

SCIENTIFIC REPORTS



OPEN

Consolidating soil carbon turnover models by improved estimates of belowground carbon input

Arezoo Taghizadeh-Toosi¹, Bent T. Christensen¹, Margaret Glendining² & Jørgen E. Olesen¹

Received: 01 June 2016
Accepted: 09 August 2016
Published: 01 September 2016

World soil carbon (C) stocks are third only to those in the ocean and earth crust, and represent twice the amount currently present in the atmosphere. Therefore, any small change in the amount of soil organic C (SOC) may affect carbon dioxide (CO₂) concentrations in the atmosphere. Dynamic models of SOC help reveal the interaction among soil carbon systems, climate and land management, and they are also frequently used to help assess SOC dynamics. Those models often use allometric functions to calculate soil C inputs in which the amount of C in both above and below ground crop residues are assumed to be proportional to crop harvest yield. Here we argue that simulating changes in SOC stocks based on C input that are proportional to crop yield is not supported by data from long-term experiments with measured SOC changes. Rather, there is evidence that root C inputs are largely independent of crop yield, but crop specific. We discuss implications of applying fixed belowground C input regardless of crop yield on agricultural greenhouse gas mitigation and accounting.

Soil carbon (C) is the third largest component of the global C cycle after the oceanic and the geologic C pools^{1,2}. The amount of soil C (including peatlands, wetlands and permafrost areas) has been estimated at 2157–2293 Pg to a depth of 1 m, comprising 1462–1545 Pg in organic forms and 695–748 Pg as carbonate¹. Soil organic matter is the source of soil organic carbon (SOC) and contains more organic carbon than global vegetation and the atmosphere combined³. In the soil, SOC is about twice the quantity of C presently in CO₂ in the atmosphere⁴. Therefore, soils are globally considered the largest terrestrial ecosystem sink or source of atmospheric CO₂ depending on land use and management practices. Globally, about 37% of land is used for agriculture and 10% of the land is under annual crops⁵. Over the last 150 years, agricultural soils have mostly been depleted in SOC due to cultivation and removal of crop residues, which suggests that agricultural land has potential to sequester SOC^{6,7}. The accurate estimation of SOC content in agricultural lands and the factors that regulate SOC accumulation are critical aspects for evaluating the extent of this potential⁴. Numerous studies have attempted to quantify changes in SOC as a key component of global climate change, i.e. as affected by elevated CO₂ or temperature using process-based models^{8,9}. The formation of soil organic matter on agricultural land depends on the C input by above- and belowground crop residues including rhizodeposition during the growth period, C input from manures or other organic by-products, and the SOC stabilization mechanisms embedded in the simulation models^{10,11}. Therefore, accurate estimations of the C input are fundamental for monitoring the development of SOC stocks of agricultural soils and also for modelling the potential for C sequestration and estimating changes in SOC content over several decades. However, reliable data on the C input of different crops is scarce and is often limited to rough estimates of above- and belowground C input using a plant C allocation approach which strongly relies on crop yield data and simplified C allocation coefficients¹².

Many major soil C simulation models are built on the premise of the metabolic theory of ecology that a number of biological properties scale allometrically with body size according to a power-law over many orders of magnitude¹³. For that, total main harvestable crop products are normally used for calculating C mass in total net primary production (NPP) assuming a concentration of 0.45 g C g⁻¹ dry matter in all crop parts and then calculating C biomass as a fraction of this value using biomass allocation functions¹⁴. A search of Web of Science in early 2016 for the combination of the words soil, carbon, model, allometric, NOT forest, NOT tree, NOT spectroscopy revealed 25 papers in the period 1990–2016. 19 of these papers use allometric functions for estimating soil C inputs. Many more studies were found when the name of specific SOC models were used instead of

¹Aarhus University, Department of Agroecology, AU-Foulum, DK-8830 Tjele, Denmark. ²Rothamsted Research, Department of Computational and Systems Biology, West Common, Harpenden, Herts AL5 2JQ, UK. Correspondence and requests for materials should be addressed to A.T.T. (email: Arezoo.Taghizadeh-Toosi@agro.au.dk)

Treatment	N rate (kg N ha ⁻¹) 1852–1985	N rate (kg N ha ⁻¹) 1985-onwards
N ₀	0	0
N ₃	144	144
N ₄	48 (year 1852–1967) & 192 (year 1968–1984)	192
N ₅	96 (year 1852–1967) & 144 (year 1968–1984)	240
N ₆	192 (year 1852–64), 0 (year 1865–83), 96 (year 1884–1984)	288

Table 1. Treatments with continuous winter wheat in the Broadbalk experiment at Rothamsted, UK, used in the current study. All plots also receive non-limiting P, K, Na and Mg.

allometric and model keywords (e.g. ICBM (30 papers), CENTURY (107 papers), CN-SIM (18 papers), DAISY (30 papers)). Most of these studies were also explicitly or implicitly applying allometric functions for estimating soil C inputs. Such methods are also commonly and increasingly used for estimating changes in SOC as part of inventories for national accounting of greenhouse gases¹⁵. Other models use alternative approaches, for example RothC (118 papers) does not attempt to compute annual returns of plant C to the soil. Instead, the model is run ‘in reverse’ to generate monthly or yearly soil C inputs from known site, soil and weather data^{16,17}.

Globally crop yields have increased greatly since the middle of the 20th century, resulting from both an increase in the total biomass production (NPP) on agricultural land and a higher proportion of the biomass in harvestable material. For example, the ratio of the grain dry matter yield to total above ground biomass (harvest index, HI) in wheat has shown a mean increase from 0.35 in 1951–1955 to 0.45 in 1995–2010 due to general progress of plant breeding, a higher nutrient availability with intensive fertilisation, and improved control of weeds, pests and diseases^{18,19}.

It has been suggested that increasing yields not only provide food for the growing population but also increase the sink of CO₂ in agroecosystems due to the accumulation of SOC or reducing the decline rate of SOC^{4,20}.

Among different crop parts which contribute to soil C input, below-ground residue inputs from roots and root exudates are of particular interest because they may contribute more to stable SOC pools than above-ground residues, partly because root derived C is more protected than other forms of C inputs due to physical protection of root materials by soil aggregates and fine pores^{12,21}. However, estimating root C input is notoriously difficult because conventional root washing misses the fine roots and provides no estimates of the amount of C exuded from roots. Tracer techniques can be used to estimate C translocation by plants such as experimental labeling with ¹³C or ¹⁴C and determination of changes in ¹³C natural abundance. However, these techniques require more sophisticated equipment and measurement approaches. Furthermore, methods based on changes in the ¹³C natural abundance are restricted to systems with a change from C₃ to C₄ plants or vice versa¹². Additionally, the estimations of total root C input from just one crop can differ due to the stage of growth of the plant, the environmental conditions, soil type, soil fertility and microbial activity^{22,23}. Therefore, estimating below-ground C input using either measurements or allometric functional relationships remains a considerable challenge.

We hypothesize that a fixed relationship between above- and belowground biomass as used in allometric functions of most soil C turnover models overestimates belowground C inputs resulting from the technological progress that has increased crop yield. We examine the impact of nitrogen (N) fertilisation on soil C development using data from a long-term fertilisation experiment and suggest an improved method for estimating below-ground soil C inputs in SOC models and in inventories of SOC change on agricultural land.

Experimental Data

Long-term field experiments provide the best foundation for the experimental verification of SOC changes and for calibrating SOC turnover models. We use data from the Broadbalk Winter Wheat Experiment, a well-managed long-term field experiment which was established in 1843 at Rothamsted Research, Harpenden, Herts, UK, and has been under winter wheat (*Triticum aestivum* L.) and known management for over 170 years. For the first few years these treatments varied a little, but in 1852 a scheme of fertiliser treatments was established that has continued, with some modifications, until today. The site is thought to have been in arable cultivation since at least 1623, and probably much earlier²⁴. The experiment does not have true, randomised replication, but has been divided into different sections over the course of the experiment²⁵. From 1852 to 1925, the plots were divided into 2 halves; therefore n = 2 for mean and s.e.m. calculations (except 1902 to 1912 and 1914 to 1915 when n = 1). In 1926, the experiment was sub-divided into five strips with generally one in fallow each year in sequence, to control weeds, crossing the treatment strips at right angles (n = 2 from 1926 to 1929, and n = 5 from 1930 to 1954). From 1955 to 1967, all plots were divided into seven strips, with one in fallow each year in sequence. The yields from the fallow plots were considered as zero. In 1968 the experiment was divided into 10 sections, in order to compare wheat grown in rotation with break crops with wheat grown continuously. For years 1968 to 2014 we have used the data from the three main continuous wheat sections (1, 6 and 9), excluding section 6 from 1968 to 1978 as it was still in a fallow rotation. The soil at Broadbalk is a well-drained clay loam to silty clay loam classified as an Aquic Paleudalf (USDA) and as a Chromic Luvisol⁵. Straw is removed at harvest each year, leaving above-ground residues of stubble, chaff and the uncollected straw. We selected the treatments from the continuous wheat sections, with a range of rates of N in mineral fertiliser (Table 1), resulting in different crop yields (Fig. 1). Austin *et al.* (1993) calculated HI based on total above-ground biomass (stubble, chaff and all of




Figure 1. Trends in grain and straw yield (85% Dry Matter) in the Broadbalk wheat experiment, Rothamsted, UK (error bars = \pm s.e.m., n = differs, see experimental data). See Table 1 for details of N treatments.

the straw) and compared an old long-strawed variety with a modern short-strawed variety, over three years and several different fertiliser treatments on Broadbalk²⁶. The varieties were typical of those which historically have been grown on the experiment. The average HI was 0.30 for the old variety, and 0.47 for the modern variety²⁶. Taking the period from 1852 to 1967 the ratio of grain yield to grain plus harvested straw yield was around 0.35 in Broadbalk. When winter wheat varieties were changed to modern short-straw varieties in 1968, the grain yield as proportion of grain plus harvested straw yield was changed substantially (less straw and more grain was produced). The average ratio of grain yield to grain plus straw yield for treatments N_3 , N_4 , N_5 , and N_6 was 0.63 during the period 1968–2013, with no difference between the treatments. The mean for N_0 was very similar to the other treatment, 0.65. Considering just the grain and harvested straw would underestimate the aboveground residues since stubble, chaff and unharvested straw was not measured. Nevertheless, it can be assumed that the ratio of grain to straw was constant, and allometric functions can be used for estimating aboveground residuals²⁷. We used harvest index (HI: the ratio of grain to total aboveground biomass) to estimate above and belowground C input (see supplementary materials).

Measured SOC was calculated from % SOC and a standard soil weight of $2.88 \times 10^6 \text{ kg ha}^{-1}$ 0–23 cm (see supplementary materials for further details).

Soil C model (C-TOOL)

For testing our hypothesis, we used a process based soil C model, C-TOOL^{11,28}. In C-TOOL, the simulated SOC stock depends on the input of organic C from crops, its rate of decomposition, soil texture, initial soil C:N ratio and temperature. In the standard version of C-TOOL, allometric relations are used to derive the C inputs from measured crop yield (see supplementary materials). Here, C input from aboveground residues was calculated based on the estimated HI and measured grain and straw yields (Fig. 1) from 1852 to 2013. We have assumed $\text{HI} = 0.35$ from 1852 to 1970 and $\text{HI} = 0.45$ from 1980 to 2013. From 1970 to 1980, we assumed a linear increase of HI from 0.35 to 0.45. For root C input estimation, three approaches were pursued: (1) the C-TOOL allometric functions (see supplementary materials) were used to calculate the root C input of winter wheat into the soil for each treatment (hereafter called allometric root C input), (2) the N_3 treatment was considered as a reference and the average N_3 root C input (for 1852 to 2013) calculated from allometric functions was used as a constant root C input in all treatments (hereafter called fixed root C input), (3) the N_3 treatment was considered as a reference




Figure 2. Trends in estimated total soil C inputs for different N fertiliser rate treatments of the Broadbalk winter wheat experiment at Rothamsted, UK. *Note:* different C calculations on each figure. See Table 1 for details of N treatments.

and N_3 root C input calculated from allometric functions for each year was applied in the corresponding year in the other treatments (hereafter called N_3 root C input) from 1852 to 2013.

All treatments were assumed to have the same SOC content prior to the start of experiment. The model initial SOC content was estimated by minimising the sum of squares between simulated and estimated SOC in 1852 of N_0 treatment using nonlinear curve-fitting function in MATLAB at the start of the model spin-up period (year 1822). That estimated initial SOC content was used for all treatments at the start of spin-up period (Spin-up period started in 1822). The spin-up period was 30 years and was used to initialise the model using similar C input calculated from allometric functions using winter wheat yield in 1852 for all treatments.

Results

Our study compared different methods for estimating soil C inputs using data from the long-term Broadbalk experiment at Rothamsted, UK, with different fertiliser rates providing different yields in winter wheat (Fig. 1). The wheat grain yields increased considerably over time in plots with the higher fertilisation rates while no increase in grain yield was observed in unfertilised plots. The estimated soil C input based on allometric root C input, fixed root C input or N_3 root C input show markedly different estimates (Fig. 2). The average measured topsoil SOC contents in each treatment did not change significantly from year 1987 onwards (See supplementary materials, Fig. S4). Therefore, we used the average for 1987–2010 of measured SOC contents for comparing with the simulated values using different approaches for estimating root C input (Fig. 3).

Discussion

The increasing wheat yields from 1960 to 1980 in all treatments receiving fertiliser was attributed to improvements in plant breeding and crop husbandry²⁹. Crop breeding, particularly cereals, has resulted in the preferential allocation of assimilated C to grains (higher HI)³⁰. However, total biomass production has also been increasing, due to a combination of improved crop varieties, better plant nutrition and improved crop protection. Increased N fertiliser rates result in increased yields; however, this relationship is non-linear with a saturation response at higher N rates (Fig. 1). The carbon input as a proportion of winter wheat yield varies in literature, but it was




Figure 3. The average of SOC contents (measured SOC and simulated SOC using allometric root C input, fixed root C input and N₃ root C input estimations for the period 1987–2010 (error bars = + s.e.m., n = 6 for measured values and n = 24 for simulated values). See Table 1 for details of N treatments.

considered 86% by Jenkinson and Rayner (1977)³¹. Chrinda *et al.* (2012) measured grain yield (9.5 t ha⁻¹), the amounts of shoot C (5.1 t C ha⁻¹) and root C (0.9 t C ha⁻¹) of winter wheat in the inorganic fertilised based rotation in Denmark³².

In SOC models that use allometric functions to estimate C inputs, higher grain yields produced by higher N fertilisation rates will result in enhanced above- and belowground biomass and thereby greater inputs of C in crop residues (stubble and roots) to the soil. As a result, higher SOC contents would be expected in high compared to low-yielding cropping systems. For the treatment with no N fertilisation (N₀), SOC was measured to be lower compared with the treatments with higher N rates. When N fertiliser was applied at 144 kg N ha⁻¹ yr⁻¹ (reference rate, N₃, since 1852), higher SOC contents were as expected obtained compared with no N application. However, rates of N fertiliser above the reference rate (N₄, N₅ and N₆) did not increase topsoil SOC contents much more, although these treatments had only been applied since 1968 (N₄) and 1985 (N₅ and N₆). In contrast, the simulations of SOC contents using the allometric function for calculating C input showed higher SOC contents in topsoil of N fertilised treatments than measured topsoil organic C contents (ranged from extra 1 to 4 t C accumulation ha⁻¹, see supplementary materials, Figure S1). We surmise that the deviation between measured and simulated SOC contents with different level of N fertiliser can be ascribed mainly to overestimation of belowground C inputs. Higher N fertiliser rates do not necessarily increase belowground biomass, possibly due to a relative decrease in cumulative fine root production as a result of fertilisation³². Root biomass has shown less response to added N than aboveground biomass causing the below- to aboveground ratio to decline with increasing N rates³³. When we used N₃ root C input for each year in different treatments, the simulated SOC contents were slightly overestimated by 0.4 to 2 t C ha⁻¹ in N₄, N₅ and N₆ treatments (see supplementary materials, Figure S2). This means that there is a need to estimate the belowground C input using an alternative to the allometric function, which could be independently derived, reliable belowground C input data. In our simulations, using fixed root C input resulted in a closer match between simulated and measured topsoil SOC (difference; 0.02 to 1 t C ha⁻¹, see supplementary materials, Figure S3). There was a slight difference in SOC contents between measured and simulated using fixed root C input in N₃ (reference N rate, Fig. 3). We believe that difference might be due to slight overestimation of aboveground C input from winter wheat residues and stubbles, possibly due to changes in the HI not accounted for in our assumptions³⁴. These results indicate that estimation of changes in SOC on arable land using allometric functions probably overestimate the ability of the land surface to accumulate C from technological progress. These findings emphasize the need to better quantify and represent shifting plant C allocation in the face of changing technologies that lead to yield improvements, but not to equivalent increases in root C inputs.

Conclusion

Use of crop yields and application of strict allometric functions for estimating C inputs from belowground crop residues and exudates in carbon models can lead to erroneous estimates of belowground C inputs in agricultural systems. The commonly applied method of assuming a fixed ratio between crop yield and root C inputs will tend to overestimate root C inputs for technologies that improve harvestable yield. However, using a process based model like C-TOOL and allometric functions in combination with fixed belowground C input value estimated from standard crop management may result in more accurate estimation of SOC trends in agricultural soils. Such an approach would contribute to an improved assessment of soil C sequestration potentials, in particular when considering measures that enhance harvestable crop yield.

References

1. Batjes, N. H. Total carbon and nitrogen in the soils of the world. *European Journal of Soil Science* **47**, 151–163 (1996).

2. Schuur, E. A. G. *et al.* Vulnerability of permafrost carbon to Climate Change: Implications for the global carbon cycle. *Bioscience* **58**, 701–714 (2008).
3. Lehmann, J. & Kleber, M. The contentious nature of soil organic matter. *Nature* **528**, 60–68 (2015).
4. Powlson, D., Whitmore, A. & Goulding, K. Soil carbon sequestration to mitigate climate change: A critical re examination to identify the true and the false. *European Journal of Soil Science* **62**, 42–55 (2011).
5. FAO. Food and Agriculture Organization of the United nations, <<http://www.fao.org/>> (2013).
6. Chapman, S. J. *et al.* Comparison of soil carbon stocks in Scottish soils between 1978 and 2009. *European Journal of Soil Science* **64**, 455–465 (2013).
7. Lal, R. Soil carbon sequestration impacts on global climate change and food security. *Science* **304**, 1623–1627 (2004).
8. Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* **440**, 165–173 (2006).
9. Van Groenigen, K. J., Xia, J., Osenberg, C. W., Luo, Y. & Hungate, B. A. Application of a two-pool model to soil carbon dynamics under elevated CO₂. *Global Change Biology* **21**, 4293–4297 (2015).
10. Andrén, O. & Kätterer, T. ICBM: The introductory carbon balance model for exploration of soil carbon balances. *Ecological Applications* **7**, 1226–1236 (1997).
11. Taghizadeh-Toosi, A. *et al.* C-TOOL: A simple model for simulating whole-profile carbon storage in temperate agricultural soils. *Ecological Modelling* **292**, 11–25 (2014).
12. Kuzyakov, Y. & Domanski, G. Carbon inputs by plants into the soil. Review. *Journal of Plant Nutrition and Soil Science* **163**, 421–431 (2000).
13. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789 (2004).
14. Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A. & VandenBygaart, A. J. An approach for estimating net primary productivity and annual carbon inputs to soil for common agricultural crops in Canada. *Agriculture, Ecosystems and Environment* **118**, 29–42 (2007).
15. Johnson, J. M. F., Allmaras, R. R. & Reicosky, D. C. Estimating source carbon from crop residues roots and rhizodeposits using the national grain-yield database. *Agronomy Journal* **98**, 622–636 (2006).
16. Coleman, K. & Jenkinson, D. S. In *Evaluation of soil organic matter models using existing long-term datasets* Vol. 38 (eds Powlson, D. S., Smith, P. & Smith, J. U.) 237–246 (Springer-Verlag, 1996).
17. Coleman, K. *et al.* Simulating trends in soil organic carbon in long-term experiments using RothC-26.3. *Geoderma* **81**, 29–44 (1997).
18. Fang, Y., Liu, L., Xu, B. C. & Li, F. M. The relationship between competitive ability and yield stability in an old and a modern winter wheat cultivar. *Plant and Soil* **347**, 7–23 (2011).
19. Shearman, V. J., Sylvester-Bradley, R., Scott, R. K. & Foulkes, M. J. Physiological processes associated with wheat yield progress in the UK. *Crop Science* **45**, 175–185 (2005).
20. Ladha, J. K., Reddy, C. K., Padre, A. T. & van Kessel, C. Role of nitrogen fertilization in sustaining organic Matter in cultivated soils. *Journal of Environmental Quality* **40**, 1756–1766 (2011).
21. Menichetti, L., Ekblad, A. & Kätterer, T. Contribution of roots and amendments to soil carbon accumulation within the soil profile in a long-term field experiment in Sweden. *Agriculture, Ecosystems and Environment* **200**, 79–87 (2015).
22. Keith, H., Oades, J. M. & Martin, J. K. Input of carbon to soil from wheat plants. *Soil Biology & Biochemistry* **18**, 445–449 (1986).
23. Saggart, S. & Hedley, C. B. Estimating seasonal and annual carbon inputs, and root decomposition rates in a temperate pasture following field ¹⁴C pulse-labelling. *Plant and Soil* **236**, 91–103 (2001).
24. Avery, B. W. & Bullock, P. The soils of Broadbalk: Morphology and classification of Broadbalk soils. *Rothamsted Experimental Station Report for 1968 Part 2*, 63–81 (1969).
25. Rothamsted. Guide to the Classical and other Long-term Experiments, Datasets and Sample Archive. (Lawes Agricultural Trust Ltd, Harpenden UK, 2006).
26. Austin, R. B., Ford, M. A., Morgan, C. L. & Yeoman, D. Old and modern wheat cultivars compared on the Broadbalk wheat experiment. *European Journal of Agronomy* **2**, 141–147 (1993).
27. Bolinder, M. A., Angers, D. A. & Dubuc, J. P. Estimating shoot to root ratios and annual carbon inputs in soils for cereal crops. *Agriculture, Ecosystems and Environment* **63**, 61–66 (1997).
28. Taghizadeh-Toosi, A. & Olesen, J. E. Modelling soil organic carbon in Danish agricultural soils suggests low potential for future carbon sequestration. *Agricultural Systems* **145**, 83–89 (2016).
29. Foulkes, M. J. *et al.* Genetic progress in yield potential in wheat: recent advances and future prospects. *Journal of Agricultural Science Cambridge* **145**, 17–29 (2007).
30. Mackay, I. *et al.* Analyses of the historical series of UK variety trials to quantify the contributions of genetic and environmental factors to trends and variability in yield over time. *Theoretical and Applied Genetics* **122**, 225–238 (2011).
31. Jenkinson, D. S. & Rayner, J. H. The turnover of soil organic matter in some of the Rothamsted classical experiments. *Soil Science* **123**, 298–305 (1977).
32. Chirinda, N., Olesen, J. E. & Porter, J. R. Root carbon input in organic and inorganic fertilizer-based systems. *Plant and Soil* **359**, 321–333 (2012).
33. Jensen, B. & Christensen, B. T. Interactions between elevated CO₂ and added N: effects on water use, biomass, and soil ¹⁵N uptake in wheat. *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science* **54**, 175–184 (2004).
34. Palosuo, T., Heikkinen, J. & Regina, K. Method for estimating soil carbon stock changes in Finnish mineral cropland and grassland soils. *Carbon Management*, doi: 10.1080/17583004.2015.1131383 (2016).

Acknowledgements

The work was funded by the Danish Ministry of Environment as part of the SINKS2 project. We thank the Lawes Agricultural Trust and Rothamsted Research for data from the Broadbalk experiment, from the e-RA database. The Rothamsted Long-term Experiments National Capability is supported by the UK Biotechnology and Biological Sciences Research Council and the Lawes Agricultural Trust.

Author Contributions

Measured soil data was provided by M.G. All the modelling simulations were performed by A.T.T. The manuscript was first drafted by A.T.T. whereas revisions were performed by all authors; A.T.T., B.T.C., M.G., J.E.O. Final preparation for submission was done by A.T.T.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Taghizadeh-Toosi, A. *et al.* Consolidating soil carbon turnover models by improved estimates of belowground carbon input. *Sci. Rep.* **6**, 32568; doi: 10.1038/srep32568 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016