

Jury is still out on the radiative forcing by black carbon

Olivier Boucher^{a,1}, Yves Balkanski^b, Øivind Hodnebrog^c, Cathrine Lund Myhre^d, Gunnar Myhre^c, Johannes Quaas^e, Bjørn Hallvard Samset^c, Nick Schutzgens^f, Philip Stier^f, and Rong Wang^b

Peng et al. (1) conclude that a fast increase in the mass absorption cross-section (MAC) of black carbon (BC) in urban environments leads to significantly increased estimates of the BC radiative forcing (RF). Their chamber measurements are highly valuable and complement observations performed in ambient conditions, but their “enhancement factor” relative to an unspecified baseline may not be directly comparable to values used or simulated in global aerosol models. MAC, a key parameter in our understanding of the net BC climate impact, is indeed a more relevant quantity to examine. A fast MAC enhancement in polluted environments as the BC gets coated with organic and inorganic species is consistent with recent findings (2, 3). Global models used in AeroCom [table S1 in Peng et al. (1), ref. 4] have an average MAC of $\sim 8 \text{ m}^2 \cdot \text{g}^{-1}$ at 550 nm. This value is reflecting reported measurements, although there is a large spatial and seasonal variability in ambient MAC for aged particles, with values of $\sim 10 \text{ m}^2 \cdot \text{g}^{-1}$ at a rural Northern Chinese site (2) (at 678 nm); $6\text{--}14 \text{ m}^2 \cdot \text{g}^{-1}$ at rural, urban, and high-altitude Indian locations (5) (at 678 nm); and $\sim 6 \text{ m}^2 \cdot \text{g}^{-1}$ at an Arctic site (6) (at 522 nm).

Coating of BC by soluble species not only enhances absorption of solar radiation but also reduces the BC atmospheric lifetime (7). Fig. 1 shows an offset between the increase in average MAC value with faster BC aging and an overall shorter BC lifetime, resulting in a near-constant BC aerosol absorption optical depth and RF with aging time. Furthermore, current global aerosol

models frequently have a too long BC lifetime and consequently overestimate BC concentrations downwind from source regions (8).

According to Peng et al. (1), their BC absorption enhancement factor of 2.4 is also an upper bound, only reached after 5 (Beijing) to 18 (Houston) h, and possibly longer in cleaner environments. Such time-scales are not small compared with the BC atmospheric lifetime of 3–5 d, especially considering that dilution effects may lengthen the aging timescale in the real atmosphere compared with the static chamber measurements performed by Peng et al. It is unclear how representative these measurements are for global and annual averages, but we know that generalizations can introduce serious errors due to spatial and temporal sampling issues (9). This implies that a simple scaling of the BC RF by the absorption enhancement factor measured by Peng et al.—as performed by the authors in their table S1 and figure 4, and extended in the commentary (10)—is overly simplistic.

In conclusion, although we welcome the advances made by Peng et al., their conclusion of a $+0.45 [0.21\text{--}0.80] \text{ W} \cdot \text{m}^{-2}$ additional RF due to a large BC enhancement factor is premature. The jury is still out on the question of the net climate impact of BC and how much climate cobenefit will result from the necessary mitigation of BC emissions. Reducing the uncertainty on the BC forcing requires better constraining BC MAC and atmospheric lifetime in global aerosol models.

^aLaboratoire de Météorologie Dynamique, Institut Pierre-Simon Laplace, Université Pierre et Marie Curie/CNRS, 75005 Paris, France; ^bLaboratoire des Sciences de l'Environnement et du Climat, Commissariat à l'Énergie Atomique et aux Énergies Alternatives/CNRS/Université de Versailles Saint-Quentin-en-Yvelines, 91198 Gif-sur-Yvette, France; ^cCenter for International Climate and Environmental Research, 0318 Oslo, Norway; ^dNorwegian Institute for Air Research, 2007 Kjeller, Norway; ^eInstitute for Meteorology, Universität Leipzig, D-04103 Leipzig, Germany; and ^fDepartment of Physics, University of Oxford, Oxford OX1 3PJ, United Kingdom

Author contributions: O.B., Y.B., Ø.H., C.L.M., G.M., J.Q., B.H.S., N.S., P.S., and R.W. wrote the paper; B.H.S. and R.W. performed the experiments. The authors declare no conflict of interest.

¹To whom correspondence should be addressed. Email: olivier.boucher@lmd.jussieu.fr.




Fig. 1. Relationship between BC aerosol absorption optical depth (AAOD) (black scale), MAC (green scale), and direct radiative effect (blue scale) as a function of the aging timescale of the BC aerosol in two different models: LMDz-OR-INCA and OsloCTM2. LMDz-OR-INCA includes all sources of BC; its MAC at 550 nm is $5.5 \text{ m}^2\text{g}^{-1}$ for fresh BC increasing to an average of $11.4 \text{ m}^2\text{g}^{-1}$ for aged BC, and the BC atmospheric residence time is about 6 d. OsloCTM2 includes fossil fuel and biofuel BC sources only; its MAC at 550 nm is $7.3 \text{ m}^2\text{g}^{-1}$ for fresh BC increasing to $11.0 \text{ m}^2\text{g}^{-1}$ for aged BC, and the BC atmospheric residence time is about 3 d (resulting in a better agreement with observations in remote regions). The BC enhancement factors are 2.1 and 1.5, respectively, but correspond to different baselines.

- 1 Peng J, et al. (2016) Markedly enhanced absorption and direct radiative forcing of black carbon under polluted urban environments. *Proc Natl Acad Sci USA* 113(16):4266–4271.
- 2 Cui X, et al. (2016) Radiative absorption enhancement from coatings on black carbon aerosols. *Sci Total Environ* 551-552:51–56.
- 3 Liu S, et al. (2015) Enhanced light absorption by mixed source black and brown carbon particles in UK winter. *Nat Commun* 6:8435.
- 4 Myhre G, et al. (2013) Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations. *Atmos Chem Phys* 13(4):1853–1877.
- 5 Ram K, Sarin MM (2009) Absorption coefficient and site-specific mass absorption efficiency of elemental carbon in aerosols over urban, rural, and high-altitude sites in India. *Environ Sci Technol* 43(21):8233–8239.
- 6 Yttri KE, et al. (2014) Quantifying black carbon from biomass burning by means of levoglucosan—A one-year time series at the Arctic observatory Zeppelin. *Atmos Chem Phys* 14(12):6427–6442.
- 7 Stier P, Seinfeld JH, Kinne S, Feichter J, Boucher O (2006) Impact of nonabsorbing anthropogenic aerosols on clear-sky atmospheric absorption. *J Geophys Res* 111(D18):D18201.
- 8 Samset BH, et al. (2014) Modelled black carbon radiative forcing and atmospheric lifetime in AeroCom Phase II constrained by aircraft observations. *Atmos Chem Phys* 14(22):12465–12477.
- 9 Schutgens NAJ, et al. (2016) Will a perfect model agree with perfect observations? The impact of spatial sampling. *Atmos Chem Phys* 16(10):6335–6353.
- 10 Gustafsson Ö, Ramanathan V (2016) Convergence on climate warming by black carbon aerosols. *Proc Natl Acad Sci USA* 113(16):4243–4245.