



Original investigation

# US Smokers' Beliefs, Experiences and Perceptions of Different Cigarette Variants Before and After the FSPTCA Ban on Misleading Descriptors Such as "Light," "Mild," or "Low"

Hua-Hie Yong PhD<sup>1</sup>, Ron Borland PhD<sup>1</sup>, K. Michael Cummings PhD<sup>2</sup>,  
Eric N. Lindblom JD<sup>3</sup>, Lin Li PhD<sup>1</sup>, Maansi Bansal-Travers PhD<sup>4</sup>,  
Richard J. O'Connor PhD<sup>4</sup>, Tara Elton-Marshall PhD<sup>5,6,7</sup>, James F. Thrasher PhD<sup>8</sup>,  
David Hammond PhD<sup>9</sup>, Mary E. Thompson PhD<sup>10</sup>, Timea R. Partos PhD<sup>11</sup>

<sup>1</sup>Nigel Gray Fellowship Group, Cancer Council Victoria, Melbourne, Australia; <sup>2</sup>Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, SC; <sup>3</sup>O'Neill Institute for National and Global Health Law, Georgetown University Law Center, Washington, DC; <sup>4</sup>Department of Health Behavior, Roswell Park Cancer Institute, Buffalo, NY; <sup>5</sup>Social and Epidemiological Research, Centre for Addiction and Mental Health, London, ON, Canada; <sup>6</sup>Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada; <sup>7</sup>School of Health Studies, Western University, London, ON, Canada; <sup>8</sup>Department of Health Promotion, Education and Behavior, Arnold School of Public Health, University of South Carolina, Columbia, SC; <sup>9</sup>School of Public Health and Health Systems, University of Waterloo, Waterloo, ON, Canada; <sup>10</sup>Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, ON, Canada; <sup>11</sup>Institute of Psychiatry, King's College London, London, UK

Corresponding Author: Hua-Hie Yong, PhD, Nigel Gray Fellowship Group, Cancer Council Victoria, 615 St Kilda Road, Melbourne, VIC 3004, Australia. Telephone: 61-3-9514-6283; Fax: 61-3-9514-6800; E-mail: [hua.yong@cancervic.org.au](mailto:hua.yong@cancervic.org.au)

## Abstract

**Introduction:** In December 2008, the Federal Trade Commission (FTC) took action that prompted the removal of nicotine and tar listings from cigarette packs and ads. As of June 2010, the US Family Smoking Prevention and Tobacco Control Act prohibited the use of explicit or implicit descriptors on tobacco packaging or in advertising that convey messages of reduced risk or exposure, specifically including terms like "light," "mild," and "low" and similar descriptors. This study evaluates the effect of these two policy changes on smokers' beliefs, experiences and perceptions of different cigarettes.

**Methods:** Using generalized estimating equations models, this study analyzed survey data collected between 2002 and 2013 by the International Tobacco Control Policy Evaluation Study regarding US smokers' beliefs, experiences, and perceptions of different cigarettes.

**Results:** Between 2002 and 2013, smoker misperceptions about "light" cigarettes being less harmful did not change significantly and remained substantial, especially among those who reported using lower-strength cigarettes. After the two policy changes, reported reliance on pack colors, color terms, and other product descriptors like "smooth" to determine cigarette strength style trended upward.

**Conclusions:** Policies implemented to reduce smoker misperceptions that some cigarettes are safer than others appear to have had little impact. Because of pack colors, color terms, descriptors such as "smooth," cigarette taste or feel, and possibly other characteristics, millions of smokers

continue to believe, inaccurately, that they can reduce their harms and risks by smoking one cigarette brand or sub-brand instead of another, which may be delaying or reducing smoking cessation.

**Implications:** *What this study adds:* This study confirms that US policies to reduce smoker misperceptions that some cigarettes are less harmful than others have not been successful. Following the removal of light/low descriptors and tar and nicotine numbers from cigarette packs and ads, pack colors, color words, other descriptors (eg, smooth), and sensory experiences of smoother or lighter taste have helped smokers to continue to identify their preferred cigarette brand styles and otherwise distinguish between which brands and styles they consider “lighter” or lower in tar and, mistakenly, less harmful than others. These findings provide additional evidence to support new enforcement or regulatory action to stop cigarettes and their packaging from misleading smokers about relative risk, which may be reducing or delaying quit attempts.

## Introduction

In December 2008, the Federal Trade Commission (FTC) issued a notice rescinding its prior guidance that listing Cambridge method or “FTC method” nicotine and tar levels on packs and in advertisements was not a violation of the FTC Act, prompting cigarette companies to remove the tar/nicotine figures from their packs and ads. In addition, the 2009 Family Smoking Prevention and Tobacco Control Act (FSPTCA) gave the US Food and Drug Administration (FDA) broad authority to regulate tobacco products and their marketing.<sup>1</sup> Section 911 of the Act specifically addressed concerns about the deceptive marketing of so-called “light” cigarettes.<sup>2</sup> As of 22 June 2010, the Act prohibited the use of explicit or implicit descriptors on tobacco packaging or in advertising that convey messages of reduced risk or exposure—specifically including the use of the words “light,” “mild,” and “low”—unless and until the manufacturer could demonstrate to the FDA that the messages conveyed by the descriptors were accurate and not misleading and that allowing the descriptors would benefit the public health.<sup>2,3</sup> Prior to the enactment of the FSPTCA, it was well understood that smokers mistakenly believed that light/low cigarettes were less harmful than regular cigarettes, which reduced cessation rates and maintained higher smoking levels and harms.<sup>4</sup>

Prior to the US ban on misleading brand descriptors in 2010, over 50 other countries had similarly banned misleading brand descriptors.<sup>5</sup> Studies evaluating the impact of the bans on misleading brand descriptors have generally concluded that such bans have not been able to substantially, much less completely, eliminate smoker misperceptions that some cigarette brands and sub-brands are less harmful than others.<sup>6,7</sup> In many of these countries, the industry anticipated or responded to the descriptor bans by introducing color-related words such as “gold,” “silver,” and “blue” on packs as replacement descriptors along with altering elements of packaging design (eg, color, brand imagery), and cigarette engineering (eg, filter venting), in an effort to differentiate brands so that smokers would continue identifying some cigarette brands and sub-brands as “light,” “mild,” or “low.”<sup>7-13</sup> Evidence from both experimental studies<sup>8,11,14,15</sup> and population-based, observational studies<sup>3,9</sup> suggests that the use of color, whether done through the use of color words, different pack colors, or different coloring of the cigarettes and filters, continues to perpetuate the myth that cigarette vary in terms of health risks and harms, which can reduce or delay smoking cessation.

Yong et al.<sup>7</sup> evaluated the impact on smokers’ misperceptions following the ban on the terms “light” and “mild” in the United Kingdom in 2003 and in Australia in 2006, comparing them with the United States and Canada, which had no such bans at the time of the study. They found that the bans were followed by only a short-term

reduction in the belief that “light/mild” cigarettes confer some health benefits, and concluded that the temporary decline in this misunderstanding was largely the result of each country complementing the bans with a temporary mass media public education campaign. They based this conclusion on the fact that such misperceptions also declined over the study period in the United States, which had not implemented a ban, following several widely publicized, high-profile court cases against the tobacco industry for misleadingly marketing cigarettes labeled as “light.”<sup>6,7</sup>

Cigarette companies have frequently argued that their use of pack colors and brand descriptors are merely intended to communicate differences between brands/sub-brands with regards to taste and texture. However, research evidence shows that consumer perceptions of product risk is also related to taste and sensory impact which can be influenced not just by the types of tobacco used or how products are engineered (eg, filter ventilation)<sup>16,17</sup> but even by the terms used to describe the brand or brand variant or the color of the cigarette, its filter or its packaging.<sup>11</sup> Ironically, increasing smoothness and lightness by filter ventilation may actually increase product risk by making the tar more mutagenic,<sup>18-20</sup> but the cooling effect of oxygen mixing with the hot smoke will prompt smokers to experience the cigarette as lighter and easier on the throat (and mistakenly think it is less harmful than other cigarettes).<sup>10,21-24</sup>

Because smokers’ belief that light/low cigarettes are smoother on the throat and chest than regular cigarettes is directly linked to their beliefs that light/low cigarettes are safer or less harmful than other types of cigarettes, monitoring trends in this sensory belief, and the factors influencing that belief, is important. Past research found smokers’ belief that “light” cigarettes are smoother on the throat and chest than regular cigarettes declined between 2002 and 2006,<sup>6</sup> but the reason for this trend and whether it would persist remained unclear.

Similar to the taste and texture research, past research also suggests that smokers trying to reduce their harms from smoking are more likely to choose cigarettes with lower strength as defined by listed tar and/or nicotine levels or by the cigarettes being labeled as light, mild, or low,<sup>24</sup> and their levels of misperception about light/low cigarettes being less harmful tend to correspond to the listed strength of the cigarettes they smoke—highest among smokers of “ultralight” cigarettes and lowest among smokers of “regular full-flavored” cigarettes.<sup>22</sup> However, no past studies have examined whether the removal of nicotine and tar numbers or “light/low” terms from cigarette packs and ads might have a differential impact on smoker misperceptions depending on whether they smoke cigarettes formerly labeled or advertised as light/low or with lower tar or nicotine numbers or that the smokers currently perceive as light/

low despite the absence of the related numbers or descriptors. One would expect the impact to be greater for smokers of those light/low cigarettes that were most directly affected by the descriptor ban than for smokers of regular cigarettes.

Since 2006, there appear to have been only two major national policy changes in the United States that may have influenced smokers' sensory and relative-harm beliefs: the previously described FTC guidance<sup>25</sup>; and the FSPTCA defacto ban on descriptors such as "light," "mild," and "low." The former prompted cigarette companies to remove the nicotine and tar listings from cigarette packs and ads while the latter resulted in cigarette companies' corresponding use of pack colors and color terms and other descriptors to distinguish different brand variants. At the same time, recent evidence suggests no significant change in the filter ventilation levels of major brands and variants following the descriptor ban in the United States.<sup>3</sup> Although it has not previously been researched, one would, therefore, anticipate that after the ban, smokers would become more reliant on package colors as an indicator of taste and texture, tar levels, whether a brand or variant was light/low or regular, and, consequently, relative harmfulness.

In the months leading up to the effective date of the FSPTCA ban on specific descriptor terms on existing cigarette packs, Philip Morris USA ran an information campaign for its leading brand Marlboro, and briefly introduced a pack insert to reassure consumers that their cigarettes remained the same despite the changed packaging (the only manufacturer known to do so), although the FDA quickly stopped this practice.<sup>26</sup> It is unclear how the information campaign might have differentially affected Marlboro smokers' misperceptions about the relative harmfulness of different cigarette variants. Marlboro has been heavily advertised and marketed in the United States and remains the most popular brand smoked by US smokers.<sup>27,28</sup>

This study extends previous research by Yong et al.<sup>7</sup> using two additional waves of data from the US arm of the ITC Four-Country (ITC4) survey, collected in 2010 and 2013, to understand the impact following the 2009 removal of tar and nicotine numbers from packs and advertisements and the 2010 defacto "light/low" descriptor ban on US smokers' beliefs, reported experiences and perceptions about different styles of cigarettes. Specifically, this study examined (1) the extent to which the removal of misleading "light/mild/low" terms from cigarette packaging affected (a) smokers' beliefs about the relative harmfulness of different cigarette variants; (b) their experiences about smoothness and taste of different variants; or (c) their perceptions about the extent to which pack color, tar/nicotine levels, and pack descriptor like "smooth" provide useful information about cigarette taste or smoothness; and (2) the extent to which any impacts varied by cigarette brand (Marlboro vs. other top-selling brands) and strength styles (ultralight, light, and regular cigarettes).

## Methods

### Sample

Data come from US arm of the ITC4 project, a cohort study of adult smokers conducted annually since 2002 in the United States, Canada, the United Kingdom, and Australia. A detailed description of the conceptual framework and methods of the ITC-4 Survey has been reported elsewhere.<sup>29,30</sup> Briefly, the ITC-4 Survey employs a prospective multi-country cohort design and involves telephone surveys of representative cohorts of adult smokers in each country using random-digit dialing (and from Wave 7 onwards web survey was also used). The sample size per country was initially around 2000 at each

wave, with replenishment sampling from the same sampling frame used to maintain sample size across waves (NB. A slightly reduced sample size was obtained at Waves 7 and 8, mainly due to budget, but the Wave 9 sample size was markedly increased through replenishment because of additional funding resources). A brief time-line of the data collection and key events related to the "light/low" labeling change in the United States is presented in [Supplementary Appendix, Supplementary Table 1](#). At the time of initial recruitment, participants were aged 18 years or more, had smoked at least 100 cigarettes over their lifetime, and had smoked at least once in the past 30 days.

## Measures

### Outcome Measures

These included beliefs, experiences and perceptions about "light/low" cigarettes, and one's own brands, relating to relative harmfulness, cigarette taste, and taste indicators (see details in [Table 1](#)). Consistent with existing research,<sup>7</sup> the "Lights are less harmful" and "Lights give less tar" belief items were combined into a scale (correlation across waves:  $r = 0.50-0.60$ , all  $P < .001$ ) by averaging the scores to form the Lights Benefit Scale (LBS). Reported usual brand and sub-brand style smoked was recorded and used to determine brand family. The reported usual sub-brand style smoked was categorized as "full flavor/regular," "light," or "ultralight" based on how the sub-brand style was labeled prior to the descriptor ban or on its current color coding, with those that could not be classified because of insufficient information coded as "other." Strength coding was based on Cornelius et al.<sup>31</sup> and guided by our US investigators. For those without a usual brand and variant, last purchase brand and variant was used.

### Covariates

These included cigarettes per day and having made a quit attempt in the prior year, as well as socio-demographics such as age, gender, annual household income (low:  $\leq \$29\,999$ ; moderate:  $\$30\,000-59\,999$ ; high:  $\geq \$60\,000$ ), education (low:  $\leq$  high school; moderate: some college/tech/trade school, no degree; high: university degree or higher), and minority ethnic status (non-white and mixed race vs. white). Survey mode (phone vs. internet) and cohort (ie, year of recruitment) were also included.

### Data Analysis

All analyses were conducted using Stata version 14. Analyses were limited to current smokers of factory-made cigarettes at each wave with sufficient data to determine brand family and the cigarette strengths of the sub-brands or brand variants smoked (about 16% were excluded because of insufficient detailed brand attribute information provided). As the number of brand families and varieties was extensive and varied across waves, brand family analyses were limited to the top 10 brands reported at each wave, which included Marlboro, Newport, Camel, Doral, Winston, Kool, Basic, Virginia Slims, Salem, Benson & Hedges, Misty, Pall Mall, Seneca, American Spirit, Maverick, and Pyramid (list and ranking of specific brands vary across waves). This approach captured the top-selling brands (ie, 73% to 80% of brands smoked by respondents across waves) while ensuring sufficient sample sizes for analyses. Marlboro was kept as a separate category for brand smoked comparisons because it was consistently the top brand (ie, reported by 26%–37% of the sample across the waves) and thus, study findings will be directly applicable to the largest share of current smokers in the United States. The remaining brands were combined into a single category.

**Table 1.** Light/Mild Related Questions Assessed in the International Tobacco Control US Survey

Survey questions	Wave (year) asked	Response options
<b>Beliefs</b>		
Light cigarettes are less harmful than regular cigarettes. Smokers of light cigarettes take in less tar than smokers of regular cigarettes.	1 to 8 (2002–2010)	Rated on a 5-point scale ranging from “strongly agree” to “strongly disagree” (NB. Responses were reversed coded and “Don’t Know” responses recoded as “neither” for analysis purpose)
Based on your experience of smoking, do you think that the brand you usually smoke, [regular brand], might be a little less harmful, no different, or a little more harmful, compared to other cigarette brands?	5 to 9 (2006–2013)	“A little less harmful,” “no different,” “a little more harmful,” or “Don’t Know” (NB. Responses were reversed coded and “Don’t Know” responses recoded as “no different” for analysis purpose)
<b>Experiences</b>		
Light cigarettes are smoother on your throat and chest than regular cigarettes.	1 to 8 (2002–2010)	Rated on a 5-point scale ranging from “strongly agree” to “strongly disagree” (NB. Responses were reversed coded and “Don’t Know” responses recoded as “neither” for analysis purpose)
Thinking about the cigarettes you are currently smoking in relation to other cigarettes, are your cigarettes... Lighter in taste or more intense in taste? Harsher or smoother on your throat?	6 to 9 (2007–2013)	“Lighter,” “about the same,” “more intense,” or “Don’t Know”; and “harsher,” “about the same,” “smoother,” or “Don’t Know,” respectively. (NB. Responses were reversed coded and “Don’t Know” responses recoded as “about the same” for analysis purpose)
<b>Perceptions</b>		
To what extent do any of the following give you useful information on how cigarettes will taste: The colors of the pack itself? The tar and nicotine levels of the brand?	6 to 9 (2007–2013)	“Not at all,” “a little,” “somewhat,” “a lot,” or “Don’t Know” (NB. Being an ordinal scale, “Don’t Know” responses were deemed to be situated somewhere between “not at all” and “a little,” thus, they were recoded as “a little” for analysis purpose)
Does the term SMOOTH on cigarette packs mean that the cigarettes are supposed to be some form of light, mild, or low-tar cigarette?	4 to 8 (2005–2010)	“Yes,” “No,” or “Don’t Know” (NB. “Don’t Know” responses were recoded as “No” for analysis purpose)

Estimates of means and proportions were computed on weighted data. In order to take into account the correlated nature of the longitudinal data, we used generalized estimating equations (GEE) to compute parameter estimates. A strength of GEE is that it allows cases with at least one wave of data to be included in analyses, thus allowing inclusion of data from replenishment samples, which helps minimize attrition bias. We assumed an unstructured working correlation structure given the large sample and used robust variance to compute the *P*-values for the parameter estimates. We tested for significant main effects of survey wave to assess change over time (both linear and quadratic trends, and also pre–post differences using simple contrast) in outcomes of interest. We also tested for significant interactions between wave and potential moderators, such as brand or strength-style smoked to assess whether the patterns of change over time in outcomes of interest differed between smokers of Marlboro versus other top-selling brands combined and between smokers of different strength styles. In all models, we included the following invariant control variables (gender, minority status, and year of recruitment) and time-varying covariates (age, education, income, cigarettes per day, any recent quit attempts, and survey mode).

## Results

### Sample Characteristics

Baseline sample characteristics are presented in Table 2. Over half of respondents were women, with the majority being white and nearly

70% aged 40 years and above. Nearly 60% had at least some college education, and nearly 60% reported having annual household income of ≥\$30 000 per year. Slightly more than one-third reported smoking 10 or fewer cigarettes a day.

### Cigarette Variant Relative Harm Beliefs Before and After the Ban

Figure 1 shows the patterns of change over time in mean level of endorsement of lights beliefs and Table 3 presents the GEE results testing for main and interaction effects. Measures of misperceptions about lights cigarettes (both individual belief items and their combined scale) showed a decline between 2002 and 2005 (significant linear trend) and then a resurgence after 2006 before plateauing (significant quadratic trend) through 2013, with no clear effect of the ban on “light/low” descriptors (nonsignificant pre–post ban effect). As expected, the overall level of endorsement of the beliefs about light cigarettes differed by brand strength style, being highest among “ultralight” smokers and lowest among “regular full-flavored” smokers ( $P < .001$ ). However, overall level of endorsement of these beliefs was lower among Marlboro smokers than among those who smoked other top-selling brands (all differences significant at  $P < .01$ ). Overall endorsement of the belief that one’s own brand is less harmful than others remained stable between 2007 and 2013 although significant differences between strength styles ( $P < .001$ ) but not between Marlboro and other top brands smoked ( $P = .439$ ) were observed. Also, differences between strength styles showed

**Table 2.** Baseline Characteristics of International Tobacco Control US Sample, 2002–2013 ( $n = 7072$ )<sup>a</sup>

Age in years (%)	
18–24	15.0
25–39	15.6
40–54	37.1
≥55	32.3
Gender (% female)	56.2
Identified minority group (%)	19.8
Education (%)	
Low	42.1
Moderate	40.1
High	17.8
Income (%)	
Low	34.8
Moderate	32.2
High	26.5
No information	5.5
Cigarettes per day (%)	
1–10	35.2
11–20	45.7
21–30	12.0
≥31	7.1
Wave (year) of recruitment ( $n$ )	
Wave 1 (2002)	2013
Wave 2 (2003)	637
Wave 3 (2004)	856
Wave 4 (2005)	687
Wave 5 (2006)	537
Wave 6 (2007)	501
Wave 7 (2008)	294
Wave 8 (2010)	271
Wave 9 (2013)	1276

NB. Percentages are based on unweighted data.

<sup>a</sup>Among current factory-made cigarette smokers with data on cigarette brand attribute.

narrowing post-ban (a significant year by strength interaction [ $P < .001$ ], [Figure 1B](#)) but such pattern of change did not differ by brand smoked (year  $\times$  strength  $\times$  brand interaction not significant). More than 3 years post-implementation of the light/low descriptor ban, 12% of current smokers still mistakenly reported that their own brands of cigarettes were less harmful than other brands.

### Cigarette Variant Sensory Beliefs/Experiences Before and After the Ban

GEE results ([Table 3](#)) revealed that overall endorsement of lights cigarettes being smoother on the throat and chest showed a gradual decline over time between 2002 and 2006, but recovered somewhat by 2007 before plateauing (53.8% endorsing this belief in 2013) with no clear effect of the 2010 ban ([Supplementary Figure 2A](#)), although the pre-ban mean endorsement was significantly higher than that of post-ban ( $P = .038$ ). Notably, the pattern of changes in this belief was similar across the different cigarette strength styles (regardless of brand) although the overall level differed by strength styles, being highest among “ultralight” smokers and lowest among “regular full-flavor” smokers ( $P < .001$ ). However, no overall differences in responses or in overall trends over time by brand smoked were observed. The trends by strength styles also did not differ by brand smoked.

Reported endorsement of one’s own brand being lighter in taste assessed between 2007 and 2013 indicated that the initial decline

between 2007 and 2008 was not sustained but increased (42.1% endorsement in 2013) after the policy change, with a greater increase among “light” smokers than among “ultralight” and “regular” smokers (year by strength interaction significant at  $P = .013$ , see [Supplementary Figure 2B](#)). The pattern of change for believing one’s own brand is smoother on the throat (with 59% endorsing it in 2013) was very similar to that of believing one’s own brand is lighter in taste but the trends did not differ by strength styles ([Supplementary Figure 2C](#)). For both measures of sensory effects of their cigarettes, no significant differences by brand smoked were observed and the trends by strength styles also did not differ by brand smoked.

### Cigarette Variant Taste Indicators Before and After the Ban

GEE results ([Table 3](#)) showed that the trend in reported utility of pack color as an indicator of taste differed by strength styles ( $P = .004$ ). Reporting pack color as an indicator of taste remained stable for regular smokers throughout the study period (36.3% reporting it in 2013). However, for both “ultra” and “light” smokers, endorsement of this taste indicator showed an initial decline between 2007 and 2008 and then an increase (with 37.9% and 45.3%, respectively, reporting it in 2013) following the 2010 ban on “light/low” descriptors (significant quadratic trend, see [Supplementary Figure 3A](#)).


The perception that nicotine and tar levels are useful indicators of taste showed a similar initial decline followed by an increase after the removal of tar/nicotine numbers in 2009 and the descriptor ban in 2010 (with 51.4% perceiving this in 2013) but this trend did not differ by strength styles ([Supplementary Figure 3B](#)). The pattern of change also did not differ by brand smoked. However, there was a clear overall difference by strength styles ( $P = .004$ ) with “ultralight” smokers being more likely to perceive nicotine and tar levels as useful taste indicators than regular smokers, with no difference between “light” and regular smokers.

Perceiving the term “smooth” on pack as indicating that the cigarette is a “light/low” cigarette showed an increase between 2005 and 2007 and subsequently plateaued over the remainder of the study period ([Supplementary Figure 3C](#), significant quadratic trend; 34.1% perceiving this in 2010) with no clear differences by strength styles in either mean level of endorsement or pattern of change over time ([Table 3](#)).

For all three measures of taste indicator perceptions, there was no evidence of a significant difference in either overall level or pattern of change between Marlboro smokers and the smokers of other brands.

## Discussion

Consistent with findings in Australia, and the United Kingdom,<sup>6,7</sup> this study confirms that the removal of misleading terms such as “light,” “mild,” and “low” in the United States, even after tar and nicotine numbers had already been removed from packs and advertising, had little impact on changing consumer misperceptions that some cigarettes are less harmful than others. A nontrivial number of current smokers (12%, which roughly translates to 5 million smokers nationwide) still reported that their own cigarette brands were less harmful than others. Also consistent with other observational studies,<sup>3,9</sup> this study shows that following the ban, smokers have increasingly relied on pack colors, color terms, and other descriptors (eg, “smooth”), as well as sensory experiences of smoother or lighter taste, to help identify their preferred cigarette brand styles and determine which brand styles are light versus regular cigarettes.



**Figure 1.** A and B: Trends in belief about the health benefits of “light” cigarettes and lower risk of one’s own brand, before and after the implementation of the removal of “light,” “mild,” and “low” descriptors in the United States. Weighted estimates adjusted for socio-demographic and smoking-related variables along with survey mode and year recruited into the survey.

Hence, the descriptor ban does not appear to have advanced the objective of eliminating smoker misperceptions that some cigarette sub-brands or variants—including those previously labeled and marketed as light, mild or low—are less harmful than others (which may lead some to smoke brands or styles they mistakenly think are less harmful instead of trying to quit smoking).

Consistent with the trend reported in Yong et al.,<sup>7</sup> this study shows that misperceptions about “light” cigarettes among current smokers declined from a high level in 2002 to a low but still significant level in 2006. This decline was possibly due to increased public awareness and understanding of the lights deception that were highlighted in several high-profile court cases about the lights fraud in the United States. By 2007, as the issue waned, misperceptions started to trend up and plateau showing little change following the 2009 removal of tar and nicotine numbers or the 2010 descriptor ban. The findings suggest that the removal of the numbers and descriptors from cigarette packs and ads had no observable impact on misperceptions. This is not surprising for two reasons. First, the policy changes were not accompanied by any public education campaign or wide media coverage in the United States, the most plausible driver of change in countries like Australia and the

United Kingdom that had implemented similar descriptor bans.<sup>6,7</sup> Second, the policy did not address other interrelated cues used by smokers as indicators of risk, including colors, color words, and other descriptors (eg, “smooth”), perceived taste or smoothness, and design features, which have been shown in past research to be potent conveyors of reduced-risk messages.<sup>3,9–11,32</sup> This study’s observed increase in the perceived utility of various indicators of cigarette taste and reported sensory experiences of smoothness and/or lightness of cigarettes further supports the role of these other cues in supporting smokers’ reduced-risk beliefs.

This study also confirms that the levels of misperceptions correspond to cigarette strength-style levels in a dose-response manner (highest among smokers of “ultralight” cigarettes and lowest among smokers of “regular” cigarettes), as found previously.<sup>6,22</sup> One interesting finding is the narrowing of the differences in misperceptions between smokers of different strength styles but only for beliefs about the harmfulness of one’s own cigarettes. It remains unclear to what extent the narrowing was due to the two policy changes as other related measures did not change in this way, as we might expect if the policy changes had any positive effect in reducing misperceptions. Whether this effect or the others will be sustained over time is unclear.

**Table 3.** Beliefs, Reported Experiences, and Perceptions of Top-Selling Cigarette Brand Variants in the United States: Results From GEE Modeling

Outcomes	n (N)	Survey year (Wave)				Overall P-value	Trend analysis			Strength style		Brand smoked	
		2007 (w6)	2008 (w7)	2010 (w8)	2013 (w9)		Linear	Quadratic	Pre vs. post-ban	Ultra vs. regular	Light vs. regular	Overall P-value	Marlboro vs. other
<b>Beliefs</b>													
Lights less harmful <sup>d</sup>													
B (SE)	5879 (10 573)	-0.08 (0.04)	Ref	-0.02 (0.06)	—	<.001	-0.17 <sup>c</sup> (0.03)	0.02 <sup>c</sup> (0.00)	-0.11 (0.08)	0.42 <sup>c</sup> (0.04)	0.16 <sup>c</sup> (0.03)	<.001	-0.11 <sup>c</sup> (0.03)
Lights less tar <sup>d</sup>													
B (SE)	5878 (10 569)	-0.04 (0.05)	Ref	-0.02 (0.06)	—	<.001	-0.13 <sup>c</sup> (0.03)	0.01 <sup>a</sup> (0.00)	-0.01 (0.08)	0.31 <sup>c</sup> (0.04)	0.10 <sup>b</sup> (0.03)	<.001	-0.08 <sup>a</sup> (0.03)
Lights benefit scale <sup>e</sup>													
B (SE)	5878 (10 567)	-0.06 (0.04)	Ref	-0.02 (0.06)	—	<.001	-0.15 <sup>c</sup> (0.02)	0.01 <sup>a</sup> (0.00)	-0.05 (0.07)	0.35 <sup>c</sup> (0.04)	0.13 <sup>c</sup> (0.03)	<.001	-0.10 <sup>a</sup> (0.03)
Own cigs less harmful <sup>e</sup>													
B (SE)	3027 (4611)	-0.01 (0.03)	Ref	-0.02 (0.03)	-0.00 (0.03)	.890	-0.00 (0.15)	0.00 (0.01)	-0.03 (0.05)	0.41 <sup>c</sup> (0.03)	0.24 <sup>c</sup> (0.02)	<.001	-0.02 (0.02)
<b>Experiences</b>													
Lights smoother <sup>d</sup>													
B (SE)	5879 (10 571)	0.06 (0.04)	Ref	-0.05 (0.07)	—	<.001	-0.13 <sup>c</sup> (0.03)	0.01 <sup>a</sup> (0.00)	-0.16 <sup>c</sup> (0.08)	0.76 <sup>c</sup> (0.03)	0.50 <sup>c</sup> (0.03)	<.001	-0.05 (0.03)
Own cigs lighter <sup>f</sup>													
OR (95% CI)	3026 (4609)	1.64 <sup>f</sup> (1.33–2.02)	Ref	1.04 (0.79–1.37)	1.94 <sup>c</sup> (1.43–2.64)	<.001	0.02 <sup>c</sup> (0.00–0.06)	1.32 <sup>c</sup> (1.21–1.45)	0.97 (0.65–1.46)	16.73 <sup>c</sup> (12.42–22.52)	6.26 <sup>c</sup> (5.20–7.55)	<.001	0.88 (0.73–1.06)
Own cigs smoother													
OR (95% CI)	3025 (4607)	1.85 <sup>c</sup> (1.51–2.26)	Ref	1.11 (0.87–1.43)	2.42 <sup>c</sup> (1.77–3.29)	<.001	0.01 <sup>c</sup> (0.00–0.02)	1.42 <sup>c</sup> (1.29–1.55)	1.03 (0.71–1.49)	3.87 <sup>c</sup> (2.96–5.06)	1.99 <sup>c</sup> (1.67–2.38)	<.001	0.89 (0.74–1.07)
<b>Perceptions</b>													
Pack color <sup>f</sup>													
B (SE)	3030 (4615)	0.10 <sup>a</sup> (0.05)	Ref	0.07 (0.06)	0.23 <sup>b</sup> (0.07)	.003	-0.94 <sup>c</sup> (0.27)	0.06 <sup>c</sup> (0.02)	0.05 (0.08)	0.01 (0.06)	-0.00 (0.04)	.970	0.03 (0.04)
Nicotine and tar levels													
B (SE)	3022 (4606)	0.11 <sup>a</sup> (0.04)	Ref	0.13 <sup>a</sup> (0.05)	0.25 <sup>c</sup> (0.07)	.001	-0.86 <sup>c</sup> (0.27)	0.06 <sup>c</sup> (0.02)	0.12 (0.08)	0.18 <sup>b</sup> (0.06)	0.02 (0.04)	.004	0.01 (0.04)
Term “smooth” <sup>a,d</sup>													
OR (95% CI)	3394 (6026)	1.39 <sup>a</sup> (1.13–1.69)	Ref	1.25 (0.99–1.59)	—	<.001	5.51 <sup>c</sup> (2.53–11.98)	0.87 <sup>c</sup> (0.81–0.93)	1.72 <sup>a</sup> (1.11–2.64)	0.99 (0.78–1.27)	1.13 (0.97–1.33)	.237	0.88 (0.75–1.04)

B = regression estimates (positive coefficients refer to higher, while negative coefficients refer to lower, mean level of endorsement of each outcome relative to the reference group); CI = confidence interval; GEE = generalized estimating equations; n = number of unique individual observations; N = number of person-wave observations; OR = odds ratios; Ref = reference group (Survey year 2008, the most proximal year prior to the ban, was set as the reference group for comparison with other survey year); SE = standard error; —, data not available as question not asked in that survey year.

<sup>a</sup>Significant (in bold) at P < .05.

<sup>b</sup>Significant (in bold) at P < .01.

<sup>c</sup>Significant (in bold) at P < .001.

<sup>d</sup>Results for this outcome with data prior to survey year 2007 are not shown (available in [Supplementary Appendix](#)).

<sup>e</sup>Data analyzed excluded wave 5 (ie, only included waves 6 to 9) because of question order and skip pattern at wave 5 being different to that of subsequent waves.

<sup>f</sup>Significant wave x strength interaction (P < .001, P = .018, and P = .004, respectively).

The additional data available from this study show that the declining trend in the belief that “light” cigarettes are smoother than regular cigarettes reported by Borland et al.<sup>6</sup> was not sustained over time as the level of this sensory belief remained relatively stable in recent years with little change following the descriptor ban. However, of concern is the upward trend observed post-ban on more recent measures of smokers’ comparative experiences of lightness and smoothness of their own brand as compared to others (reported by 42% and 59%, respectively, in 2013), although the reason for this uptick is unclear. Nevertheless, our data show a clear correspondence between strength-style of cigarettes and their reported sensory effects, suggesting that the design features of cigarettes—in particular filter ventilation<sup>33</sup> but also including known and perceived historical labeling, as well as current color coding, coloring, color terms, and other descriptors—will continue to produce and reinforce smoker misperceptions of product characteristics and product safety<sup>10</sup> regardless of how these brand variant differences are described.<sup>34</sup>

This study also shows that a significant number of smokers (as high as one in two), across all strength styles and irrespective of brand family, appear to have come to understand that, post ban, they can use pack color, color terms, and other descriptors such as “smooth” to identify cigarettes they mistakenly believe to be less harmful/risky. The significantly greater use of nicotine and tar yield information among “ultralight” smokers (presumably through color coding and other proxies following the removal of tar numbers) is consistent with the manufacturer’s more common use of such numbers on these brand variants and in their advertising in the past. Indeed, following the descriptor ban color has become a key visual signifier differentiating one variant from another and the reliance on this strategy post-ban is what was helping to maintain the misperceptions about product risks among US consumers.<sup>3,9,11,14,15</sup>

Data from this study did not reveal any clear evidence of trend differences in beliefs, experiences and perceptions about different cigarettes between the dominant brand Marlboro and the other top-selling brands. This finding is rather surprising given the known efforts made by Philip Morris USA just prior to the ban to educate consumers (via pack inserts and onsets) on how to identify particular brands/sub-brands based on new color coding.<sup>3,11</sup> Nevertheless, the overall level of misperceptions was significantly lower among Marlboro smokers than that among smokers of other top-selling brands, possibly due to the effect of the information campaigns by Philip Morris USA to inform its customers via pack inserts that lights cigarettes are not less dangerous as part of their efforts to mitigate the negative impacts of the publicity surrounding the court cases regarding the lights deception.<sup>35</sup>

A few study limitations warrant some discussion. First, effects found may be underestimated due to the use of self-report data which may be affected by social desirability biases (eg, the discussion of equivalent harmfulness of so-called “light” cigarettes might have inhibited some people’s preparedness to report differences) and/or misclassification errors. Second, our sample excluded those with missing data on brand and brand varieties which could limit the generalizability of our findings. Third, our study only evaluated the relatively short-term impact of the descriptor ban. Longer term trend and impact analyses await future study. That said, it seems unlikely that differences will emerge with time, unless the ban has a much larger influence on those taking up smoking than it has had on existing smokers, and we can see no good reason why that might happen. Because of limitations inherent in the survey questions and answers, this study was also unable to determine what, exactly, smokers meant when they reported that the cigarettes they smoked were less

harmful than others. Further research would be needed to identify which specific brands, variants or types of cigarettes those smokers think are more harmful (eg, other variants of the same brand, other brands of the same variant, other strength styles, or possibly some other specific cigarette brand they think are the most harmful).

In conclusion, this study confirms that the removal of “light,” “mild,” and “low” descriptors from cigarette packaging and advertising pursuant to the related ban in the Tobacco Control Act—following the removal of nicotine and tar numbers from cigarette packs and ads—has not corrected consumers’ misperceptions that some cigarettes are safer than others. The defacto light/low descriptor ban in the United States has also led to an increase in the number of consumers who report relying upon on other brand descriptors (eg, smooth) and other features of the package and product to differentiate brands by (inaccurately) perceived differences in harmfulness. These findings provide further support for FDA action to remove the elements of product packaging and engineering (such as color coding and the descriptor “smooth”) that contribute to consumer misperceptions regarding product risk or to enforce against manufacturers marketing cigarettes with those misleading elements. The existing FSPTCA clearly prohibits manufacturers from having misleading labeling or ads or making explicit or implicit reduced-risk or reduced-exposure claims about any cigarette brand or sub-brand unless the claims are not false or misleading and the manufacturer has first obtained a modified risk tobacco product order from FDA. These findings also support the introduction of standardized packaging and other standardized product characteristics, including the regulation of product engineering such as filter ventilation, as additional strategies to help minimize consumer misperceptions that some cigarettes are safer, which can delay or prevent smoking cessation.

## Supplementary Material

Supplementary Appendix, Figures 1–3 and Tables 1 and 2 can be found online at <http://www.ntr.oxfordjournals.org>

## Funding

The ITC US Survey is supported by multiple grants including R01 CA 100362 and P50 CA111236 and also in part from grant P01 CA138389 and an ITC pilot study grant (Medical University of South Carolina, Charleston, South Carolina), all funded by the National Cancer Institute of the United States, and Robert Wood Johnson Foundation (045734).

## Declaration of Interests

*KMC has received grant funding from Pfizer, Inc to study the impact of a hospital-based tobacco cessation intervention and also has served as an expert witness in litigation filed against the tobacco industry. ENL is a former Director of the Office of Policy at Food and Drug Administration’s Center for Tobacco Products but ENL’s participation in this article is entirely independent of his FDA affiliation. RJO has served as a consultant to the FDA on tobacco regulation as a member of the Tobacco Products Scientific Advisory Committee. JFT and DH have both received payment as an expert witness in litigation involving the tobacco industry. All other authors have no conflicts of interest to declare.*

## Acknowledgments

All waves of the ITC US study have received ethical approval from the relevant research ethics committee at the Medical University of South Carolina, United States and Roswell Park Cancer Institute, United States.



## References

- Family Smoking Prevention and Tobacco Control Act. 2009. [www.fda.gov/TobaccoProducts/GuidanceComplianceRegulatoryInformation/ucm261829.htm](http://www.fda.gov/TobaccoProducts/GuidanceComplianceRegulatoryInformation/ucm261829.htm). Accessed September 14, 2015.
- US Department of Health and Human Services. Guidance for industry and FDA staff: Use of “light”, “mild”, “low”, or similar descriptors in the label, labeling, or advertising of tobacco products. 2010. [www.fda.gov/tobaccoproducts/guidancecomplianceregulatoryinformation/ucm214597.htm](http://www.fda.gov/tobaccoproducts/guidancecomplianceregulatoryinformation/ucm214597.htm). Accessed September 14, 2015.
- Connolly GN, Alpert HR. Has the tobacco industry evaded the FDA's ban on 'Light' cigarette descriptors? *Tob Control*. 2014;23(2):140–145. doi:10.1136/tobaccocontrol-2012-050746.
- Tindle HA, Shiffman S, Hartman AM, Bost JE. Switching to “lighter” cigarettes and quitting smoking. *Tob Control*. 2009;18(6):485–490. doi:10.1136/tc.2008.029314.
- Hammond D. Tobacco packaging and labeling policies under the U.S. Tobacco Control Act: research needs and priorities. *Nicotine Tob Res*. 2012;14(1):62–74. doi:10.1093/ntr/ntr182.
- Borland R, Fong GT, Yong HH, et al. What happened to smokers' beliefs about light cigarettes when “light/mild” brand descriptors were banned in the UK? Findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2008;17(4):256–262. doi:10.1136/tc.2007.023812.
- Yong HH, Borland R, Cummings KM, et al. Impact of the removal of misleading terms on cigarette pack on smokers' beliefs about 'light/mild' cigarettes: cross-country comparisons. *Addiction*. 2011;106(12):2204–2213. doi:10.1111/j.1360-0443.2011.03533.x
- Bansal-Travers M, Hammond D, Smith P, Cummings KM. The impact of cigarette pack design, descriptors, and warning labels on risk perception in the U.S. *Am J Prev Med*. 2011;40(6):674–682. doi:10.1016/j.amepre.2011.01.021.
- Mutti S, Hammond D, Borland R, Cummings MK, O'Connor RJ, Fong GT. Beyond light and mild: cigarette brand descriptors and perceptions of risk in the International Tobacco Control (ITC) Four Country Survey. *Addiction*. 2011;106(6):1166–1175. doi:10.1111/j.1360-0443.2011.03402.x.
- O'Connor RJ, Caruso RV, Borland R, et al. Relationship of cigarette-related perceptions to cigarette design features: findings from the 2009 ITC U.S. Survey. *Nicotine Tob Res*. 2013;15(11):1943–1947. doi:10.1093/ntr/ntt075.
- Bansal-Travers M, O'Connor R, Fix BV, Cummings KM. What do cigarette pack colors communicate to smokers in the U.S.? *Am J Prev Med*. 2011;40(6):683–689. doi:10.1016/j.amepre.2011.01.019.
- King B, Borland R. What was “light” and “mild” is now “smooth” and “fine”: new labelling of Australian cigarettes. *Tob Control*. 2005;14(3):214–215. doi:10.1136/tc.2005.011692.
- Thrasher JF, Hammond D, Arillo-Santillan E. The alchemy of Marlboro: transforming 'light' into 'gold' in Mexico. *Tob Control*. 2010;19(4):342–343. doi:10.1136/tc.2010.037044.
- Hammond D, Dockrell M, Arnott D, Lee A, McNeill A. Cigarette pack design and perceptions of risk among UK adults and youth. *Eur J Public Health*. 2009;19(6):631–637. doi:10.1093/eurpub/ckp122.
- Hammond D, Parkinson C. The impact of cigarette package design on perceptions of risk. *J Public Health*. 2009;31(3):345–353. doi:10.1093/pubmed/fdp066.
- Rees VW, Kreslake JM, Cummings KM, et al. Assessing consumer responses to potential reduced-exposure tobacco products: a review of tobacco industry and independent research methods. *Cancer Epidem Biomar*. 2009;18(12):3225–3240. doi:10.1158/1055-9965.EPI-09-0946.
- Kozlowski LT, Goldberg ME, Yost BA, Ahern FM, Aronson KR, Sweeney CT. Smokers are unaware of the filter vents now on most cigarettes: results of a national survey. *Tob Control*. 1996;5(4):265–270. [www.ncbi.nlm.nih.gov/pmc/articles/PMC1759525/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1759525/). Accessed September 14, 2015.
- Pages RA. *Research Report: 6906 Annual Report-Biological Effects of Smoke*. Philip Morris; 1978. Report No.: Bates No. 2001243600-2001243673. Legacy Tobacco Documents Library, UCSF. <https://industrydocuments.library.ucsf.edu/tobacco/docs/smky0116>. Accessed December 21, 2015.
- Rapp KE. *Memorandum to R.A. Pages: Model III WSC Activities in the Salmonella/Microsome Assay: WSC Activity as a Function of Cigarette Filter Dilution*. Philip Morris; 1979. Report No.: Bates No. 2501507859-2501507866. Legacy Tobacco Documents Library, UCSF. <https://industrydocuments.library.ucsf.edu/tobacco/docs/hrgh0115>. Accessed December 21, 2015.
- Levins RJ. *Research Report: 6908 Contribution of the Blend Components of Low Tar Reference Cigarette to CCSC, PAH III Yield and Salmonella Activity*. Philip Morris; 1984. Report No.: Bates No. 2001291173-2001291198. Legacy Tobacco Documents Library, UCSF. <https://industrydocuments.library.ucsf.edu/tobacco/docs/lxpd0123>. Accessed December 21, 2015.
- Kozlowski LT, O'Connor RJ. Cigarette filter ventilation is a defective design because of misleading taste, bigger puffs, and blocked vents. *Tob Control*. 2002;11(suppl 1):i40–50. doi:10.1136/tc.11.suppl\_1.i40.
- Shiffman S, Pillitteri JL, Burton SL, Rohay JM, Gitchell JG. Smokers' beliefs about “Light” and “Ultra Light” cigarettes. *Tob Control*. 2001;10(suppl 1):i17–23. doi:10.1136/tc.10.suppl\_1.i17.
- Elton-Marshall T, Fong GT, Yong H-H, et al. Smokers' sensory beliefs mediate the relation between smoking a 'light/low tar' cigarette and perceptions of harm. *Tob Control*. 2015;24(suppl 4):iv21–27. doi:10.1136/tobaccocontrol-2014-051977.
- Borland R, Yong HH, King B, et al. Use of and beliefs about light cigarettes in four countries: findings from the International Tobacco Control Policy Evaluation Survey. *Nicotine Tob Res*. 2004;6(suppl 3):S311–321. doi:10.1080/1462220412331320716.
- U.S. Federal Trade Commission. *Rescission of FTC Guidance Concerning the Cambridge Filter Method*. Federal Register; 2008;74500–74505. [www.gpo.gov/fdsys/pkg/FR-2008-12-08/pdf/E8-28969.pdf](http://www.gpo.gov/fdsys/pkg/FR-2008-12-08/pdf/E8-28969.pdf). Accessed January 22, 2016.
- FDA. Letter to Philip Morris USA, Inc. Marketing Marlboro Lights cigarettes with an onsert. 2010. [www.fda.gov/downloads/TobaccoProducts/Labeling/RegulationsGuidance/UCM284131.pdf](http://www.fda.gov/downloads/TobaccoProducts/Labeling/RegulationsGuidance/UCM284131.pdf). Accessed September 14, 2015.
- O'Connor RJ. What brands are US smokers under 25 choosing? *Tob Control*. 2005;14(3):213. doi:10.1136/tc.2004.010736.
- CDC. Cigarette Brand Preference Among Middle and High School Students Who Are Established Smokers - United States, 2004 and 2006. *MMWR*. 2009;58(5):112–115. [www.cdc.gov/mmwr/preview/mmwrhtml/mm5805a3.htm](http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5805a3.htm). Accessed September 14, 2015.
- Fong GT, Cummings KM, Borland R, et al. The conceptual framework of the International Tobacco Control (ITC) Policy Evaluation Project. *Tob Control*. 2006;15(suppl 3):iii3–11. doi:10.1136/tc.2005.015438.
- Thompson ME, Fong GT, Hammond D, et al. Methods of the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2006;15(suppl 3):iii12–18. doi:10.1136/tc.2005.013870.
- Cornelius ME, Cummings KM, Fong GT, et al. The prevalence of brand switching among adult smokers in the USA, 2006–2011: findings from the ITC US surveys. *Tob Control*. 2015;24(6):609–615. doi:10.1136/tobaccocontrol-2014-051765.
- Cohen JE, Yang J, Donaldson EA. Impact of the removal of light and mild descriptors from cigarette packages in Ontario, Canada: switching to “light replacement” brand variants. *Prev Med*. 2014;69:120–125. doi:10.1016/j.ypmed.2014.08.037.
- King B, Borland R. The “low-tar” strategy and the changing construction of Australian cigarettes. *Nicotine Tob Res*. 2004;6(1):85–94. doi:10.1080/14622200310001656907.
- Borland R, Savvas S. The effects of variant descriptors on the potential effectiveness of plain packaging. *Tob Control*. 2014;23(1):58–63. doi:10.1136/tobaccocontrol-2012-050736. doi:10.1136/tobaccocontrol-2012-050736.
- Cummings KM, Hyland A, Bansal MA, Giovino GA. What do Marlboro Lights smokers know about low-tar cigarettes? *Nicotine Tob Res*. 2004;6(suppl 3):S323–332. doi:10.1080/14622200412331320725.