

Middle East Consensus Statement on the Diagnosis and Management of Functional Gastrointestinal Disorders in <12 Months Old Infants

Yvan Vandenplas, Muath Abdurrahman Alturaiki^{*}, Wafaa Al-Qabandi[†], Fawaz AlRefaee[‡], Ziad Bassil[§], Bassam Eid^{||}, Ahmed El Beleidy[¶], Ali Ibrahim Almehaidib^{**}, Pierre Mouawad^{††}, and Maroun Sokhn^{§,††}

Department of Pediatrics, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Brussels, Belgium, ^{}Department of Pediatrics, King Salman Hospital, Riyadh, Saudi Arabia, [†]Faculty of Medicine, Kuwait University, Kuwait City, Kuwait, [‡]Department of Pediatrics, Al Adan Hospital, Kuwait City, Kuwait, [§]Department of Pediatric Gastroenterology, Hepatology and Pediatric Nutrition, St. Joseph Hospital, Beirut, Lebanon, ^{||}Department of Pediatric Gastroenterology, Hotel Dieu de France, Beirut, Lebanon, [¶]Department of Pediatrics, Cairo University, Cairo, Egypt, ^{**}Department of Pediatric Gastroenterology, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia, ^{††}Department of Pediatric Gastroenterology, St. Georges Orthodox, Beirut, Lebanon*

This paper covers algorithms for the management of regurgitation, constipation and infantile colic in infants. Anti-regurgitation formula may be considered in infants with troublesome regurgitation, while diagnostic investigations or drug therapy are not indicated in the absence of warning signs. Although probiotics have shown some positive evidence for the management of functional gastrointestinal disorders (FGIDs), the evidence is not strong enough to make a recommendation. A partially hydrolyzed infant formula with prebiotics and β -palmitate may be considered as a dietary intervention for functional constipation in formula fed infants. Lactulose has been shown to be effective and safe in infants younger than 6 months that are constipated. Macrogol (polyethylene glycol, PEG) is not approved for use in infants less than 6 months of age. However, PEG is preferred over lactulose in infants >6 months of age. Limited data suggests that infant formula with a partial hydrolysate, galacto-oligosaccharides/fructo-oligosaccharides, added β -palmitate may be of benefit in reducing infantile colic in formula fed infants in cases where cow's milk protein allergy (CMPA) is not suspected. Evidence suggests that the use of extensively hydrolyzed infant formula for a formula-fed baby and a cow's milk free diet for a breastfeeding mother may be beneficial to decrease infantile colic if CMPA is suspected. None of the FGIDs is a reason to stop breastfeeding.

Key Words: Breast feeding, Colic, Constipation, Diarrhea, Formula feeding, Gastrointestinal diseases, Regurgitation

Received : June 24, 2016, Revised : August 22, 2016, Accepted : August 31, 2016

Corresponding author: Yvan Vandenplas, Department of Pediatrics, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Laarbeeklaan 101, 1090 Brussels, Belgium. Tel: +32-2-477-57-81, Fax: +32-2-477-57-83, E-mail: yvan.vandenplas@uzbrussel.be

Copyright © 2016 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Functional gastrointestinal disorders (FGIDs) such as regurgitation, constipation and infantile colic are extremely common among infants across the globe with an estimated global incidence ranging from 40-60%, thus affecting approximately half the population of infants [1]. Increasing awareness towards the importance of these functional disorders has led to recent publications of consensus statements and international guidelines towards appropriate management of these conditions. An expert review of global data on various FGIDs estimated a prevalence of 10% to 25% for functional constipation, colic and regurgitation, while functional diarrhea has a prevalence below 5% [1]. There are some data available on the incidence of FGIDs in the Middle East Region. This region is known as a highly diversified region in the world in terms of ecology (green valleys and dry yellow deserts), political structures and stability and economic diversity (including countries that classify as among the world's richest and poorest). Health and nutritional status of this region are determined by all these factors.

Although the different FGIDs are discussed separately with independent management algorithms, in real-life they are inter-related and many times present in combination. For e.g., the incidence of other FGIDs is two times higher in infants with colic vs. those without colic [2]. Up to now, unified recommendations for the management of FGIDs in infants, specifically tailored for the Middle East region, do not exist.

METHODS

A panel of experts met on September 5, 2015 in Dubai to develop a primary care Middle East Consensus for the management of common FGIDs in < 12 months old infants, covering regurgitation, functional constipation, colic and functional diarrhea. The panel included nine experts representing four countries in the Middle East countries "Egypt,

Kuwait, Lebanon, and Saudi Arabia" with expertise in general pediatrics, gastroenterology, and nutrition. The panel was led by an international chair with extensive experience on the subject of GI disorders in infants. Each of the statements was discussed within the group before voting. In order to reach consensus, a structured method, previously shown to be efficacious, was utilized [3]. Consensus was formally achieved through a nominal group technique, a structured quantitative method. The group consisted of 10 members voting anonymously. Before the voting took place, the statements were reviewed by each co-author until agreement was reached. A nine-point scale was used (one for strongly disagree to nine for fully agree) [4]. It was decided from beforehand that consensus was reached if over 75% of the votes were "six, seven, eight, or nine". A vote of six and above meant "agreement", nine being an expression of stronger agreement than six. Each statement reached a maximal consensus with all panel members voting 9.

REGURGITATION

There is scarcity of data from Middle East region on the prevalence of regurgitation. Available data from global studies show considerable variability in estimates, with prevalence figures ranging from 6% to 61%, with a median prevalence of 26% [1]. Regurgitation in infants, although considered a self-limiting and benign condition, can pose considerable inconvenience for parents, with increased parental stress and potential impact on quality of life (QoL). The perception of mothers tends to exaggerate the symptoms, and as a result, clear and objective documentation of symptoms and reassurance is the cornerstone of regurgitation management. Regurgitation is not a reason to stop breastfeeding. Feeding techniques and volumes are extremely important in preventing and managing regurgitation. In formula fed infants, anti-regurgitation (AR)-formula could be recommended for infants with troublesome regurgitation. There is no evidence to recommend mixing formula with breast milk, or mixing a feed thickener

with breast milk, or mixing of different formulas. Appropriate feeding is key, irrespective of whether the baby is breastfed or formula-fed, as infants with regurgitation are often overfed (causing the regurgitation) or sometimes underfed (as a consequence of the regurgitation). Regurgitation shows a typical regression pattern of decreasing incidence with age, and thickened formulas 'accelerate' this process of regression. Although some available data suggesting benefits of pre- and probiotics, the evidence is not strong enough for active recommendation of prebiotics or probiotics in this indication (Table 1) [5]. Commercial, readily available AR-formula is preferable over ad hoc thickening at home, due to the negative effects of extra calories and osmolarity changes in home preparations (e.g., preparations using cereals/starch). A practical algorithm for the management of regurgitation in breast- and formula-fed infants was developed (Fig. 1).

CONSTIPATION

There is scarcity of data from the Middle East region on constipation. Available data from global studies show considerable variability in estimates, with prevalence figures ranging from 2.3% to 61%, with a median prevalence of 8% [1]. A study in the Kingdom of Saudi Arabia (KSA) showed that 23.6% of mothers changed the infant formula due to constipation [6]. There are studies that indicate some benefits of prebiotics (galacto-oligosaccharides [GOS]/fructo-oligosaccharides [FOS]) [7,8], and/or prebiotics, partially hydrolyzed formula (pHF), beta-palmitate [9,10], and probiotic strains such as *Lactobacillus reuteri* DSM 17938 and *Bifidobacterium longum* [1,5,6, 11,12]. Probiotics may be considered as an option, although there is not enough evidence to recommend them. Prebiotic oligosaccharides in combination with other ingredients such as beta-palmitate and

Table 1. Consensus Statement on Regurgitation

Consensus statements
<p>Otherwise healthy infants, who present with four or more episodes of regurgitation per day during at least two weeks, and in whom the onset is either after the age of one week or before the age of six months, should be considered as presenting probably with physiologic regurgitation and should be managed by parental reassurance and AR-formula can be used if additional intervention to reduce regurgitation is felt needed.</p> <p>Key features</p> <ul style="list-style-type: none"> • ≥ 4 episodes/day • > 2 weeks duration • Starts > 1 week of age or < 6 months of age <p>Diagnostic investigations are not indicated for the diagnosis and management of "troublesome regurgitation" in the absence of warning signals*</p> <p>*Warning signals requiring investigation in infants with regurgitation or vomiting [5]</p> <ul style="list-style-type: none"> • Bilious vomiting • Gastrointestinal bleeding • Apnea, chronic cough, recurrent aspiration • Consistently forceful vomiting • Onset of vomiting after 6 months of life • Failure to thrive • Diarrhea • Constipation • Fever • Lethargy • Hepatosplenomegaly • Bulging fontanelle • Macro/microcephaly • Seizures • Abdominal tenderness or distension • Documented or suspected genetic/metabolic syndrome




Fig. 1. Algorithm for regurgitation. BF: breastfed, CMPA: cow’s milk protein allergy, AR-formula: anti-regurgitation formula, FF: fomula fed, eHF: extensively hydrolyzed formula, GERD: gastro-esophageal reflux disease.

partial protein hydrolysates have been shown to soften the stools in constipated infants [13,14]. There is no evidence to recommend change in diet e.g., extra fiber and/or extra fluids above the recommended amount during weaning. Macrogol (polyethylene glycol, PEG) is not approved for use in infants less than 6 months of age. However, PEG is preferred over lactulose in infants >6 months of age. Lactulose and PEG dosages are outlined in Table 2.

Duration of PEG/lactulose treatment should be minimal two weeks with good compliance. If symptoms do not improve, compliance should be checked before considering another intervention. Rectal treatment should not be used as a routine and should be restricted to providing acute relief only. Types of rectal treatment are glycerine suppositories, enemas; rectal stimulation should be discouraged (Table 3,

Table 2. Lactulose and PEG Dosage

Lactulose	- 1-2 g/kg, (1-3 mL/kg), once or twice/day
PEG 3350 maintenance	- 0.2-0.8 g/kg/day - 1 to 1.5 g/kg/day for 3 days if disimpaction is needed

PEG: polyethylene glycol.

Fig. 2). Although magnesium is used a lot, there is no data to support its use. The use of castor oil and mineral oil is discouraged, especially in this age group.

INFANTILE COLIC

There is also scarcity of data from the Middle East region on colic. Small-scale epidemiological studies in Egypt showed an estimate ranging from 20% to

Table 3. Consensus Statements on Constipation

Consensus statements
Constipation is defined as “difficult and/or rare defecation (less than one stool per week in breastfed, and less than one stool per three days in formula fed infants), lasting for at least two weeks”.
In an infant presenting with constipation, each of the following manifestations indicates possible organic disease:
1) The absence of meconium within the first 24 hours of life in term infants
2) Abdominal distention
3) Bilious vomiting
4) Failure to thrive
5) Bloody stools
6) Neurodevelopmental delay
7) Anal/sacral abnormalities
Lactulose may be considered as an intervention for functional constipation.
Macrogol (polyethylene glycol, PEG) is preferred over lactulose as an intervention for functional constipation for infants over six months of age.
A partially hydrolysed infant formula with prebiotics and β -palmitate may be considered as a dietary intervention for functional constipation in formula fed infants.
Breastfed babies that have less than one stool per week should be referred, except if there are asymptomatic.
When cow’s milk protein allergy is suspected, a cow’s milk elimination diet may be considered as an intervention for constipation for two to four weeks followed by a challenge.
Rectal treatment should not be used as a routine and should be restricted to providing acute relief only.




Fig. 2. Algorithm for functional constipation. FF: formula fed, BF: breastfed, pHF: partially hydrolyzed formula, GOS: galacto-oligosaccharides, FOS: fructo-oligosaccharides, L: *Lactobacillus*, B: *Bifidobacterium*, Mg: magnesium, PEG: polyethylene glycol, eHF: extensively hydrolyzed formula, CMPA: cow’s milk protein allergy.

37% [15,16], while a study in KSA showed that 32% of mothers changed the infant formula due to infantile colic [6]. Available data from global studies also show considerable variability in estimates, with prevalence figures ranging from 4% to 64%, with a median prevalence of 18% [1]. Changing of feeding techniques may not be effective in the management of true infantile colic. Feeding issues such as under-feeding, overfeeding, infrequent burping, swallowing too much air should be ruled out before initiating other management. Feeding the baby in a vertical position and frequent burping may reduce swallowed air, but there is no evidence that feeding in vertical position and special bottles reduce colic. Colic is often confused with simple fussiness, and adding unsoothable or inconsolable crying as an additional criterion will help identify 'true' colic from simple fussiness.

Five randomized controlled trials assessing two strains of probiotic *L. reuteri* in mostly breastfed infants were identified. Analysis of response rates showed that infants receiving probiotics had a 2.3-fold greater chance of having a 50% or greater decrease in crying/fussing time compared to controls ($p=0.01$). Probiotic supplementation was not associated with any adverse events [17]. Supplementation with the probiotic *L. reuteri* DSM 17938 in breastfed infants appears to be safe and effective for the management of infantile colic. Further research is needed to determine the role of probiotics in infants who are formula-fed [17]. Harb and co-workers also performed a meta-analysis on the efficacy of different therapeutic options in the treatment of infantile colic [18]. They concluded that: six studies included for subgroup meta-analysis on probiotic treatment, notably *L. reuteri*, demonstrated that probiotics appear an effective treatment, with an overall mean difference in crying time at day 21 of -55.8 min/day (95%

confidence interval [CI], -64.4 to -47.3 ; $p=0.001$) [18]. The 3 studies included for subgroup meta-analysis on preparations containing fennel suggest it to be effective, with an overall mean difference of -72.1 min/day (95% CI, -126.4 to -17.7 ; $p<0.001$) [18]. Xu et al. [19] came to a similar conclusion: *L. reuteri* increased colic treatment effectiveness at two weeks (relative risk [RR], 2.84; 95% CI, 1.24-6.50; $p=0.014$) and three weeks (RR, 2.33; 95% CI, 1.38-3.93; $p=0.002$) but not at four weeks (RR, 1.41; 95% CI, 0.52-3.82; $p=0.498$). *L. reuteri* decreased crying time (min/day) at two weeks (weighted mean difference [WMD], -42.89 ; 95% CI, -60.50 to -25.29 ; $p<0.001$) and three weeks (WMD, -45.83 ; 95% CI, -59.45 to -32.21 ; $p<0.001$) (Table 4) [18-21].

In a double-blind, placebo-controlled trial, a pHF, with high beta-palmitate, and a specific prebiotics mixture with GOS/FOS also resulted in a significant reduction in colic within one and two weeks of intervention [22]. Limited evidence suggests that the use of fermented formula with Lactofidus™ (Danone [Nutricia Early Life Nutrition], Amsterdam, The Netherlands) can be effective in treatment of infantile colic [23-25] (Table 5, Fig. 3).

FUNCTIONAL DIARRHEA & DYSCHYZIA

Functional diarrhea and dyschezia are not discussed in this consensus statement. Functional diarrhea is a rare condition in the <1 year age group with an incidence of about 2% (Table 6). The presence of prolonged diarrhea lasting >2 weeks in infants <1 year of age should not be managed by primary care, and mandates a referral to a specialist. Since there is no management for dyschezia except reassurance, this entity is not discussed either.

Table 4. Caution When Assessing Evidence with Probiotics

- On an average, the probiotic *Lactobacillus reuteri* decrease crying time with only one hour approximately [18-21].
- The published evidence on *L. reuteri* DSM 17938 is limited to the drops containing the bacilli, and there are no existing PubMed published studies that show positive results on *L. reuteri*-containing infant formula.

Table 5. Consensus Statements on Infantile Colic

Consensus statements
Otherwise healthy infants younger than three months of age, presenting with fussiness, irritability or frequent unsoothable crying lasting for three or more hours per day, for three or more days per week, for at least one week may be suffering from infantile colic.
Evidence suggests that the use of extensively hydrolysed infant formula in a formula-fed baby and a cow's milk free diet for a breastfeeding mother may be beneficial to decrease infantile colic if CMPA is suspected. However, these data reflects selected infant populations.
Limited data suggests that infant formula with partial hydrolysate, GOS/FOS added β -palmitate may be of some benefit in reducing infantile colic in cases where cow's milk allergy is not suspected in formula fed infants.
There is convincing data that <i>Lactobacillus reuteri</i> DSM 17938 drops decreases infantile colic in selected breastfed infants. There is no evidence that addition <i>L. reuteri</i> DSM 17938 to infant formula alleviates infantile colic.
There is insufficient data to recommend <i>L. reuteri</i> DSM 17938 in all colicky infants.
Anti-acid medications is not a treatment for colicky infants
The evidence of benefit is too limited to recommend herbal medicine such as fennel and peppermint for alleviating infantile colic. Decreasing stimulation and white noise may help in colic. Swaddling may also be effective.




Fig. 3. Algorithm for infantile colic. GI: gastrointestinal, CM-free: cow's milk-free, BF: breastfed, eHF: extensively hydrolyzed formula, FF: formula fed, CMPA: cow's milk protein allergy, L: *Lactobacillus*, pHF: partially hydrolyzed formula, GOS: galacto-oligosaccharides, FOS: fructo- oligosaccharides.

Table 6. Consensus Statement Functional Diarrhea and Dyschezia

Consensus statement
A separate algorithm is not required for functional diarrhea and dyschezia in infants <1 year of age.

DISCUSSION

FGIDs such as regurgitation, infantile colic and constipation are benign conditions, they have a negative impact on the QoL of parents and infants. Although "time is the cure, their immediate management is often frustrating for health care professional because lack of scientific evidence for effective treatment. Because in principle all FGIDs disappear over time, management should be safe and devote of adverse effects. Therefore, reassurance is the cornerstone of the management, endorsed by nutritional therapy. This paper is the result of a consensus discussion among key-opinion leaders in the Middle East region, starting from the recently published international recommendations [6,7,26]. Unneeded investigations and medical treatments are often prescribed in these infants presenting with FGIDs. Medication has failed to improve significantly these conditions. Therefore, the algorithms focus on reassurance, education and dietary intervention. Although the evidence of the data does not change, meaning that guidelines should be followed as much as possible worldwide, there is at the same time the possibility and need for adaptations to local situations. The scientific background of these algorithms was largely discussed in the other papers [6,7,26], and will not be repeated here. These guidelines, adapted to the specific needs of the Middle East region, are in line with previous algorithms [6,7,26]. They are never in contradiction. Local data on the epidemiology of FGIDs were added when available. However, some further changes were incorporated. A relevant and important addition is the recommendation in the regurgitation algorithm for the breastfed baby. FGIDs are almost never a reason to stop breastfeeding. It was felt necessary to add time frames in the constipation algorithm and specific-

ities on the pre- and probiotics were added. Also rectal treatment should be restricted to provide acute relief only. The possible benefit of prebiotics and/or probiotics in combination with a partial whey hydrolysate with beta-palmitate and probiotics were added to the algorithm of infantile colic. The possible benefits of a fermented formula were also added.

CONCLUSION

This Middle-East consensus guideline endorses the previously published recommendations [6,7,26]. However, reviewing the previous publications allowed to highlight some limitations of the previous guidelines. These guidelines are intended to be widespread in the Middle East region.

ACKNOWLEDGEMENTS

The authors thank Nutricia Early Life Nutrition for bringing them together and to provide an unrestricted grant for this consensus.

REFERENCES

1. Vandenplas Y, Abkari A, Bellaiche M, Benninga M, Chouraqui JP, Çokura F, et al. Prevalence and health outcomes of functional gastrointestinal symptoms in infants from birth to 12 months of age. *J Pediatr Gastroenterol Nutr* 2015;61:531-7.
2. Vandenplas Y, Ludwig T, van Elburg R, Bouritius H, Huet F. Association of infantile colic and functional gastrointestinal disorders and symptoms. Paper presented at: ESPGHAN 48th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition; 2015 May 6-9; Amsterdam, The Netherlands. PO-G-016.
3. Vandenplas Y, Dupont C, Eigenmann P, Host A, Kuitunen M, Ribes-Koninckx C, et al. A workshop report on the development of the cow's milk-related symptom score awareness tool for young children. *Acta Paediatr* 2015;104:334-9.
4. McMurray AR. Three decision-making aids: brainstorming, nominal group, and Delphi technique. *J Nurs Staff Dev* 1994;10:62-5.
5. Vandenplas Y, Rudolph CD, Di Lorenzo C, Hassall E, Liptak G, Mazur L, et al. Pediatric gastroesophageal re-

- flux clinical practice guidelines: joint recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition (NASPGHAN) and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition (ESPGHAN). *J Pediatr Gastroenterol Nutr* 2009;49:498-547.
6. Alfaleh K, Alluwaimi E, Aljefri S, Alosaimi A, Behaisi M. Infant formula in Saudi Arabia: a cross sectional survey. *Kuwaiti Med J* 2014;46:328-32.
 7. Moro G, Minoli I, Mosca M, Fanaro S, Jelinek J, Stahl B, et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr* 2002;34:291-5.
 8. Moro GE, Boehm G. Clinical outcomes of prebiotic intervention trials during infancy: a review. *Funct Food Rev* 2012;4:101-13.
 9. Vandenplas Y, Alarcon P, Alliet P, De Greef E, De Ronne N, Hoffman I, et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Paediatr* 2015;104:449-57.
 10. Vandenplas Y, Gutierrez-Castrellon P, Velasco-Benitez C, Palacios J, Jaen D, Ribeiro H, et al. Practical algorithms for managing common gastrointestinal symptoms in infants. *Nutrition* 2013;29:184-94.
 11. Coccorullo P, Strisciuglio C, Martinelli M, Miele E, Greco L, Staiano A. *Lactobacillus reuteri* (DSM 17938) in infants with functional chronic constipation: a double-blind, randomized, placebo-controlled study. *J Pediatr* 2010;157:598-602.
 12. Indrio F, Di Mauro A, Riezzo G, Civardi E, Intini C, Corvaglia L, et al. Prophylactic use of a probiotic in the prevention of colic, regurgitation, and functional constipation: a randomized clinical trial. *JAMA Pediatr* 2014;168:228-33.
 13. Bongers ME, de Lorijn F, Reitsma JB, Groeneweg M, Taminiou JA, Benninga MA. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.
 14. Savino F, Cresi F, Maccario S, Cavallo F, Dalmasso P, Fanaro S, et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolysed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003;91:86-90.
 15. Ali AM. *Helicobacter pylori* and infantile colic. *Arch Pediatr Adolesc Med* 2012;166:648-50.
 16. Ali ASA, Elhady MAA. Prevalence and risk factors for infantile colic in Egyptian infants. *J Am Sci* 2013;9:340-3.
 17. Schreck Bird A, Gregory PJ, Jalloh MA, Risoldi Cochrane Z, Hein DJ. Probiotics for the treatment of infantile colic: a systematic review. *J Pharm Pract* 2016. pii: 0897190016634516. [Epub ahead of print]
 18. Harb T, Matsuyama M, David M, Hill RJ. Infant colic-what works: a systematic review of interventions for breast-fed infants. *J Pediatr Gastroenterol Nutr* 2016;62:668-86.
 19. Xu M, Wang J, Wang N, Sun F, Wang L, Liu XH. The efficacy and safety of the probiotic bacterium *Lactobacillus reuteri* dsm 17938 for infantile colic: a meta-analysis of randomized controlled trials. *PLoS One* 2015; 10:e0141445.
 20. Chau K, Lau E, Greenberg S, Jacobson S, Yazdani-Brojeni P, Verma N, et al. Probiotics for infantile colic: a randomized, double-blind, placebo-controlled trial investigating *Lactobacillus reuteri* DSM 17938. *J Pediatr* 2015;166:74-8.
 21. Savino F, Cordisco L, Tarasco V, Palumeri E, Calabrese R, Oggero R, et al. *Lactobacillus reuteri* DSM 17938 in infantile colic: a randomized, double-blind, placebo-controlled trial. *Pediatrics* 2010;126:e526-33.
 22. Savino F, Palumeri E, Castagno E, Cresi F, Dalmasso P, Cavallo F, et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006;60:1304-10.
 23. van de Heijning BJ, Berton A, Bouritius H, Goulet O. GI symptoms in infants are a potential target for fermented infant milk formulae: a review. *Nutrients* 2014;6:3942-67.
 24. Kearney PJ, Malone AJ, Hayes T, Cole M, Hyland M. A trial of lactase in the management of infant colic. *J Hum Nutr Diet* 1998;11:281-5.
 25. Kanabar D, Randhawa M, Clayton P. Improvement of symptoms in infant colic following reduction of lactose load with lactase. *J Hum Nutr Diet* 2001;14:359-63.
 26. Vandenplas Y, Benninga M, Broekaert I, Falconer J, Gottrand F, Guarino A, et al. Functional gastrointestinal disorder algorithms focus on early recognition, parental reassurance and nutritional strategies. *Acta Paediatr* 2016;105:244-52.