

# Corrigendum: Deactivation of excitatory neurons in the prelimbic cortex via Cdk5 promotes pain sensation and anxiety

Guo-Qiang Wang, Cheng Cen, Chong Li, Shuai Cao, Ning Wang, Zheng Zhou, Xue-Mei Liu, Yu Xu, Na-Xi Tian, Ying Zhang, Jun Wang, Li-Ping Wang & Yun Wang

*Nature Communications* 6:7660 doi: 10.1038/ncomms8660 (2015); Published 16 Jul 2015; Updated 10 Oct 2016

In Fig. 7i of this Article, a different section of the control trace was used to represent the Cdk5 shRNA trace. The correct version of Fig. 7, including a new representative Cdk5 shRNA trace in panel i, appears below as Fig. 1. The original conclusions are not affected by this change and the raw data files from which these traces were selected are now available as Supplementary Data 1 and 2.




Figure 1

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016