

“Theory Based Health Education: Application of Health Belief Model for Iranian Obese and Overweight Students about Physical Activity” in Urmia, Iran

Baratali Rezapour^{1,2}, Firoozeh Mostafavi², Hamidreza Khalkhali³

¹Department of Public Health, School of Health, Urmia University of Medical Sciences, Urmia, Iran, ²Department of Health Education and Health Promotion, Isfahan University of Medical Sciences, Isfahan, Iran, ³Department of Epidemiology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran

Correspondence to:

Dr. Firoozeh Mostafavi, Department of Health Education and Health Promotion, Isfahan University of Medical Sciences, Isfahan, Iran.

E-mail: f_mostafavi@yahoo.com

How to cite this article: Rezapour B, Mostafavi F, Khalkhali H. "Theory based health education: Application of health belief model for Iranian obese and overweight students about physical activity" in Urmia, Iran. *Int J Prev Med* 2016;7:115.

ABSTRACT

Background: Obesity is a major problem in both developed and underdeveloped countries. The aim of this study is to investigate the effects of a physical education program (PEP) on promoting health belief model (HBM) scores, increasing physical activity (PA), and reducing obesity among Iranian high school students.

Methods: This quasi-experimental study was accomplished at four high schools that were randomly divided into two groups of experiment (forty) and control (forty) male students in junior high schools in Urmia, Iran. Students in the experimental group received a PEP during 6 months. The essential parameters were used for evaluating the effects of educational program on HBM, PA, and body mass index (BMI) of students.

Results: After the intervention of 3 and 6 months, the experimental group showed a significant difference on the results of HBM constructs. According to the result of repeated-measures ANOVA, there is a significant difference between the experimental and control groups about the components of PA constructs. Analysis of covariance showed that although BMI reduced in 6 months after intervention, there was no significant difference in BMI.

Conclusions: Results of the study revealed that implementation of PEP was effective on increasing the score of the components of HBM and PA of students.

Keywords: Health belief model, obesity, physical activity, physical education program

INTRODUCTION

Obesity is an important factor underlying chronic diseases, and health workers need to act to deal with it.^[1] Governments have enacted policies to increase physical activity (PA) in schools as a way to fight

childhood obesity.^[2] The worldwide prevalence of obesity has approximated that 1.12 billion adults will be obese globally in 2030.^[3] The prevalence of overweight and obesity in children under 18 years in national Iranian studies was 5.0–13.5% and 3.2–11.9%, respectively.^[1] Obesity has grown modestly in the United States, Canada, England, Italy, Korea, and Spain, but has increased by a further 2–3% in Australia, France, Mexico, and Switzerland.^[4] Sedentary life style (SLS) causes a

Access this article online

Quick Response Code:



Website: www.ijpvmjournal.net/www.ijpm.ir

DOI:
10.4103/2008-7802.191879

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

rapid rise in obesity and overweight over the past few years in Iran. Theory-based health education (TBHE) with health belief model (HBM) and duration of PA can be increased to reduce SLS and the prevalence of obesity and overweight.^[5] This research used physical education program (PEP) to interfere in the current status of students and schools to modify the rates of HBM, PA, and body mass index (BMI) of participants. HBM is one of the behavior change theories that has been widely used in public health for understanding health behavior.^[6] HBM can explain the change in health behavior to be used as a basis for educational interventions.^[7] HBM is a health-specific social cognition model whose key components and constructs are perceived susceptibility (PSU), perceived severity (PSE), perceived benefits (PBE), perceived barriers (PBA), and self-efficacy (SE).^[8] Behavioural obesity interventions are based on the assumptions that physical activity is the proximal mediator of body weight.^[9]

This type of education increases students' awareness of opportunities for increasing PA, expresses methods for removing barriers and negative attitudes about PA, and increases PA.

The results of this article showed that the application of PEP is useful to increase the scores of the components of HBM, perception of the first junior high school students about PA and work activity (WA), sports activity (SA), leisure activity (LA), and SLS to manage overweight and obesity. Hence, the present study hypothesizes that PEP can be effective in improving the HBM and PA and reducing the overweight and obesity of the students.

METHODS

Study design and participants

This study was a quasi-experimental, TBHE that took place in educational districts of the city of Urmia. After obtaining ethical clearance (ethical code number 393,847), the researcher entered the selected schools. Based on the sample size formula for *t*-tests, the probability of making Type I error was 0.05, the power of the hypothesis test was 0.8%, Cohen's *d* effect size was 0.7, and the minimum required sample size in each group was 32. A total of 80 junior high school male students whose age varied between 12 and 14 years old were studied.

Four high schools were randomly sampled from two districts in Urmia [Figure 1], and divided into two experimental high schools ($n = 40$ obese or overweight students) and two control high schools ($n = 40$ obese or overweight students) from the selected high schools participated in this study. The inclusion criteria were BMI ≥ 25 and/or no limitation on exercising. The students that were used medications for 6 months




Figure 1: Study design and participant flow in the intervention and control groups of students

known to affect body weight before the study and/or the participants who did not intend to continue the study were excluded from the study. The goal, its importance, and benefits of the study were explained to the participants. Students and their parents were told that participants' profile will be used anonymously, and hence personal information will be strictly confidential. The participants and their parents were informed of the confidentiality of the information and they consented to participate in the study.

Study instruments and variables assessment

A hybrid questionnaire was prepared by the researcher which includes variables such as father's education, mother's education, father's occupation, mother's occupation, family size, and rates of weight and height to determine BMI.

Overweight and obesity prevalence was determined by the International Obesity Task Force, Centers for Disease Control and Prevention and World Health Organization standard criteria. Weight and Height properties have been determined on the basis of age and gender. Overweight and obesity are defined as follows for children aged between 5–19 years: overweight is BMI-for-age greater than 1 standard deviation above the WHO Growth Reference median; and obesity is greater than 2 standard deviations above the WHO Growth Reference median.^[10]

The components of HBM construct and Baecke *et al.* questionnaire for measuring PA habits in epidemiological

studies, which in most studies of obesity and PA, have been used.^[11] The PA was divided into three distinct dimensions: WA, SA, and LA. Weight was taken with light clothing with the aid of a weighing Seca 750 scale with the weights measured to the nearest 1 kg, and height was measured using a stadiometer. Then, BMI was calculated from weight (in kg) divided by a square of the height (in meters). Information was collected by a three-part questionnaire. The first part includes demographic data, the second part was a HBM-based questionnaire that included PSU ($\alpha = 0.836$), PSE ($\alpha = 0.737$), PBE ($\alpha = 0.773$), and PBA ($\alpha = 0.885$), and the third part was the Baecke *et al.* questionnaire that included WA ($\alpha = 0.726$), SA ($\alpha = 0.675$), and LA ($\alpha = 0.881$). The questionnaire included 16 items, with proven validity which had been studied in previous studies.^[11] Interventional education in the intervention group conducted with lecture, participatory method, group discussion and role playing in six sessions, each session was 1 h.

HBM constructs were utilized to formulate the educational strategies. To assess the impact of educational

interventions to increase PA and reduce overweight of the participants, the data related to the participants' HBM and PA were collected by the same questionnaire at three times (before, 3 months, and 6 months as follow-up) after the TBHE.

Statistical analysis

The results are expressed as percentages for qualitative variables and mean and standard deviation (SD) for quantities variables in two groups. Chi-square test was used to compare demographical variables and HBM components in baseline, 3, and 6 months after the intervention between the two groups. The repeated-measures (RM) ANOVA was used to assess intervention and time effect on the components of PA constructs. Mauchly's test was used to investigate sphericity, and Kolmogorov-Smirnov test was used to investigate the normality of the distribution for each quantity variable. Furthermore, analysis of covariance (ANCOVA) was used to compare the mean of BMI reduced during 6 months after the intervention. The selected high school students' data were analyzed using the SPSS statistic software version 16, Urmia-Iran.

Table 1: Frequency of demographical characteristics of students at baseline in experimental and control groups

Variables	Answer	Experimental group, n (%)	Control group, n (%)	P (χ^2)
Father's literacy	High literate	23 (57.5)	19 (47.5)	0.370
	Low literate	17 (42.5)	21 (52.5)	
Mother's literacy	High literate	21 (52.5)	16 (40.0)	0.262
	Low literate	19 (47.5)	24 (60.0)	
Father' job	Employment	16 (40.0)	18 (45.0)	0.651
	Self-employed	24 (60.0)	22 (55.0)	
Mother' job	Housewife	31 (77.5)	33 (82.5)	0.576
	Employment	9 (22.5)	7 (17.5)	
Family size	3-4	34 (85.0)	32 (80.0)	0.556
	5+	6 (15.0)	8 (20.0)	
Father's obesity	Yes	21 (52.5)	23 (57.5)	0.653
	No	19 (47.5)	17 (42.5)	
Mother's obesity	Yes	26 (65.0)	31 (77.5)	0.217
	No	14 (35.0)	9 (22.5)	
Brother's obesity	Yes	33 (82.5)	35 (87.5)	0.531
	No	7 (17.5)	5 (12.5)	
Sister's obesity	Yes	39 (97.5)	36 (93.8)	0.166
	No	1 (2.5)	4 (10)	
Uncle's obesity	Yes	32 (80)	27 (67.5)	0.204
	No	8 (20)	13 (32.5)	
Aunt's obesity	Yes	31 (77.5)	35 (87.5)	0.239
	No	9 (22.5)	5 (17.5)	
Grandmother's obesity	Yes	25 (62.5)	30 (75.0)	0.228
	No	15 (37.5)	10 (25.0)	
Grandfather's obesity	Yes	29 (72.5)	33 (82.5)	0.284
	No	11 (27.5)	7 (17.5)	

RESULTS

In this study, eighty high school students from Urmia cooperated with the researchers.

Table 1 describes the frequency of demographical characteristics (parent's literacy, their job, family size, and obesity in family) of the students at baseline compared with two groups (experimental and control) of students by analyzing the data using Chi-square test.

Table 2 describes the application of the components of HBM about PA at baseline, 3, and 6 months after intervention by analyzing the data using Chi-square test.

Table 3 shows the effects of PEP on WA, SA, LA, and BMI in the experimental group in the baseline, 3, and 6 months after intervention by analyzing the data using RM ANOVA.

Table 4 demonstrates the BMI of students at 6 months after the intervention compared with baseline by analyzing the data using ANCOVA.

Table 1 shows the demographic characteristics of the two groups of students. There was no significant statistical difference between demographic variables at baseline.

Table 2 shows the effect of TBHE on the components of HBM constructs 3 and 6 months after the intervention in the experimental and control groups. Chi-square test shows that there are significant differences in many components of HBM 3 and 6 months after intervention in the experimental group in comparison with the baseline (i.e., after 3 months, PSU 2; $P < 0.0001$, PSE

Table 2: The components of health belief model in the groups (experimental and control) in the baseline, 3, and 6 months after the intervention

Variables	Answer	Baseline			3 months after intervention			6 months after intervention		
		Experimental ^f , n (%)	Control ^g , n (%)	Significant*	Experimental, n (%)	Control, n (%)	Significant	Experimental	Control	Significant
PSU^a										
1	Agree	10 (25)	5 (12.5)	0.232	5 (12.5)	5 (12.5)	0.251	5 (12.5)	6 (15)	0.217
	No idea	7 (17.5)	5 (12.5)		9 (23.1)	16 (40)		6 (15)	12 (30)	
	Disagree	23 (57.5)	30 (75)		25 (64.1)	19 (47.5)		29 (72.5)	22 (55)	
2	Agree	9 (22.5)	7 (17.5)	0.348	6 (15)	12 (30)	0.007	6 (15)	10 (25)	0.040
	No idea	9 (22.5)	5 (12.5)		8 (20)	16 (40)		5 (12.5)	12 (30)	
	Disagree	22 (55)	28 (70)		26 (65)	12 (30)		29 (72.5)	18 (45)	
3	Agree	9 (22.5)	14 (35)	0.466	6 (15)	17 (42.5)	0.0001	5 (12.5)	11 (27.5)	0.001
	No idea	6 (15)	5 (12.5)		7 (17.5)	16 (40)		7 (17.5)	19 (47.5)	
	Disagree	25 (62.5)	21 (52.5)		27 (67.5)	7 (17.5)		28 (70)	10 (25)	
4	Agree	13 (32.5)	14 (35)	0.945	10 (25)	13 (32.5)	0.003	6 (15)	12 (30)	0.004
	No idea	7 (17.5)	6 (15)		7 (17.5)	18 (45)		7 (17.5)	16 (40)	
	Disagree	20 (50)	20 (50)		23 (57.5)	9 (23.1)		27 (67.5)	12 (30)	
PSE^b										
1	Agree	16 (40)	13 (32.5)	0.753	5 (12.5)	16 (40)	0.0001	6 (15)	12 (30)	0.046
	No idea	8 (20)	8 (20)		11 (27.5)	17 (42.5)		8 (20)	13 (32.5)	
	Disagree	16 (40)	19 (47.5)		24 (60)	7 (17.5)		26 (65)	15 (37.5)	
2	Agree	10 (25)	7 (17.5)	0.686	8 (20)	14 (35)	0.014	6 (15)	18 (45)	0.006
	No idea	9 (22.5)	11 (27.5)		8 (20)	15 (37.5)		7 (17.5)	8 (20)	
	Disagree	21 (52.5)	22 (55)		24 (60)	11 (27.5)		27 (67.5)	14 (35)	
3	Agree	13 (32.5)	14 (35)	0.866	11 (27.5)	8 (20)	0.201	7 (17.5)	6 (15)	0.953
	No idea	10 (25)	8 (20)		7 (17.5)	14 (35)		8 (20)	8 (20)	
	Disagree	17 (42.5)	18 (45)		22 (55)	18 (45)		25 (62.5)	26 (65)	
PBE^c										
1	Agree	8 (20)	9 (22.5)	0.632	6 (15)	8 (20)	0.827	5 (12.5)	7 (17.5)	0.363
	No idea	15 (37.5)	11 (27.5)		12 (30)	12 (30)		6 (15)	10 (25)	
	Disagree	17 (42.5)	20 (50)		22 (55)	20 (50)		29 (72.5)	23 (57.5)	
2	Agree	7 (17.5)	8 (20)	0.657	6 (15)	8 (20)	0.028	5 (12.5)	7 (17.5)	0.164
	No idea	10 (25)	13 (32.5)		5 (12.5)	14 (35)		6 (15)	12 (60)	
	Disagree	23 (57.5)	19 (47.5)		29 (72.5)	18 (45)		29 (72.5)	21 (52.5)	
3	Agree	9 (22.5)	9 (22.5)	0.876	8 (20)	9 (22.5)	0.367	5 (12.5)	8 (20)	0.043
	No idea	12 (30)	14 (35)		10 (25)	15 (37.5)		7 (17.5)	15 (37.5)	
	Disagree	19 (47.5)	17 (42.5)		22 (55)	16 (40)		28 (70)	17 (42.5)	
PBA^d										
1	Agree	16 (40)	11 (27.5)	0.386	26 (65)	7 (17.5)	0.0001	27 (67.5)	18 (45)	0.040
	No idea	7 (17.5)	6 (15)		7 (17.5)	10 (25)		7 (17.5)	6 (15)	
	Disagree	17 (42.5)	23 (57.5)		7 (17.5)	23 (57.5)		6 (15)	16 (40)	
2	Agree	26 (65)	22 (55)	0.446	28 (70)	7 (17.5)	0.0001	25 (62.5)	7 (17.5)	0.001
	No idea	6 (15)	5 (12.5)		6 (15)	10 (25)		8 (20)	17 (42.5)	
	Disagree	8 (20)	13 (32.5)		6 (15)	23 (57.5)		7 (17.5)	16 (40)	
3	Agree	27 (67.5)	21 (52.5)	0.182	28 (70)	8 (20)	0.0001	28 (70)	8 (20)	0.001
	No idea	7 (17.5)	6 (15)		7 (17.5)	20 (50)		7 (17.5)	16 (40)	
	Disagree	6 (15)	13 (32.5)		5 (12.5)	12 (30)		5 (12.5)	16 (40)	
4	Agree	13 (32.5)	15 (37.5)	0.879	24 (60)	8 (20)	0.001	26 (65)	12 (30)	0.003
	No idea	6 (15)	5 (12.5)		10 (25)	16 (40)		7 (17.5)	8 (20)	
	Disagree	21 (25.5)	20 (50)		6 (15)	16 (40)		7 (17.5)	20 (50)	

Contd...

Table 2: Contd...

Variables	Answer	Baseline			3 months after intervention			6 months after intervention		
		Experimental ^f , n (%)	Control ^g , n (%)	Significant*	Experimental, n (%)	Control, n (%)	Significant	Experimental	Control	Significant
5	Agree	27 (67.5)	26 (65)	0.953	29 (72.5)	27 (67.5)	0.817	30 (75)	28 (70)	0.882
	No idea	6 (15)	7 (17.5)		6 (15)	6 (15)		5 (12.5)	6 (15)	
	Disagree	7 (17.5)	7 (17.5)		5 (12.5)	7 (17.5)		5 (12.5)	6 (15)	
SE ^e										
1	Agree	17 (42.5)	21 (52.5)	0.642	6 (15)	24 (60)	0.0001	5 (12.5)	10 (25)	0.0001
	No idea	7 (17.5)	5 (12.5)		12 (30)	10 (25)		11 (27.5)	25 (62.5)	
	Disagree	16 (40)	14 (35)		22 (55)	6 (15)		24 (60)	5 (12.5)	
2	Agree	6 (15)	13 (32.5)	0.043	6 (15)	15 (37.5)	0.0001	5 (12.5)	7 (17.5)	0.004
	No idea	5 (12.5)	9 (22.5)		7 (17.5)	20 (50)		5 (12.5)	17 (42.5)	
	Disagree	29 (72.5)	18 (45)		27 (67.5)	5 (12.5)		30 (75)	16 (40)	
3	Agree	5 (12.5)	6 (15)	0.913	5 (12.5)	6 (15)	0.480	5 (12.5)	6 (15)	0.042
	No idea	6 (15)	5 (12.5)		7 (17.5)	11 (27.5)		5 (12.5)	14 (35)	
	Disagree	29 (72.5)	29 (72.5)		28 (70)	23 (57.5)		30 (75)	20 (50)	
4	Agree	5 (12.5)	10 (66.7)	0.208	5 (12.5)	10 (25)	0.014	5 (12.5)	13 (32.5)	0.003
	No idea	11 (27.5)	6 (15)		7 (17.5)	15 (37.5)		6 (15)	13 (32.5)	
	Disagree	24 (60)	24 (60)		28 (70)	15 (37.5)		29 (72.5)	14 (35)	

*Significant, ^aPSU=Perceived susceptibility, ^bPSE=Perceived severity, ^cPBE=Perceived benefits, ^dPBA=Perceived barriers, ^eSE=Self-efficacy, ^fExperimental, ^gControl

1; $P < 0.0001$, PBE 2; $P = 0.028$, PBA; $P < 0.0001$, and SE 1, 2; $P < 0.0001$).

Table 3 shows the mean and SD in two groups (experimental and control) about the components of PA constructs (i.e., WA, SA, and LA) in baseline, 3, and 6 months after the intervention.

The Kolmogorov–Smirnov test prove the normality of the distribution for components of PA in each time ($P > 0.05$). The intervention and time effect were asses on components of PA constructs. The result of using R.M.ANOVA showed that the main effect of time, the interaction between the time with intervention is significant ($P = 0.023$), and there is a significant difference between the experiment and control groups in terms of the mean score of work Activity ($P < 0.001$). Also the result showed that the main effect of time ($P < 0.001$), and the interaction between the time with intervention is significant $P = 0.047$, and there is a significant difference between the experiment and control groups in terms of the mean score of Sport Activity ($P < 0.001$). Also the results showed that the main effect of time $P = 0.004$, and the interaction between the time with intervention is significant ($P = 0.092$), and there is a significant difference between the experiment and control groups in terms of the mean score of Leisure Activity ($P < 0.001$).

Table 4 shows the mean and SD of BMI at the baseline 6 months after intervention. The baseline BMI of participants in the study was 27.20 ± 3.04 for the experimental group and 27.66 ± 3.08 for the control group. The Kolmogorov–Smirnov test showed

Table 3: Comparison of work activity, sports activity, and leisure activity in the experimental and control groups in the baseline, 3, and 6 months after the intervention (follow-up)

PA*	Group	Mean ± SD		
		Baseline	3 months after intervention	6 months after intervention
WA ^c	Experimental ^a	3.21 ± 0.54	3.43 ± 0.50	3.47 ± 0.47
	Control ^b	2.70 ± 0.40	2.99 ± 0.34	3.19 ± 0.41
SA ^d	Experimental	3.06 ± 1.36	3.13 ± 0.98	4.75 ± 2.20
	Control	3.15 ± 1.59	3.28 ± 1.52	3.76 ± 2.44
LA ^e	Experimental	12.42 ± 3.44	10.85 ± 3.94	11.37 ± 3.36
	Control	11.07 ± 3.13	10.75 ± 3.45	10.85 ± 3.15

*Physical activity, ^aExperimental, ^bControl, ^cWork activity, ^dSports activity, ^eLeisure activity. SD=Standard deviation

Table 4: Mean and standard deviation of body mass index in the experiment and control groups at the baseline and 6 months after the intervention

Group	Mean ± SD	
	Baseline	6 months after the intervention
Experimental	27.70 ± 3.42	27.198 ± 3.04
Control	27.94 ± 3.08	27.65 ± 3.37

SD=Standard deviation

that the BMI had a normal distribution ($P > 0.05$). The result of ANCOVA showed that although BMI reduced during 6 months after the intervention, there was no significant difference in BMI 6 months

after the intervention, after baseline's BMI was adjusted $P = 0.57$.

CONCLUSIONS

Increasing the PA is a practical strategy for students to optimize their BMI and reduce their overweight and obesity. The most effective overweight and obesity reduction strategies for students at schools include prevention through theoretical health education and health promotion.^[12] This finding supports the hypothesis that education based on HBM can be effective in increasing PA and weight loss in participants. Some studies determined educational need to increase the health belief model structures, lead to promoting preventive behaviors about overweight and obesity.^[13]

The findings of the present study indicate that the components of HBM (PSU, PSE, PBE, PBA, and SE) scores of the participants to reduce their overweight and obesity by the intervention in the intervention group significantly increased PA while little change in the behavior of the control group was created. These findings are consistent with the results of Masumeh *et al*^[15] and Jennifer Robertson-Wilson *et al*^[16] and Robertson-Wilson *et al*.^[14] Due to the prevalence of overweight and its complications, increasing rates of HBM is important. Thus, activities that encourage students to increase their PA causes to reduce overweight. The results of this study are consistent with the results of the TBHE programs based on HBM interventions which are focused on the prevention and treatment of overweight and obesity that are appropriate to achieve the objectives of increasing rates of the HBM about PA. There was a significant difference between the mean scores in the intervention group compared with the control group about PA, and the interaction between the time trends is significant ($P < 0.001$). The increase in the performance of the participants in this study is also consistent with the findings of Jafari-Adli *et al.*,^[1] Tell *et al*^[17] Killen *et al*^[18] Franco *et al*^[4] Epstein *et al*^[19,20] and Ford *et al*.^[21]

Considering the positive effects of HBM and physical education program on increasing physical activity and reducing the body mass index of students, and also considering that today's students would be the adults of the future, and given the low cost of prevention of obesity and overweight in comparison with the high cost of treatment, it is essential that educational efforts at the country level based on HBM implemented for all student groups.

Financial support and sponsorship

This study was part of a PhD thesis supported by IUMS (grant No: 393847).

Conflicts of interest

There are no conflicts of interest.

Received: 05 Dec 15 Accepted: 09 Aug 16

Published: 10 Oct 16

REFERENCES

- Jafari-Adli S, Jouyandeh Z, Qorbani M, Soroush A, Larjani B, Hasani-Ranjbar S. Prevalence of obesity and overweight in adults and children in Iran: A systematic review. *J Diabetes Metab Disord* 2014;13:121.
- Heidari K, Minasian V, Marandi SM, Kelishadi R, Khalighinejad P, Davari H. Prevalence of obesity and overweight in 12-14-year-old students in Isfahan-Iran. *Int J Prev Med* 2014;5 Suppl 2:S120-5.
- Fallahi E. Vitamin D and obesity: Which one is affected by the other? *Nutr Food Sci Res* 2015;2:1-3.
- Franco SS, Wilson S, Devaux M, Cecchini M, Analyst HP. *OECD Obesity Update*. 2014.
- Yarhamadi SH, Etemad K, Hazaveh AM, Azhang N. Urbanization and non-communicable risk factors in the capital city of 6 big provinces of Iran. *Iran J Public Health* 2013;42 Suppl 1:113-8.
- Baghianimoghadam MH, Shogafard G, Sanati HR, Baghianimoghadam B, Mazloomi SS, Askarshahi M. Application of the health belief model in promotion of self-care in heart failure patients. *Acta Med Iran* 2013;51:52-8.
- Baghianimoghadam MH, Shogafard G, Sanati HR, Baghianimoghadam B, Mazloomi SS, Askarshahi M. Application of the health belief model in promotion of self-care in heart failure patients. *Acta Med Iran* 2013;51:53.
- Taylor D, Bury M, Campling N, Carter S, Garfield S, Newbould J, *et al*. A review of the use of the Health Belief Model (HBM), the Theory of Reasoned Action (TRA), the Theory of Planned Behaviour (TPB) and the Trans-Theoretical Model (TTM) to study and predict health related behaviour change. The School of Pharmacy, University of London, London WC1N 1AX: National Institute for Clinical Excellence Department of Health; 2007. p. 33.
- Kelly AS, Barlow SE, Rao G, Inge TH, Hayman LL, Steinberger J, *et al*. Obesity in children and adolescents: Identification, associated health risks, and treatment. *Am Heart Assoc* 2013;128:1697.
- Strong WB, Malina RM, Blimkie CJ, Daniels SR, Dishman RK, Gutin B, *et al*. Evidence based physical activity for school-age youth. *J Pediatr* 2005;146:732-7.
- Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982;36:936-42.
- Katz DL, O'Connell M, Yeh MC, Nawaz H, Njike V, Anderson LM, *et al*. Public Health Strategies for Preventing and Controlling Overweight and Obesity in School and Worksite Settings. A Report on Recommendations of the Task Force on Community Preventive Services. CDC; 2005.
- Ebadi Fard Azar F, Solhi M, Yekaninejad M, Pakpour A. The impact of health education through Health Belief Model (H.B.M) on mother's perceptions about obesity children obese male student in elementary schools. *J Health Adm* 2006;8:13.
- Robertson-Wilson J, Lévesque L, Richard L. Using an analytic framework to identify potential targets and strategies for ecologically based physical activity interventions in middle schools. *Health Promot Pract* 2009;10:232-43.
- Masumeh S, Yusef M, Javad N, Maryam M. Obesity Among Female Adolescents of Tehran Schools. *Journal of Premedical School, Tehran University of Medical Sciences, Peyavard Salamat* 1391;6:403-11.
- Robertson-Wilson J, Lévesque L, Richard L. Using an Analytic Framework to Identify Potential Targets and Strategies for Ecologically Based Physical Activity Interventions in Middle Schools. *Health Promotion Practice* 2009;10:240-1.
- Tell GS, Vellar OD. Physical fitness, physical activity, and cardiovascular disease risk factors in adolescents, the Oslo Youth Study. *Prev Med* 1988;17:12-24.
- Killen JD, Telch MJ, Robinson TN, Maccoby N, Taylor CB, Farquhar JW. Cardiovascular disease risk reduction for tenth graders. A multiple-factor school-based approach. *JAMA* 1988;260:1728-33.
- Epstein LH, Smith JA, Vara LS, Rodefer JS. Behavioral economic analysis of activity choice in obese children. *Health Psychol* 1991;10:311-6.
- Epstein LH, Saelens BE, Myers MD, Vito D. Effects of decreasing sedentary behaviors on activity choice in obese children. *Health Psychol* 1997;16:107-13.
- Ford BS, McDonald TE, Owens AS, Robinson TN. Primary care interventions to reduce television viewing in African-American children. *Am J Prev Med* 2002;22:106-9.