

Non-diluted seawater enhances nasal ciliary beat frequency and wound repair speed compared to diluted seawater and normal saline

Arnaud Bonnomet, PhD^{1,2}, Emilie Luczka, PhD¹, Christelle Coraux, PhD¹ and Ludovic de Gabory, MD, PhD³

Background: The regulation of mucociliary clearance is a key part of the defense mechanisms developed by the airway epithelium. If a high aggregate quality of evidence shows the clinical effectiveness of nasal irrigation, there is a lack of studies showing the intrinsic role of the different irrigation solutions allowing such results. This study investigated the impact of solutions with different pH and ionic compositions, eg, normal saline, non-diluted seawater and diluted seawater, on nasal mucosa functional parameters.

Methods: For this randomized, controlled, blinded, *in vitro* study, we used airway epithelial cells obtained from 13 nasal polyps explants to measure ciliary beat frequency (CBF) and epithelial wound repair speed (WRS) in response to 3 isotonic nasal irrigation solutions: (1) normal saline 0.9%; (2) non-diluted seawater (Physiomer®); and (3) 30% diluted seawater (Stérimar). The results were compared to control (cell culture medium).

Results: Non-diluted seawater enhanced the CBF and the WRS when compared to diluted seawater and to normal saline. When compared to the control, it significantly enhanced CBF and slightly, though nonsignificantly, improved the WRS. Interestingly, normal saline markedly reduced the number of epithelial cells and ciliated cells when compared to the control condition.

Conclusion: Our results suggest that the physicochemical features of the nasal wash solution is important because it determines the optimal conditions to enhance CBF and epithelial WRS thus preserving the respiratory mucosa in pathological conditions. Non-diluted seawater obtains the best results on CBF and WRS vs normal saline showing a deleterious effect on epithelial cell function. © 2016 The Authors International Forum of Allergy & Rhinology, published by ARSAAOA, LLC.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Key Words:

seawater; saline; ciliary beat frequency; wound repair; nasal mucosa

How to Cite this Article:

Bonnomet A, Luczka E, Coraux C, de Gabory L. Non-diluted seawater enhances nasal ciliary beat frequency and wound repair speed compared to diluted seawater and normal saline. *Int Forum Allergy Rhinol.* 2016;6:1062-1068.

¹National Institute of Health and Medical Research (Institut National de la Santé et de la Recherche Médicale [INSERM]) Scientific Mixed Research unit UMRS-S 903, Federative Structure Health Research Champagne-Ardenne-Picardie (SFR CAP-Santé): FED 4231, University of Reims-Champagne-Ardenne, Reims, France; ²Cellular and Tissular Imaging Platform (PICT), University of Reims-Champagne-Ardenne, Reims, France; ³Ear Nose and Throat (ENT) Department, University Hospital of Bordeaux, Hôpital Pellegrin, Bordeaux, France

Correspondence to: Ludovic de Gabory, MD, PhD, Department of Otorhinolaryngology, University Hospital of Bordeaux, Hôpital Pellegrin, Centre F-X Michelet, Place A. Raba-Léon, F-33076, Bordeaux Cedex, France; e-mail: ludovic.de-gabory@chu-bordeaux.fr

Additional Supporting Information may be found in the online version of this article.

Funding sources for the study: Laboratoire de la Mer to INSERM UMRS-S 903.

Nasal mucosa plays a particularly important protective role. The mucociliary clearance mechanism acts as a highly effective, nonspecific waste disposal system that is sometimes insufficient to prevent allergic response or microbial infection to airborne allergens, pollutants or pathogens.

In vitro and *in vivo* studies have revealed the air pollutants attenuating properties on ciliary beat frequency (CBF).¹⁻³ Other studies have shown an impaired CBF in patients with allergic rhinitis or asthma.⁴⁻⁷ Certain topical antibiotics have been shown to reduce the CBF

Potential conflict of interest: None provided.

Received: 23 November 2015; Revised: 17 February 2016; Accepted: 8 March 2016

DOI: 10.1002/alr.21782

View this article online at wileyonlinelibrary.com.

and thus cannot be recommended in treatment of local infections.⁸ Pathogens targeting the airway like *Pseudomonas aeruginosa*, *Streptococcus pneumoniae*, or *Haemophilus influenzae* produce various cilio-inhibitory factors.^{9–12} This is clearly illustrated in cystic fibrosis or primary ciliary dyskinesia in which the impaired ciliary beat and/or thicker mucus result in a defective mucociliary clearance that may participate in repeated lung infections and pathophysiology.^{13–16} It is thus important to restore an efficient ciliary beat to ensure the protective role of the respiratory mucosa.

Furthermore, contact prolongation of airborne allergens, pollutants, and pathogens with the nasal mucosa worsen the inflammatory reaction leading to epithelial lesions that threaten mucosal integrity. Mucosal surface wounds become entry points for bacteria and viruses that target airway epithelial cells and may result in respiratory infections. In order to restore its functionality, the airway epithelium enters in turn in a process of repair and regeneration.¹⁷ In cases of inflammatory respiratory diseases such as asthma,¹⁷ it is becoming apparent that normal airway epithelium repair is compromised. Ensuring the proper conditions of wound repair process represents therefore a great therapeutic interest for any respiratory conditions associated with mucosal inflammation.

Nasal irrigation with saline solutions is commonly used as adjunctive treatment in upper respiratory conditions as well as in postsurgical follow-up.¹⁸ A 2012 systematic review and meta-analysis by Hermelingmeier et al.¹⁹ showed the benefit of nasal irrigation in symptom relief and drug reduction in cases of allergic rhinitis. It has also been recommended as add-on therapy in pediatric allergic rhinitis by European experts.²⁰ In addition, nasal irrigation was helpful in improving the nasal peak expiratory flow rates and quality of life scores in children with atopic²¹ and nonatopic²² acute sinusitis. A Cochrane review by Kassel et al.²³ of data in adults with acute upper respiratory tract infections (URTIs) revealed benefit in quicker symptom resolution and return to work, though the results were not statistically significant. Recently, several international expert groups recommended nasal saline irrigation in chronic rhinosinusitis with or without nasal polyps,^{24–26} and after endonasal surgery.²⁴

If the efficacy of nasal irrigation is no longer questioned *in vivo*, the literature on nasal irrigation composition specifically, remains limited and does not provide any evidence on its potential impact. The aim of this study is to investigate and compare the functional impact of 3 commonly used isotonic nasal irrigation solutions, reflecting the diversity of products available on the market: (1) normal saline (0.9%); (2) isotonic non-diluted seawater solution; and (3) isotonic 30% diluted seawater solution. We hypothesize that the non-diluted seawater solution is more effective on the CBF and wound repair speed (WRS) vs other solutions.

Patients and methods

Study design

Prospective, randomized, controlled, and blinded *in vitro* study using biological materials from nasal polyp explants.

Material

The bioethical law N° 2004–800 of the French Public Health Code authorizes the use of human tissues. Local Institutional Review Board approved this study and informed consent was obtained from all patients. Three centers provided nasal polyps explants: University Hospital Hautepierre (Strasbourg, France); University Hospital Robert Debré (Paris, France); and private clinic Courlancy (Reims, France). Nasal polyps were removed from patients diagnosed with sinonasal polyposis without any other comorbidities (asthma or nonsteroidal anti-inflammatory drugs allergies). Age, sex, and clinical characteristics are summarized in Table 1. Some of the patients received corticoids and/or antibiotics. Treatments are described in Table 1.

Randomization

Biological materials were randomly assigned (using random letters A, B, C) into either the normal saline group, the non-diluted seawater group, and the 30% diluted seawater group.

Blinding

Blinding of tested solutions except control medium was strictly maintained for researchers including laboratory staff. The composition of the tested solutions was revealed by the sponsor after study completion. Study solutions were provided and prepackaged by the sponsor's head pharmacist (located at a distant site) into anonymous kits and lettered accordingly.

Tested solutions

Non-diluted seawater (Physiomer[®], Laboratoire de la Mer, Saint-Malo, France) is a sterile, isotonic 100% non-diluted seawater solution. Isotonicity is achieved by selective electro dialysis that removes NaCl ions while preserving seawater full content in other minerals. Physiomer[®] is sterilized aseptically through microfiltration at 0.2 μm .

Diluted seawater (Stérimar[®], Laboratoires Fumouze, Levallois-Perret, France) is a sterile, isotonic, seawater solution diluted at 30%. It is rendered isotonic through dilution with purified water. Its minerals content is reduced by two-thirds vs seawater. Stérimar[®] is sterilized by gamma irradiation.

Normal saline 0.9% (CDM Lavoisier, Paris, France) is a sterile, ready-for-injection solution. It contains 3500 mg/L of sodium ions and 5500 mg/L of chloride ions. Normal saline conforms to pharmaceutical quality standards.

Table 2 summarizes the different test solutions and their physicochemical characteristics.

TABLE 1. Demographic and clinical data of patients

Patient number	Age (years)	Sex	Allergy history	Intolerance history	Treatment
01	75	M	None	None	None
02	34	M	None	None	None
03	69	F	None	None	None
04	44	F	None	None	Beclomethasone
05	53	M	Latex	None	Beclomethasone
06	52	M	None	None	Pristinamycin
07	77	M	None	None	None
08	63	M	None	None	Prednisolone; pristinamycin
09	36	M	None	None	Beclomethasone
10	67	F	None	None	None
11	56	M	None	None	Beclomethasone
12	50	M	None	None	None
13	39	M	None	None	None
14	51	M	None	None	Mometasone furoate

TABLE 2. pH and osmolarity of solutions

Tested solution	Batch number	Measured pH	Measured osmolarity (mOsm/kg)
Physiomer®	P1201024A	7.9	308
Stérimar®	FE2234	7.28	302
Lavoisier normal saline (0.9%)	2F258; 2F220	5.21	308

Cell cultures

For the measurement of CBF, nasal epithelial tissue from nasal polyps from 10 patients (1 to 2 mm² in size) were seeded on 12-well culture plates (BD Falcon, Franklin Lakes, NJ, USA) coated with type IV collagen (Sigma Aldrich, St. Louis, MO, USA) and incubated in Bronchial Epithelial cell Growth Medium™ (BEGM; Lonza, Verviers, Belgium) at 37°C for 4 to 6 days until explants were surrounded by a cell outgrowth that contained well-visible ciliated cells.

BEGM™ was used for cell cultures as control. Saline solution is usually considered as a reference product for nasal irrigation; however, it is not used and dedicated for cell culture. BEGM™ contains equal proportions of Bronchial Epithelial Basal Medium (Lonza) and Dulbecco's Modified Eagle Medium (Lonza). The medium is supplemented with 0.1 mM of retinoic acid, 0.5 mg/L of human epidermal growth factor, 5 mg/L of epinephrine, 0.13 g/L of bovine pituitary extract, 0.5 mg/L of hydrocortisone, 5 mg/L of insulin, 6.5 mg/L of triiodothyronine, 0.5 mg/L of transferrin (all from Lonza, Verviers, Belgium),

200 U/mL of penicillin, 200 mg/L of streptomycin, and 1.5 mg/L of bovine serum albumin (all from Sigma-Aldrich, Lyon, France).

For the WRS assay, nasal epithelial cells were isolated from nasal polyps from 13 patients by incubation with 0.1% type XIV collagenase (Sigma-Aldrich) in Roswell Park Memorial Institute (RPMI) 1640 culture medium medium (Life Technologies, Saint Aubin, France) supplemented with 200 U/mL penicillin and 200 µg/mL streptomycin (Life Technologies) overnight at 4°C.

Isolated epithelial cells were washed, suspended in BEGM medium, counted, and then seeded on 12-well culture plates coated with type IV collagen at a density of 6×10^4 cells/cm² and cultured in BEGM medium at 37°C until confluence was reached.

CBF measurement

After a culture period of 4 to 6 days, once explants were surrounded by a cell outgrowth and contained well-visible ciliated cells, wells were rinsed with either BEGM medium,

TABLE 3. Ciliary beat frequency for each explants and solutions

Explant number	Control (Hz)	Physiomer® (Hz)	Stérimar® (Hz)	0.9% NaCl (Hz)
1	13.69	15.31	14.16	13.14
2	12.98	13.21	12.90	11.32
3	12.49	15.44	13.37	nd
4	14.09	15.04	12.61	nd
5	13.07	13.58	13.54	nd
6	10.74	12.11	11.36	nd
7	11.79	13.12	11.75	nd
8	14.78	15.92	13.51	9.59
9	12.28	12.12	12.37	12.67
10	12.10	12.47	10.50	9.08
Mean	12.80	13.83**	12.61	11.16**
SD	1.18	1.46	1.13	1.80

***p* < 0.01, comparison of Physiomer® and 0.9% NaCl vs CTRL. CTRL = control; nd = not determined (death cell); SD = standard deviation.




FIGURE 1. CBF: mean and distribution of the values of CBF for the control (•), Physiomer® (■), Stérimar® (▲), and normal saline 0.9% (▼) conditions. CBF is expressed in hertz (Hz). ***p* = 0.0039, ***p* = 0.0059, CBF = ciliary beat frequency; CTRL = control; ns = nonsignificant.

non-diluted seawater, diluted seawater, or normal saline. Plates were then incubated in the corresponding solutions for 30 minutes, under an inverted microscope (Axiovert 200; Zeiss, Germany) equipped with an environmental chamber maintained at 37°C and 5% CO₂ and a CoolSNAP CCD camera (Roper Scientific, Inc., Tucson, AZ, USA) allowing live-cell imaging. For each condition, 20 areas with ciliated cells were recorded over 10 seconds with an acquisition frequency of 50 images/second (×32 magni-

fication). CBF was quantified using an in-house-developed plug-in on the ImageJ software (NIH, Bethesda, MD; <http://imagej.nih.gov/ij/index.html>).²⁷ The control condition was achieved by incubating the cells with BEGM medium. Values were expressed as mean frequency in hertz (Hz) for each condition.

WRS measurement

At confluence, cultures were rinsed and incubated with BEGM medium, non-diluted seawater, diluted seawater, or normal saline. The control condition was achieved by incubating cells with BEGM medium. After a 4-hour incubation period, an epithelial wound was realized by scratching each epithelial cell monolayer in a linear pattern with a pipette tip (100 μL).

Wells were then rinsed with BEGM medium to eliminate debris and cells were incubated with fresh culture medium. Three areas per condition and culture were controlled by live-cell imaging (×10 magnification) until wound closure. Phase contrast images were captured every 10 minutes and then analyzed with an in-house-developed plug-in on ImageJ software. WRS was expressed as mean speed for each condition and culture in μm²/hour.

Statistical analysis

Statistical analysis was carried out independently by the research unit INSERM UMRS-S 903 using the nonparametric Wilcoxon test in Prism 5©. CBF and WRS were expressed as median ± standard errors of the mean. Statistical significance was defined as *p* < 0.05.

TABLE 4. Wound repair speed for each explant and solution

Explant number	Control ($\mu\text{m}^2/\text{hour}$)	Physiomer [®] ($\mu\text{m}^2/\text{hour}$)	Stérimar [®] ($\mu\text{m}^2/\text{hour}$)	0.9% NaCl ($\mu\text{m}^2/\text{hour}$)
1	42068.83	76953.81	37498.06	-3707.19
2	53085.60	72525.33	52519.79	125.43
3	108927.93	90072.95	100699.66	8232.33
4	59098.62	62144.68	48757.18	2013.14
5	35406.18	20563.92	18061.37	3243.46
6	61025.22	70991.49	30805.66	1283.73
7	84179.58	63026.18	18131.67	9497.80
8	40599.71	62023.00	23785.74	15311.40
9	106692.41	114806.43	39197.86	-6140.88
10	59580.00	75823.75	45010.63	1961.25
11	105500.62	116436.27	97488.66	-3296.17
12	94326.33	79993.78	85877.69	3243.46
13	57236.36	66952.39	42256.76	33702.66
Mean	69825.18	74793.38	49237.74 ^{***}	5036.18 ^{***}
SD	26634.36	24298.26	28228.66	10374.19

^{***} $p < 0.001$, comparison of Stérimar[®] and 0.9% NaCl vs CTRL. CTRL = control; SD = standard deviation.

Results

CBF in response to non-diluted seawater (Physiomer[®]), diluted seawater (Stérimar[®]), and normal saline

After a 4-day to 6-day culture period, explant outgrowths show visible and functional ciliated cells. Supporting Video 1 presents CBF in the 3 different nasal irrigation solutions compared to control. Mean values and standard deviations for each culture and condition are presented in Table 3. The CBF mean was 12.80 ± 1.18 Hz for the control condition, 13.83 ± 1.46 Hz for cells exposed to non-diluted seawater, 12.61 ± 1.13 Hz for cells exposed to diluted seawater, and 11.16 ± 1.80 Hz for cells exposed to normal saline. Figure 1 shows the means and the distribution of the measured CBF values for each condition. Incubation of ciliated cells with non-diluted seawater significantly increased the CBF compared to the control condition ($p = 0.0039$). Diluted seawater, on the other hand, did not modify the CBF compared to the control ($p > 0.05$). Surprisingly, incubation with normal saline induced greater ciliated cell death compared with control, non-diluted seawater, and diluted seawater ($p = 0.0039$). Moreover, after 30 minutes, 5 of the 10 nasal explant cultures incubated with normal saline exhibited no ciliated cells and CBF could not be determined (Table 3). Finally, ciliated cells incubated with non-diluted seawater exhibited a significantly better CBF compared to cells incubated with diluted seawater ($p = 0.0059$).

WRS in response to non-diluted seawater (Physiomer[®]), diluted seawater (Stérimar[®]), and normal saline

Our WRS model exposed to different conditions is shown in Supporting Video 2. Mean values and standard deviations for each culture and conditions are presented in Table 4. The mean WRS was 69825.18 ± 26634.36 $\mu\text{m}^2/\text{hour}$ for the control condition, 74793.38 ± 24298.26 $\mu\text{m}^2/\text{hour}$ for cells exposed to non-diluted seawater, 49237.74 ± 28228.66 $\mu\text{m}^2/\text{hour}$ for cells exposed to diluted seawater, and 5036.18 ± 10374.19 $\mu\text{m}^2/\text{hour}$ for cells exposed to normal saline. Figure 2 shows the WRS means and the distribution of the measured values for each condition.

Incubation of epithelial cells with non-diluted seawater slightly raised the WRS compared to the control condition, but the results did not reach significance. Diluted seawater on the other hand, reduced the WRS compared to the control ($p = 0.0002$). Nasal epithelial cells exposed to non-diluted seawater exhibited a significantly better WRS compared to cells exposed to diluted seawater ($p = 0.0024$) and normal saline ($p = 0.0002$). Interestingly, incubation with normal saline greatly reduced the WRS, showing a deleterious effect on epithelial cell function ($p = 0.0002$). In fact, the reduced WRS was exacerbated by a high cell death rate in normal saline (data not shown).




FIGURE 2. WRS: mean and distribution of the values of WRS for the control (•), Physiomer® (■), Stérimar® (▲), and normal saline 0.9% (▼) conditions. WRS is expressed in μm^2 per hour. $**p = 0.0024$, $***p = 0.0002$. CTRL = control; WRS = wound repair speed.

Discussion

Our *in vitro* results showed that non-diluted seawater markedly enhanced the CBF and slightly but nonsignificantly raised the WRS when compared to the control condition. Both enhancements were significant when compared to normal saline and diluted seawater. Moreover, our results confirmed previous findings^{28,29} demonstrating that normal saline is potentially deleterious for nasal mucosa as it induced nasal epithelial cell death *in vitro*. A recent review of the literature showed the effects of selected ions on epithelial cells such as magnesium on the control of local inflammation resulting from allergy, calcium in the regulation of the ciliary beat frequency, and the implication of potassium on healing.³⁰

Many models have been described for the *in vitro* evaluation of the CBF and other features of the nasal mucosa. In addition to cell lines, primary cells have been extensively used. They are isolated from different regions of the nasal mucosa, grown as explant outgrowth cultures or dissociated tissue cultures, with coated or uncoated supports, in a perfusion system or CO_2 incubator. The various methods and time of CBF recording result in a stable, coordinated CBF ranging from 7 to 13 Hz according to the literature,^{31–34} 12 to 13 Hz being the CBF recorded *in vivo*.^{35,36} In the present study, the mean baseline CBF (12.80 Hz) from the nasal polyps explants outgrowth is in accordance with previous published reports using similar methods *in vitro* and *in vivo*.^{34,37,38} As for the wound repair capacity of the nasal mucosa, *in vitro* studies using a similar model are lacking.

The difficulty of this kind of investigation lies in the high interpatient variability, which may explain why the difference between non-diluted seawater and the control medium

did not reach statistical significance. Our results confirm that non-diluted seawater unlike normal saline and other diluted seawater solutions, does not inhibit the physiological process of wound repair.

Previous studies^{28,29} demonstrated *in vitro* that the viability of bronchial epithelial cells (BECs) incubated in normal saline decreases by 40% and 20% after 2-hour and 4-hour incubation periods, respectively. In contrast, incubation in non-diluted seawater (Physiomer®) maintains a healthy BEC culture with adherent cells and intercellular junctions. Protein contents are also higher in BEC cultures with non-diluted seawater (Physiomer®) compared to normal saline, or compared to diluted seawater (Stérimar®). Our results show that, *in vitro*, non-diluted seawater gives better results than normal saline and diluted seawater solutions on CBF and WRS. This suggests that *in vivo* superiority of mineral-rich solutions could rely on such an *in vitro* effect on CBF and WRS.

Nasal irrigation with normal saline has been recommended and performed for many years. Although there has not been reported any deleterious effect on the nasal mucosa like in our model, it shows the impact of nasal solutions specifically in terms of composition. This corroborates recent findings showing that nasal irrigation was proven more effective when performed with seawater-derived or mineral-rich solutions compared to a normal saline 0.9% solution in allergic rhinitis,³⁹ chronic rhinosinusitis,^{40,41} and post-endonasal surgery.^{42–44} Recently, Low et al.⁴⁴ showed *in vivo* moderate, though statistically better and faster symptom resolution with nasal irrigation with lactated Ringer's solution after endoscopic sinus surgery when compared with normal saline. Lactated Ringer's solution is an isotonic solution, composed of a sodium lactate as buffer (2500 mg/L), sodium (3000 mg/L), chloride (3900 mg/L), calcium (120 mg/L), and potassium (150 mg/L).

The effect of non-diluted seawater can be explained by the preservation of seawater mineral composition. Indeed, calcium ions are well known for their implication in ciliary beat regulation.^{45–47} Potassium, magnesium, and zinc ions have been shown to assist in epithelial wound repair.^{48–50} Diluted seawater solutions contain far lower mineral content due to dilution. Moreover non-diluted seawater alkaline pH (7.9) is more favorable to the ciliary beat, as previously shown *in vitro*,⁵¹ whereas normal saline acidic pH (5.21) is deleterious for the *in vitro* ciliary beat.⁵¹

Conclusion

Our results suggest that the physicochemical features of the nasal wash solution is important because it determines the optimal conditions to enhance CBF and epithelial WRS, thus preserving the respiratory mucosa in pathological conditions. Further *in vivo* studies will be needed to confirm the superiority of non-diluted seawater vs normal saline and diluted seawater in pathological conditions such as chronic rhinosinusitis and/or allergic rhinitis. 🌊

Acknowledgments

We thank Pr Christian Debry, Dr Jean-Claude Mérol and Dr. Talal Nasser for the delivery of polyeps, as well

as Pr. Philippe Birembaut for his general support in the implementation of the study as former head of INSERM UMRS-S 903.

References

- Bayram H, Devalia JL, Sapsford RJ, et al. The effect of diesel exhaust particles on cell function and release of inflammatory mediators from human bronchial epithelial cells in vitro. *Am J Respir Cell Mol Biol*. 1998;18:441–448.
- Calderon-Garciduenas L, Valencia-Salazar G, Rodriguez-Alcaraz A, et al. Ultrastructural nasal pathology in children chronically and sequentially exposed to air pollutants. *Am J Respir Cell Mol Biol*. 2001;24:132–138.
- Sakai N, Tamaoki J, Chiyotani A, et al. [Inhibitory effect on sulfur dioxide on ciliary motility in rabbit tracheal epithelium and its prevention by intracellular cyclic AMP]. *Nihon Kyobu Shikkan Gakkai Zasshi*. 1993;31:733–737. Japanese
- Holmström M, Lund V, Scadding G. Nasal ciliary beat frequency after nasal allergen challenge. *Am J Rhinol*. 1992;6:101–105.
- Maurizi M, Paludetti G, Todisco T, et al. Ciliary ultrastructure and nasal mucociliary clearance in chronic and allergic rhinitis. *Rhinology*. 1984;22:233–240.
- Mezey RJ, Cohn MA, Fernandez RJ, et al. Mucociliary transport in allergic patients with antigen-induced bronchospasm. *Am Rev Respir Dis*. 1978;118:677–684.
- Ohashi Y, Nakai Y, Kihara S, et al. Ciliary activity in patients with nasal allergies. *Arch Otorhinolaryngol*. 1985;242:141–147.
- Gosepath J, Grebneva N, Mossikhin S, Mann WJ. Topical antibiotic, antifungal, and antiseptic solutions decrease ciliary activity in nasal respiratory cells. *Am J Rhinol*. 2002;16:25–31.
- Janson H, Carlén B, Cervin A, et al. Effects on the ciliated epithelium of protein D-producing and non-producing nontypeable *Haemophilus influenzae* in nasopharyngeal tissue cultures. *J Infect Dis*. 1999;180:737–746.
- Read RC, Roberts P, Munro N, et al. Effect of *Pseudomonas aeruginosa* rhamnolipids on mucociliary transport and ciliary beating. *J Appl Physiol*. 1992;72:2271–2277.
- Read RC, Rutman AA, Jeffery PK, et al. Interaction of capsulate *Haemophilus influenzae* with human airway mucosa in vitro. *Infect Immun*. 1992;60:3244–3252.
- Steinfurt C, Wilson R, Mitchell T, et al. Effect of *Streptococcus pneumoniae* on human respiratory epithelium in vitro. *Infect Immun*. 1989;57:2006–2013.
- Sagel SD, Davis SD, Campisi P, Dell SD. Update of respiratory tract disease in children with primary ciliary dyskinesia. *Proc Am Thorac Soc*. 2011;8:438–443.
- Worlitzsch D, Tarran R, Ulrich M, et al. Effects of reduced mucus oxygen concentration in airway *Pseudomonas* infections of cystic fibrosis patients. *J Clin Invest*. 2002;109:317–325.
- Knowles MR, Boucher RC. Mucus clearance as a primary innate defense mechanism for mammalian airways. *J Clin Invest*. 2002;109:571–577.
- Livraghi A, Randell SH. Cystic fibrosis and other respiratory diseases of impaired mucus clearance. *Toxicol Pathol*. 2007;35:116–129.
- Grainge CL, Davies DE. Epithelial injury and repair in airways diseases. *Chest*. 2013;144:1906–1912.
- Tomooka LT, Murphy C, Davidson TM. Clinical study and literature review of nasal irrigation. *Laryngoscope*. 2000;110:1189–1193.
- Hermelingmeier KE, Weber RK, Hellmich M, et al. Nasal irrigation as an adjunctive treatment in allergic rhinitis: a systematic review and meta-analysis. *Am J Rhinol Allergy*. 2012;26:e119–e125.
- Roberts G, Xatzipsalti M, Borrego LM, et al. Paediatric rhinitis: position paper of the European Academy of Allergy and Clinical Immunology. *Allergy*. 2013;68:1102–1116.
- Wang YH, Ku MS, Sun HL, Lue KH. Efficacy of nasal irrigation in the treatment of acute sinusitis in atopic children. *J Microbiol Immunol Infect*. 2014;47:63–69.
- Wang YH, Yang CP, Ku MS, et al. Efficacy of nasal irrigation in the treatment of acute sinusitis in children. *Int J Pediatr Otorhinolaryngol*. 2009;73:1696–1701.
- Kassel JC, King D, Spurling GK. Saline nasal irrigation for acute upper respiratory tract infections. *Cochrane Database Syst Rev*. 2010;(3):CD006821.
- Fokkens WJ, Lund VJ, Mullol J, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2012. *Rhinol Suppl*. 2012;(23):3 p preceding table of contents, 1–298.
- Rosenfeld RM, Andes D, Bhattacharyya N, et al. Clinical practice guideline: adult sinusitis. *Otolaryngol Head Neck Surg*. 2007;137(3 Suppl):S1–S31.
- Wei CC, Adappa ND, Cohen NA. Use of topical nasal therapies in the management of chronic rhinosinusitis. *Laryngoscope*. 2013;123:2347–2359.
- Zahm JM, Dionisius JP, Pierrot D, et al. [Measuring the frequency of ciliary beating on video screen.] Mesure de la fréquence de battement ciliaire sur écran vidéo. *Innov Tech Biol Med*. 1990;2:118–129.
- Tabary O, Muselet C, Miesch MC, et al. Reduction of chemokine IL-8 and RANTES expression in human bronchial epithelial cells by a sea-water derived saline through inhibited nuclear factor-kappaB activation. *Biochem Biophys Res Commun*. 2003;309:310–316.
- Traissac L, Bordenave L, Bareille R, et al. In vitro study of the effect of sea water by-products on the respiratory mucosa. *Rev Soc Fr ORL*. 1995;32:43–49.
- Bastier PL, Lehot A, Bordenave L, et al. Nasal irrigation: From empiricism to evidence-based medicine. A review. *Eur Ann Otorhinolaryngol Head Neck Dis*. 2015;132:281–285.
- Birk R, Aderhold C, Stern-Strater J, et al. Polyhexanide-containing solution reduces ciliary beat frequency of human nasal epithelial cells in vitro. *Eur Arch Otorhinolaryngol*. 2015;272:377–383.
- Carson JL, Lu TS, Brighton L, et al. Phenotypic and physiologic variability in nasal epithelium cultured from smokers and non-smokers exposed to second-hand tobacco smoke. *In Vitro Cell Dev Biol Anim*. 2010;46:606–612.
- Jiao J, Meng N, Wang H, Zhang L. Comparison of human nasal epithelial cells grown as explant outgrowth cultures or dissociated tissue cultures in vitro. *Front Med*. 2013;7:486–491.
- Shaari J, Palmer JN, Chiu AG, et al. Regional analysis of sinonasal ciliary beat frequency. *Am J Rhinol*. 2006;20:150–154.
- Rutland J, Cole PJ. Nasal mucociliary clearance and ciliary beat frequency in cystic fibrosis compared with sinusitis and bronchiectasis. *Thorax*. 1981;36:654–658.
- Stanley PJ, Wilson R, Greenstone MA, et al. Effect of cigarette smoking on nasal mucociliary clearance and ciliary beat frequency. *Thorax*. 1986;41:519–523.
- Chevillard M, Hinrasky J, Zahm JM, et al. Proliferation, differentiation and ciliary beating of human respiratory ciliated cells in primary culture. *Cell Tissue Res*. 1991;264:49–55.
- Zahm JM, Pierrot D, Hinrasky J, et al. Functional activity of ciliated outgrowths from cultured human nasal and tracheal epithelia. *Biorheology*. 1990;27:559–565.
- Cordray S, Harjo JB, Miner L. Comparison of intranasal hypertonic dead sea saline spray and intranasal aqueous triamcinolone spray in seasonal allergic rhinitis. *Ear Nose Throat J*. 2005;84:426–430.
- Friedman M, Hamilton C, Samuelson CG, et al. Dead Sea salt irrigations vs saline irrigations with nasal steroids for symptomatic treatment of chronic rhinosinusitis: a randomized, prospective double-blind study. *Int Forum Allergy Rhinol*. 2012;2:252–257.
- Friedman M, Vidyasagar R, Joseph N. A randomized, prospective, double-blind study on the efficacy of dead sea salt nasal irrigations. *Laryngoscope*. 2006;116:878–882.
- Keerl R, Weber R, Muller C, Schick B. [Effectiveness and tolerance of nasal irrigation following paranasal sinus surgery]. *Laryngorhinootologie*. 1997;76:137–141.
- Michel O, Charon J. [Postoperative inhalation treatment after paranasal sinus interventions. A placebo-controlled, double-blind and randomized study]. *HNO*. 1991;39:433–438.
- Low TH, Woods CM, Ullah S, Carney AS. A double-blind randomized controlled trial of normal saline, lactated Ringer's, and hypertonic saline nasal irrigation solution after endoscopic sinus surgery. *Am J Rhinol Allergy*. 2014;28:225–231.
- Di Benedetto G, Magnus CJ, Gray PT, Mehta A. Calcium regulation of ciliary beat frequency in human respiratory epithelium in vitro. *J Physiol*. 1991;439:103–113.
- Schmid A, Salathe M. Ciliary beat co-ordination by calcium. *Biol Cell*. 2011;103:159–169.
- Zsembery A, Boyce AT, Liang L, et al. Sustained calcium entry through P2X nucleotide receptor channels in human airway epithelial cells. *J Biol Chem*. 2003;278:13398–13408.
- Bardou O, Trinh NT, Brochiero E. Molecular diversity and function of K⁺ channels in airway and alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol*. 2009;296:L145–L155.
- Tesfaigzi Y. Roles of apoptosis in airway epithelia. *Am J Respir Cell Mol Bio*. 2006;34:537–547.
- Zsembery A, Fortenberry JA, Liang L, et al. Extracellular zinc and ATP restore chloride secretion across cystic fibrosis airway epithelia by triggering calcium entry. *J Biol Chem*. 2004;279:10720–10729.
- Luk CK, Dulfano MJ. Effect of pH, viscosity and ionic-strength changes on ciliary beating frequency of human bronchial explants. *Clin Sci (Lond)*. 1983;64:449–451.