

Molecular Characterization of Extended-Spectrum-Cephalosporin-Resistant *Enterobacteriaceae* from Wild Kelp Gulls in South America

Apostolos Liakopoulos,^a Björn Olsen,^{b,c} Yvon Geurts,^a Karin Artursson,^d Charlotte Berg,^e Dik J. Mevius,^{a,f} Jonas Bonnedahl^g

Department of Bacteriology and Epidemiology, CVI of Wageningen University, Lelystad, the Netherlands^a; Section of Clinical Microbiology and Infectious Diseases, Department of Medical Sciences, Uppsala University, Uppsala, Sweden^b; Zoonosis Science Center, IMBIM, Uppsala University, Uppsala, Sweden^c; National Veterinary Institute, SVA, Uppsala, Sweden^d; Department of Animal Environment and Health, Swedish University of Agricultural Sciences, Uppsala, Sweden^e; Faculty of Veterinary Medicine, Department of Infectious Diseases and Immunology, Utrecht University, Utrecht, the Netherlands^f; Department of Infectious Diseases, Kalmar County Hospital, Kalmar, Sweden^g

Extended-spectrum-cephalosporin-resistant *Enterobacteriaceae* are a public health concern due to limited treatment options. Here, we report on the occurrence and the molecular characteristics of extended-spectrum-cephalosporin-resistant *Enterobacteriaceae* recovered from wild birds (kelp gulls). Our results revealed kelp gulls as a reservoir of various extended-spectrum cephalosporinase genes associated with different genetic platforms. In addition, we report for the first time the presence of a known epidemic clone of *Salmonella enterica* serotype Heidelberg (JF6X01.0326/XbaI.1966) among wild birds.

Extended-spectrum-cephalosporinase-producing *Enterobacteriaceae* have been reported worldwide among isolates obtained from humans and from food-producing and companion animals, as well as from environmental sources (1). In spite of the limited number of studies regarding the occurrence of antibiotic resistance in natural environments, where animals do not naturally come into contact with antibiotics, the occurrence of extended-spectrum-cephalosporin-resistant (ESC^r) *Enterobacteriaceae* has been detected lately in wild birds, especially in populations of gulls (*Laridae*) (2–7). The kelp gull (*Larus dominicanus*) is a large gull species distributed in coastal areas through much of the Southern Hemisphere and is the only gull species inhabiting the Antarctic continent. It is known to be a food generalist, regularly feeding on food resulting from human activities (abattoirs, garbage, sewage outfalls, etc.) (8). This behavior makes it an interesting sentinel species for the study of the environmental spread of antibiotic-resistant bacteria. Our aim was to determine the occurrence and the molecular characteristics of ESC^r *Enterobacteriaceae* isolates recovered from kelp gulls, as this species could favor the dissemination of ESC^r *Enterobacteriaceae* in human populations and in the pristine Antarctic environment.

(Preliminary results from this study were presented as an oral presentation at the 26th European Congress of Clinical Microbiology and Infectious Diseases [ECCMID], 9 to 12 April 2016, Amsterdam, the Netherlands.)

During November 2012, fresh fecal specimens ($n = 50$) were collected from a flock of approximately 500 kelp gulls on a sandy beach where they were roosting in Ushuaia, in Argentina. All samples were enriched either in brain heart infusion broth (Becton-Dickinson, Franklin Lakes, NJ, USA), supplemented with 16 mg/liter vancomycin, or in buffered peptone water (SVA, Uppsala, Sweden) for 18 to 24 h in 37°C and subsequently inoculated on ChromID ESBL (bioMérieux, Solna, Sweden) for the selective isolation of extended-spectrum- β -lactamase (ESBL)-producing *Enterobacteriaceae* or on modified semisolid Rappaport Vassiliadis agar (SVA, Uppsala, Sweden) for the selective isolation of *Salmonella* species, respectively. Presumptive extended-spectrum-cephalosporinase-producing isolates were identified using

(MALDI-TOF) mass spectrometry (Brucker, Coventry, United Kingdom), while *Salmonella* isolates were further serotyped by the microtitration method. Antibiotic susceptibility of the isolates was assessed by broth microdilution and interpreted according to the epidemiologic cutoff values recommended by the European Committee on Antimicrobial Susceptibility Testing (<http://mic.eucast.org>), whereas ESBL and/or AmpC production was evaluated by a combined disc test, as previously described (9).

Genes conferring the ESC^r phenotype were sought and their genetic location on either the chromosome or a plasmid was determined as previously described (9). Standard methods (PCR-based replicon [rep] typing, plasmid multilocus sequence typing [pMLST]/plasmid double-locus sequence typing [pDLST]/replicon sequence typing [RST], and S1 nuclease pulsed-field gel electrophoresis [PFGE]) were applied for further plasmid analysis, while the conjugal transferability of the extended-spectrum cephalosporinase genes and the presence of known insertion sequences (ISs) upstream of them were examined (9). Genetic relatedness among *Escherichia coli* and *Salmonella enterica* serotype Heidelberg isolates was assessed by MLST and XbaI-PFGE typing, respectively, as previously described (9, 10).

Overall, we recovered 37 nonduplicate ESC^r *Enterobacteriaceae* isolates from 34 of the fecal samples included in the study. Among them, 91.9% ($n = 34$) were identified as *E. coli* and 8.1% ($n = 3$) as *S. Heidelberg*. The copresence of ESC^r *E. coli* and *S. Heidelberg* was documented in three fecal samples. The recovered isolates exhibited non-wild-type MICs mainly for ciprofloxacin ($n = 27$; 73.0%), nalidixic acid ($n = 25$; 67.6%), tetracycline ($n = 22$;

Received 25 May 2016 Returned for modification 5 July 2016

Accepted 23 August 2016

Accepted manuscript posted online 29 August 2016

Citation Liakopoulos A, Olsen B, Geurts Y, Artursson K, Berg C, Mevius DJ, Bonnedahl J. 2016. Molecular characterization of extended-spectrum-cephalosporin-resistant *Enterobacteriaceae* from wild kelp gulls in South America. *Antimicrob Agents Chemother* 60:6924–6927. doi:10.1128/AAC.01120-16.

Address correspondence to Apostolos Liakopoulos, apostolos.liakopoulos@wur.nl.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

TABLE 1 Characteristics of ESC^c Enterobacteriaceae isolates recovered from kelp gulls (*Larus dominicanus*) in Ushuaia, Argentina, in 2012^a

| Species (no. of isolates) | ST/PFGE type (no. of isolates) | Resistance phenotypes (no. of isolates) | Plasmid characteristic(s) | | | | | Upstream region |
|------------------------------|-----------------------------------|--|---|--|---|-----------------------------|---|-------------------------|
| | | | ESBL/AmpC gene (no. of isolates) | Location (no. of isolates) | Plasmid type/subtype (no. of isolates) | Size (kb) | Transferability | |
| <i>E. coli</i> (34) | ST744 (6) | AMP, CAZ, CHL, CIP, CST, CTX, NAL, SMX, TET, TMP (1) AMP, AZM, CAZ, CHL, CIP, CTX, NAL, SMX, TET, TMP (2) AMP, CHL, CIP, CST, CTX, NAL, SMX, TET, TMP (3) AMP, CAZ, CHL, CIP, CTX, NAL, TET (4) | <i>bla</i> _{CTX-M-14} (6) | Plasmid (6) | InclII/ST80 (6) | 105 | Conjugative | — ^e |
| | ST617 (5) | AMP, CAZ, CIP, CTX, NAL, TET (1) | <i>bla</i> _{SHV-2A} (4) | Plasmid (4) | InclII/ST187 (3) InclF/Fl:A1:B1 (1) | 125 70 | Conjugative Conjugative | IS26 IS26 |
| | ST57 (3) | AMP, CAZ, CIP, CTX, GEN, NAL, SMX, TET, TMP (3) | <i>bla</i> _{CTX-M-14} (1) | Plasmid (1) | InclII/ST80 (1) | 105 | Conjugative | — ^e |
| | ST93 (3) | AMP, CAZ, CHL, CTX, SMX, TMP (1) AMP, CAZ, CIP, CTX, GEN, SMX (1) AMP, CAZ, CTX, SMX, TET (1) | <i>bla</i> _{CTX-M-2} (3) <i>bla</i> _{CTX-M-2} (3) | Plasmid (3) Plasmid (1) Chromosome (2) | InclF/Fl:8-A-B1 (3) InclII/ST2 (1) NA | 208 202 NA | Conjugative Conjugative Nonconjugative ^d | ISCR1 ISCR1 ISCR1 |
| | ST4038 (3) | AMP, CAZ, CTX (3) | <i>bla</i> _{SHV-2} (3) | Plasmid (3) | InclII/ST12 (3) | 145 | Conjugative | IS26 |
| | ST10 (2) | AMP, CAZ, CTX, TET (1) AMP, CHL, CIP, CTX (1) | <i>bla</i> _{CTX-M-14} (2) | Plasmid (2) | InclII/ST80 (2) | 105 | Conjugative | — ^e |
| | STNew1 (2) ^b | AMP, CAZ, CIP, CTX, NAL (1) AMP, CAZ, CHL, CIP, CTX, GEN, NAL, SMX (1) | <i>bla</i> _{CTX-M-14} (1) <i>bla</i> _{CTX-M-2} (1) | Plasmid (1) Chromosome (1) | InclII/ST80 (1) NA | 105 NA | Conjugative Nonconjugative ^d | — ^e ISCR1 |
| | ST69 (1) | AMP, CAZ, CTX, SMX, TET (1) | <i>bla</i> _{CTX-M-2} (1) | Chromosome (1) | NA | NA | Nonconjugative ^d | ISCR1 |
| | ST88 (1) | AMP, CTX, TET (1) | <i>bla</i> _{CTX-M-14} (1) | Plasmid (1) | InclII/ST80 (1) | 105 | Conjugative | — ^e |
| | ST101 (1) | AMP, CAZ, CIP, CST, CTX, NAL, SMX, TET (1) | <i>bla</i> _{CTX-M-2} (1) | Plasmid (1) | InclA/C (1) | 100 | Nonconjugative ^d | ISCR1 |
| | ST117 (1) | AMP, CAZ, CIP, CTX, NAL, SMX (1) | <i>bla</i> _{CTX-M-2} (1) | Plasmid (1) | NT (1) | 132 | Conjugative | ISCR1 |
| | ST212 (1) | AMP, CAZ, CTX (1) | <i>bla</i> _{SHV-2} (1) | Plasmid (1) | InclII/ST12 (1) | 121 | Conjugative | IS26 |
| | ST359 (1) | AMP, CAZ, CHL, CIP, CTX, NAL, SMX, TET (1) | <i>bla</i> _{CTX-M-2} (1) | Plasmid (1) | InclF/F24-A-B1 (1) | 170 | Conjugative | ISCR1 |
| | ST1011 (1) | AMP, CAZ, CIP, CTX, GEN, NAL, SMX (1) | <i>bla</i> _{CTX-M-2} (1) | Chromosome (1) | NA | NA | Nonconjugative ^d | ISCR1 |
| | ST1193 (1) | AMP, CAZ, CIP, CTX, NAL, TET (1) | <i>bla</i> _{CTX-M-15} (1) | Plasmid (1) | InclF/Fl:A1:B1 (1) | 87 | Nonconjugative ^d | ISFcp1 |
| ST2485 (1) | AMP, CAZ, CHL, CTX, SMX, TET (1) | <i>bla</i> _{CTX-M-2} (1) | Plasmid (1) | InclF/F24-A-B10 (1) | 205 | Conjugative | ISCR1 | |
| STNew2 (1) ^b | AMP, CAZ, CIP, CTX, NAL, SMX (1) | <i>bla</i> _{CTX-M-2} (1) | Chromosome (1) | NA | NA | Nonconjugative ^d | ISCR1 | |
| 5. Heidelberg (3) | JF6X01.0326 (3) ^c | AMP, CAZ, CIP, CTX, NAL (3) | <i>bla</i> _{CMV-2} (3) | Plasmid (3) | InclII/ST12 (3) | 110 | Conjugative | ISFcp1 |

^a ST₁ sequence type; AMP, ampicillin; AZM, azithromycin; CAZ, ceftazidime; CIP, ciprofloxacin; CHL, chloramphenicol; CST, colistin; CTX, cefotaxime; GEN, gentamicin; MEMF, meropenem; NAL, nalidixic acid; SMX, sulfamethoxazole; TET, tetracycline; TGC, tigecycline; TMP, trimethoprim; NA, not applicable.

^b Assignment to a specific ST could not be performed, as uploading new sequences and STs based on ABI files is no longer supported by the MLST database (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>).

^c PFGE pattern numbers correspond to the PulseNet database.

^d No transconjugants were obtained after liquid mating experiments, suggesting either the presence of nonconjugative plasmids or conjugation frequencies below the detection limit ($\leq 1 \times 10^{-9}$).

^e ISFcp1, ISCR1, or IS26 insertion sequences were not found upstream of the ESBL genes for these STs.

59.5%), sulfamethoxazole ($n = 20$; 54.0%), and chloramphenicol ($n = 15$; 40.5%). All isolates were susceptible to meropenem and tigecycline, whereas they exhibited non-wild-type MICs for the remaining tested agents (ranging from 5.4% to 27.0%). All *E. coli* isolates exhibited an ESBL phenotype and carried $bla_{CTX-M-2}$ ($n = 14$; 41.2%), $bla_{CTX-M-14}$ ($n = 11$; 32.3%), bla_{SHV-2} ($n = 4$; 11.8%), bla_{SHV-2A} ($n = 4$; 11.8%), and $bla_{CTX-M-15}$ ($n = 1$; 2.9%) genes, whereas all *S. Heidelberg* isolates exhibited an AmpC phenotype and carried the bla_{CMY-2} gene.

The broad-host-range IncI1 ($n = 21$; 67.7%) and narrow-host-range IncF ($n = 7$; 22.6%) plasmids were by far the most common rep types accounted for the ESC^r phenotype among the recovered isolates. The $bla_{CTX-M-2}$ gene was found mainly on the chromosome ($n = 6$; 42.9%) or on plasmids of different replicon types, including IncF plasmids with fused FIB-FII replicons ($n = 5$; 35.7%), IncHI2 ($n = 1$; 7.1%), IncA/C ($n = 1$; 7.1%), and non-typeable ones ($n = 1$; 7.14%). The $bla_{CTX-M-14}$ and $bla_{CTX-M-15}$ genes were identified exclusively on IncI1/sequence type 80 (ST80) and IncFIA-FIB plasmids, respectively. The bla_{SHV-2} gene was associated with IncI1/ST12 and bla_{SHV-2A} with IncI1/ST187 and IncFIA-FIB plasmids, whereas the bla_{CMY-2} gene was located on IncI1/ST12 plasmids. Detailed results regarding the subtyping, the size, and the transferability of the plasmids are summarized in Table 1.

Three insertion sequence elements previously associated with the mobilization and support of extended-spectrum cephalosporinase genes were identified. Briefly, in all isolates carrying $bla_{CTX-M-2}$, the gene was accompanied upstream by a copy of ISCR1 in the same orientation as the resistance gene, regardless of the plasmid replicon type or whether the gene was chromosomally located (Table 1). Similarly, ISEcp1 was found upstream of the $bla_{CTX-M-15}$ and bla_{CMY-2} genes, while IS26 was found upstream of the bla_{SHV-2} and bla_{SHV-2A} genes. ISEcp1, ISCR1, or IS26 insertion sequences were not found upstream of $bla_{CTX-M-14}$ gene (Table 1).

High diversity of genotypes was observed among the *E. coli* isolates, resulting in 17 different STs, each comprised of one to six isolates. The most predominant genotypes were ST744 ($n = 6$; 17.6%), ST617 ($n = 5$; 14.7%), ST57 ($n = 3$; 8.8%), ST93 ($n = 3$; 8.8%), and ST4038 ($n = 3$; 8.8%), while isolates belonging to ST10, ST69, ST88, ST101, ST117, ST212, ST359, ST1011, ST1193, ST2485, STNew1, and STNew2 were also identified. All *S. Heidelberg* isolates belonged to epidemic clone JF6X01.0326/XbaI.1966 (PulseNet database). Different ESBL determinants were found among isolates with the same genotype; conversely, different genotypes carrying the same ESBL determinants were identified (Table 1).

Several studies have documented the occurrence of ESC^r *Enterobacteriaceae* isolates among wild birds at prevalences ranging from 0% to 37% (4, 11–15). However, our study revealed a higher occurrence among kelp gulls in accordance with studies regarding Brown-headed gulls and Franklin's gulls (5). Although the resistance gene families described in this study are similar to those reported previously (2, 4, 5, 12, 14–19), we documented for the first time the presence of bla_{SHV-2A} and the predominance of $bla_{CTX-M-2}$ among wild birds. The latter mirrors the situation observed for nosocomial infections in Argentinian hospitals (20, 21), confirming the endemicity of $bla_{CTX-M-2}$ within this area and its potential transmission from humans to wild birds and/or vice versa. Of note was the association of $bla_{CTX-M-2}$ gene with ISCR1 on four different plasmid replicon types associated with six differ-

ent *E. coli* STs and on the chromosome of five other different *E. coli* STs, underscoring that ISCR1 has probably played a significant role in the capture of this gene by conjugative plasmids and in its further interreplicon and interclone dissemination. Moreover, our data suggest the horizontal transfer of a conjugative IncI1/ST80 plasmid (105 kb) carrying $bla_{CTX-M-14}$ among five different *E. coli* STs, underscoring the dissemination of this gene owing to a successful plasmid-gene combination.

Among the 17 different STs detected here, we identified several, namely, ST10, ST69, ST101, ST117, ST167, ST617, and ST744, that have been previously reported from ESC^r *E. coli* isolates of human and animal origin (1, 5, 12, 15). Interestingly, some of the identified STs (ST10, ST117, ST157, ST359, ST617, and ST744) have been previously reported among wild birds as well, but they have been found to harbor different extended-spectrum cephalosporinase genes, suggesting that avian commensal *E. coli* strains play a role in the maintenance and dissemination of these genes (1, 5, 12, 15). In contrast with the solely ESC^r *S. Heidelberg* isolate carrying bla_{CMY-2} on a 97-kb IncN plasmid reported previously from an Argentinian adult inpatient (22), here we documented for the first time the presence in wild birds of a known epidemic ESC^r *S. Heidelberg* clone (JF6X01.0326/XbaI.1966), carrying bla_{CMY-2} on a 110-kb IncI1/ST12 plasmid. This PFGE type, circulating in the United States and recently introduced to Europe (9), has been documented to cause outbreaks and exhibit potency for bloodstream infections (23).

In conclusion, although there are few studies on the presence of resistance genes conferring the ESC^r phenotype among *Enterobacteriaceae* from wild birds, to our knowledge this is the first report presenting a detailed characterization of ESC^r *Enterobacteriaceae*, including the underlying antibiotic resistance gene content and its genetic support (plasmids and IS elements). Our data imply that kelp gulls act as reservoirs of a variety of extended-spectrum cephalosporinase genes associated with different genetic platforms that could facilitate their horizontal transfer. In addition, our findings underscore the potential role of kelp gulls as a bridge species for transfer of ESC^r *Enterobacteriaceae* between humans and wildlife and as a spreader of these isolates among human populations and naturally antibiotic-resistant-bacterium-free environments (Antarctic continent) via their movement and migration.

ACKNOWLEDGMENTS

We gratefully acknowledge Kees Veldman, Joop Testerink, and Marga Japing for the antimicrobial susceptibility testing of the isolates and Quark Expeditions for supporting the field trip.

FUNDING INFORMATION

This work was supported by the Dutch Ministry of Economic Affairs (BO-22.04-008-001).

REFERENCES

- Ewers C, Bethe A, Semmler T, Guenther S, Wieler LH. 2012. Extended-spectrum beta-lactamase-producing and AmpC-producing *Escherichia coli* from livestock and companion animals, and their putative impact on public health: a global perspective. *Clin Microbiol Infect* 18:646–655. <http://dx.doi.org/10.1111/j.1469-0691.2012.03850.x>.
- Bonnedahl J, Drobni M, Gauthier-Clerc M, Hernandez J, Granholm S, Kayser Y, Melhus A, Kahlmeter G, Waldenström J, Johansson A, Olsen B. 2009. Dissemination of *Escherichia coli* with CTX-M type ESBL between humans and yellow-legged gulls in the South of France. *PLoS One* 4:e5958. <http://dx.doi.org/10.1371/journal.pone.0005958>.

3. Bonnedahl J, Drobni P, Johansson A, Hernandez J, Melhus A, Stedt J, Olsen B, Drobni M. 2010. Characterization, and comparison, of human clinical and black-headed gull (*Larus ridibundus*) extended-spectrum beta-lactamase-producing bacterial isolates from Kalmar, on the southeast coast of Sweden. *J Antimicrob Chemother* 65:1939–1944. <http://dx.doi.org/10.1093/jac/dkq222>.
4. Bonnedahl J, Hernandez J, Stedt J, Waldenstrom J, Olsen B, Drobni M. 2014. Extended-spectrum beta-lactamases in *Escherichia coli* and *Klebsiella pneumoniae* in gulls, Alaska, USA. *Emerg Infect Dis* 20:897–899.
5. Báez J, Hernández-García M, Guamparito C, Díaz S, Olave A, Guerrero K, Cantón R, Baquero F, Gahona J, Valenzuela N, Del Campo R, Silva J. 2015. Molecular characterization and genetic diversity of ESBL-producing *Escherichia coli* colonizing the migratory Franklin's gulls (*Leucophaeus pipixcan*) in Antofagasta, North of Chile. *Microb Drug Resist* 21:111–116. <http://dx.doi.org/10.1089/mdr.2014.0158>.
6. Bonnedahl J, Stedt J, Waldenstrom J, Svensson L, Drobni M, Olsen B. 2015. Comparison of extended-spectrum beta-lactamase (ESBL) CTX-M genotypes in Franklin gulls from Canada and Chile. *PLoS One* 10: e0141315.
7. Stedt J, Bonnedahl J, Hernandez J, Waldenström J, McMahon BJ, Tolf C, Olsen B, Drobni M. 2015. Carriage of CTX-M type extended spectrum beta-lactamases (ESBLs) in gulls across Europe. *Acta Vet Scand* 57:74. <http://dx.doi.org/10.1186/s13028-015-0166-3>.
8. Bertellotti M, Yorio P. 1999. Spatial and temporal patterns in the diet of the kelp gull in Patagonia. *Condor* 101:790–798. <http://dx.doi.org/10.2307/1370066>.
9. Liakopoulos A, Geurts Y, Dierikx CM, Brouwer MSM, Kant A, Wit B, Heymans R, van Pelt W, Mevius DJ. July 2016. Introduction of extended-spectrum cephalosporin-resistant *Salmonella enterica* serotype Heidelberg strains in the Netherlands. *Emerg Infect Dis* <http://dx.doi.org/10.3201/eid2207.151377>.
10. Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, Karch H, Reeves PR, Maiden MC, Ochman H, Achtman M. 2006. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 60: 1136–1151. <http://dx.doi.org/10.1111/j.1365-2958.2006.05172.x>.
11. Guenther S, Ewers C, Wieler LH. 19 December 2011. Extended-spectrum beta-lactamases producing E-coli in wildlife, yet another form of environmental pollution? *Front Microbiol* <http://dx.doi.org/10.3389/fmicb.2011.00246>.
12. Guenther S, Aschenbrenner K, Stamm I, Bethe A, Semmler T, Stubbe A, Stubbe M, Batsajkhan N, Glupczynski Y, Wieler LH, Ewers C. 2012. Comparable high rates of extended-spectrum-beta-lactamase-producing *Escherichia coli* in birds of prey from Germany and Mongolia. *PLoS One* 7:e53039. <http://dx.doi.org/10.1371/journal.pone.0053039>.
13. Jarhult JD, Stedt J, Gustafsson L. 25 July 2013. Zero prevalence of extended spectrum beta-lactamase-producing bacteria in 300 breeding collared flycatchers in Sweden. *Infect Ecol Epidemiol* <http://dx.doi.org/10.3402/iee.v3i0.20909>.
14. Veldman K, van Tulden P, Kant A, Testerink J, Mevius D. 2013. Characteristics of cefotaxime-resistant *Escherichia coli* from wild birds in the Netherlands. *Appl Environ Microbiol* 79:7556–7561. <http://dx.doi.org/10.1128/AEM.01880-13>.
15. Alcalá L, Alonso CA, Simón C, González-Esteban C, Orós J, Rezusta A, Ortega C, Torres C. 21 December 2015. Wild birds, frequent carriers of extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* of CTX-M and SHV-12 types. *Microb Ecol* <http://dx.doi.org/10.1007/s00248-015-0718-0>.
16. Hasan B, Melhus A, Sandegren L, Alam M, Olsen B. 2014. The gull (*Chroicocephalus brunnicephalus*) as an environmental bioindicator and reservoir for antibiotic resistance on the coastlines of the Bay of Bengal. *Microb Drug Resist* 20:466–471. <http://dx.doi.org/10.1089/mdr.2013.0233>.
17. Ewers C, Bethe A, Stamm I, Grobbel M, Kopp PA, Guerra B, Stubbe M, Doi Y, Zong Z, Kola A, Schaufler K, Semmler T, Fruth A, Wieler LH, Guenther S. 2014. CTX-M-15-D-ST648 *Escherichia coli* from companion animals and horses: another pandemic clone combining multiresistance and extraintestinal virulence? *J Antimicrob Chemother* 69:1224–1230. <http://dx.doi.org/10.1093/jac/dkt516>.
18. Literak I, Dolejska M, Janoszowska D, Hrusakova J, Meissner W, Rzycka H, Bzoma S, Cizek A. 2010. Antibiotic-resistant *Escherichia coli* bacteria, including strains with genes encoding the extended-spectrum beta-lactamase and QnrS, in waterbirds on the Baltic Sea Coast of Poland. *Appl Environ Microbiol* 76:8126–8134. <http://dx.doi.org/10.1128/AEM.01446-10>.
19. Hernandez J, Bonnedahl J, Eliasson I, Wallensten A, Comstedt P, Johansson A, Granholm S, Melhus A, Olsen B, Drobni M. 2010. Globally disseminated human pathogenic *Escherichia coli* of O25b-ST131 clone, harbouring blaCTX-M-15, found in Glaucous-winged gull at remote Commander Islands, Russia. *Environ Microbiol Rep* 2:329–332. <http://dx.doi.org/10.1111/j.1758-2229.2010.00142.x>.
20. Sennati S, Santella G, Di Conza J, Pallecchi L, Pino M, Ghiglione B, Rossolini GM, Radice M, Gutkind G. 2012. Changing epidemiology of extended-spectrum beta-lactamases in Argentina: emergence of CTX-M-15. *Antimicrob Agents Chemother* 56:6003–6005. <http://dx.doi.org/10.1128/AAC.00745-12>.
21. Quinteros M, Radice M, Gardella N, Rodriguez MM, Costa N, Korbenfeld D, Couto E, Gutkind G; Microbiology Study Group. 2003. Extended-spectrum beta-lactamases in enterobacteriaceae in Buenos Aires, Argentina, public hospitals. *Antimicrob Agents Chemother* 47:2864–2867. <http://dx.doi.org/10.1128/AAC.47.9.2864-2867.2003>.
22. Cejas D, Vignoli R, Quinteros M, Marino R, Callejo R, Betancor L, Gutkind GO, Radice MA. 2014. First detection of CMY-2 plasmid mediated beta-lactamase in *Salmonella* Heidelberg in South America. *Rev Argent Microbiol* 46:30–33. [http://dx.doi.org/10.1016/S0325-7541\(14\)70044-6](http://dx.doi.org/10.1016/S0325-7541(14)70044-6).
23. Centers for Disease Control and Prevention. 2011. Investigation update: multistate outbreak of human *Salmonella* Heidelberg infections linked to “kosher broiled chicken livers” from Schreiber Processing Corporation. Centers for Disease Control and Prevention, Atlanta, GA.