

Type I γ Phosphatidylinositol Phosphate 5-Kinase i5 Controls the Ubiquitination and Degradation of the Tumor Suppressor Mitogen-inducible Gene 6*

Received for publication, May 4, 2016, and in revised form, August 11, 2016. Published, JBC Papers in Press, August 24, 2016, DOI 10.1074/jbc.M116.736041

Ming Sun[‡], Jinyang Cai[‡], Richard A. Anderson[§], and Yue Sun^{‡1}

From the [‡]Philips Institute for Oral Health Research, School of Dentistry and Massey Cancer Center, Virginia Commonwealth University, Richmond, Virginia 23298 and the [§]Molecular and Cellular Pharmacology Program, University of Wisconsin, School of Medicine and Public Health, Madison, Wisconsin 53706

Mitogen-inducible gene 6 (Mig6) is a tumor suppressor, and the disruption of Mig6 expression is associated with cancer development. Mig6 directly interacts with epidermal growth factor receptor (EGFR) to suppress the activation and downstream signaling of EGFR. Therefore, loss of Mig6 enhances EGFR-mediated signaling and promotes EGFR-dependent carcinogenesis. The molecular mechanism modulating Mig6 expression in cancer remains unclear. Here we demonstrate that type I γ phosphatidylinositol phosphate 5-kinase i5 (PIPKI γ i5), an enzyme producing phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂), stabilizes Mig6 expression. Knockdown of PIPKI γ i5 leads to the loss of Mig6 expression, which dramatically enhances and prolongs EGFR-mediated cell signaling. Loss of PIPKI γ i5 significantly promotes Mig6 protein degradation via proteasomes, but it does not affect the Mig6 mRNA level. PIPKI γ i5 directly interacts with the E3 ubiquitin ligase neuronal precursor cell-expressed developmentally down-regulated 4-1 (NEDD4-1). The C-terminal domain of PIPKI γ i5 and the WW1 and WW2 domains of NEDD4-1 are required for their interaction. The C2 domain of NEDD4-1 is required for its interaction with PtdIns(4,5)P₂. By binding with NEDD4-1 and producing PtdIns(4,5)P₂, PIPKI γ i5 perturbs NEDD4-1-mediated Mig6 ubiquitination and the subsequent proteasomal degradation. Thus, loss of NEDD4-1 can rescue Mig6 expression in PIPKI γ i5 knockdown cells. In this way, PIPKI γ i5, NEDD4-1, and Mig6 form a novel molecular nexus that controls EGFR activation and downstream signaling.

EGF receptor (EGFR)² is a critical receptor tyrosine kinase that controls cell growth and differentiation during embryo-

genesis and adult homeostasis (1). After stimulation by its agonists, EGFR mediates multiple downstream signaling cascades that control cell proliferation, apoptosis, migration, survival, angiogenesis, and other biological processes (2). The appropriate regulation of EGFR activation is critical to maintain normal physiology, and loss or overactivation of EGFR can lead to multiple diseases (2, 3). The overexpression or aberrant kinase activity of EGFR could result in unregulated growth stimulation, tumorigenesis, and metastasis in various tumor types (4). Therefore, targeted therapies directed against EGFR have been a primary focus of antitumor drug development over the past 10 years (5). An understanding of the molecular mechanisms controlling EGFR activation is required to design efficient anti-EGFR therapies and to identify biomarkers predicting higher sensitivity to anti-EGFR drugs.

Mig6 (also known as RALT or Gene 33) is a widely expressed adaptor protein that directly binds to the kinase domain of EGFR to block its activation (6–8). Agonist binding stimulates the formation of an asymmetric EGFR dimer by the intracellular kinase domains in which the carboxy-terminal lobe (C lobe) of one kinase domain induces an active conformation in the other (9). Mig6 inhibits EGFR activation by blocking formation of the activating EGFR dimer interface (6). Furthermore, Mig6 controls the endosomal trafficking of EGFR (10). Following agonist stimulation, activated EGFR is internalized, and subsequent trafficking events determine the fate of internalized EGFR, including recycling back to the plasma membrane or trafficking to the lysosomes for degradation (11, 12). Mig6 ensures the sorting of internalized EGFR to late endosomes and the subsequent sorting to lysosomes for degradation (10). Mig6 can be transcriptionally induced by growth factors, cytokines, and cell stress (13–16). Deletion of Mig6 leads to hyperactivated EGFR signaling and causes carcinogenesis in multiple organs (14, 17). The disruption of Mig6 in mice accelerates initiation and progression of mutant EGFR-driven lung adenocarcinoma (18). Thus, Mig6 is a tumor suppressor, and its expression is down-regulated in various cancers (14). The molecular mechanisms controlling Mig6 expression in cancer remain to be clarified.

Type I γ phosphatidylinositol phosphate kinase (PIPKI γ) is an enzyme that generates PtdIns(4,5)P₂ (19). The PIPKI γ gene

* This work was supported by Scientist Development Grant 12SDG11950022 from the American Heart Association and Institutional Research Grant IRG-14-192-40 from the American Cancer Society. The authors declare that they have no conflicts of interest with the contents of this article.

¹ To whom correspondence should be addressed: Philips Institute for Oral Health Research, School of Dentistry and Massey Cancer Center, Virginia Commonwealth University, 4154 Perkinson Bldg., 1101 E. Leigh St., P.O. Box 980566, Richmond, VA 23298-0566. Tel.: 804-827-6094; Fax: 804-828-0150; E-mail: ysun4@vcu.edu.

² The abbreviations used are: EGFR, EGF receptor; PIPKI γ i5, type I γ phosphatidylinositol phosphate 5-kinase i5; PtdIns(4,5)P₂, phosphatidylinositol 4,5-bisphosphate; KD, kinase-dead; CRISPR, clustered regularly interspaced short palindromic repeats; sgRNA, single guide RNA; PAM, protospacer adjacent motif; Ni²⁺-NTA, nickel-nitrilotriacetic acid; HECT, the homologous to the E6AP carboxyl terminus; PtdIns(3,4,5)P₃, phosphatidylinositol 3,4,5-trisphosphate; PTEN, phosphatase and tensin homolog; ILV, intraluminal vesicle; ESCRT, endosomal sorting complex required for trans-

port; Hrs, hepatocyte growth factor-regulated tyrosine kinase substrate; IP, immunoprecipitation; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PIP, phosphatidylinositol phosphate.

PIP2KIγ5 Controls Mig6 Ubiquitination and Degradation




FIGURE 1. PIP2KIγ5 controls Mig6 expression and EGFR signaling. MDA-MB-231 cells were transfected with control siRNA, PIP2KIγ5 (*lγi5*) siRNA, or Mig6 siRNA separately. 72 h after siRNAs transfection, cells were stimulated with TGFα (10 nM) for the times indicated. *A*, the EGFR protein level, EGFR activation, AKT activation, and Mig6 expression were detected. The following were quantified: Mig6 protein level in TGFα-untreated cells (*B*), EGFR activation detected by Tyr(P)-1068 antibody (*C*), and AKT activation detected by Ser(P)-473 antibody (*D*). Quantification of the Mig6 protein level was normalized with the GAPDH level. Quantification of EGFR or AKT activation was normalized with the total EGFR or AKT level. GFP, HA-tagged wild-type PIP2KIγ5, or kinase dead-mutant PIP2KIγ5 was constructed into PWPT (Addgene) lentivirus vectors. GFP, HA-PIP2KIγ5 WT, or HA-PIP2KIγ5 KD was then expressed by lentivirus transfection in PIP2KIγ5 knockdown MDA-MB-231 cells, and the effects on Mig6 expression were detected by immunoblotting (*E*) and quantified (*F*). Error bars indicate mean ± S.E. from three independent experiments. *, *p* < 0.05.

is alternatively spliced, resulting in at least six protein variants expressed in humans, known as PIP2KIγ1-i6 (20). These splicing variants have the same N terminus and kinase domain, and they only differ by different extensions at the C terminus (20, 21). These extensions allow for association with unique binding partners and allow for different PIP2KIγ splice variants to produce PtdIns(4,5)P₂ with distinct subcellular distributions necessary to perform specific biological functions. Distinct roles of different PIP2KIγ splice variants have been found in mediating or controlling EGFR signaling. For instance, PIP2KIγ2 specifically binds to talin at cell adhesions, and the production of PtdIns(4,5)P₂ modulates talin-integrin interaction to regulate inside-out integrin signaling and cell adhesion turnover (22). EGFR phosphorylates PIP2KIγ2 to control local PtdIns(4,5)P₂ levels at adhesions. In this way, PIP2KIγ2 is required for nascent adhesion formation at the leading edge to facilitate EGFR-induced cell directional migration (22). Unlike PIP2KIγ2, PIP2KIγ5 is localized at endosomes and is required for EGFR sorting from endosomes to lysosomes for degradation (23). Therefore, the loss of PIP2KIγ5 blocks the lysosomal trafficking and degradation of activated EGFR, which dramatically enhances and prolongs EGFR activation and downstream signaling such as AKT and ERK MAPK activation (23). These results indicate that PIP2KIγ5 is a critical regulator of EGFR.

Here we show that PIP2KIγ5 directly interacts with an E3 ubiquitin ligase, NEDD4-1, to block NEDD4-1-mediated ubiquitination of Mig6. In this way, PIP2KIγ5 protects Mig6 from

being degraded by the proteasome. Therefore, PIP2KIγ5 ensures Mig6 expression and down-regulates EGFR activation. The data uncover a novel molecular nexus composed of PIP2KIγ5, Mig6, and NEDD4-1 that controls EGFR signaling.

Results

Loss of PIP2KIγ5 Dramatically Decreases Mig6 Expression— PIP2KIγ5 is required for down-regulating EGFR activation and signaling (23). However, the molecular mechanism by which PIP2KIγ5 controls EGFR is not fully understood. Mig6 is a suppressor of EGFR, and knockdown of PIP2KIγ5 by siRNA in MDA-MB-231 cells dramatically decreased Mig6 protein expression levels (Fig. 1, *A* and *B*). Consistently, loss of PIP2KIγ5 significantly enhanced and prolonged TGFα-induced autophosphorylation of EGFR on tyrosine 1068 (Fig. 1, *A* and *C*), indicating increased levels of EGFR activation. Downstream AKT activation was also enhanced and prolonged by PIP2KIγ5 knockdown (Fig. 1, *A* and *D*). Knockdown of Mig6 by siRNA similarly promoted EGFR activation and AKT activation as the effects induced by PIP2KIγ5 knockdown (Fig. 1, *A*, *C*, and *D*). These results indicate that PIP2KIγ5 can regulate EGFR signaling by modulating Mig6 expression.

To determine whether PIP2KIγ5 lipid kinase activity was required for modulation of Mig6 expression, a knockdown-rescue approach was developed. siRNA was used to knock down endogenous PIP2KIγ5, and then wild-type PIP2KIγ5 or a kinase-dead mutant (PIP2KIγ5KD) construct containing siRNA-




FIGURE 2. **CRISPR-mediated genome disruption of PIPKI γ 5 leads to loss of Mig6 expression.** Two different guide RNA sequences were designed to induce genome editing of PIPKI γ 5 (γ 5) in MDA-MB-231 or H-1975 cells separately. Single clones were selected and validated by sequencing the targeted PIPKI γ 5 loci. The sequences of two mutant alleles in MDA-MB-231 single cell clone 4 are shown in (A). PIPKI γ 5 and Mig6 were detected from MDA-MB-231 parental (Par) or single clone 4 by immunoblotting (B), and Mig6 expression levels were quantified (C). The sequences of two mutant alleles in H-1975 single cell clone 2 are shown in (D). PIPKI γ 5 and Mig6 were detected from H-1975 parental or single clone 2 by immunoblotting (E), and Mig6 expression levels were quantified (F). Protospacer adjacent motif (PAM) sequences are underlined. Error bars indicate mean \pm S.E. from three independent experiments. *, $p < 0.05$.

resistant silent mutations was re-expressed using lentivirus-mediated infection. Expression of wild-type PIPKI γ 5 but not PIPKI γ 5KD significantly rescued Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 1, E and F). These results confirm the roles of PIPKI γ 5 in Mig6 expression and indicate that kinase activity with resulting production of PtdIns(4,5)P₂ is required for PIPKI γ 5 control of Mig6 expression.

To rule out possible PIPKI γ 5 siRNA off-target effects, another approach, clustered regularly interspaced short palindromic repeats (CRISPR)-cas9 genome editing (24, 25), was used to disrupt PIPKI γ 5 expression. Two different sgRNA sequences were designed to mediate the genome editing of PIPKI γ 5 in a breast cancer cell line, MDA-MB-231, and a lung cancer cell line, H-1975. Single clones with CRISPR-cas9-mediated PIPKI γ 5 genome editing were selected from MDA-MB-231 and H-1975 cells. The single clone selected from MDA-MB-231 has an in-frame 28-bp deletion at the targeted PIPKI γ 5 locus in one allele, leading to full disruption, and an in-frame 9-bp deletion at another allele, leading to a deletion of three amino acids at the C terminus of PIPKI γ 5 (Fig. 2A). Although it is not a full disruption of PIPKI γ 5, PIPKI γ 5 protein expression is dramatically decreased and Mig6 expression is significantly decreased in this clone, similar to the siRNA-induced PIPKI γ 5 knockdown cells (Fig. 2, B and C). The H-1975 single clone has an in-frame 8-bp and an in-frame 59-bp deletion at the PIPKI γ 5 locus in two different alleles separately (Fig. 2D). This is a full disruption of PIPKI γ 5, and the Mig6 expression in this clone is significantly decreased (Fig.

2, E and F). These results further confirm that PIPKI γ 5 controls Mig6 expression.

To demonstrate whether PIPKI γ 5 regulates steady-state Mig6 mRNA levels, a real-time PCR assay was performed to detect the mRNA levels of Mig6 in control or PIPKI γ 5 knockdown cells. As shown in Fig. 3A, knockdown of PIPKI γ 5 does not significantly change Mig6 mRNA levels. This suggests that PIPKI γ 5 may modulate Mig6 protein degradation but not the steady-state Mig6 mRNA levels. The efficiency of PIPKI γ 5 knockdown was confirmed by Western blotting (Fig. 3B). To define whether PIPKI γ 5-regulated Mig6 degradation is dependent on lysosomes or proteasomes, the lysosome inhibitor chloroquine or the proteasome inhibitor MG132 was used to treat the cells. Only inhibition of the proteasome function increased Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 3C). This shows that PIPKI γ 5-regulated Mig6 degradation is dependent on proteasomes but not lysosomes. To confirm the efficiency of chloroquine in inhibiting lysosome function, the LC3A/BII level was detected because its degradation is dependent on lysosomes (26). To test the efficiency of MG132 in blocking proteasome function, the levels of SMAD ubiquitination regulatory factor 1 (Smurf1) and TGF β receptor I (TGF β RI) were detected because their degradation is dependent on proteasomes (27, 28). Chloroquine treatment increased the LC3A/BII level (Fig. 3D), and MG132 treatment enhanced Smurf1 and TGF β RI levels (Fig. 3E). This validates the function of these two inhibitors. Exogenously expressed HA-tagged Mig6 was also degraded when endogenous PIPKI γ 5 was lost

PIPKI γ 5 Controls Mig6 Ubiquitination and Degradation




FIGURE 3. PIPKI γ 5 controls the protein degradation of Mig6. *A*, control siRNA or PIPKI γ 5 (*l* γ 5) siRNA was transfected into MDA-MB-231 cells. 72 h after siRNAs transfection, the Mig6 mRNA levels in these cells were quantified by real-time PCR. *B*, PIPKI γ 5 expression in control or PIPKI γ 5 knockdown cells was detected by immunoblotting. *C*, control or PIPKI γ 5 knockdown MDA-MB-231 cells were treated with the lysosome inhibitor chloroquine (80 μ M) or the proteasome inhibitor MG132 (50 μ M) for 4 h, and then Mig6 expression was detected. *D*, MDA-MB-231 cells were treated with chloroquine (80 μ M) for the times indicated, and then the LC3A/BII protein level was detected. *E*, MDA-MB-231 cells were treated with MG132 (50 μ M) for the times indicated, and then the Smurf1 or TGF β R1 protein levels were detected. *F*, HA-Mig6 was transfected into control or PIPKI γ 5 knockdown MDA-MB-231 cells, and then the expression of HA-Mig6 was detected by anti-HA antibody or anti-Mig6 antibody via immunoblotting. Error bars indicate mean \pm S.E. from three independent experiments.

(Fig. 3*F*). This further supports that PIPKI γ 5 controls Mig6 protein degradation but not the mRNA level.

The E3 Ubiquitin Ligase NEDD4-1 Is Responsible for PIPKI γ 5-regulated Mig6 degradation—Mig6 possesses two peptide sequences that are rich in proline, glutamic acid, serine, and threonine (PEST sequences). These PEST sequences promote Mig6 ubiquitination and proteasomal degradation (29). However, the E3 ubiquitin ligase responsible for Mig6 ubiquitination has not been defined. PIPKI γ 5 is involved in the modulation of E3 ligase NEDD4-1-mediated ubiquitination of the hepatocyte growth factor-regulated tyrosine kinase substrate (Hrs) (23). Therefore, the effect of NEDD4-1 on PIPKI γ 5-regulated Mig6 expression was detected. As shown in Fig. 4, *A* and *B*, knockdown of NEDD4-1 rescued Mig6 expression in PIPKI γ 5 knockdown cells. NEDD4-1 belongs to the NEDD4 family E3 ligases. Knockdown of other NEDD4 family E3 ligases such as NEDD4-2, Smurf1, or SMAD ubiquitination regulatory factor 2 (Smurf2) cannot rescue Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 4, *A* and *B*). These results indicate that NEDD4-1 is the E3 ubiquitin ligase responsible for PIPKI γ 5-regulated Mig6 expression. To rule out siRNA off-target effects, three different NEDD4-1 siRNAs were used. All of these siRNAs similarly knocked down NEDD4-1 expression and rescued Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 4, *C* and *D*). These results suggest that PIPKI γ 5 may modulate Mig6 ubiquitination and degradation by regulating NEDD4-1.

To determine whether NEDD4-1 can use Mig6 as a substrate and mediate Mig6 ubiquitination, Mig6-NEDD4-1 interaction was first investigated. Endogenous Mig6 was immunoprecipitated from cell lysates and examined by Western blotting for association of endogenous NEDD4-1. As shown in Fig. 5*A*, NEDD4-1 was detected with the Mig6 complex. Their direct binding was confirmed using *in vitro* pulldown assays with purified GST-NEDD4-1 and His $_6$ -Mig6 (Fig. 5*B*). An *in vivo* ubiquitination assay was then performed to demonstrate whether NEDD4-1 can promote Mig6 ubiquitination. Myc-Mig6 and His $_6$ -ubiquitin were co-transfected into HEK-293 cells, and the ubiquitinated proteins were pulled down by Ni $^{2+}$ -NTA beads. Unubiquitinated proteins may also be pulled down by NTA beads via nonspecific binding or via interaction with ubiquitinated protein. To remove the unubiquitinated proteins, the NTA beads were washed under strong denaturing conditions as described under "Experimental Procedures." The amount of Mig6 in total ubiquitinated proteins was then detected by anti-Myc antibody via Western blotting. NEDD4-1 expression enhanced the Mig6 poly-ubiquitination level (Fig. 5*C*). To further confirm that NEDD4-1 directly ubiquitinates Mig6, an *in vitro* ubiquitination assay was used with purified NEDD4-1 and GST-Mig6 proteins. Addition of Mig6 and NEDD4-1 dramatically enhanced total ubiquitinated protein levels (Fig. 5*D*). Mig6 in the reaction system was immunoprecipitated by anti-Mig6 antibody, and then the levels of ubiquitinated Mig6 were detected by anti-ubiquitin antibody (Fig. 5*E*) via Western blotting. Addition of NEDD4-1 induced Mig6 ubiquitination (Fig. 5*E*). These results indicate that NEDD4-1 ubiquitinates Mig6.

PIPKI γ 5 Directly Interacts with NEDD4-1—The *in vitro* binding assay by purified His $_6$ -PIPKI γ 5 and GST-NEDD4-1 indicated the direct interaction of PIPKI γ 5 with NEDD4-1 (Fig. 6*A*). To determine whether the PIPKI γ 5-NEDD4-1 interaction is specific, the interaction of NEDD4-1 with other PIPKI γ splicing variants was examined by co-IP assay. As shown in Fig. 6*B*, PIPKI γ 1 and PIPKI γ 2 do not have the ability to bind with NEDD4-1. This result demonstrated that the unique C terminus of PIPKI γ 5 is required for its association with NEDD4-1. Consistently, knockdown of PIPKI γ 2 does not affect Mig6 expression (Fig. 6*C*). This suggests that binding with NEDD4-1 is required for PIPKI γ 5 to modulate Mig6 expression. PIPKI γ 5KD still interacts with NEDD4-1 (Fig. 6*B*). This demonstrates that the kinase activity producing PtdIns(4,5)P $_2$ is not required for PIPKI γ 5-NEDD4-1 interaction. A series of truncations in the PIPKI γ 5-specific C terminus were made (Fig. 7*A*), and their ability to interact with NEDD4-1 was examined. The truncation mutant PIPKI γ 5 1–652 (the C terminus after amino acid 652 was deleted) lost the ability to interact with NEDD4-1 (Fig. 7*B*). This further supports that the C terminus of PIPKI γ 5 is required for NEDD4-1 interaction. NEDD4-1 is composed of three functional regions: an N-terminal C2 domain for membrane binding, a central region containing four WW domains for protein-protein interactions, and a C-terminal homologous to the E6AP carboxyl terminus (HECT) domain for ubiquitin protein ligation (30). A series of NEDD4-1 truncation mutants were made (Fig. 7*C*), and the deletion of WW1 and WW2 decreased




FIGURE 4. Loss of the E3 ubiquitin ligase NEDD4-1 rescues Mig6 expression in PIP1 γ 5 knockdown cells. Smurf1 siRNA, Smurf2 siRNA, NEDD4-1 siRNA, or NEDD4-2 siRNA were co-transfected with PIP1 γ 5 (γ 5) siRNA into MDA-MB-231 cells as indicated. 72 h after siRNA transfection, the Mig6 and different E3 ubiquitin ligase expression levels were detected by immunoblotting (A), and the Mig6 expression levels were quantified (B). Three different siRNAs targeting NEDD4-1 were separately transfected into PIP1 γ 5 knockdown MDA-MB-231 cells. The Mig6 and NEDD4-1 expression levels were detected by immunoblotting (C), and the Mig6 expression levels were quantified (D). The NEDD4-1 siRNA used in A and B was NEDD4-1 siRNA1. Quantification of Mig6 protein levels was normalized with actin levels. Error bars indicate mean \pm S.E. from three independent experiments. *, $p < 0.05$.

NEDD4-1 association with PIP1 γ 5 (Fig. 7, D and E). This indicates that the WW1 and WW2 domains are required for NEDD4-1-PIP1 γ 5 interaction.

PIP1 γ 5 Blocks NEDD4-1-mediated Mig6 Ubiquitination—The effect of PIP1 γ 5 on NEDD4-1-mediated ubiquitination was tested by *in vitro* ubiquitination assays. Addition of purified wild-type His₆-PIP1 γ 5 or kinase-dead mutant His₆-PIP1 γ 5 proteins dramatically decreased NEDD4-1-mediated ubiquitination *in vitro* (Fig. 8A). Both wild-type and kinase-dead mutant PIP1 γ 5 interact with NEDD4-1 (Fig. 6B). This suggests that PIP1 γ 5-NEDD4-1 interaction blocks NEDD4-1-mediated ubiquitination in the *in vitro* system.

Unlike the *in vitro* system, although kinase-dead mutant PIP1 γ 5 was expressed at a similar level as wild-type PIP1 γ 5, it was less efficient to decrease NEDD4-1-mediated Mig6 polyubiquitination compared with wild-type PIP1 γ 5 in the *in vivo* system (Fig. 8B). This indicates that kinase activity to produce PtdIns(4,5)P₂ is required for PIP1 γ 5 to efficiently modulate NEDD4-1-mediated Mig6 ubiquitination *in vivo*. This is consistent with the result that only wild-type PIP1 γ 5 but not the kinase-dead mutant can rescue Mig6 expression in PIP1 γ 5 knockdown cells (Fig. 1, E and F). Expression of wild-type PIP1 γ 5 but not the KD mutant decreased endogenous Mig6-NEDD4-1 interaction (Fig. 8, C and D). This suggests that PtdIns(4,5)P₂ produced by PIP1 γ 5 can disturb Mig6-NEDD4-1 interaction *in vivo*.

PtdIns(4,5)P₂ Binds to NEDD4-1 and Decreases NEDD4-1-mediated Mig6 Ubiquitination—The C2 domain of the yeast NEDD4 homolog could bind phosphoinositides (31). To determine whether NEDD4-1 is a PtdIns(4,5)P₂ binding protein, a protein-lipid overlay assay was performed. As shown in Fig. 9A, purified GST-NEDD4-1 can bind multiple species of phospholipids, including PtdIns(4,5)P₂. The C2 domain deletion mutant GST-NEDD4-1 lost the ability to interact with PtdIns(4,5)P₂ (Fig. 9A). This indicates that the C2 domain of NEDD4-1 is required for binding with PtdIns(4,5)P₂.

To further confirm the NEDD4-1-PtdIns(4,5)P₂ interaction, a liposome binding assay was performed. Adding PtdIns(4,5)P₂ into the liposomes significantly enhanced liposome binding with GST-NEDD4-1 (Fig. 9, B and C), although adding PtdIns(4,5)P₂ did not increase liposome binding with C2 domain deletion mutant GST-NEDD4-1 (Fig. 9, D and E). These results further support that NEDD4-1 binds PtdIns(4,5)P₂ and that the C2 domain is critical for this interaction.

To determine whether PtdIns(4,5)P₂ binding can affect NEDD4-1-mediated ubiquitination, an *in vitro* NEDD4-1 E3 ubiquitin ligase activity assay was performed. As shown in Fig. 9F, addition of PtdIns(4,5)P₂-containing liposomes into the system decreased NEDD4-1-mediated ubiquitination. Liposomes not containing PtdIns(4,5)P₂ did not affect NEDD4-1-mediated




FIGURE 6. PIPKI γ 5 directly interacts with NEDD4-1. A, recombinant GST-NEDD4-1 (1 μ g) and His₆-PIPKI γ 5 (1 μ g) were purified from *E. coli* and subjected to pull-down assays with His₆-affinity Ni²⁺-NTA beads. The quality of purified proteins was verified by Coomassie staining. IB, immunoblot. B, HA-NEDD4-1 was co-transfected with Myc-PIPKI γ 5, Myc-PIPKI γ 5 kinase dead mutant, Myc-PIPKI γ 1, or Myc-PIPKI γ 2 separately into HEK-293T cells, subjected to immunoprecipitation with anti-HA antibody, and then immunoblotted with antibodies as indicated. C, control, PIPKI γ 5-, or PIPKI γ 2-specific siRNAs were transfected into MDA-MB-231 cells, and the Mig6 expression levels were detected by Western blotting 72 h after siRNA transfection.

increase Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 1, E and F). Furthermore, disturbing PIPKI γ 5 expression by another approach, CRISPR-cas9 genome editing, similarly decreases the Mig6 level as using PIPKI γ 5 siRNA (Fig. 2). These results clearly indicate that PIPKI γ 5 controls Mig6 expression.

Kinase-dead mutant PIPKI γ 5 interacts with NEDD4-1 to the same extent as wild-type PIPKI γ 5 (Fig. 6B), and it can block NEDD4-1-mediated ubiquitination similarly as wild-type PIPKI γ 5 in the *in vitro* system (Fig. 8A). But in the *in vivo* system, kinase-dead mutant PIPKI γ 5 cannot decrease NEDD4-1-mediated Mig6 ubiquitination as efficiently as wild-type PIPKI γ 5 (Fig. 8B), and only wild-type PIPKI γ 5 can significantly rescue Mig6 expression in PIPKI γ 5 knockdown cells (Fig. 1F). These results suggest that PtdIns(4,5)P₂ plays an important role in modulating Mig6 degradation. PtdIns(4,5)P₂ binds to NEDD4-1 and decreases NEDD4-1-mediated ubiquitination (Fig. 9). Furthermore, PtdIns(4,5)P₂ can modulate NEDD4-1-Mig6 interaction *in vivo*, as only wild-type PIPKI γ 5 but not kinase-dead PIPKI γ 5 decreased endogenous NEDD4-1-Mig6 interaction (Fig. 8, C and D). In the *in vivo* system, PtdIns(4,5)P₂ may also modulate NEDD4-1 interaction with

other components to affect Mig6 ubiquitination, and this cannot be reflected in the *in vitro* ubiquitination assay system.

Although all PIPKI γ splicing variants produce the same product, PtdIns(4,5)P₂, they play distinct biological functions by interacting with specific effector proteins (39). This work identified NEDD4-1 as a novel specific effector binding with PIPKI γ 5 (Fig. 6). Consistently, PIPKI γ 5 specifically controls Mig6 expression; other PIPKI γ splicing variants such as PIPKI γ 2 do not play the similar role (Fig. 6C). PIPKI γ 5 and PIPKI γ 2 may differentially modulate cancer progression. PIPKI γ 5 down-regulates EGFR signaling and can function as a tumor suppressor, whereas PIPKI γ 2 promotes anchorage-independent cancer growth, cancer cell migration, and invasion (22, 40–43). NEDD4-1 is an oncoprotein and has been found to be frequently overexpressed in many types of human cancers such as non-small cell lung cancer (44), gastric carcinomas (45), and breast cancer (46). Suppression of NEDD4-1 expression can significantly reduce tumor cell growth (47, 48). Our research indicates that NEDD4-1 mediates degradation of the tumor suppressor Mig6 (Fig. 4 and 5), which provides a novel mechanism for NEDD4-1 promotion of tumor progression. This work also identified PIPKI γ 5 as a suppressor of NEDD4-1 function.

PIPKI γ 5 is required for EGFR sorting from the limiting membrane of endosomes into intraluminal vesicles (ILVs) of the multivesicular bodies, which is a key step for EGFR sorting to lysosomes for degradation (23). Hrs is a key component of the endosomal sorting complexes required for transport (ESCRT)-0 (49), which binds to ubiquitinated EGFR and recruits additional ESCRT components to mediate EGFR sorting into ILVs (50). PIPKI γ 5 facilitates Hrs interaction with Sorting Nexin 5 (SNX5), which blocks NEDD4-1-mediated Hrs ubiquitination and promotes Hrs association with EGFR to mediate EGFR sorting into ILV (23). Here we found that PIPKI γ 5 directly interacts with NEDD4-1 to modulate Mig6 ubiquitination and degradation. As such, the ubiquitination events are critical for PIPKI γ 5 in modulating EGFR. PIPKI γ 2 can be ubiquitinated by E3 ligase HECT domain E3 ubiquitin protein ligase 1 (HECTD1) in the N terminus at lysine 97 and resulted in PIPKI γ 2 degradation (51). Blocking PIPKI γ 2 ubiquitination by mutating lysine 97 to arginine perturbs focal adhesion assembly and cancer cell migration (51). PIPKI γ 5 has exactly the same N terminus and lysine 97 as PIPKI γ 2. Therefore, PIPKI γ 5 itself has the potential to be modified by ubiquitination, and this can modulate its function.

EGFR mutations have been identified in lung cancer, including the point mutation L858R and an in-frame short deletion mutation, del746–750 (52–55). These mutations render EGFR with constitutive kinase activity, and they are deficient for being sorted to the lysosomes for degradation (56, 57). Mig6 still interacts with these EGFR mutants and plays essential roles in down-regulating signaling mediated by them in cancer. Loss of Mig6 dramatically accelerates the initiation and progression of EGFR L858R-driven lung adenocarcinomas (18). By modulating Mig6 expression, PIPKI γ 5 can regulate EGFR mutant-initiated cancer progression.

Mig6 not only interacts with EGFR but also interacts with other members of ErbB family receptors such as ErbB2 (7).

PIPK1 γ 5 Controls Mig6 Ubiquitination and Degradation




FIGURE 7. Identification of the regions responsible for PIPK1 γ 5-NEDD4-1 interaction. *A*, schematic of a series of PIPK1 γ 5 C terminus truncation mutants. *B*, V5-NEDD4-1 was co-transfected with HA-tagged wild type PIPK1 γ 5 (wt) or a series of PIPK1 γ 5 C terminus truncation mutants into HEK-293T cells, subjected to immunoprecipitation with anti-V5 antibody, and then immunoblotted (IB) with antibodies as indicated. *C*, schematic of the domains of NEDD4-1 and a series of NEDD4-1 deletion mutants. *D*, HA-PIPK1 γ 5 was co-transfected with V5-tagged wild-type NEDD4-1 (wt) or a series of NEDD4-1 deletion mutants into HEK-293T cells, subjected to immunoprecipitation with anti-V5 antibody, and then immunoblotted with antibodies as indicated. *E*, the amount of PIPK1 γ 5 co-immunoprecipitated with wild-type NEDD4-1 or NEDD4-1 deletion mutants was quantified. Error bars indicate mean \pm S.E. from three independent experiments. *, $p < 0.05$.

Thus, Mig6 suppresses the signaling from ErbB2 (29). As an adaptor protein, Mig6 also interacts and modulates the function of many important components of different signaling pathways, including cdc42, $\text{I}\kappa\text{B}\alpha$, GRB2, Src, PI3K p85, and 14-3-3 ζ (7, 58–60). This suggests that PIPK1 γ 5 can function in multiple signaling pathways by controlling the expression of Mig6.

Experimental Procedures

Cell Cultures and Transfection—The breast cancer cell line MDA-MB-231 was cultured using DMEM supplemented with 10% FBS. The lung cancer cell line H-1975 was cultured using 1640 medium supplemented with 10% FBS. For plasmid transfection, cells were transfected using Lipofectamine 2000 (Invitrogen) following the instructions of the manufacturer. For siRNA transfection, cells were transfected with Oligofectamine (Invitrogen) for 72 h following the instructions of the manufacturer.

Reagents—TGF α (catalog no. GF022) was purchased from EMD Millipore (Billerica, MA). Antibodies to EGFR (catalog no. sc-03), Mig6 (catalog no. sc-137154), Myc-tag (catalog no. sc-40), NEDD4-2 (catalog no. sc-514954), and GAPDH (catalog no. sc-32233) were from Santa Cruz Biotechnology (Santa Cruz, CA). Antibodies to NEDD4-1 (catalog no. 2740), total AKT (catalog no. 4685), phosphorylated AKT (catalog no.

4060), and phosphorylated EGFR (catalog no. 3777) were from Cell Signaling Technology (Danvers, MA). Anti-HA antibody (catalog no. 901502) was from BioLegend (San Diego, CA). Anti-FLAG antibody (catalog no. F3165) was from Sigma-Aldrich (St. Louis, MO). Anti-GFP (catalog no. 11814460001) antibody was from Roche Life Science (Indianapolis, IN). Anti-PIP1 γ 2- and -i5-specific antibodies were generated as described previously (21). Secondary antibodies were obtained from Jackson ImmunoResearch Laboratories (West Grove, PA).

Constructs—PIP1 γ 5 (GenBank accession no. NM_001300849.1) was amplified by PCR, ligated into the pcDNA3.0 vector (Invitrogen) with an HA tag at the N terminus, and subcloned into the PWPT lentivirus vector (Addgene) through MluI/SalI sites. Mig6 (GenBank accession no. NM_018948.3) was subcloned into the pcDNA3.0 vector via BamHI/XhoI sites. PIPK1 γ 5 C terminus truncation mutants were generated by regular PCR, and NEDD4-1 truncation mutants were generated using PCR primer overlap extension with primers containing the desired mutations. To generate V5-tag-NEDD4-1, NEDD4-1 (GenBank accession no. NM_006154.3) was amplified by PCR and inserted into the pcDNA3.1 vector (Invitrogen) through BamHI/XhoI sites. For expression in *Escherichia coli*, PIPK1 γ 5 was subcloned into pET28c(+) (Novagen), Mig6 was subcloned




FIGURE 8. PIPKI γ 5 blocks NEDD4-1-mediated Mig6 ubiquitination. *A*, the ubiquitination reaction of GST-Mig6 was carried out in the presence of E1, UbcH5b, NEDD4-1, ubiquitin, and ATP with or without purified wild-type His₆-PIPKI γ 5 or kinase dead mutant His₆-PIPKI γ 5 *in vitro*, and the whole ubiquitination level was detected by immunoblotting. *B*, Myc-Mig6 was co-transfected with His₆-ubiquitin, NEDD4-1, wild-type HA-PIPKI γ 5, or kinase dead mutant PIPKI γ 5 into HEK-293T cells as indicated, and Myc-Mig6 ubiquitination was detected by an *in vivo* ubiquitination assay. *IB*, immunoblot. *C*, GFP, HA-tagged wild-type PIPKI γ 5 or kinase-dead mutant PIPKI γ 5 was constructed into PWPT lentivirus vectors. GFP, HA-PIPKI γ 5 WT, or HA-PIPKI γ 5 KD was then expressed via lentivirus transfection in MDA-MB-231 cells, and then cells were subjected to immunoprecipitation with Mig6 antibody. The amount of endogenous NEDD4-1 co-immunoprecipitated with endogenous Mig6 was detected by immunoblotting. *D*, quantification of the amount of endogenous NEDD4-1 co-immunoprecipitated with endogenous Mig6 as in *C*. Error bars indicate mean \pm S.E. from three independent experiments. *, $p < 0.05$.

into pET52b(+) (Novagen) or pGEX 5x-2 (GE Healthcare), and NEDD4-1 were subcloned into pGEX 5x-2.

Recombinant Protein Purification—*E. coli* strain BL21 (DE3) was transformed with pET28c-PIPKI γ 5 (His₆-tagged), pET52b-Mig6 (His₆-tagged), pGEX 5x-2-Mig6 (GST-tagged) or pGEX 5x-2-NEDD4-1 (GST-tagged), respectively. Single clones were picked, and 0.1 mM isopropyl β -D-thiogalactopyranoside was added when A₆₀₀ reached 0.4–0.6. The culture was then further incubated at 30 °C overnight. The cells were harvested and sonicated in buffer (50 mM Tris-HCl, 150 mM NaCl, 1% Nonidet P-40, and 1 mM DTT). The GST-tagged proteins were purified by glutathione Sepharose™ 4B resin (GE Healthcare). The His₆-tagged proteins were purified by Ni²⁺-NTA-agarose resin (Thermo Fisher Scientific).

siRNA—The sequence of control scrambled siRNA was 5'-GUACCUGUACUUAUCAGCAG-3'. The siRNA sequence for human PIPKI γ 5 was 5'-GGAUGGGAGGUACUGGAU-3'. The siRNA sequence specific targeting human PIPKI γ 2 was 5'-GAGCGACACAUAUUUCUA-3'. The siRNA sequence for Smurf1 was 5'-GCAUCGAAGUGUCCAGAGAAG-3'. The siRNA sequence for Smurf2 was 5'-CCUUCUGUGUUGAACAUAA-3'. The NEDD4-2 siRNA sequence was 5'-AAC-

CACAACACAAAGUCACAC-3'. The NEDD4-1 siRNA1 sequence was 5'-UCCAUGAAUCUAGAAGAACA-3'. The NEDD4-1 siRNA2 sequence was 5'-UAGAGCCUGGCUGGUUGUUU-3'. The NEDD4-1 siRNA3 sequence was 5'-CAGCAUCGAGCUCAAAUCA-3'. Mig6 siRNA was purchased from Santa Cruz Biotechnology (catalog no. sc-45704).

CRISPR-cas9 Genome Editing—Genomic disruptions were created in MDA-MB-231 and H-1975 cells using the CRISPR/Cas9 system as described previously (61). The lentiCRISPR v2 plasmid (Addgene, plasmid no. 52961) was used to mediate the expression of the single guide RNA (sgRNA) and cas9 protein by lentivirus infection. The sgRNAs targeting the PIPKI γ 5-specific sequence were designed using online software. Two different sgRNA-targeting sequences were chosen: 5'-TCA-CGGCCCCGAGTCGGCGA-3' and 5'-GCGGTCACTTACAGTCCCCG-3'. Lentivirus-infected cells were cultured in medium containing 1.0 μ g/ml puromycin for 2 days for selection. Surviving cells were then trypsinized, and a single cell was isolated by limited dilution of the selected cells in 96-well plates. Each single clone was passaged and expanded. The total DNA of each colony was extracted and genotyped. The genomic region surrounding the CRISPR/Cas9 target site for

PIPK1 γ 5 Controls Mig6 Ubiquitination and Degradation




FIGURE 9. PtdIns(4,5)P₂ binds NEDD4-1 and decreases NEDD4-1-mediated ubiquitination. Wild-type GST-NEDD4-1 and C2 domain-deletion mutant GST-NEDD4-1 (*Del C2*) were purified from *E. coli*. **A**, binding of GST-NEDD4-1 (WT) or GST-NEDD4-1 (*Del C2*) with different lipids was detected by protein-lipid overlay assay with PIP strips. *PI*, phosphatidylinositol. **B** and **C**, the binding of GST-NEDD4-1 (WT) with PtdIns(4,5)P₂-containing or non-containing liposomes (*control*) was detected (**B**) and quantified (**C**). **D** and **E**, the binding of GST-NEDD4-1 with PtdIns(4,5)P₂-containing or non-containing liposomes was detected (**D**) and quantified (**E**). **F**, the ubiquitination reaction of GST-Mig6 was carried out in the presence of E1, UbcH5b, NEDD4-1, ubiquitin, and ATP with or without PtdIns(4,5)P₂-containing or non-containing liposomes *in vitro*, and the whole ubiquitination level was detected by immunoblotting. *LPA*, lysophosphatidic acid; *LPC*, lysophosphocholine; *SIP*, sphingosine 1-phosphate.




FIGURE 10. Model for PIPK1 γ 5 regulation of Mig6 expression and EGFR signaling. By directly interacting with NEDD4-1 and producing PtdIns(4,5)P₂, PIPK1 γ 5 blocks NEDD4-1-mediated Mig6 ubiquitination. This protects Mig6 from being degraded by proteasomes. As a result, increased Mig6 protein levels inhibit EGFR activation and signaling. Loss of PIPK1 γ 5 promotes NEDD4-1-mediated Mig6 ubiquitination and the subsequent Mig6 degradation by proteasomes. This enhances EGFR signaling.

PIPK1 γ 5 was PCR-amplified. The amplicons were cloned into the pCRTM 2.1 Vector (Invitrogen) and validated by sequencing.

RNA Extraction and Real-time PCR—MDA-MB-231 cells were harvested, and total RNA was extracted using the

ISOLATE RNA mini kit (Bioline, catalog no. BIO-52072). 2 μ g of total RNA was used for cDNA reverse transcription with the kit (Applied Biosystems, catalog no. 4368814). SYBR Green master mix (Applied Biosystems, catalog no. A25742) was used for real-time PCR, and the results were analyzed by QUANT-STUDIO 3 software. The Mig6 mRNA abundance was normalized to the expression of GAPDH. Primers used for the PCR were as follows: 5'-CTACTGGAGCAGTCGCAGTG-3' (forward) and 5'-CCTCTTCATGTGGTCCCAAG-3' (reverse) for Mig6 and 5'-AATCCCATCACCATCTTCCA-3' (forward) and 5'-TGGACTCCACGACGTACTCA-3' (reverse) for GAPDH.

Immunoprecipitation and Immunoblotting—Immunoprecipitation was performed as described previously (62). Briefly, cells were harvested and lysed in 25 mM HEPES (pH 7.2), 150 mM NaCl, 0.5% Nonidet P-40, 1 mM MgCl₂, and protease inhibitor mixture and then centrifuged and incubated with protein G-Sepharose and 2 μ g of antibody as indicated at 4 °C for 4 h. The immunocomplexes were separated by SDS-PAGE and analyzed as indicated.

In Vivo Ubiquitination Assay—The ubiquitination of Mig6 was evaluated as described previously (63). His₆-ubiquitin-conjugated Mig6 in MDA-MB-231 cells was purified by Ni²⁺-NTA beads (Qiagen). MDA-MB-231 cell was lysed in IP buffer (25 mM HEPES (pH 7.2), 150 mM NaCl, 0.5% Nonidet P-40, 1 mM MgCl₂, and protease inhibitor mixture) and incubated with Ni²⁺-NTA beads for 2 h at 4 °C. The beads were washed with IP buffer, buffer A (8 M urea, 0.1 M Na₂PO₄/NaH₂PO₄, 0.01 M Tris-HCl (pH 8.0), and 10 mM β -mercaptoethanol), and buffer B (8 M urea, 0.1 M Na₂PO₄/NaH₂PO₄, 0.01 M Tris-HCl (pH 5.3),

and 10 mM β -mercaptoethanol), and bound proteins were eluted with buffer C (200 mM imidazole, 0.15 M Tris-HCl (pH 6.7), 30% glycerol, 0.72 M β -mercaptoethanol, and 5% SDS). The eluted proteins were analyzed by Western blotting for the presence of His₆-ubiquitin-conjugated Mig6 by anti-Mig6 antibody.

In Vitro E3 Ubiquitin Ligase Activity Assay—The E3 ubiquitin ligase activity assay was based on a method described in previous studies (64, 65) with minor modifications. The ubiquitination reaction mixture contained 0.2 μ M E1, 1 μ M UBCH5C (E2), 0.5 μ M NEDD4-1, and 10 μ M ubiquitin in buffer (50 mM Tris-HCl (pH 7.5), 2 mM ATP, and 5 mM MgCl₂). After incubation at 30 °C for 1 h, the reaction was terminated by SDS loading sample buffer. Mig6 poly-ubiquitination chains were prepared as above. To isolate Mig6 from the mixture, SDS was added to the reaction mixture to a final concentration of 1%, and the sample was incubated at 90 °C for 10 min. After a 10-fold dilution of SDS with buffer (50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1% Nonidet P-40, and 10% glycerol), Mig6 antibody (1.0 μ g, Santa Cruz Biotechnology) and protein G-Sepharose beads (5 μ l) were added. The mixture was incubated for 4 h at room temperature and spun at 1000 \times g to collect protein G-Sepharose beads. The beads were washed three times with buffer plus 0.1% SDS, and then proteins were eluted with SDS loading sample buffer and detected by immunoblotting with ubiquitin antibody or Mig6 antibody.

Protein-Lipid Overlay Assay—PIP strips (catalog no. P-6001, Echelon) are 2 \times 6 cm hydrophobic membranes that have been spotted with 100 picomoles of all eight phosphoinositides and seven other biological important lipids. PIP strips were first blocked with PBS including 3% BSA for 1 h and then incubated with GST-NEDD4-1 (WT) or GST-NEDD4-1 (Del C2) (1 μ g/ml) in PBS with 3% BSA. Subsequently, the PIP strips were washed extensively in PBS containing 0.1% Tween 20, and the bound protein was detected with anti-GST antibody by immunoblotting.

Liposome Binding Assay—Liposome binding assays were carried out essentially as described previously (66, 67). PtdIns(4,5)P₂-containing PolyPIPosomes (catalog no. Y-P045, Echelon) are composed of 65% phosphatidylcholine, 29% phosphatidylethanolamine (PE), 1% biotin-PE, and 5% PtdIns(4,5)P₂. Control PolyPIPosomes (catalog no. Y-P000, Echelon) are composed of 65% phosphatidylcholine, 29% PE, 1% biotin-PE, and 5% phosphatidylinositol. 15 μ mol of PolyPIPosomes were diluted in 100 μ l of binding buffer (50 mM Tris (pH 7.8), 150 mM NaCl, 0.05% Nonidet P-40 (v/v), and 20 mM iodoacetamide) and incubated for 15 min at room temperature with 10 μ g of GST-NEDD4-1 (WT) or GST-NEDD4-1 (Del C2). Liposome-protein complexes were recovered by 10-min centrifugation at room temperature (11,356 \times g), and pellets were resuspended in 500 μ l of binding buffer and further incubated with streptavidin-agarose (Invitrogen) for 1 h at room temperature. Biotinylated liposomes bound to GST proteins were washed with binding buffer and eluted with Laemmli sample buffer, separated by SDS-PAGE, and immunoblotted with anti-GST antibody.

Statistics—All data analyses were performed using SigmaPlot. Bar graphs represent mean \pm S.E. as indicated. Statistical significance was assessed using Student's *t* test.

Author Contributions—M. S. conducted most of the experiments and analyzed the results. J. C. conducted the CRISPR-cas9 genome editing and selected single clones. R. A. A. provided assistance with the preparation of the manuscript. Y. S. conceived the idea for the project, designed the experiments, and wrote the paper with M. S.

Acknowledgments—We thank Prof. W. A. Yeudall for the HN31 cells. We also thank Prof. X. Jiang for the gift of the NEDD4-1 construct. We thank Prof. I. Morgan for discussions of the project.

References

- Schlessinger, J. (2002) Ligand-induced, receptor-mediated dimerization and activation of EGF receptor. *Cell* **110**, 669–672
- Hynes, N. E., and MacDonald, G. (2009) ErbB receptors and signaling pathways in cancer. *Curr. Opin. Cell Biol.* **21**, 177–184
- Casalini, P., Iorio, M. V., Galmozzi, E., and Ménard, S. (2004) Role of HER receptors family in development and differentiation. *J. Cell. Physiol.* **200**, 343–350
- Yarden, Y., and Pines, G. (2012) The ERBB network: at last, cancer therapy meets systems biology. *Nat. Rev. Cancer* **12**, 553–563
- Chong, C. R., and Jänne, P. A. (2013) The quest to overcome resistance to EGFR-targeted therapies in cancer. *Nat. Med.* **19**, 1389–1400
- Zhang, X., Pickin, K. A., Bose, R., Jura, N., Cole, P. A., and Kuriyan, J. (2007) Inhibition of the EGF receptor by binding of MIG6 to an activating kinase domain interface. *Nature* **450**, 741–744
- Fiorentino, L., Pertica, C., Fiorini, M., Talora, C., Crescenzi, M., Castellani, L., Alemà, S., Benedetti, P., and Segatto, O. (2000) Inhibition of ErbB-2 mitogenic and transforming activity by RALT, a mitogen-induced signal transducer which binds to the ErbB-2 kinase domain. *Mol. Cell. Biol.* **20**, 7735–7750
- Frosi, Y., Anastasi, S., Ballarò, C., Varsano, G., Castellani, L., Maspero, E., Polo, S., Alemà, S., and Segatto, O. (2010) A two-tiered mechanism of EGFR inhibition by RALT/MIG6 via kinase suppression and receptor degradation. *J. Cell Biol.* **189**, 557–571
- Zhang, X., Gureasko, J., Shen, K., Cole, P. A., and Kuriyan, J. (2006) An allosteric mechanism for activation of the kinase domain of epidermal growth factor receptor. *Cell* **125**, 1137–1149
- Ying, H., Zheng, H., Scott, K., Wiedemeyer, R., Yan, H., Lim, C., Huang, J., Dhakal, S., Ivanova, E., Xiao, Y., Zhang, H., Hu, J., Stommel, J. M., Lee, M. A., Chen, A. J., et al. (2010) Mig-6 controls EGFR trafficking and suppresses gliomagenesis. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 6912–6917
- Sorkin, A., and Goh, L. K. (2009) Endocytosis and intracellular trafficking of ErbBs. *Exp. Cell Res.* **315**, 683–696
- Carpenter, G., and Liao, H. J. (2009) Trafficking of receptor tyrosine kinases to the nucleus. *Exp. Cell Res.* **315**, 1556–1566
- Wick, M., Bürger, C., Funk, M., and Müller, R. (1995) Identification of a novel mitogen-inducible gene (mig-6): regulation during G₁ progression and differentiation. *Exp. Cell Res.* **219**, 527–535
- Ferby, I., Reschke, M., Kudlacek, O., Knyazev, P., Pantè, G., Amann, K., Sommergruber, W., Kraut, N., Ullrich, A., Fässler, R., and Klein, R. (2006) Mig6 is a negative regulator of EGF receptor-mediated skin morphogenesis and tumor formation. *Nat. Med.* **12**, 568–573
- Xu, D., Makkinje, A., and Kyriakis, J. M. (2005) Gene 33 is an endogenous inhibitor of epidermal growth factor (EGF) receptor signaling and mediates dexamethasone-induced suppression of EGF function. *J. Biol. Chem.* **280**, 2924–2933
- Segatto, O., Anastasi, S., and Alemà, S. (2011) Regulation of epidermal growth factor receptor signalling by inducible feedback inhibitors. *J. Cell Sci.* **124**, 1785–1793
- Anastasi, S., Sala, G., Huiping, C., Caprini, E., Russo, G., Iacovelli, S., Lucini, F., Ingvarsson, S., and Segatto, O. (2005) Loss of RALT/MIG-6 ex-

PIPK1 γ 5 Controls Mig6 Ubiquitination and Degradation

- pression in ERBB2-amplified breast carcinomas enhances ErbB-2 oncogenic potency and favors resistance to Herceptin. *Oncogene* **24**, 4540–4548
18. Maity, T. K., Venugopalan, A., Linnoila, I., Cultraro, C. M., Giannakou, A., Nemati, R., Zhang, X., Webster, J. D., Ritt, D., Ghosal, S., Hoschuetzky, H., Simpson, R. M., Biswas, R., Politi, K., Morrison, D. K., *et al.* (2015) Loss of MIG6 accelerates initiation and progression of mutant epidermal growth factor receptor-driven lung adenocarcinoma. *Cancer Discov.* **5**, 534–549
 19. Heck, J. N., Mellman, D. L., Ling, K., Sun, Y., Wagoner, M. P., Schill, N. J., and Anderson, R. A. (2007) A conspicuous connection: structure defines function for the phosphatidylinositol-phosphate kinase family. *Crit. Rev. Biochem. Mol. Biol.* **42**, 15–39
 20. Xia, Y., Irvine, R. F., and Giudici, M. L. (2011) Phosphatidylinositol 4-phosphate 5-kinase I γ _v6, a new splice variant found in rodents and humans. *Biochem. Biophys. Res. Commun.* **411**, 416–420
 21. Schill, N. J., and Anderson, R. A. (2009) Two novel phosphatidylinositol-4-phosphate 5-kinase type I γ splice variants expressed in human cells display distinctive cellular targeting. *Biochem. J.* **422**, 473–482
 22. Sun, Y., Ling, K., Wagoner, M. P., and Anderson, R. A. (2007) Type I γ phosphatidylinositol phosphate kinase is required for EGF-stimulated directional cell migration. *J. Cell Biol.* **178**, 297–308
 23. Sun, Y., Hedman, A. C., Tan, X., Schill, N. J., and Anderson, R. A. (2013) Endosomal type I γ PIP 5-kinase controls EGF receptor lysosomal sorting. *Dev. Cell* **25**, 144–155
 24. Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., and Zhang, F. (2013) Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**, 2281–2308
 25. Shen, B., Zhang, W., Zhang, J., Zhou, J., Wang, J., Chen, L., Wang, L., Hodgkins, A., Iyer, V., Huang, X., and Skarnes, W. C. (2014) Efficient genome modification by CRISPR-Cas9 nickase with minimal off-target effects. *Nat. Methods* **11**, 399–402
 26. Tanida, I., Minematsu-Ikeguchi, N., Ueno, T., and Kominami, E. (2005) Lysosomal turnover, but not a cellular level, of endogenous LC3 is a marker for autophagy. *Autophagy* **1**, 84–91
 27. Yang, Y. L., Liu, Y. S., Chuang, L. Y., Guh, J. Y., Lee, T. C., Liao, T. N., Hung, M. Y., and Chiang, T. A. (2009) Bone morphogenetic protein-2 antagonizes renal interstitial fibrosis by promoting catabolism of type I transforming growth factor- β receptors. *Endocrinology* **150**, 727–740
 28. Nie, J., Wu, M., Wang, J., Xing, G., He, F., and Zhang, L. (2010) REG γ proteasome mediates degradation of the ubiquitin ligase Smurf1. *FEBS Lett.* **584**, 3021–3027
 29. Fiorini, M., Ballarò, C., Sala, G., Falcone, G., Alemà, S., and Segatto, O. (2002) Expression of RALT, a feedback inhibitor of ErbB receptors, is subjected to an integrated transcriptional and post-translational control. *Oncogene* **21**, 6530–6539
 30. Yang, B., and Kumar, S. (2010) Nedd4 and Nedd4-2: closely related ubiquitin-protein ligases with distinct physiological functions. *Cell Death Differ.* **17**, 68–77
 31. Dunn, R., Klos, D. A., Adler, A. S., and Hicke, L. (2004) The C2 domain of the Rsp5 ubiquitin ligase binds membrane phosphoinositides and directs ubiquitination of endosomal cargo. *J. Cell Biol.* **165**, 135–144
 32. Balla, T. (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol. Rev.* **93**, 1019–1137
 33. Bunney, T. D., and Katan, M. (2010) Phosphoinositide signalling in cancer: beyond PI3K and PTEN. *Nat. Rev. Cancer* **10**, 342–352
 34. Yuan, T. L., and Cantley, L. C. (2008) PI3K pathway alterations in cancer: variations on a theme. *Oncogene* **27**, 5497–5510
 35. Samuels, Y., Wang, Z., Bardelli, A., Silliman, N., Ptak, J., Szabo, S., Yan, H., Gazdar, A., Powell, S. M., Riggins, G. J., Willson, J. K., Markowitz, S., Kinzler, K. W., Vogelstein, B., and Velculescu, V. E. (2004) High frequency of mutations of the PIK3CA gene in human cancers. *Science* **304**, 554
 36. Emerling, B. M., Hurov, J. B., Poulgiannis, G., Tsukazawa, K. S., Choo-Wing, R., Wulf, G. M., Bell, E. L., Shim, H. S., Lamia, K. A., Rameh, L. E., Bellinger, G., Sasaki, A. T., Asara, J. M., Yuan, X., Bullock, A., *et al.* (2013) Depletion of a putatively druggable class of phosphatidylinositol kinases inhibits growth of p53-null tumors. *Cell* **155**, 844–857
 37. Mendelsohn, J., and Baselga, J. (2006) Epidermal growth factor receptor targeting in cancer. *Semin. Oncol.* **33**, 369–385
 38. Kittler, R., Pelletier, L., Ma, C., Poser, I., Fischer, S., Hyman, A. A., and Buchholz, F. (2005) RNA interference rescue by bacterial artificial chromosome transgenesis in mammalian tissue culture cells. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 2396–2401
 39. Sun, Y., Thapa, N., Hedman, A. C., and Anderson, R. A. (2013) Phosphatidylinositol 4,5-bisphosphate: targeted production and signaling. *BioEssays* **35**, 513–522
 40. Sun, Y., Turbin, D. A., Ling, K., Thapa, N., Leung, S., Huntsman, D. G., and Anderson, R. A. (2010) Type I γ phosphatidylinositol phosphate kinase modulates invasion and proliferation and its expression correlates with poor prognosis in breast cancer. *Breast Cancer Res.* **12**, R6
 41. Thapa, N., Choi, S., Tan, X., Wise, T., and Anderson, R. A. (2015) Phosphatidylinositol phosphate 5-kinase I γ and phosphoinositide 3-kinase/Akt signaling couple to promote oncogenic growth. *J. Biol. Chem.* **290**, 18843–18854
 42. Thapa, N., Choi, S., Hedman, A., Tan, X., and Anderson, R. A. (2013) Phosphatidylinositol phosphate 5-kinase I γ 2 in association with Src controls anchorage-independent growth of tumor cells. *J. Biol. Chem.* **288**, 34707–34718
 43. Chen, C., Wang, X., Xiong, X., Liu, Q., Huang, Y., Xu, Q., Hu, J., Ge, G., and Ling, K. (2015) Targeting type I γ phosphatidylinositol phosphate kinase inhibits breast cancer metastasis. *Oncogene* **34**, 4635–4646
 44. Amodio, N., Scrima, M., Palaia, L., Salman, A. N., Quintiero, A., Franco, R., Botti, G., Pirozzi, P., Rocco, G., De Rosa, N., and Viglietto, G. (2010) Oncogenic role of the E3 ubiquitin ligase NEDD4-1, a PTEN negative regulator, in non-small-cell lung carcinomas. *Am. J. Pathol.* **177**, 2622–2634
 45. Sun, A., Yu, G., Dou, X., Yan, X., Yang, W., and Lin, Q. (2014) Nedd4-1 is an exceptional prognostic biomarker for gastric cardia adenocarcinoma and functionally associated with metastasis. *Mol. Cancer* **13**, 248
 46. Jung, S., Li, C., Jeong, D., Lee, S., Ohk, J., Park, M., Han, S., Duan, J., Kim, C., Yang, Y., Kim, K. I., Lim, J. S., Kang, Y. S., and Lee, M. S. (2013) Oncogenic function of p34SEI-1 via NEDD4-mediated PTEN ubiquitination/degradation and activation of the PI3K/AKT pathway. *Int. J. Oncol.* **43**, 1587–1595
 47. Wang, X., Trotman, L. C., Koppie, T., Alimonti, A., Chen, Z., Gao, Z., Wang, J., Erdjument-Bromage, H., Tempst, P., Cordon-Cardo, C., Pandolfi, P. P., and Jiang, X. (2007) NEDD4-1 is a proto-oncogenic ubiquitin ligase for PTEN. *Cell* **128**, 129–139
 48. Zou, X., Levy-Cohen, G., and Blank, M. (2015) Molecular functions of NEDD4 E3 ubiquitin ligases in cancer. *Biochim. Biophys. Acta* **1856**, 91–106
 49. Henne, W. M., Buchkovich, N. J., and Emr, S. D. (2011) The ESCRT pathway. *Dev. Cell* **21**, 77–91
 50. Eden, E. R., White, I. J., and Futter, C. E. (2009) Down-regulation of epidermal growth factor receptor signalling within multivesicular bodies. *Biochem. Soc. Trans.* **37**, 173–177
 51. Li, X., Zhou, Q., Sunkara, M., Kutys, M. L., Wu, Z., Rychahou, P., Morris, A. J., Zhu, H., Evers, B. M., and Huang, C. (2013) Ubiquitylation of phosphatidylinositol 4-phosphate 5-kinase type I γ by HECTD1 regulates focal adhesion dynamics and cell migration. *J. Cell Sci.* **126**, 2617–2628
 52. Eberhard, D. A., Johnson, B. E., Amler, L. C., Goddard, A. D., Heldens, S. L., Herbst, R. S., Ince, W. L., Jänne, P. A., Januario, T., Johnson, D. H., Klein, P., Miller, V. A., Ostland, M. A., Ramies, D. A., Sebanovic, D., *et al.* (2005) Mutations in the epidermal growth factor receptor and in KRAS are predictive and prognostic indicators in patients with non-small-cell lung cancer treated with chemotherapy alone and in combination with erlotinib. *J. Clin. Oncol.* **23**, 5900–5909
 53. Paez, J. G., Jänne, P. A., Lee, J. C., Tracy, S., Greulich, H., Gabriel, S., Herman, P., Kaye, F. J., Lindeman, N., Boggon, T. J., Naoki, K., Sasaki, H., Fujii, Y., Eck, M. J., Sellers, W. R., *et al.* (2004) EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* **304**, 1497–1500
 54. Sakurada, A., Shepherd, F. A., and Tsao, M. S. (2006) Epidermal growth factor receptor tyrosine kinase inhibitors in lung cancer: impact of primary or secondary mutations. *Clin. Lung Cancer* **7**, S138–144
 55. Siegelin, M. D., and Borczuk, A. C. (2014) Epidermal growth factor receptor mutations in lung adenocarcinoma. *Lab. Invest.* **94**, 129–137

56. Grandal, M. V., Zandi, R., Pedersen, M. W., Willumsen, B. M., van Deurs, B., and Poulsen, H. S. (2007) EGFRvIII escapes down-regulation due to impaired internalization and sorting to lysosomes. *Carcinogenesis* **28**, 1408–1417
57. Shtiegman, K., Kochupurakkal, B. S., Zwang, Y., Pines, G., Starr, A., Vexler, A., Citri, A., Katz, M., Lavi, S., Ben-Basat, Y., Benjamin, S., Corso, S., Gan, J., Yosef, R. B., Giordano, S., and Yarden, Y. (2007) Defective ubiquitinylation of EGFR mutants of lung cancer confers prolonged signaling. *Oncogene* **26**, 6968–6978
58. Makkinje, A., Quinn, D. A., Chen, A., Cadilla, C. L., Force, T., Bonventre, J. V., and Kyriakis, J. M. (2000) Gene 33/Mig-6, a transcriptionally inducible adapter protein that binds GTP-Cdc42 and activates SAPK/JNK. A potential marker transcript for chronic pathologic conditions, such as diabetic nephropathy: possible role in the response to persistent stress. *J. Biol. Chem.* **275**, 17838–17847
59. Tsunoda, T., Inokuchi, J., Baba, I., Okumura, K., Naito, S., Sasazuki, T., and Shirasawa, S. (2002) A novel mechanism of nuclear factor κ B activation through the binding between inhibitor of nuclear factor- κ B α and the processed NH(2)-terminal region of Mig-6. *Cancer Res.* **62**, 5668–5671
60. Pante, G., Thompson, J., Lamballe, F., Iwata, T., Ferby, I., Barr, F. A., Davies, A. M., Maina, F., and Klein, R. (2005) Mitogen-inducible gene 6 is an endogenous inhibitor of HGF/Met-induced cell migration and neurite growth. *J. Cell Biol.* **171**, 337–348
61. Sanjana, N. E., Shalem, O., and Zhang, F. (2014) Improved vectors and genome-wide libraries for CRISPR screening. *Nat. Methods* **11**, 783–784
62. Ling, K., Doughman, R. L., Iyer, V. V., Firestone, A. J., Bairstow, S. F., Mosher, D. F., Schaller, M. D., and Anderson, R. A. (2003) Tyrosine phosphorylation of type I γ phosphatidylinositol phosphate kinase by Src regulates an integrin-talin switch. *J. Cell Biol.* **163**, 1339–1349
63. Pan, Y., and Chen, J. (2003) MDM2 promotes ubiquitination and degradation of MDMX. *Mol. Cell. Biol.* **23**, 5113–5121
64. Wang, J., Peng, Q., Lin, Q., Childress, C., Carey, D., and Yang, W. (2010) Calcium activates Nedd4 E3 ubiquitin ligases by releasing the C2 domain-mediated auto-inhibition. *J. Biol. Chem.* **285**, 12279–12288
65. Xia, Z. P., Sun, L., Chen, X., Pineda, G., Jiang, X., Adhikari, A., Zeng, W., and Chen, Z. J. (2009) Direct activation of protein kinases by unanchored polyubiquitin chains. *Nature* **461**, 114–119
66. Elkin, S. K., Ivanov, D., Ewalt, M., Ferguson, C. G., Hyberts, S. G., Sun, Z. Y., Prestwich, G. D., Yuan, J., Wagner, G., Oettinger, M. A., and Gozani, O. P. (2005) A PHD finger motif in the C terminus of RAG2 modulates recombination activity. *J. Biol. Chem.* **280**, 28701–28710
67. Naslavsky, N., Rahajeng, J., Chenavas, S., Sorgen, P. L., and Caplan, S. (2007) EHD1 and Eps15 interact with phosphatidylinositols via their Eps15 homology domains. *J. Biol. Chem.* **282**, 16612–16622