

# Endothelin and Angiotensin II Stimulation of $\text{Na}^+\text{-H}^+$ Exchange Is Impaired in Cardiac Hypertrophy

Nobuhiko Ito,\* Yutaka Kagaya,\* Ellen O. Weinberg,\* William H. Barry,† and Beverly H. Lorell\*

\*The Charles A. Dana Research Institute and the Harvard-Thorndike Laboratory of Beth Israel Hospital and the Department of Medicine, Cardiovascular Division, Beth Israel Hospital, and Harvard Medical School, Boston, Massachusetts, 02215; and †Division of Cardiology, Department of Medicine, University of Utah School of Medicine, Salt Lake City, Utah 84132

## Abstract

We compared the effects of endothelin-1 (ET-1) on intracellular pH, intracellular  $[\text{Ca}^{2+}]_i$ , and cell contraction in hypertrophied adult ventricular myocytes from ascending aortic banded rats and age-matched controls. Intracellular pH ( $\text{pH}_i$ ) was measured in individual myocytes with SNARF-1, and  $[\text{Ca}^{2+}]_i$  was measured with indo-1, simultaneous with cell motion. Experiments were performed at  $36^\circ\text{C}$  in myocytes paced at 0.5 Hz in Hepes-buffered solution ( $\text{pH}_o$  7.40) containing 1.2 mM  $\text{CaCl}_2$ . At baseline, calibrated  $\text{pH}_i$ , diastolic and systolic  $[\text{Ca}^{2+}]_i$  values, and the amplitude of cell contraction were similar in hypertrophied and control myocytes. Exposure of the control myocytes to 10 nM ET-1 caused an increase in the amplitude of cell contraction to  $163 \pm 22\%$  of baseline ( $P < 0.05$ ), associated with intracellular alkalinization ( $\text{pH}_i + 0.08 \pm 0.02$  U,  $P < 0.05$ ) and a slight increase in peak systolic  $[\text{Ca}^{2+}]_i$  ( $104 \pm 1\%$  of baseline,  $P < 0.05$ ). In contrast, in the hypertrophied myocytes, exposure to ET-1 did not increase the amplitude of cell contraction or cause intracellular alkalinization ( $-0.01 \pm 0.02$  U, NS). Similar effects were observed in the hypertrophied and control myocytes in response to exposure to 10 nM angiotensin II. ET-1 also increased the rate of recovery from intracellular acidosis induced by the washout of  $\text{NH}_4\text{Cl}$  in the control cells, but did not do so in the hypertrophied cells. In the presence of 10  $\mu\text{M}$  5-(*N*-ethyl-*N*-isopropyl)-amiloride, which inhibits  $\text{Na}^+\text{-H}^+$  exchange, ET-1 did not cause a positive inotropic effect or intracellular alkalinization in control cells. The activation of protein kinase C by exposure to phorbol ester caused intracellular alkalinization and it increased the rate of recovery from intracellular acidification induced by an  $\text{NH}_4\text{Cl}$  pulse in control cells but not in hypertrophied cells. ET-1, as well as angiotensin II, and phorbol ester, fail to stimulate forward  $\text{Na}^+\text{-H}^+$  exchange in adult hypertrophied myocytes. These data suggest a defect in the coupling of protein kinase C signaling with  $\text{Na}^+\text{-H}^+$  exchange in adult hypertrophied myocytes. (*J. Clin. Invest.* 1997; 99: 125–135.) Key words: hypertrophy • myocytes • endothelin-1 • angiotensin II

## Introduction

Cardiac pressure overload hypertrophy is associated with changes in gene expression and function that recapitulate a fetal pattern of cellular biology (1, 2). For example, we have shown that hypertrophied myocytes from aortic banded rats show a fetal pattern of gene programming and metabolism (3, 4). Recent studies suggest that the vasoactive peptides endothelin-1 (ET-1)<sup>1</sup> and angiotensin II may play an important role as trophic growth factors during the hypertrophic growth response in immature myocytes (5, 6). We have shown that the effects of these vasoactive peptides on immediate early cardiac gene induction and protein synthesis differ in normal and hypertrophied adult hearts (7–9), suggesting the potential for differing intracellular signal transduction pathways in adult hypertrophied myocytes.

Previous studies have demonstrated that ET-1, as well as angiotensin II, cause a positive inotropic effect in isolated adult ventricular myocytes from rabbits, which appears to be mediated by intracellular alkalinization which sensitizes the myofilaments to  $\text{Ca}^{2+}$  rather than by changes in cytosolic  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ) (10, 11). In adult rabbit and rat myocytes, this intracellular alkalosis appears to be due in part to the stimulation of the  $\text{Na}^+\text{-H}^+$  exchanger (12–14). In contrast, both ET-1 and angiotensin II fail to increase contractility and induce a slight intracellular acidosis in immature cultured ventricular myocytes from both neonatal rat and embryonic chick hearts (10).

The objective of this study was to test the hypothesis that the effects of ET-1 on cell contraction and intracellular pH in adult hypertrophied myocytes differ from those in adult normal myocytes. Therefore, we compared the effects of ET-1, as well as angiotensin II, on cell contraction, intracellular pH, and  $[\text{Ca}^{2+}]_i$  in isolated normothermic adult ventricular myocytes from aortic-banded rats and normal age-matched controls. We also examined the effects of ET-1 and angiotensin II on the recovery from intracellular acidification and the potential mechanistic role of the stimulation of forward  $\text{Na}^+\text{-H}^+$  exchange. Our findings indicate that there is a defect in the coupling of protein kinase C activation with enhanced  $\text{Na}^+\text{-H}^+$  exchange in hypertrophied myocytes.

## Methods

*Preparation of aortic-banded rats.* Weanling male Wistar rats (Charles River Breeding Laboratories, Wilmington, DE) were banded at the age of 3–4 wk (body weight, 75–100 grams) by placing a stainless steel

Address correspondence to Beverly H. Lorell, M.D., Cardiovascular Division, Beth Israel Hospital, 330 Brookline Avenue, Boston, MA 02215. Phone: 617-667-8727; FAX: 617-667-4833.

Received for publication 8 June 1995 and accepted in revised form 1 November 1996.

*J. Clin. Invest.*

© The American Society for Clinical Investigation, Inc.

0021-9738/97/01/0125/11 \$2.00

Volume 99, Number 1, January 1997, 125–135

1. *Abbreviations used in this paper:* 2DG, 2-deoxyglucose;  $[\text{Ca}^{2+}]_i$ , intracellular calcium; EIPA, 5-(*N*-ethyl-*N*-isopropyl)-amiloride; ET-1, endothelin-1; TMA-Cl, trimethylamine hydrochloride; TPA, 12-*O*-tetradecanoylphorbol-13-acetate.

clip of 0.6 mm internal diameter on the ascending aorta via a thoracic incision. Age-matched control rats underwent a sham-operation. The rats were fed normal rat chow and water ad libitum, and were used 12 wk after banding. We have shown at this stage after aortic banding this model of pressure-overload hypertrophy is characterized by an increase in left ventricular weight relative to sham-operated controls, concentric hypertrophy, and the absence of chamber dilatation (4, 15–17).

**Simultaneous measurement of  $[Ca^{2+}]_i$  and cell contraction.** Left ventricular myocyte dissociation was performed by a modification of methods of Capogrossi et al. (18) and Haddad et al. (19) and has been described in detail elsewhere (3).  $[Ca^{2+}]_i$  was measured with  $Ca^{2+}$ -sensitive fluorescence indicator indo-1 AM (20) (Molecular Probes, Inc., Eugene, OR) as described previously (3, 11) using a modification of the method of duBell et al. (21) and Ikenouchi et al. (22). Myocytes were attached on coverslips with cell adhesive (Cell-Tak; Collaborative Research, Inc., Waltham, MA) and loaded with 5  $\mu$ M indo-1 AM in Hepes-buffered solution at room temperature for 30 min. The coverslip was rinsed with indo-1 AM free buffer solution and placed in a flow-through heated (36°C) cell superfusion chamber on the stage of an inverted microscope (Nikon, Tokyo, Japan). The instrumentation for fluorescence measurement has been described in detail elsewhere (3, 11, 22–24). The excitation source was a high-pressure Hg-arc lamp which provides an intense emission peak at 360 nm. Further selection of this excitation was made with narrow band width interference filters. The excitation beam was chopped at 360 Hz to reduce bleaching and the myocyte was illuminated via epifluorescent optics using Fluor  $\times$ 40 objective lens (Nikon). The fluorescence light was collected by the objective lens and transmitted to a custom-modified spectrofluorometer (FM-1000; Rincon Scientific Instruments, Santa Barbara, CA) for simultaneous measurement of both 400 and 500 nm wavelengths using two separate photomultiplier tubes. The spectrofluorometer provided analogue signals representing the fluorescence intensity at both wavelengths and the ratio of emitted fluorescence (400/500 nm). The subtraction of background autofluorescence was done by offsetting the photomultiplier tube outputs during the measurement of fluorescence from an unloaded myocyte at the beginning of each experiment. An adjustable iris was used to restrict the optical image to only one myocyte of interest in each experiment to minimize background fluorescence from other myocytes. The image of the beating myocyte was obtained by illumination via the 50-W standard microscope light source passed through a 645-nm band-pass filter. This wavelength was long enough not to interfere with the fluorescence detection at 400 and 500 nm. The motion of the myocyte was monitored using a solid state camera (GP-CD60; Panasonic, Secaucus, NJ) and a custom-modified video detector system (24) (Crescent Electronics, Sandy, UT). The analogue output of the cell motion signal was monitored and recorded continuously with the analogue signal of the  $[Ca^{2+}]_i$ -sensitive fluorescence ratio (F400/500 nm). Two platinum electrodes placed in the bathing fluid were connected to a stimulator (SD9G; Grass Instruments, Quincy, MA) and used to stimulate the myocyte at 0.5 Hz with 3-ms pulses.

In these experiments, myocytes from 51 hearts from sham-operated rats and 40 hearts from aortic-banded rats were studied. Two to four experiments were performed in sequence from separate coverslips of myocytes isolated from one heart. We have documented recently the stability of both cell motion and  $[Ca^{2+}]_i$  for the 15-min duration of the protocols used in this study and demonstrated minimal compartmentation of indo-1 in intracellular organelles in chemically skinned myocytes (3). To determine the absolute values of peak-systolic and end-systolic  $[Ca^{2+}]_i$  under baseline conditions in hypertrophied and control myocytes, we performed calibration studies described (3) by a modification of the methods of Cheung et al. (25) and Borzak et al. (26). The calibration methods have been described in detail elsewhere (3).

**Measurement of intracellular pH.** Intracellular pH ( $pH_i$ ) was measured with the pH-sensitive indicator SNARF-1 AM (Molecular probes, Inc.) as described recently (10–12, 22). First, 50  $\mu$ g of

SNARF-1 AM was added to 50  $\mu$ l of dimethyl sulfoxide, which was mixed with 450  $\mu$ l of fetal bovine serum. Loading of SNARF-1 AM was done by exposing the myocytes on coverslips of a final concentration of 8  $\mu$ M SNARF-1 AM for 15 min at 36°C. The coverslip was rinsed with SNARF-1 AM free solution and placed on a flow-through cell chamber as described above. Excitation was performed at 540 nm, and fluorescence emission was collected simultaneously at 580 and 640 nm using the same optics system as described above except for the substitution of different dichroic mirror and interference filters. To improve the signal-to-noise ratio, the 640/580 fluorescence ratio signal was filtered by a high cutoff filter at 1.0 Hz. At the end of the experiment, the emission ratio from each cell was calibrated in situ by exposing cells to solutions of varying pH. Each solution contained (mM):  $K^+$  140 (adjusted to keep  $[K^+]$  constant),  $MgCl_2$  1.0, Hepes 4.0, EGTA 2.0, 2,3-butanedione monoxime 30, BAPTA-AM 50  $\mu$ M (Molecular Probes Inc.) and nigericin 14  $\mu$ M (Sigma Chemical Co., St. Louis, MO), and was titrated to various pH values (6.7, 6.85, 7.0, 7.2, 7.4) using 1.0 N KOH. Since changes in pH are not linear to the ratio of fluorescence emission (23),  $pH_i$  was calculated by the equation,  $pH_i = (ax + c)/(1 + bx)$ ; where x is the measured emission ratio, and a, b, and c are constant parameters (12, 23).

**Experimental protocol: effects of ET-1 on cell contraction and  $[Ca^{2+}]_i$ .** Hypertrophied ( $n = 21$ ) and control ( $n = 17$ ) myocytes were superfused with oxygenated Hepes-buffered normal Tyrode solution of the following composition (mM): NaCl 137, KCl 3.7,  $MgCl_2$  0.5, Hepes (free acid) 4.0,  $CaCl_2$  1.2, glucose 11, and probenecid 0.5 with a final pH of 7.40. Probenecid, a blocker of organic anion transport, was added as it has been shown to inhibit secretion of both indo-1 and fura-2 free acids from loaded cells (27, 28). The temperature of the myocytes was maintained at 36–37°C. The myocytes were placed at 0.5 Hz. After recording baseline data, the cells were superfused with oxygenated Hepes-buffered solution containing ET-1 (Sigma) over a concentration range of  $10^{-10}$  to  $10^{-6}$  M. Each concentration was studied in a separate group of myocytes (5–7 cells per group) for 10 min. The analogue cell motion signals and the F400:500 nm analogue signals were recorded simultaneously. The F400/500 values were converted to  $[Ca^{2+}]_i$  using the procedure described in the previous section.

In separate experiments, we also studied the effects of angiotensin II (Sigma) on cell motion and intracellular calcium handling in additional control ( $n = 23$ ) and hypertrophied myocytes ( $n = 17$ ). First, myocytes were superfused with the same Hepes-buffered solution used for baseline superfusion in the ET-1 protocol, then with the same solution containing angiotensin II over a concentration range of  $10^{-10}$  to  $10^{-6}$  M for 10 min. Each concentration was studied in separate groups of myocytes (4–7 cells per group). The temperature, pH, and the pacing rate were the same as those in the ET-1 protocol.

**Effects of ET-1 on intracellular pH.** To investigate whether the increased amplitude of cell contraction in myocytes exposed to 10 nM ET-1 and the lack of an inotropic effect in hypertrophied control myocytes is due to changes in the  $pH_i$ , separate experiments were performed in hypertrophied ( $n = 12$ ) myocytes and control myocytes ( $n = 11$ ) loaded with SNARF-1. Myocytes were first superfused with the same Hepes-buffered solution used for baseline superfusion to eliminate effects of  $Cl^-/HCO_3^-$  exchange and  $Na^+/HCO_3^-$  exchange on  $pH_i$ . The superfusate was then switched to solution with 10 nM ET-1 for 10 min. In additional experiments in hypertrophied ( $n = 8$ ) and control ( $n = 8$ ) myocytes, the effects of an  $NH_4Cl$  pulse in the presence and absence of 10 nM ET-1 was examined by abrupt exposure and washout of Hepes-buffered solution containing 10.0 mM  $NH_4Cl$  prepared by equimolar replacement of NaCl with  $NH_4Cl$ . As previously described (12, 23), the abrupt exposure to  $NH_4Cl$  initially increases  $pH_i$  as basic  $NH_3$  rapidly enters the myocyte. The  $pH_i$  then falls as charged  $NH_4^+$  enters the cell via  $K^+$  channels and dissociates. Upon washout of extracellular  $NH_4Cl$ , intracellular acidosis is created as internal  $NH_3$  leaves the cell, causing the intracellular retention of  $H^+$ . Previous studies have clarified that the initial rate of recovery from this intracellular acidosis depends predominantly on forward

Na<sup>+</sup>-H<sup>+</sup> exchange (12, 29). In additional experiments, the same protocol was done in SNARF-1 loaded hypertrophied ( $n = 4$ ) and control ( $n = 8$ ) myocytes in the presence and absence of 10 nM angiotensin II.

**Contribution of Na<sup>+</sup>-H<sup>+</sup> exchange.** To further assess the contribution of Na<sup>+</sup>-H<sup>+</sup> exchange to ET-1-induced alkalosis, additional SNARF-1-loaded control ( $n = 9$ ) and hypertrophied myocytes ( $n = 6$ ) were perfused with 10 μM 5-(*N*-ethyl-*N*-isopropyl)-amiloride (EIPA) (Sigma), which inhibits Na<sup>+</sup>-H<sup>+</sup> exchange. As described above, all experiments were performed using Hepes-buffered solutions with p*H*<sub>o</sub> of 7.40. Intracellular pH was measured at baseline and after 6 min of exposure to 10 μM EIPA, followed by 6 min of exposure to 10 nM ET-1 in the presence of EIPA. The identical protocol was performed in additional control ( $n = 4$ ) and hypertrophied ( $n = 4$ ) myocytes to examine effects of exposure to 10 nM angiotensin II in the presence of EIPA. The same protocols were performed in an additional series of myocytes to measure cell contractions, since optical considerations do not permit simultaneous measurement of SNARF-1 fluorescence and detection of cell motion. Because ET-1 causes p*H*<sub>i</sub> to fall in both control and hypertrophied myocytes during inhibition of Na<sup>+</sup>-H<sup>+</sup> exchange, the possibility of metabolic acid production was evaluated by preincubating cells in glucose-free solution containing 15.0 mM 2-deoxyglucose (2DG) to inhibit glycolysis with 5 mM acetate as substrate for mitochondrial oxidative phosphorylation (3, 12) and 10 μM EIPA. The effects of exposure to 10 nM ET-1 in the presence of 2DG and 10 μM EIPA were examined in SNARF-1-loaded control ( $n = 4$ ) and hypertrophied ( $n = 4$ ) myocytes.

**Calculation of intrinsic intracellular buffering power (B<sub>i</sub>).** B<sub>i</sub> was experimentally estimated using step reductions in trimethylamine hydrochloride (TMA-Cl; Sigma) to cause stepwise changes in p*H*<sub>i</sub> as previously described in isolated myocytes (12, 30, 31). To eliminate transsarcolemmal flux of acid equivalents via Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchange, Na<sup>+</sup>-HCO<sub>3</sub><sup>-</sup> transport, and Na<sup>+</sup>/H<sup>+</sup> exchange, all cell bathing solutions used for B<sub>i</sub> measurements contained no added HCO<sub>3</sub><sup>-</sup>, CO<sub>2</sub>, or sodium, and were buffered with Hepes. Sodium-free solutions were prepared by equal replacement of NaCl with *N*-methyl-D-glucamine (Sigma), using 1.0 N HCl to adjust pH to 7.40. To minimize [Ca<sup>2+</sup>]<sub>i</sub> accumulation via Na<sup>+</sup>/Ca<sup>2+</sup> exchange, the solutions also contained no added calcium. The final composition of the sodium-free solutions (mM) was: *N*-methyl-D-glucamine 137, KCl 3.7, glucose 11, Hepes, 4.0, MgCl 0.5. These solutions were gassed with 100% O<sub>2</sub> through a column of Ba(OH)<sub>2</sub> crystals to remove any residual bicarbonate. The solutions containing TMA-Cl of 15, 10, 5, 1, and 0 mM were prepared by equimolar replacement of *N*-methyl-D-glucamine with TMA-Cl, respectively. p*H*<sub>i</sub> was allowed to stabilize in sodium-free solution before applying TMA-Cl. The relationship between B<sub>i</sub> and p*H*<sub>i</sub>, where p*H*<sub>i</sub> was taken at the midpoint of each p*H*<sub>i</sub> step, was plotted as line fit by least squares, linear regression for the control myocytes (27 measurement points from 7 myocytes), and hypertrophied myocytes (31 measurement points from 8 myocytes). B<sub>i</sub> (mM/pH) was calculated as described previously (12, 30, 31):

$$B_i = \Delta [TMAH^+]_i / \Delta pH_i;$$

where Δp*H*<sub>i</sub> is the measured change in p*H*<sub>i</sub>, and Δ[TMAH<sup>+</sup>]<sub>i</sub> is the change in intracellular concentration of trimethylamine ions, calculated according to:

$$[TMAH^+]_i = \frac{[TMA]_o 10^{pH_o - pH_i}}{1 + 10^{pH_o - pK}};$$

where [TMA]<sub>o</sub> is the total extracellular concentration of trimethylamine, p*H*<sub>o</sub> is extracellular pH, with the dissociation constant (p*K*) taken as 9.8 (30). Since all experiments were conducted in the nominal absence of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>, intracellular HCO<sub>3</sub><sup>-</sup> buffering was negligible and total intracellular buffering (B<sub>T</sub>) was equal to B<sub>i</sub>.

**Effects of protein kinase C on Na<sup>+</sup>-K<sup>+</sup> exchange.** To examine the potential role of the activation of protein kinase C in control and hy-

perrophied myocytes, the effects of 80 nM 12-*O*-tetradecanoylphorbol-13-acetate (TPA; Sigma) on p*H*<sub>i</sub> were compared in control ( $n = 7$ ) and hypertrophied myocytes ( $n = 5$ ) loaded with SNARF-1. In addition, the effect of preincubation with 80 nM TPA on the rate of recovery from intracellular acidification induced by an NH<sub>4</sub>Cl pulse was studied in control ( $n = 5$ ) and hypertrophied ( $n = 5$ ) cells. Following the observation that TPA caused intracellular alkalinization in control myocytes, the effect of the inactive phorbol ester 4-α phorbol 12,13-didecanoate (100 nM, Research Biochemicals Inc., Natick, MA) on p*H*<sub>i</sub> was also studied ( $n = 4$  cells). In addition, we studied the effects of 10 nM ET-1 ( $n = 6$ ) or 10 nM angiotensin II ( $n = 5$ ) on p*H*<sub>i</sub> in the presence of 30 nM GF-109203X, a selective protein kinase C inhibitor (Research Biochemicals Inc.), in control cells. Intracellular pH was measured at baseline and after 6 min of exposure to 30 nM GF-109203X, followed by 6 min of exposure to 10 nM ET-1 or 10 nM angiotensin II in the presence of GF-109203X.

To determine if protein kinase C also mediates the intracellular acidification that is seen when Na<sup>+</sup>-H<sup>+</sup> exchange is blocked, the effect of TPA on p*H*<sub>i</sub> after inhibition of Na<sup>+</sup>-H<sup>+</sup> exchange with 10 μM EIPA was studied in control ( $n = 5$ ) and hypertrophied ( $n = 6$ ) myocytes.

**Statistical analysis.** Two-way ANOVA with repeated measures was used for the comparisons between the control versus hypertrophied myocytes, or control versus hypertrophied perfused hearts, of values measured over time in response to exposure to ET-1, or to angiotensin II. Unpaired Student's *t* test was used for comparisons of the values between the groups at baseline.  $P < 0.05$  was considered significant. Results are expressed as mean ± SEM.

## Results

**Characteristics of isolated hypertrophied and normal myocytes.** Both left ventricular weight and left ventricular weight-to-body weight ratio were significantly increased in left ventricles from aortic-banded rats compared with those from sham-operated rats (Table I). Table II shows baseline function of left ventricular control myocytes from sham-operated rats and hypertrophied myocytes from the aortic-banded rats used in the experimental protocols. Myocytes were superfused with oxygenated Hepes-buffered solution (pH 7.40) containing 1.2 mM CaCl<sub>2</sub> and were paced at 0.5 Hz at 36–37°C. There was a significant increase in myocyte cell length in hypertrophied myocytes compared with controls. The amplitude of cell contraction and percent fractional shortening were not statistically different between the two groups. Both time to peak shortening and the time to 50% relengthening were slightly prolonged in the hypertrophied myocytes relative to the control myocytes. Table II shows the results of the intracellular Ca<sup>2+</sup> calibration. Under the baseline experimental conditions of this study, systolic and diastolic [Ca<sup>2+</sup>]<sub>i</sub> levels were similar between control and hypertrophied myocytes. The calibrated values of

**Table I. Body Weight and Left Ventricular Weight of Control and Hypertrophied Rats**

	BW	LVW	LVW/BW
	grams	mg	
Controls ( $n = 51$ )	535 ± 12	1895 ± 45	3.44 ± 0.06
LVH ( $n = 40$ )	520 ± 13	2410 ± 50*	4.71 ± 0.10*

Mean ± SEM. \* $P < 0.05$  versus control rats. BW, body weight; LVW, left ventricular wet weight; LVW/BW, the ratio of left ventricular wet weight to body weight; LVH, hypertrophied aortic-banded rats.

Table II. Baseline Characteristics of Myocyte Function and Intracellular  $Ca^{2+}$

	Controls (n = 51)	Hypertrophy (n = 43)
Myocyte length ( $\mu\text{m}$ )	136 $\pm$ 2	149 $\pm$ 2*
Fractional cell shortening (%)	4.46 $\pm$ 0.30	4.52 $\pm$ 0.31
Amplitude cell shortening ( $\mu\text{m}$ )	6.2 $\pm$ 0.3	6.8 $\pm$ 0.3
Peak positive first derivative of cell motion ( $\mu\text{m/s}$ )	169 $\pm$ 10	158 $\pm$ 15
Peak negative first derivative of cell motion ( $\mu\text{m/s}$ )	-141 $\pm$ 11	-136 $\pm$ 13
Time to peak shortening (ms)	54 $\pm$ 2	75 $\pm$ 2*
Time to 50% relengthening (ms)	33 $\pm$ 1	43 $\pm$ 2*
Peak-systolic $[Ca^{2+}]_i$ (nm)	659 $\pm$ 71	738 $\pm$ 40
End-diastolic $[Ca^{2+}]_i$ (nm)	96 $\pm$ 14	99 $\pm$ 12
Time to peak $[Ca^{2+}]_i$ (ms)	33 $\pm$ 1	37 $\pm$ 1*
Time to 50% decline in $[Ca^{2+}]_i$ from peak (ms)	95 $\pm$ 3	111 $\pm$ 3*
Intracellular pH (pH <sub>i</sub> )	(n = 40) 7.07 $\pm$ 0.02	(n = 25) 7.03 $\pm$ 0.02

Mean $\pm$ SEM. \* $P < 0.05$  versus control myocytes. All data were obtained under baseline perfusion conditions of 36°C, perfusate  $[Ca^{2+}]$  1.2 mM, and paced rate of 0.5 Hz.  $[Ca^{2+}]_i$  was measured using cells loaded with the  $Ca^{2+}$ -sensitive indicator indo-1. Intracellular pH was measured using the pH-sensitive indicator SNARF-1.

diastolic and systolic  $[Ca^{2+}]_i$  in the normothermic paced rat myocytes are similar to those previously reported by us and others (3, 26, 32). The time to peak  $[Ca^{2+}]_i$  and time to 50% decline in  $[Ca^{2+}]_i$  from peak were slightly prolonged in the hypertrophied myocytes (12 and 17% above control myocyte values, respectively). As shown in Table II, the calibrated values of pH<sub>i</sub> using the indicator SNARF-1 were similar in control (n = 20) and hypertrophied myocytes (n = 19), and these val-

ues are comparable with measurements reported by others in isolated ventricular myocytes (22, 23).

**Effects of ET-1 in isolated hypertrophied and normal myocytes.** ET-1 caused a maximal positive inotropic effect at a concentration of 10 nM in the control myocytes, manifest as an increase in the amplitude of contraction to 163 $\pm$ 22% of baseline ( $P < 0.05$ ). However, ET-1 did not increase the amplitude of cell contraction at any concentration in the hypertrophied myocytes (102 $\pm$ 7% of baseline, NS, in response to 10 nM ET-1). The effects of 10 nM ET-1 on fractional cell shortening and peak systolic  $[Ca^{2+}]_i$ , relative to baseline are illustrated in Fig. 1. Perfusion with 10 nM ET-1 caused a slight but consistent increase in peak systolic  $[Ca^{2+}]_i$  in the control and hypertrophied myocytes (104 $\pm$ 1%,  $P < 0.05$ ; and 103 $\pm$ 1% of baseline,  $P < 0.05$ , respectively). There was no change in diastolic  $[Ca^{2+}]_i$  or diastolic cell length in either group in response to 10 nM ET-1.

The effects of angiotensin II were studied in additional control and hypertrophied myocytes. Angiotensin II also caused a maximal positive inotropic effect at a concentration of 10 nM in the control myocytes, manifest as an increase in the amplitude of cell contraction to 119 $\pm$ 4% of baseline ( $P < 0.05$ ). However, angiotensin II did not increase the amplitude of cell contraction at any concentration in the hypertrophied myocytes (105 $\pm$ 4% of baseline, NS, in response to 10 nM angiotensin II). Similar to the effects of ET-1, angiotensin II caused only a slight increase in peak systolic  $[Ca^{2+}]_i$  in the control and hypertrophied myocytes (105 $\pm$ 0.6% of baseline,  $P < 0.05$ ; and 108 $\pm$ 3.0% of baseline,  $P < 0.05$ , respectively) without any change in diastolic  $[Ca^{2+}]_i$  levels. In addition, exposure to 10<sup>-6</sup> M BMS 186-295, a highly specific angiotensin AT<sub>1</sub> receptor subtype blocker (Bristol-Myers Squibb), completely inhibited the positive inotropic effect of 10 nM angiotensin II in control cells (n = 8, 108 $\pm$ 9% of baseline, NS) and had no effect in the hypertrophied cells (n = 5, 94 $\pm$ 4% of baseline, NS).

**Effects of ET-1 and angiotensin II on intracellular pH.** As illustrated in Fig. 2, 10 nM ET-1 caused a significant increase in pH<sub>i</sub> in the control myocytes (+0.08 $\pm$ 0.02 U,  $P < 0.05$ ). How-




Figure 1. Representative traces of cell motion and  $[Ca^{2+}]_i$ -sensitive ratio of fluorescence intensities of 400 to 500 nm (F400/500) during baseline condition and in response to 10 nM ET-1 in a control myocyte (left) and in a hypertrophied myocyte (right). These representative recordings were selected from the control (n = 7) and hypertrophied (n = 7) myocyte studies. Tracings are displayed with the convention that systolic myocyte shortening is displayed as an upward deflection of the cell motion trace. Note that ET-1 increased the amplitude of cell shortening in the control myocyte, with minimal change in  $[Ca^{2+}]_i$ , but did not cause a positive inotropic effect in the hypertrophied myocyte.




Figure 2. Representative traces of the effects of 10 nM ET-1 and 10 nM angiotensin II on  $pH_i$  in a control myocyte (left) and on hypertrophied myocyte (right) loaded with the pH-sensitive fluorescence indicator, SNARF-1. ET-1 and angiotensin II caused intracellular alkalization in the control myocyte. In contrast, there was no change in  $pH_i$  in the hypertrophied myocyte in response to ET-1 and angiotensin II.

ever, the hypertrophied myocytes showed no increase or a slight fall in  $pH_i$  in response to 10 nM ET-1 ( $-0.01 \pm 0.0$  U, NS). The average effects of ET-1 on  $pH_i$  are shown in Fig. 3 A. The effect of 10 nM angiotensin II on  $pH_i$  was studied in additional control and hypertrophied cells (Fig. 3 B). Although the effect was not as marked as that observed with ET-1, angiotensin II also caused an intracellular alkalization in  $pH_i$  in control cells ( $+0.02 \pm 0.01$  U,  $P < 0.05$ ) and no change in hypertrophied myocytes ( $+0.001 \pm 0.01$  U, NS).

To assess the potential contribution of ET-1 to the recovery from acidosis, additional control and hypertrophied myocytes were exposed to an abrupt pulse and washout of  $NH_4Cl$  in the absence and in the presence of 10 nM ET-1. As illus-

trated in Fig. 4, *aA* and *bA*, the  $NH_4Cl$  pulse caused an initial intracellular alkalosis and secondary intracellular acidosis. The magnitude and time course of recovery from intracellular acidification was similar in the control and hypertrophied myocytes. In the presence of 10 nM ET-1, the recovery from intracellular acidification and from the associated negative inotropic effect was enhanced in the control myocytes but not in the hypertrophied myocytes (Fig. 4). The average effects of ET-1 on the increase in  $pH_i$  during the initial 3 min of recovery from maximal intracellular acidification are shown in Fig. 5 and Table III. The effect of 10 nM angiotensin II on the initial rate of recovery from intracellular acidification was also studied in additional control and hypertrophied myocytes (Fig. 5 and Table





Figure 3. Average effects of 10 nM ET-1 (A) and 10 nM angiotensin II (B) on intracellular pH in hypertrophied and control myocytes. Values are normalized relative to baseline for each myocyte. At baseline,  $pH_i$  was similar in both groups. Both ET-1 and angiotensin II caused intracellular alkalization in the control cells, whereas no change or a slight fall in  $pH_i$  was observed in the hypertrophied myocytes. Values are reported under steady state conditions 6 min after exposure to drug.




**Figure 4.** Representative effects of exposure and washout of NH<sub>4</sub>Cl in the absence (*aA* and *bA*) and presence (*aB* and *bB*) of 10 nM ET-1 in control (*a*) and hypertrophied (*b*) myocytes on myocyte shortening (cell length) and p<sub>H</sub><sub>i</sub>. In the absence of ET-1 (*aA* and *bA*), the NH<sub>4</sub>Cl pulse caused an initial intracellular alkalinization and secondary acidification which was of similar time course and magnitude in the control and hypertrophied myocytes. Arrows denote the onset of recovery from acidification. Note that alkalinization was accompanied by an increase in the amplitude of cell contraction and incomplete relaxation, whereas acidification caused a negative inotropic effect. Panel *aB* illustrates that in the control myocytes, the presence of ET-1 during the NH<sub>4</sub>Cl pulse was associated with an exaggerated initial alkalinization, and acceleration of the recovery of p<sub>H</sub><sub>i</sub> from the secondary phase of acidification, implicating the stimulation of Na<sup>+</sup>-H<sup>+</sup> exchange. In contrast, in the hypertrophied myocytes (*bB*), ET-1 did not modify the changes in p<sub>H</sub><sub>i</sub> induced by the NH<sub>4</sub>Cl pulse.

III). Exposure to angiotensin II enhanced the rate of recovery from intracellular acidification induced by an NH<sub>4</sub>Cl pulse in control myocytes but not in the hypertrophied myocytes.

In these experiments using the NH<sub>4</sub>Cl pulse, differences in the effects of ET-1 and angiotensin II on recovery from intracellular acidosis in HEPES-buffered control and hypertrophied cells could be related to differences in the stimulation of H<sup>+</sup> flux via Na<sup>+</sup>-H<sup>+</sup> exchange or to differences in intrinsic buffering power (B<sub>i</sub>). As shown in Fig. 6, the intrinsic buffering

power of hypertrophied and control myocytes was similar under the experimental conditions of this study.

**Na<sup>+</sup>-H<sup>+</sup> exchange in hypertrophied myocytes.** To assess the contribution of Na<sup>+</sup>-H<sup>+</sup> exchange to the effects of ET-1, control and hypertrophied myocytes were exposed with 10  $\mu\text{M}$  EIPA for 6 min to inhibit Na<sup>+</sup>-H<sup>+</sup> exchange and then exposed to the 10 nM ET-1 for 6 min during continuous exposure to EIPA. EIPA alone did not modify p<sub>H</sub><sub>i</sub> in control or hypertrophied cells (Table IV). In the presence of EIPA, ET-1 did not



**Figure 5.** The time course of recovery from maximal intracellular acidosis elicited by washout of a 10.0 mM  $\text{NH}_4\text{Cl}$  pulse in control and hypertrophied cells. The final recovery value 6 min after washout of  $\text{NH}_4\text{Cl}$  is shown. The left panel shows data for control myocytes at baseline in the absence of drug and in the presence of 10 nM ET-1, 10 nM angiotensin II, and 80 nM TPA. The right panel shows data for hypertrophied myocytes. Five to eight cells were studied per group. For clarity, the standard error bars are not shown. In comparison with no drug, ET-1, angiotensin II, and TPA each increased the rate of recovery from intracellular acidification in the control cells ( $P < 0.05$ ). In contrast, these interventions had no effect on the rate of recovery from intracellular acidification in the hypertrophied cells ( $P = \text{NS}$ ), suggesting a failure of the interventions to stimulate  $\text{Na}^+\text{-H}^+$  exchange.

cause intracellular alkalinization in the control myocytes; instead, ET-1 caused intracellular acidification in both control and hypertrophied myocytes (Table IV). The effects of 10 nM angiotensin II on  $\text{pH}_i$  during inhibition of  $\text{Na}^+\text{-H}^+$  exchange with 10  $\mu\text{M}$  EIPA were also studied in additional control and hypertrophied myocytes. Angiotensin II in the presence of EIPA caused intracellular acidification in both control and hypertrophied myocytes (Table IV). To examine the possibility that the intracellular acidification induced by ET-1 in the presence of the inhibition of  $\text{Na}^+\text{-H}^+$  exchange is in part related to metabolic acid production (12), the effects of 10 nM ET-1 on  $\text{pH}_i$  were studied in control and hypertrophied cells in the presence of solution containing 10  $\mu\text{M}$  EIPA and 15 mM 2DG to inhibit glycolysis with 5.0 mM acetate as substrate for oxidative phosphorylation. The presence of 2DG completely prevented the ET-1-induced intracellular acidification during exposure to EIPA in both control ( $-0.03 \pm 0.02$  U, NS) and hypertrophied ( $-0.02 \pm 0.02$  U, NS) cells.

**Activation of protein kinase C.** Krämer et al. (13) and Kohmoto et al. (10) have suggested previously that the effects of ET-1 on intracellular alkalinization may be mediated in part by activation of protein kinase C. Therefore, we compared the effects of the phorbol ester TPA on  $\text{pH}_i$  in additional control and hypertrophied cells. Exposure to 80 nM TPA increased

**Table III.** Effects of Interventions on the Initial Recovery of Intracellular pH from  $\text{NH}_4\text{Cl}$ -induced Acidification

	No drug	ET-1	A II	TPA
Controls	$\Delta 0.12 \pm 0.01$	$\Delta 0.20 \pm 0.02^{**}$	$\Delta 0.19 \pm 0.02^{**}$	$\Delta 0.18 \pm 0.02^{**}$
LVH	$\Delta 0.12 \pm 0.01$	$\Delta 0.11 \pm 0.02$	$\Delta 0.11 \pm 0.02$	$\Delta 0.11 \pm 0.01$

Mean  $\pm$  SEM.  $^{*}P < 0.05$  versus no drug,  $^{**}P < 0.05$  versus LVH. The  $\Delta\text{pH}_i$  (pH units) was measured during the first 3 min of recovery from maximal acidification induced by washout of  $\text{NH}_4\text{Cl}$  pulse in presence and absence of drug. Each intervention was performed in separate groups of cells (4–8 cells/group). ET-1, exposure to 10 nM ET-1; A II, exposure to 10 nM angiotensin II; TPA, exposure to 80 nM TPA.

$\text{pH}_i$  in control cells ( $+0.06 \pm 0.01$  U,  $P < 0.05$ ) but not in hypertrophied cells ( $+0.01 \pm 0.01$  U, NS). Exposure to the inactive phorbol ester 100 nM 4- $\alpha$  phorbol 12,13-didecanoate had no effect on  $\text{pH}_i$  in the control cells ( $+0.01 \pm 0.01$  U, NS). In addition, we studied the effects of 10 nM ET-1 and 10 nM angiotensin II on  $\text{pH}_i$  in the presence of specific inhibition of protein kinase C with 30 nM GF-109203X in control cells. In the presence of GF-109203X, ET-1 ( $-0.01 \pm 0.01$  U, NS) and angiotensin II ( $-0.03 \pm 0.02$  U, NS) had no effect on  $\text{pH}_i$ . To examine the effects of TPA on  $\text{Na}^+\text{-H}^+$  exchange, the rate of recovery from intracellular acidification induced by an  $\text{NH}_4\text{Cl}$  pulse in the presence of 10 nM TPA was compared in additional control and hypertrophied cells. The results are shown in Fig. 5 and Table III. In the control cells, 80 nM TPA enhanced the initial rate of recovery from intracellular acidification; in contrast, TPA had no effect on the initial rate of recovery from intracellular acidification in the hypertrophied cells.

The effect of TPA on  $\text{pH}_i$  was also examined after inhibition of  $\text{Na}^+\text{-H}^+$  exchange with EIPA to determine if protein kinase C also mediates the intracellular acidification that is seen when  $\text{Na}^+\text{-H}^+$  exchange is blocked (Table IV). Following inhibition of forward  $\text{Na}^+\text{-H}^+$  exchange, TPA caused intracellular acidification rather than alkalinization in both control cells and hypertrophied cells. This suggests that the defect in protein kinase C signaling in hypertrophied cells is not generalized beyond the defect in coupling between protein kinase C and  $\text{Na}^+\text{-H}^+$  exchange. Thus, it is apparent that the effects of ET-1, angiotensin II, and TPA on the stimulation of  $\text{Na}^+\text{-H}^+$  exchange are similar in control cells, but differ in adult hypertrophied cells.

## Discussion

In this study, exposure of normal adult rat ventricular myocytes to ET-1 caused an increase in the amplitude of contraction which was mediated by intracellular alkalinization rather than by a predominant increase in  $[\text{Ca}^{2+}]_i$ . The responses of adult hypertrophied myocytes were strikingly different since




Figure 6. The intrinsic buffering power ( $B_i$ ) relative to  $pH_i$  measured in control and hypertrophied ventricular myocytes ( $pH_o = 7.40$ ). All cell bathing solutions contained *N*-methyl-D-glucamine as replacement for sodium, zero sodium, and were buffered with Hepes. As illustrated in the top panel,  $B_i$  was calculated during stepwise changes in  $pH_i$  that were elicited by application and removal of TMA-Cl. The line fits for data obtained from the control and hypertrophied cells are displayed and are virtually superimposed, showing that  $B_i$  is similar in control and hypertrophied cells over the range of  $pH_i$  shown (bottom). Line fits by least squares, linear regression:  $B_i$  (mM/pH) =  $63.14 - 6.39 pH_i$  ( $n = 27$  measurements from seven control myocytes);  $B_i = 65.55 - 6.72 pH_i$  ( $n = 31$  measurements from eight LVH myocytes).

ET-1 did not increase contractility or cause intracellular alkalization. Similar effects on contractility, intracellular pH, and  $[Ca^{2+}]_i$  were observed in response to angiotensin II as in response to ET-1. It is well established that the effects of the vasoactive peptides ET-1 and angiotensin II may vary with the preparation which is studied and with the stage of development. In intact isolated heart preparations, both positive and negative inotropic effects of ET-1 and angiotensin II have been reported in adult rabbits and rat hearts (10, 11, 16, 29, 33). These conflicting responses may be related to the effects of these vasoactive peptides on coronary vasomotor tone, thus confounding the assessment of direct myocardial actions. Recent studies on isolated normothermic paced adult rabbit myocytes have demonstrated that ET-1 and angiotensin II cause a

Table IV. Effects of EIPA on Intracellular pH

	Controls	LVH
Baseline	( $n = 18$ ) $7.07 \pm 0.02$	( $n = 17$ ) $7.05 \pm 0.02$
EIPA alone	( $n = 18$ ) $7.05 \pm 0.02$	( $n = 17$ ) $7.02 \pm 0.03$
EIPA with ET-1	( $n = 9$ ) $6.96 \pm 0.03^*$	( $n = 8$ ) $6.89 \pm 0.02^*$
EIPA with AT II	( $n = 4$ ) $6.92 \pm 0.04^*$	( $n = 4$ ) $6.89 \pm 0.03^*$
EIPA with TPA	( $n = 5$ ) $6.97 \pm 0.03^*$	( $n = 5$ ) $6.94 \pm 0.02^*$

Mean  $\pm$  SEM. \* $P < 0.05$  versus baseline. EIPA, exposure to 10 mM EIPA, an inhibitor of  $Na^+ - H^+$  exchange for 6 min; ET-1, exposure to 10 nM ET-1 for 6 min during continued perfusion with EIPA; A II, exposure to 10 nM angiotensin II for 6 min during continued perfusion with EIPA. TPA, exposure to 80 nM TPA for 6 min during continued perfusion with EIPA. Intracellular pH was measured at baseline, 6 min after exposure to EIPA, and 6 min after the addition of ET-1, AT II, or TPA. In the presence of EIPA, ET-1, AT II, and TPA each caused intracellular acidification in both the control and LVH cells.

minimal change in intracellular  $Ca^{2+}$  transients and promote an increase in the amplitude of cell contraction which is mediated predominantly by intracellular alkalization (10, 11). In contrast, the exposure of immature embryonic or neonatal chick and rat ventricular myocytes to ET-1, as well as angiotensin II, has been reported to cause an absent or negative inotropic effect and a slight intracellular acidosis (10, 34). Thus, it appears that the intracellular signal transduction pathways of these vasoactive peptides are strikingly heterogeneous in myocytes of the same species at different stages of development. This has implications for understanding the cellular mechanisms of these peptides as trophic growth factors in immature versus adult myocardium.

In this study, we used adult dissociated myocytes from ascending aortic-banded rats killed 12 wk after banding. At this stage of chronic pressure-overload, the animals are characterized by concentric left ventricular hypertrophy with preserved systolic shortening and indices of contractile function, the absence of chamber dilation, a predominant increase in myocyte width, and slight increase in myocyte length (4, 15, 17). At this stage of pressure overload, the animals are characterized by a fetal pattern of gene reprogramming including the enhanced expression of multiple genes characteristic of the immature program including atrial natriuretic factor,  $\alpha$ -skeletal actin,  $\beta$  myosin heavy chain, as well as cardiac ACE and angiotensinogen (4, 16, 35), with preservation of normal message levels of SR  $Ca^{2+}$  ATPase (4). Our interest in examining the signaling of ET-1 and angiotensin II was stimulated by our recent observations that the acute induction of protooncogenes and new protein synthesis in response to these vasoactive peptides are impaired in adult hypertrophied hearts in comparison with normal hearts (7–9).

Under the basal normothermic perfusion conditions of this study, both intracellular pH and diastolic and systolic  $[Ca^{2+}]_i$  levels were similar in normal and hypertrophied myocytes. In these experiments, we observed a slight increase in peak systolic  $[Ca^{2+}]_i$  induced by both ET-1 and angiotensin II in control and hypertrophied myocytes. A small and somewhat inconsistent increase in adult myocyte peak systolic  $[Ca^{2+}]_i$  has also been found after exposure to angiotensin II or ET-1 in previous studies (13, 36, 37) and may be due to a stimulatory effect of intracellular alkalosis on the L-channel  $Ca^{2+}$  current



(38). It is not clear if this mechanism could account for the effect of these peptides on systolic  $\text{Ca}^{2+}$  in hypertrophied myocytes, in which an effect of intracellular pH was not observed. In immature cultured neonatal rat ventricular myocytes, angiotensin II induces an immediate but transient increase in diastolic  $[\text{Ca}^{2+}]$  (6, 39) which appears due to an inositol triphosphate-mediated release of  $\text{Ca}^{2+}$  from sarcoplasmic reticulum stores (39). This is followed by a reduction in the amplitude of the  $\text{Ca}^{2+}$  transient, an effect ascribed to diacylglycerol-induced activation of protein kinase C (39). A similar effect of angiotensin II on release of  $\text{Ca}^{2+}$  from the sarcoplasmic reticulum was not found in embryonic chick myocytes, where only a steady state reduction in the amplitude of the  $\text{Ca}^{2+}$  transient was observed after exposure to ET-1, angiotensin II, or phorbol ester (10). In the present experiments, the effects of these peptides on the steady state  $[\text{Ca}^{2+}]$  transient (slight increase or no change) in hypertrophied myocytes resembled those seen in normal adult myocytes rather than immature cultured myocytes.

Our experiments in normal adult rat ventricular myocytes confirmed prior observations that ET-1 and angiotensin II induce sensitization of the myofilaments to calcium by intracellular alkalinization which is mediated in part by activation of  $\text{Na}^+\text{-H}^+$  exchange (12–14) and provided the novel observation that these effects do not occur in adult hypertrophied myocytes. Exposure of the normal myocytes to ET-1, as well as to angiotensin II, in the presence of EIPA to inhibit  $\text{Na}^+\text{-H}^+$  exchange prevented the expected intracellular alkalinization and increase in contraction amplitude. In the presence of EIPA, ET-1 and angiotensin II caused an intracellular acidosis and depression of contractility which could be blocked by 2DG, which inhibits glycolysis (3, 22, 40). This finding is congruent with recent observations in adult rabbit myocytes by Matsui et al. (12) which suggest that angiotensin II may stimulate metabolic acid production, an effect that is unmasked when  $\text{Na}^+\text{-H}^+$  exchange is blocked.

To gain further understanding of the regulation of intracellular pH in normal and hypertrophied adult myocytes and its modification by ET-1 and angiotensin II, we measured changes in  $\text{pH}_i$  and cell contraction in response to a sudden intracellular alkalosis and subsequent acidosis induced by abrupt exposure to and subsequent removal of  $\text{NH}_4\text{Cl}$  (12, 30, 31). These experiments yielded two insights. First, in the absence of ET-1 or angiotensin II, exposure to  $\text{NH}_4\text{Cl}$  caused a similar magnitude and time course of initial alkalinization and subsequent acidosis in control and hypertrophied myocytes. Second, in the presence of ET-1, as well as angiotensin II, control myocytes exhibited an exaggerated initial alkalosis and faster rate of recovery from the secondary acid load at the same level of  $\text{pH}_i$ . In contrast, in the hypertrophied myocytes, we observed no effect of either ET-1 or angiotensin II on the speed of recovery from intracellular acidification induced by  $\text{NH}_4\text{Cl}$  challenge.

In these experiments using washout of the  $\text{NH}_4\text{Cl}$  pulse in Hepes-buffered cells to induce an intracellular acid load, transsarcolemmal flux of acid equivalents via  $\text{Cl}^-/\text{HCO}_3^-$  transport was eliminated because the cell bathing solutions contained no added  $\text{HCO}_3^-$  and were buffered with Hepes. The rate of switching of solutions was a constant in all experiments. Thus, the rate of recovery from an intracellular acidosis depends on both the net  $\text{H}^+$  efflux via  $\text{Na}^+\text{-H}^+$  exchange as well as the intracellular buffering capacity. Therefore, we measured intrinsic buffering power ( $B_i$ ) in control and hypertrophied cells. Cell

bathing solutions contained no added sodium and were buffered with Hepes to eliminate transsarcolemmal flux of acid equivalents via  $\text{Cl}^-/\text{HCO}_3^-$  exchange,  $\text{Na}^+/\text{HCO}_3^-$  or  $\text{Na}^+\text{-H}^+$  exchange.  $B_i$  tended to increase slightly as  $\text{pH}_i$  declined, and values of  $B_i$  were similar to determinations by others in ventricular myocytes (12, 30). We observed that intrinsic buffering power was similar in the control and hypertrophied myocytes. Thus, it is reasonable to conclude that the faster rate of recovery from intracellular acidification which was observed in control cells but not in hypertrophied cells in response to ET-1 and angiotensin II was related to the stimulation of  $\text{H}^+$  efflux via  $\text{Na}^+\text{-H}^+$  exchange.

Differences in the stimulation of  $\text{Na}^+\text{-H}^+$  exchange in response to ET-1 and angiotensin II in hypertrophied and normal myocytes could be related to differences at the level of the receptor or downstream signaling. We and others have observed a reduction in angiotensin II receptor density in hypertrophied rat myocardium (33, 36) and no change in ET-1 receptor density (36). Prior studies in isolated normal adult rat ventricular myocytes have supported the notion that protein kinase C activation mediates the stimulation of  $\text{Na}^+\text{-H}^+$  exchange by ET-1 and angiotensin II (12–14). In the present experiments, ET-1 and angiotensin II failed to cause intracellular alkalinization in the presence of a selective protein kinase C inhibitor in normal cells. MacLeod and Harding (29) have reported previously that the recovery from an acute acid load in isolated rat myocytes is partially inhibited by amiloride or by removing extracellular  $\text{Na}^+$ . They observed that exposure to phorbol ester 12-myristate 13-acetate (which activates protein kinase C) exaggerated the changes in intracellular pH and associated changes in cell shortening in normal rat and guinea pig myocytes in response to  $\text{NH}_4\text{Cl}$  challenge, which suggests the stimulation of  $\text{Na}^+\text{-H}^+$  exchange. We observed that rapid application of the phorbol ester TPA in normal myocytes increased the rate of recovery from intracellular acidification induced by an  $\text{NH}_4\text{Cl}$  pulse, consistent with the stimulation of  $\text{Na}^+\text{-H}^+$  exchange by the activation of protein kinase C. Exposure to an inactive phorbol ester did not cause intracellular alkalinization in normal myocytes. The failure of active phorbol ester, as well as ET-1 and angiotensin II, to cause intracellular alkalinization in hypertrophied myocytes suggests that down-regulation of ET-1 and angiotensin II receptors does not account for the results; instead, the defect lies in the coupling between activation of protein kinase C and stimulation of  $\text{Na}^+\text{-H}^+$  exchange.

The physiologic significance of the abnormal signaling of protein kinase C and its coupling to  $\text{Na}^+\text{-H}^+$  exchange in hypertrophied hearts is not yet known. Angiotensin II is locally released during ischemia, and we have shown that low-flow ischemia causes greater contractile dysfunction in hypertrophied hearts than in normal hearts (41). This may be related in part to differences in the stimulation of  $\text{Na}^+\text{-H}^+$  exchange by angiotensin II during ischemia. In addition, we have shown that the early growth responses (protooncogene induction and protein synthesis) are blunted in adult hypertrophied hearts in response to stimulation with angiotensin II,  $\alpha$ -adrenergic agonists, as well as mechanical load (7, 8). The present investigation raises the possibility that defective coupling of protein kinase C and  $\text{Na}^+\text{-H}^+$  exchange may contribute to the alteration of growth responses in the hypertrophied heart.

*Summary.* We have confirmed that ET-1, as well as angiotensin II, promotes an increase in contractility in normal adult

rat myocytes via intracellular alkalinization which is partly mediated by activation of  $\text{Na}^+\text{-H}^+$  exchange. This study shows that the effects of ET-1, as well as angiotensin II, differ in adult hypertrophied myocytes and simulate the fetal response of the absence of a positive inotropic effect and the failure to develop intracellular alkalinization. In contrast with normal adult myocytes, exposure to ET-1, angiotensin II, and the phorbol ester TPA fails to stimulate  $\text{Na}^+\text{-H}^+$  exchange in hypertrophied myocytes. These studies suggest that intracellular signal transduction pathways of ET-1 and angiotensin II which stimulate  $\text{Na}^+\text{-H}^+$  exchange via activation of protein kinase C differ in normal and hypertrophied adult myocytes.

## Acknowledgments

We greatly appreciate the expert surgical assistance of Mr. Soeun Ngoy and Mrs. Megumi Ito in the preparation of the aortic-banded rat colonies. We also thank Dr. Kenneth W. Spitzer for his suggestions regarding the determination of intracellular buffering power. We also appreciate the assistance of Tony L. Baptista and John Jaster in preparation of the manuscript.

This work was supported in part by National Heart, Lung, and Blood Institute grants HL-38189 (B.H. Lorell and Y. Kagaya), HL-44431 (B.H. Lorell), and HL-30478 and P50-53773 (W.H. Barry).

## References

- Izumo, S., B. Nadal-Ginard, and V. Mahdavi. 1988. Protooncogene induction and reprogramming of cardiac gene expression produced by pressure overload. *Proc. Natl. Acad. Sci. USA.* 85:339–343.
- Chien, K.R., K.V. Knowlton, H. Zhu, and S. Chien. 1991. Regulation of cardiac gene expression during myocardial growth and hypertrophy: molecular studies of an adaptive physiologic response. *FASEB (Fed. Am. Soc. Exp. Biol.) J.* 5:3037–3046.
- Kagaya, Y., E.O. Weinberg, N. Ito, T. Mochizuki, W.H. Barry, and B.H. Lorell. 1995. Glycolytic inhibition: effects on diastolic relaxation and intracellular calcium handling in hypertrophied rat ventricular myocytes. *J. Clin. Invest.* 95:2766–2776.
- Feldman, A.M., E.O. Weinberg, P.E. Ray, and B.H. Lorell. 1993. Selective changes in cardiac gene expression during compensated hypertrophy and the transition to cardiac decompensation in rats with chronic aortic banding. *Circ. Res.* 73:184–192.
- Shubeita, H.E., P.M. McDonough, A.N. Harris, K.U. Knowlton, C.C. Glembotski, J.H. Brown, and K.R. Chien. 1990. Endothelin induction of inositol phospholipid hydrolysis, sarcomere assembly, and cardiac gene expression in ventricular myocytes. A paracrine mechanism for myocardial cell hypertrophy. *J. Biol. Chem.* 265:20555–20562.
- Sadoshima, J., Y. Xu, S. Slayter, and S. Izumo. 1993. Autocrine release of angiotensin II mediates stretch-induced hypertrophy of cardiac myocytes in vitro. *Cell.* 75:977–984.
- Schunkert, H., E.O. Weinberg, G. Bruckschlegel, A.J.G. Reigger, and B.H. Lorell. 1995. Alteration of growth responses in established cardiac pressure overload hypertrophy in rats with aortic banding. *J. Clin. Invest.* 96:2768–2774.
- Schunkert, H., J.-I. Sadoshima, T. Cornelius, Y. Kagaya, E.O. Weinberg, S. Izumo, G. Riegger, and B.H. Lorell. 1995. Angiotensin II-induced growth responses in isolated adult rat hearts: evidence for load-independent induction of cardiac protein synthesis by angiotensin II. *Circ. Res.* 76:489–497.
- Schunkert, H., L. Jahn, S. Izumo, C.S. Apstein, and B.H. Lorell. 1991. Localization and regulation of *c-fos* and *c-jun* protooncogene induction by systolic wall stress in normal and hypertrophied rat hearts. *Proc. Natl. Acad. Sci. USA.* 88:11480–11484.
- Kohmoto, O., H. Ikenouchi, Y. Harata, S. Momomura, T. Serizawa, and W.H. Barry. 1993. Variable effects of endothelin-1 on  $[\text{Ca}^{2+}]_i$  transients,  $\text{pH}_i$ , and contraction in ventricular myocytes. *Am. J. Physiol.* 265:H793–H800.
- Ikenouchi, H., W.H. Barry, J.H.B. Bridge, E.O. Weinberg, C.S. Apstein, and B.H. Lorell. 1994. Effects of angiotensin II on contractility, intracellular  $\text{Ca}^{2+}$ ,  $\text{I}_{\text{Ca}}$ , and  $\text{pH}$ : studies in isolated beating hearts and myocytes loaded with the indicator Indo 1. *J. Physiol. (Lond.)* 480:203–215.
- Matsui, H., W.H. Barry, C. Livsey, and K.W. Spitzer. 1995. Angiotensin II stimulates  $\text{Na}^+\text{-H}^+$  exchange in adult rabbit ventricular myocytes. *Cardiovasc. Res.* 29:215–221.
- Krämer, B.K., T.W. Smith, and R.A. Kelly. 1991. Endothelin and increased contractility in adult rat myocytes. Role of intracellular alkalosis induced by activation of the protein kinase C-dependent  $\text{Na}^+\text{-H}^+$  exchanger. *Circ. Res.* 68:269–279.
- Wang, J., G. Pail, and J.P. Morgan. 1991. Endothelin 1 enhances myofilament  $\text{Ca}^{2+}$  responsiveness in aequorin-loaded ferret myocardium. *Circ. Res.* 69:582–589.
- Weinberg, E.O., F.J. Schoen, D. George, Y. Kagaya, C.R. Benedict, and B.H. Lorell. 1994. Angiotensin-converting enzyme inhibition prolongs survival and modifies the transition to heart failure in rats with pressure overload hypertrophy due to aortic stenosis. *Circulation.* 90:1410–1422.
- Schunkert, H., V.J. Dzau, S.S. Tang, A.T. Hirsch, C. Apstein, and B.H. Lorell. 1990. Increased rat cardiac angiotensin-converting enzyme activity and mRNA levels in pressure overload left ventricular hypertrophy: effects on coronary resistance, contractility and relaxation. *J. Clin. Invest.* 86:1913–1920.
- Litwin, S.E., S.E. Katz, E.O. Weinberg, B.H. Lorell, G.P. Aurigemma, and P.S. Douglas. 1995. Serial echocardiographic-Doppler assessment of left ventricular geometry and function in rats with pressure-overload hypertrophy: chronic angiotensin-converting enzyme inhibition attenuates the transition to heart failure. *Circulation.* 91:2642–2654.
- Capogrossi, M.C., A.A. Kort, H.A. Spurgeon, and E.F. Lakatta. 1986. Single adult rabbit and rat cardiac myocytes retain the  $\text{Ca}^{2+}$ - and species-dependent systolic and diastolic contractile properties of the intact muscle. *J. Gen. Physiol.* 88:589–613.
- Haddad, J., M.L. Decker, L. Hsieh, M. Lesch, A.M. Samarel, and R.S. Docker. 1988. Attachment and maintenance of adult rabbit cardiac myocytes in primary cell culture. *Am. J. Physiol.* 255:C19–C27.
- Grynkiewicz, G., M. Poenie, and R.Y. Tsien. 1995. A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J. Biol. Chem.* 260:3440–3450.
- duBell, W.H., C. Philips, and S.R. Houser. 1988. A technique for measuring systolic free  $\text{Ca}^{2+}$  with indo 1 in feline myocytes. *In* Biology of Isolated Adult Cardiac Myocytes. W. Clark, editor. Elsevier Science Publishing Co., New York. 187–201.
- Ikenouchi, H., O. Kohmoto, M. McMillan, and W.H. Barry. 1991. The contributions of  $[\text{Ca}^{2+}]_i$ ,  $[\text{Pi}]_i$ , and  $\text{pH}_i$  to altered diastolic myocyte tone during partial metabolic inhibition. *J. Clin. Invest.* 88:55–61.
- Spitzer, K.W., and J.H.B. Bridge. 1992. Relationship between intracellular pH and tension development in resting ventricular muscle and myocytes. *Am. J. Physiol.* 262:C317–C327.
- Steadman, B.W., K.B. Moore, K.W. Spitzer, and J.H.B. Bridge. 1988. A video system for measuring motion in contracting heart cells. *IEEE (Inst. Electr. Electron Eng.) Trans. Biomed. Eng.* 35:264–272.
- Cheung, J.Y., D.L. Tillotson, R.V. Yelamarty, and R.C. Scaduto, Jr. 1989. Cytosolic free calcium concentration in individual cardiac myocytes in primary culture. *Am. J. Physiol.* 256:C1120–C1130.
- Borzak, S., R.A. Kelly, B.K. Kramer, Y. Matoba, J.D. Marsh, and M. Reers. 1990. In situ calibration of fura-2 and BCECF fluorescence in adult rat ventricular myocytes. *Am. J. Physiol.* 259:H973–H981.
- Arkhammer, P., T. Nilsson, and P.O. Berggren. 1990. Glucose-stimulated efflux of indo-1 from pancreatic beta-cells is reduced by probenecid. *FEBS Lett.* 273:182–184.
- DiVirgilio, F., T.H. Steinberg, J.A. Swanson, and S.C. Silverstein. 1988. Fura-2 secretion and sequestration in macrophages. A blocker of organic anion transport reveals that these processes occur via a membrane transport system for organic anions. *J. Immunol.* 140:915–920.
- MacLeod, K. T., and S.E. Harding. 1991. Effect of phorbol ester on contraction, intracellular pH, and intracellular  $\text{Ca}^{2+}$  in isolated mammalian ventricular myocytes. *J. Physiol. (Lond.)* 444:481–498.
- Xu, P., and K.W. Spitzer. 1994.  $\text{Na}^+$ -dependent  $\text{Cl}^-$ - $\text{HCO}_3^-$  exchange mediates recovery from alkalosis in guinea pig ventricular myocyte. *Am. J. Physiol.* 267:H85–H91.
- Lagadic-Gossman, D., K.J. Buckler, and R.D. Vaughan-Jones. 1992. role of bicarbonate in pH recovery from intracellular acidosis in the guinea pig myocyte. *J. Physiol. (Lond.)* 458:361–384.
- Sollott, S.J., B.D. Siman, and E.G. Lakatta. 1992. Novel technique to load indo-1 free acid into single adult cardiac myocytes to assess cytosolic  $\text{Ca}^{2+}$ . *Am. J. Physiol.* 262:H1941–H1949.
- Lopez, J.J., B.H. Lorell, J.R. Ingelfinger, E.O. Weinberg, H. Schunkert, D. Diamant, and S.-S. Tang. 1994. Distribution and function of cardiac angiotensin  $\text{AT}_1$  and  $\text{AT}_2$  receptor subtypes: an analysis of hypertrophied rat hearts. *Am. J. Physiol.* 267(Heart Circ. Physiol. 36):H844–H852.
- Allen, I.S., N.M. Cohen, R.S. Dhallen, S.T. Gaa, W.J. Lederer, and T.B. Rogers. 1988. Angiotensin II increases spontaneous contractile frequency and stimulates calcium current in cultured neonatal rat heart myocytes: insights into the underlying biochemical mechanisms. *Circ. Res.* 62:524–534.
- Schunkert, H., B. Jackson, S.-S. Tang, F.J. Schoen, J.F.M. Smits, C.S. Apstein, and B.H. Lorell. 1993. Distribution and functional significance of cardiac angiotensin converting enzyme in hypertrophied rat hearts. *Circulation.* 87:1328–1339.
- Farch, J., R.M. Touyz, E.L. Schiffrin, and G. Thibault. 1996. Endothelin-1 and angiotensin II receptors in cells from rat hypertrophied heart. *Circ. Res.* 78:302–311.
- Barry, W.H., H. Matsui, J.H.B. Bridge, and K.W. Spitzer. 1995. Excita-

tion-contraction coupling in ventricular myocytes: effects of angiotensin II. *In* Molecular and Subcellular Cardiology: Effects on Structure and Function. S. Sideman and R. Beyar, editors. Plenum Publishing Corp., New York.

38. Karibara, M., S. Mitarai, K. Yano, and M. Kameyama. 1994. Involvement of Na<sup>+</sup>-H<sup>+</sup> antiporter in regulation of L-type Ca<sup>2+</sup> channel current by angiotensin II in rabbit ventricular myocytes. *Circ. Res.* 75:1121–1125.

39. Kem, D.C., E.I.M. Johnson, A.M. Capponi, D. Chardonnens, U. Lang, B. Blondel, H. Koshida, and M.B. Valotton. 1991. Effects of angiotensin II on

cytosolic free calcium in neonatal rat cardiomyocytes. *Am. J. Physiol.* 261:C77–C85.

40. Pirollo, J.S., and D.G. Allen. 1986. Assessment of techniques for preventing glycolysis in cardiac muscle. *Cardiovasc. Res.* 20:837–844.

41. Eberli, F.R., C.S. Apstein, S. Ngoy, and B.H. Lorell. 1992. Exacerbation of ischemic diastolic dysfunction by pressure-overload hypertrophy. Modification by specific inhibition of cardiac angiotensin converting enzyme. *Circ. Res.* 70:931–943.