

RESEARCH ARTICLE

In Vitro Antioxidant versus Metal Ion Chelating Properties of Flavonoids: A Structure-Activity Investigation

Sabri Ahmed Cherrak¹, Nassima Mokhtari-Soulimane^{1*}, Farid Berroukeche¹, Bachir Bensenane¹, Angéline Cherbonnel², Hafida Merzouk¹, Mourad Elhabiri^{2*}

1 Laboratory of Physiology, Pathophysiology and Biochemistry of Nutrition, Department of Biology, Faculty of Natural and Life Sciences, Earth and Universe, Abou Bekr Belkaid University, 13000 Tlemcen, Algeria,

2 Laboratory of Bioorganic and Medicinal Chemistry, UMR 7509 CNRS, ECPM, 25 rue Becquerel, 67200 Strasbourg, France

* nassima_amel@yahoo.fr (NMS); elhabiri@unistra.fr (ME)



OPEN ACCESS

Citation: Cherrak SA, Mokhtari-Soulimane N, Berroukeche F, Bensenane B, Cherbonnel A, Merzouk H, et al. (2016) *In Vitro* Antioxidant versus Metal Ion Chelating Properties of Flavonoids: A Structure-Activity Investigation. PLoS ONE 11(10): e0165575. doi:10.1371/journal.pone.0165575

Editor: Jamshidkhan Chamani, Islamic Azad University Mashhad Branch, ISLAMIC REPUBLIC OF IRAN

Received: August 12, 2016

Accepted: October 13, 2016

Published: October 27, 2016

Copyright: © 2016 Cherrak et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information file.

Funding: This work was supported by the French-Algerian Cooperation Program PHC Tassili International Research Extension Grant TASSILI 13MDU892 and the French Foreign Office (Campus France). This work was partly supported by the CNRS (UMR 7509) and the University of Tlemcen (University Abou Bekr Belkaid).

Abstract

Natural flavonoids such as quercetin, (+)catechin and rutin as well as four methoxylated derivatives of quercetin used as models were investigated to elucidate their impact on the oxidant and antioxidant status of human red blood cells (RBCs). The impact of these compounds against metal toxicity was studied as well as their antiradical activities with DPPH assay. Antihemolytic experiments were conducted on quercetin, (+)catechin and rutin with excess of Fe, Cu and Zn (400 μM), and the oxidant (malondialdehyde, carbonyl proteins) and antioxidant (reduced glutathione, catalase activity) markers were evaluated. The results showed that Fe and Zn have the highest prooxidant effect (37 and 33% of hemolysis, respectively). Quercetin, rutin and (+)catechin exhibited strong antioxidant properties toward Fe, but this effect was decreased with respect to Zn ions. However, the Cu showed a weak antioxidant effect at the highest flavonoid concentration (200 μM), while a prooxidant effect was observed at the lowest flavonoid concentration (100 μM). These results are in agreement with the physico-chemical and antiradical data which demonstrated that binding of the metal ions (for FeNTA: (+)Catechin, $K_{\text{LFeNTA}} = 1.6(1) \times 10^6 \text{ M}^{-1}$ > Rutin, $K_{\text{LFeNTA}} = 2.0(9) \times 10^5 \text{ M}^{-1}$ > Quercetin, $K_{\text{LFeNTA}} = 1.0(7) \times 10^5 \text{ M}^{-1}$ > Q35OH, $K_{\text{LFeNTA}} = 6.3(8.7) \times 10^4 \text{ M}^{-1}$ > Quercetin3'4'OH and Quercetin 3OH, $K_{\text{LFeNTA}} \sim 2 \times 10^4 \text{ M}^{-1}$) reflects the (anti)oxidant status of the RBCs. This study reveals that flavonoids have both prooxidant and antioxidant activity depending on the nature and concentration of the flavonoids and metal ions.

Introduction

Flavonoids that belong to the polyphenols family are secondary plant metabolites and one the most occurring groups of phytochemicals. They occur in fruits, seeds, flowers and vegetables among others and are of important in the human diet. These compounds are of high physiological and morphological importance in plants [1, 2]. Protection of the plant from UV radiation, pigmentation in fruits and flowers or signalling properties (allelochemicals) can be cited.

Competing Interests: The authors have declared that no competing interests exist.

Flavonoids have an important function in plant defence. They can be transferred to the soil and inhibit the growth of the competitors. They can also act as signalling compounds for microbes or bacteria [3, 4]. Many flavonoids have also antifungal [5] and insecticide properties [6]. Last but not least, these phenolic compounds mainly regulate the antioxidant balance of the plants and prevent the plant from mutagenesis.

In humans, consumption of plant polyphenolic compounds (from fruit, vegetables, tea, red wine. . .) was also shown to contribute to numerous health benefits (pleiotropic action) [7]. Polyphenols display a broad spectrum of physiological activities such as anti-allergenic, anti-atherogenic, antimicrobial or cardioprotective effects, to cite a few [8]. For instance, polyphenols-rich diets have been associated with a lower risk of cancer and coronary heart disease mortality in a number of epidemiological studies [9, 10]. These beneficial effects were proposed to be related to reduced exposure to oxidative stress [11, 12].

Even though flavonoids share a common C₆-C₃-C₆ (benzo- γ -pyrone) carbon skeleton (Fig 1), a wide range of structural diversity exists. They are usually found with various degrees of hydroxylation, methoxylation, glycosylation or glucuronidation [13, 14], which contributes to the great variety in biological properties and form the largest group in the polyphenol family with over 6,000 compounds identified so far [13].

The main biological activity of flavonoids thoroughly studied so far is their antioxidant activity. During normal metabolism, reactive oxygen species (ROS/RNS, e.g. hydroxyl OH[•], superoxide O₂^{•-}, nitric oxide NO[•], nitrogen dioxide NO₂[•], peroxy ROO[•] and lipid peroxy LOO[•] to cite a few) are constantly formed and are believed to contribute to cellular aging, [15, 16] mutagenesis or carcinogenesis [17]. It is indeed thought that ROS produce these effects by DNA damages or LDL oxidation [18]. Flavonoids have been demonstrated to be essential radical scavengers [19, 20] because of their ability to stabilize free radicals and other active species. These antiradical/antioxidant capacities are intimately related to the redox properties of their phenolic hydroxyl groups, which can be easily oxidized [21]. As a consequence, their conjugated rings and hydroxyl groups allow them to act as radical scavengers, reducing the effect of ROS in the body. Structure-reactivity studies have demonstrated that the antiradical/antioxidant activities are related to structural criteria such as: (i) presence of an ortho-hydroxyl on the B-ring, (ii) presence of one or several free hydroxyl groups, (iii) presence of a C2-C3 double bond in the C-ring, or (iv) presence of a 3-hydroxyl group (Fig 1) [22–24]. Another potential mechanism by which flavonoids act as antioxidants relates to their interactions with redox enzymes. Flavonoids were shown to activate detoxifying enzymes such as NAD(P)H-quinone oxidoreductase, glutathione S-transferase or UDP-glucuronosyl transferase, which all belong to the defence arsenal towards oxidative stress [25].

The ability of flavonoids to prevent or scavenge the toxicity of redox active metal ions such as Fe or Cu has been less considered. These cations are believed to catalyze the production of oxidant species leading to lipid peroxidation, DNA and protein damages [26, 27]. Fe(II), even in tiny amounts, catalyzes, in the presence of hydrogen peroxide, the formation of hydroxyl radical [•]OH by the well known redox cycling process known as Fenton cycle [28, 29]. Cu displays a more intricate behaviour [30]. Reaction of Cu(II) with H₂O₂ can occur via a “free radical” mechanism in which Cu(II) oxidizes H₂O₂ to O₂^{•-} with Cu(I) being formed and reacting with excess H₂O₂ to form HO[•] (Fenton-like reaction). It was also suggested that rather than changing its oxidation state, Cu(II) forms a complex with peroxide with no formation of radical species (*i.e.*, “complex” mechanism) [31]. Other studies [30] suggested that a higher oxidation state of copper (*i.e.*, Cu(III) formed from Cu(I)+H₂O₂ + 2H⁺ ↔ Cu(III) + 2H₂O) may be a reactive oxidizing intermediate that reacted with various substrates at rate constants that are, however, several orders of magnitude lower than the corresponding reactions with [•]OH. Given these facts, the ability of flavonoids to strongly chelate prooxidant metal ions such as Cu or Fe




Fig 1. Basic structure of flavonoids and general chemical structure of different flavonoids.

doi:10.1371/journal.pone.0165575.g001

that are involved in oxidants/radicals formation contributes as well to their antioxidant properties. A large number of plant flavonoids are capable to form stable metal complexes through their multiple OH groups and the carbonyl moiety, whenever present [32]. For instance, quercetin, that is characterized by three potential bidentate binding sites (α -hydroxy-carbonyl, β -hydroxy-carbonyl or catechol), can lead to stable metallic complexes [33]. Complexation of metal cations by quercetin has been already reported for a large number of metal ions such as Mo(VI), Fe(II)/Fe(III), Cu(II), Zn(II), Al(III), Tb(III), Pb(II), Co(II) [34, 35].

Red blood cells (RBCs) membrane is the widest and most used model for studying biomembrane oxidative damage [36]. As most of the membranes, it plays a fundamental role in maintaining cellular homeostasis of RBCs. RBCs are indeed submitted to continuous fluxes of oxidative stress during their normal aerobic functions. But, in healthy subjects, this stress is balanced out by a powerful and efficient enzymatic and non enzymatic antioxidant network [37]. RBCs are especially susceptible to oxidation due to their high content of polyunsaturated lipids, their rich oxygen supply and the presence of transition metals such as Fe and Cu [38]. The aim of the present study was therefore to investigate, using physico-chemical tools, the metal binding capacities of selected flavonoids of interest that are commonly found in the human diet (*e.g.*, quercetin, rutin and (+)catechin) as well as synthetic models of quercetin (*i.e.*, four polymethylated derivatives of quercetin) that retain one or two bidentate binding sites (Fig 2). The antiradical properties (as scavengers of the 2,2-diphenyl-1-picrylhydrazyl DPPH radical) of these latter natural or synthetic flavonoids were then evaluated *in vitro*. Finally, the resistance of human RBCs to metal-induced hemolysis in the absence or in the presence of flavonoids was assessed *in vitro* as well as their oxidant/antioxidant status. This study was not devoted to mimic the *in vivo* conditions since the bioavailability of flavonoids is rather low [39, 40] and they would be found in low concentrations in human plasma. This fundamental approach allowed highlighting relationships between the chemical structure of the flavonoids and their effectiveness and ability to complex metal transition ions and to prevent RBCs hemolysis.

Methods

Solvent and Materials for the Physico-chemistry Investigations

(+)-Catechin ($C_{15}H_{14}O_6$, MW = 290.27 g mol⁻¹) was extracted from green tea by the group of Dr. A. P. Davies (Unilever, Bedford, UK) and was used without further purification [41].




Fig 2. Chemical structures of quercetin, rutin, (+)-catechin and of the four polymethylated analogues of quercetin (the thick grey colour points out the potential bidentate binding sites).

doi:10.1371/journal.pone.0165575.g002

Quercetin dihydrate ($C_{15}H_{10}O_7 \cdot 2H_2O$, MW = 338.27 g mol⁻¹, Sigma-Aldrich, 98%) and rutin trihydrate ($C_{27}H_{30}O_{16} \cdot 3H_2O$, MW = 664.56 g mol⁻¹, Sigma-Aldrich, 95%) were purchased from Sigma-Aldrich and were used without further purification. NTA (nitrilotriacetate trisodium salt, Fluka, purum) was used as received. Quercetin derivatives such as 3,5,7-tri-O-methyl-quercetin (noted hereafter quercetin-3'4'OH), 5,7,3',4'-tetra-O-methyl-quercetin (noted hereafter quercetin-3OH), 7,3',4'-tri-O-methyl-quercetin (noted hereafter quercetin-35OH) and 3,7,3',4'-tetra-O-methyl-quercetin (noted hereafter quercetin-5OH) were prepared according literature procedures [42–44].

For the sake of solubility, the investigated flavonoids and models were dissolved in a mixed solvent made of 80% of methanol (Merck) and 20% of water by weight. Distilled water was purified by passing it through a mixed bed of ion-exchanger (Bioblock Scientific R3-83002, M3-83006) and activated carbon (Bioblock Scientific ORC-83005) and was de-oxygenated by CO₂- and O₂-free argon (Sigma Oxiclear cartridge) before use. Spectrophotometric grade methanol (Merck, p.a.) was also de-oxygenated by CO₂- and O₂-free argon (Sigma Oxiclear cartridge). All the stock solutions were prepared by weighing solid products using an AG 245 Mettler Toledo analytical balance (precision 0.01 mg).

Copper(II) perchlorate hexahydrate ($Cu(ClO_4)_2 \cdot 6H_2O$, MW = 370.54 g mol⁻¹, reagent grade), Zinc(II) perchlorate hexahydrate ($Zn(ClO_4)_2 \cdot 6H_2O$, MW = 327.38 g mol⁻¹, reagent grade) and Fe(III) perchlorate hydrate ($Fe(ClO_4)_3 \cdot xH_2O$, MW = 354.20 g mol⁻¹ anhydrous basis, reagent grade) were purchased from Alfa Aesar and their stock solutions ($\sim 5\text{--}8 \times 10^{-2}$ M) were prepared from their solid salts in water saturated with argon. The metal contents of the solutions were determined according to the classical colorimetric titrations [45]. The cupric solutions ($Cu(ClO_4)_2 \cdot 6H_2O$) were acidified with 0.1 M HClO₄ to avoid hydroxide precipitation and their concentrations were ascertained by colorimetric titrations with standardized

Na₂H₂EDTA solution (Merck, Titriplex III, 0.1 M) using ammonium acetate (Prolabo, Rectapur) as buffer and PAR (4-(2-(2-Pyridylazo)resorcinol monosodium salt monohydrate) as indicator. The concentrations of the Zn(II) solutions (Zn(ClO₄)₂·6H₂O) stock solutions were ascertained by colorimetric titrations with Na₂H₂EDTA solution (Merck, Titriplex III, 0.1 M), ammoniac (Prolabo, Rectapur) and buffer tablet indicator (Merck). The concentration of Fe(III) stock solutions was ascertained by UV-Vis. absorption spectrophotometry ($\epsilon^{240} = 4.16 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ and $\epsilon^{260} = 2.88 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ in diluted aqueous perchloric acid solution; 4% v/v of HClO₄ at 70%) [46].

CAUTION! Perchlorate salts combined with organic ligands are potentially explosive and should be handled in small quantities and with the adequate precautions [47].

The FeNTA stock solutions ($\sim 10^{-3} \text{ M}$) were prepared by mixing equimolar amounts of Fe(III) perchlorate and NTA. A little excess of NTA was used to ensure complete Fe(III) binding and avoid formation of insoluble ferric tris(hydroxide) at pH 7.4. A Hepes buffer solution (0.1 M) was prepared at pH 7.4 in methanol/water mixture (80/20 w/w).

Absorption spectrophotometric Titrations of the Polyphenols by FeNTA, Cu(II) and Zn(II) at pH 7.4

Microvolumes of concentrated FeNTA, Cu(II) or Zn(II) salts were added to 2 mL of the flavonoid solution ($2-5 \times 10^{-5} \text{ M}$) in a 1 cm path length optical Hellma cell. The corresponding UV-Vis. spectra were recorded from 230 nm to 800 nm to follow the evolution of complexation on a Cary 50 (Varian) or a Cary 5000 (Agilent) spectrophotometer maintained at $(25.0 \pm 2^\circ\text{C})$. The spectrophotometric data were then analyzed with Specfit [48] program which adjusts the absorptivities and the stability constants of the species formed at equilibrium. Specfit uses factor analysis to reduce the absorbance matrix and to extract the eigenvalues prior to the multi-wavelength fit of the reduced data set according to the Marquardt algorithm [49, 50].

Antioxidant Activities Assessed with DPPH Radicals

The antioxidant/antiradical properties of the flavonoids were estimated by the radical scavenging activity method using 2,2-diphenyl-1-picrylhydrazyl radical (DPPH[•]) and compared to that of a standard (*i.e.*, ascorbic acid). Microvolumes of methanolic stock solutions of the samples or the standard were added to 2 mL of a 125 μM of DPPH[•] methanolic solution. The DPPH[•] stock solutions were prepared daily and kept at 4°C until it was used. The absorbance at 515 nm was measured over time (every 30 second) on a Varian Cary 50 UV-Vis. Spectrophotometer until the reaction time reached 16 minutes. The percentage of remaining DPPH[•] as assessed by the absorbance at 515 nm was plotted to evaluate the EC₅₀ (concentration of the antioxidant required to scavenge the initial DPPH[•] concentration by 50%) [51–53]. Low EC₅₀ values reflect high antioxidant capacities of the considered systems.

Preparation of the Flavonoids and Metal Solutions for Biochemical Analyses

Solutions of flavonoids and metals were prepared immediately before use. Quercetin, (+)-catechin and rutin were first dissolved in DMSO and then diluted to a final DMSO concentration of 0.1% in the reaction tube. At this concentration, DMSO had no appreciable effect on RBC hemolysis. Flavonoids were used at two different final concentrations in the reaction tube (200 μM and 100 μM). Fe, Cu and Zn stock solutions were first prepared in distilled water, and then diluted to a final concentration of 400 μM in the reaction tubes.

Ethics Statement

This study was carried out following the Algerian law (25/2006, Resolution No. 387). The donor gave his written informed consent and the ethic committee at the University of Tlemcen approved the study.

Preparation of Human Blood Samples

Human RBCs were separated from heparinized blood that was drawn from a healthy donor. The blood was centrifuged at 2000 rpm for 10 min to separate the RBCs from plasma, and then the RBCs were washed three times with phosphate-buffered saline solution (PBS, pH 7.4) until the color of the supernatant turned clear [54].

Hemolysis Assay

The 5% suspension of RBCs in PBS (pH 7.4) was incubated under air atmosphere at 37°C for 30 minutes with or without the flavonoid, into which an aqueous solution of the metals was added to initiate hemolysis. The reaction mixture was shaken gently while being incubated at 37°C. The extent of hemolysis was determined spectrophotometrically as described elsewhere [55]. Briefly, aliquots of the reaction mixture were taken out after 4 hours of incubation, diluted with PBS, and centrifuged at 2000 rpm for 10 min to separate the RBCs. The percentage of hemolysis was determined by measuring the absorbance of the supernatant at 540 nm and compared with that of complete hemolysis by treating the same RBC suspension with distilled water.

$$\%_{hemolysis} = \left(\frac{A_{sample}}{A_{total}} \right) \times 100$$

Where: A_{sample} is the absorbance of the test sample collected 4 hours from the reaction mixture, and A_{total} is the total hemoglobin content of the cells.

Determination of Markers of the Oxidant/Antioxidant Status

Hemolysate reduced glutathione (GSH) levels were assayed by a colorimetric method of Ellman [56] in which the reduction of Ellman's reagent (5,5-dithiobis-(2-nitrobenzoic) acid DTNB) by thiol groups of GSH to generate 2-nitro-5-thiobenzoic acid which has yellow colour, according to a Sigma Aldrich Kit (Saint Louis, MO, USA). The absorbance at 412 nm was measured, and the GSH concentration was then determined with the GSH standard curve.

The catalase activity (CAT, EC 1.11.1.6) was measured by spectrophotometric analysis of the decomposition rate of hydrogen peroxide according to the method of Aebi [57]. The results were expressed as unit of catalase per mg of Hg.

Carbonyl proteins (marker of proteins oxidation) were assayed by the derivatisation of carbonyl protein groups with 2,4-dinitrophenylhydrazine (DNPH) leading to the formation of stable dinitrophenyl (DNP) hydrazone adducts, which can be detected spectrophotometrically at 375 nm (Sigma Aldrich Kit Saint Louis, MO, USA). Oxidized BSA standard was used for the standard curve [58].

Malondialdehyde (MDA, marker of lipid peroxidation) was estimated by the method of Draper and Hadley et al. [59] using thiobarbituric acid (TBA). Absorbance was measured at 532 nm. The results were expressed as nmol per g of Hb of MDA, using the molar extinction coefficient of chromophore ($1.56 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$).

Statistical Methods

The results are presented as means \pm standard deviations of at least three repetitions. Significant differences ($P < 0.05$) or ($P < 0.01$) were determined by paired Student t-test to compare means between controls and treated RBC's. All tests were performed using STATISTICA-8 program (StatSoft, Tulsa, OK).

Results

Absorption and Stabilities of metal Complexes

In the human body, Fe(III) is mainly exposed to neutral pH (pH \sim 7.4). To maintain its solubility under physiological conditions, Fe is therefore constantly bound to proteins (hemoglobin, transferrin) and to low molecular weight moderate chelators such as citrates. The non-protein bounded iron is called "*labile iron pool*" or "*chelatable iron pool*", and could be the target of exogenous chelators such as polyphenols [60, 61]. Polyaminocarboxylate-type ligands such as citrate and nitrilotriacetic acid (NTA) are currently used in model systems to ensure Fe(III) solubility [62]. For instance, they have been used to study Fe(III) uptake processes by siderophores [63], polyphenols [64, 65], hydroxamic acids [66] or transferrin [67] at physiological pH. The aminotricarboxylate chelator NTA can be considered as a model of citric acid and is one of the most studied organic chelators. The speciation of its ferric complexes is well known [68, 69].

Absorption spectrophotometric titrations of quercetin and its synthetic methylated analogues (Fig 2) with FeNTA were therefore undertaken at pH 7.4 in CH₃OH/H₂O solvent in order to enlighten which chelation site is the favourable one (catecholate *versus* α -hydroxycarbonyl *versus* β -hydroxycarbonyl) and to compare their stability with the other flavones such as rutin (catecholate *versus* β -hydroxycarbonyl) or (+)-catechin (catecholate), two other widely distributed flavonoids in the plants kingdom. As exemplified in Fig 3, the absorption titration of the 3,5,7-tri-O-methyl-quercetin (quercetin3'4'OH) with FeNTA revealed, in addition to the bathochromic shift of the π - π^* transitions lying at \sim 350 nm, the formation of a broad absorption band (ligand-to-metal Charge Transfer LMCT) in the visible region that is characteristic of Fe(III) binding [70] by catecholate based systems.

These absorption transitions lying in the visible region can be indeed attributed to charge transfers from the p_{π} orbitals of catechol or α -hydroxy-carbonyl oxygens to the d_{π} and d_{σ} orbitals of iron(III) and are a clear spectrophotometric signature of the ferric metal complexes. The red shift of the π - π^* transitions of the polyphenolic compounds can be ascribed to electronic effects induced by Fe(III) complexation. Similar effects are indeed observed upon deprotonation of the phenolic sites. LMCT transitions as well as bathochromic shifts of the flavonoids π - π^* transitions were also observed for the other ferric-complexes with FeNTA (Table 1). The absorption spectrophotometric data sets of quercetin and its methylated analogues, (+)-catechin and rutin have been statistically processed [48] to determine the successive stability constants (Table 1). Stoichiometric 1:1:1 FeNTA complexes were systematically evidenced and confirmed the previous studies on catecholate [65] or hydroxamate [71] based systems with FeNTA. The ternary complexes with FeNTA were also further characterized by ESI-MS which reveals the presence of monocharged (positively or negatively) isotopic patterns that agree with the simulated data (*i.e.*, Quercetin3'4'OH in Fig 3). Quercetin3'4'OH (Fig 3), quercetin, rutin and (+)catechin were shown to be the best Fe(III) chelators examined herein, suggesting that the catecholate is a more efficient binding site for FeNTA. In the presence of FeNTA, quercetin3OH and quercetin3,5OH were also shown to display LMCT transitions in the visible region (but lying at much higher energies), [72] which indicates that this molecule, through its α -hydroxycarbonyl unit, also binds FeNTA. However, this binding site afforded less stable ferric




Fig 3. Absorption spectrophotometric titration of 3,5,7-tri-O-methyl-quercetin (quercetin3'4'OH). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the [FeNTA]₀ of the FeNTA complex with quercetin3'4'OH. Solvent: CH₃OH/H₂O (80/20 by weight); pH = 7.4 (Hepes buffer); T = 25.0(2°C); l = 1 cm. [Quercetin 3'4'OH]₀ = 2.89 × 10⁻⁵ M. (D) Electrospray mass spectra of quercetin3'4'OH ferric complex (noted LH₂) in the presence of NTA. Solvent: CH₃OH, capillary voltage = 4000 V. [L.FeNTA]_{tot} = 5 × 10⁻⁵ M; Negative mode; Fragmentor = -100 V.

doi:10.1371/journal.pone.0165575.g003

complexes with respect to catecholate (quercetin3'4'OH, rutin or quercetin). Last but not the least, the absorption titration of the quercetin5OH with FeNTA demonstrates that the β-hydroxycarbonyl, within this flavonol series, doesn't bind Fe(III).

This absorption spectrophotometric investigation of this flavonoids series thus enables us to establish the binding sequence with respect to Fe(III): quercetin3'4'OH > quercetin3OH >> quercetin5OH. The LMCT absorption band measured for quercetin 3'4'OH was found to be centred at about 600 nm as observed for the quercetin LMCT absorption band (λ_{max} LCMT = 608 nm). Assuming these facts, we can propose that the complexation of the quercetin first proceeds through its catechol binding site.

Table 1. Stability constants of the FeNTA, Cu and Zn complexes with rutin, (+)-catechin and quercetin and its O-methylated analogues^a.

Flavonoid	Q	Q3OH	Q5OH	Q35OH	Q3'4'OH	(+)-Catechin	Rutin	
Fe	$K_{L,FeNTA}(\sigma)$ [M ⁻¹]	1.0(7) × 10 ⁵	2.5(1.1) × 10 ⁴	nc	6.3(8.7) × 10 ⁴	2.0(1.4) × 10 ⁴	1.6(1) × 10 ^{6b}	2.0(9) × 10 ⁵
	$K_{L,FeNTA}(\sigma)$ [M ⁻¹]	620			425/515	600	570	598
	$\epsilon^{\lambda_{max}}$ (10 ³ M ⁻¹ cm ⁻¹)	3.7	4.8		4.8/1.1	4.05	2.4	2.7
Cu	$K_{L,Cu}(\sigma)$ [M ⁻¹]	1.0(5) × 10 ⁶	nd	nd	nd	nd	1.0(0.5) × 10 ⁴	4.0(9) × 10 ⁴
	$K_{L2,Cu}(\sigma)$ [M ⁻¹]	6.3(4.4) × 10 ⁴					-	-
Zn	$K_{L,Zn}(\sigma)$ [M ⁻¹]	1.70(8) × 10 ⁴	nd	nd	nd	nd	~ 10 ²	4.2(3) × 10 ²

Q = quercetin, Q3'4'OH = Quercetin3'4'OH, Q35OH = Quercetin35OH, Q3OH = Quercetin3OH, Q5OH = Quercetin5OH. log K_{FeNTA} = 6.0 at pH = 7.4.

^a Solvent: CH₃OH/H₂O (80/20 by weight); pH = 7.4 (Hepes buffer); l = 0.1 M (Hepes); T = 25.0(2°C).

^b Solvent: H₂O; pH = 7.4 (Hepes buffer 0.05 M); T = 25.0(2°C); nd: not determined; nc: no to weak complexation. σ = standard deviation.

doi:10.1371/journal.pone.0165575.t001

The Cu(II) and Zn(II) complexation properties of quercetin, rutin and (+)catechin were then examined using the same analytical approach. The stability sequence satisfactorily follows the Irving-Williams order [73] with the Cu(II) complexes being the more stable with respect to Zn(II). In addition, cupric monochelate LCu and bischelates L₂Cu were characterized for quercetin in agreement with reported data [74].

Antioxidant Properties

Table 2 summarizes the results of the antiradical activities obtained using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay.

Ascorbic acid was used for comparisons purposes. All the compounds investigated in this work exhibited potent antiradical activities. The values of the EC₅₀ (effective concentration of the antioxidant by which the initial DPPH concentration is decreased by 50%) are inversely proportional to the antioxidant activity (*i.e.*, lower the EC₅₀ is, higher the antioxidant activity of a compound is). Among the tested compounds, quercetin (12.02 ± 1.08 μM) and rutin (12.62 ± 0.78 μM) exhibited the highest radical scavenging activity (Fig 4). It was found to be two times more efficient as the standard ascorbic acid. Generally speaking, compounds bearing a catechol B subunit (quercetin, rutin, (+)catechin and quercetin3'4'OH) were found to be more efficient than ascorbic acid. The antiradical activity sequence was established as follows: quercetin > rutin > quercetin3'4'OH > (+)catechin > ascorbic acid > quercetin3OH > quercetin3,5OH > quercetin5OH and perfectly matches with the one established for Fe(III) binding. It thus demonstrates that, within the flavonoids sub-family, an efficient antiradical/antioxidant compound is also an efficient Fe(III) chelator.

In vitro Activities

We then investigated the effect of metal addition on the hemolysis of RBC membranes (Fig 5). For the sake of solubility, Fe was added as its ferrous state. It is anticipated that Fe(II) will be readily oxidized to the Fe(III) state [75]. Addition of 400 μM of Fe²⁺, Zn²⁺ and Cu²⁺ led to 37%, 22% and 32% of RBC hemolysis, respectively.

We have shown above that quercetin, rutin and (+)catechin, that display a B-catecholate subunit are strong radical scavengers and good Fe(III) chelators. It is therefore not surprising to observe that these three natural flavonoids are able due to their intrinsic properties to markedly prevent RBC hemolysis in the presence of Fe overload. Quercetin displays potent RBC anti-hemolytic activity and decreases the hemolysis percentage induced by metal supply (37% at 400 μM Fe²⁺) to very low levels (4.18% and 7.45% at 200 and 100 μM of quercetin, respectively). However, quercetin did not show the same extent of protection in the presence

Table 2. Effective concentration (EC₅₀±SD) of the investigated flavonoids and standard (*i.e.*, ascorbic acid). SD: Standard Deviation.

Compound	EC ₅₀ (μM)	Slope/R ²
Quercetin	12.02±1.08	7.67/0.998
Rutin	12.62±0.78	7.13/0.999
Quercetin3',4'OH	18.16±0.65	6.61/0.998
catechin	21.11±0.86	4.62/0.986
Ascorbic acid	24.84±0.12	3.96/0.998
Quercetin3OH	25.72±0.46	3.24/0.999
Quercetin3,5OH	37.86±0.22	2.11/0.999
Quercetin5OH	50.27±0.08	1.79/0.999

doi:10.1371/journal.pone.0165575.t002




Fig 4. DPPH scavenging activity of studied flavonoids and the standard: ○-Quercetin, ■-Rutin, ▲-Quercetin 3',4'-OH, ▼-Catechin, □-Ascorbic acid, ◀-Quercetin 3 OH, ▶-Quercetin 3,5 OH, ●-Quercetin 5 OH. The average error on the inhibition percentage was estimated to be 4% for all the examined concentrations.

doi:10.1371/journal.pone.0165575.g004

of Zn^{2+} ions. It decreases the initial hemolysis from 33% (400 μM of Zn^{2+}) to only 28% and 22% at 100 μM and 200 μM of quercetin, respectively. This is in agreement with the weaker chelating capacities of catecholate-based ligands toward Zn^{2+} (Table 1). Strikingly, quercetin used at both concentrations (100 μM and 200 μM) shows a prooxidant activity in the presence of Cu^{2+} . RBC Hemolysis increased from 22% (400 μM of Cu^{2+}) to 35% (200 μM) and 42% (100 μM), respectively. These results are consistent with previous works that demonstrate Cu (II) reduction by catechols to more toxic Cu(I) ions.




Fig 5. Hemolysis of a 5% human RBCs treated with metal ions (400 μM) in the absence or the presence of flavonoids (200 μM and 100 μM) under air atmosphere at 37 $^{\circ}C$. The significance of the differences between treated RBCs and control was determined by the Student t-test: *P < 0.05. **P < 0.01.

doi:10.1371/journal.pone.0165575.g005




Fig 6. Oxidant status of a 5% human RBCs treated with metal ions (400 μM) in the absence or the presence of flavonoids (200 μM and 100 μM) under air atmosphere at 37°C. The significance of the differences between treated RBCs and control was determined by the Student t-test: *P < 0.05. **P < 0.01.

doi:10.1371/journal.pone.0165575.g006

(+)Catechin and rutin, both displaying a B-catecholate subunit, led to comparable results. Their firm Fe binding capacities significantly reduce the RBC hemolysis induced by Fe²⁺ supply. Rutin showed the same anti-hemolytic activities with Zn²⁺ than quercetin, while catechin displayed a higher anti-hemolytic effect in the presence of Zn²⁺. In the presence of Cu²⁺, catechin and rutin had a weak protective effect at 200 μM, whereas this moderate protective effect turns to a prooxidant effect at 100 μM raising the RBC hemolysis percentage to ~ 29% for both catechin and rutin.

We then turned our attention to the oxidant status (*i.e.*, by measuring carbonyl proteins and MDA) of the treated RBCs under different experimental conditions (Fig 6).

Levels of carbonyl proteins were raised from 2.66 nmol/mg (control) to 4.67 (Fe²⁺), 3.78 (Cu²⁺) and 4.46 (Zn²⁺) nmol/mg of proteins, respectively, which confirmed the prooxidant effect of the metal ions. In the presence of Fe²⁺, the three flavonoids turned back the carbonyl proteins to their normal levels (*i.e.*, quercetin is the most potent). When treated with Zn²⁺ ions, carbonyl proteins levels of the treated RBCs also decreased for the three flavonoids; nonetheless this effect was found to be weaker by comparison with Fe²⁺. For the RBCs treated with Cu²⁺, the carbonyl proteins levels further increased whatever the flavonoid considered (100 μM) and exemplify the prooxidant properties of the cuprous catecholate complexes.

With respect to another oxidant marker such as MDA (Fig 6), addition of excess of Fe²⁺ and Zn²⁺ to 5% human RBCs strongly raised its levels from 0.29 nmol/g Hb (control) to 0.44 (Fe²⁺) and 0.49 nmol/g Hb (Zn²⁺), respectively. With Cu²⁺, this effect was much weaker




Fig 7. Antioxidant status of a 5% human RBCs treated with metal ions (400 μM) in the absence or the presence of flavonoids (200 μM and 100 μM) under air atmosphere at 37°C. The significance of the differences between treated RBCs and control was determined by the Student t-test: *P < 0.05. **P < 0.01.

doi:10.1371/journal.pone.0165575.g007

(increase from 0.29 nmol/g Hb for the control to 0.35 nmol/g Hb for 400 μM Cu²⁺). The antioxidant and chelating properties of the three flavonoids are clearly in agreement with the decrease of the MDA marker that was induced by Fe²⁺ and Zn²⁺ overload. By contrast, formation of the prooxidant Cu(I) complexes with catechol-based flavonoids (100 μM, see above) significantly increases the MDA concentrations and reflects altered oxidant status. The use of higher concentrations of flavonoids (200 μM) slightly reverses this deleterious state.

With respect to the antioxidant markers (GSH and catalase), the glutathione GSH levels (7.12 mmol/g Hb for the control) decreased for all the metals considered. This decrease is marked in the case of Fe²⁺ and Zn²⁺ (4.25 and 4.97 mmol/g Hb, respectively) in agreement with the RBCs hemolysis and the markers of the oxidant status. As seen previously, the GSH levels also met the levels of the controls when the RBCs supplied with Fe²⁺ were treated with the three flavonoids. The same situation stands for Zn²⁺ but the GSH levels were not restored as observed for Fe²⁺. Once again, a clear prooxidant effect can be seen when the Cu²⁺ treated RBCs were supplied with 100 μM of the flavonoids (Fig 7).

As for GSH, catalase levels (237 U/mg Hb for the control) significantly decreased for the redox active metal ions Fe²⁺ and Cu²⁺, confirming that the antioxidant status of the RBCs was considerably altered. The effect is less obvious for Zn²⁺. As seen previously, supplying flavonoids to the treated RBCs increases the catalase levels in particular in the case of Fe²⁺, the marked effect being obtained for rutin. By contrast with the other (anti)oxidant markers, the catalase activity of the Cu²⁺-treated RBCs was almost not affected in the presence of catechin and increases for rutin. However, in the presence of quercetin (100 μM) and Cu²⁺, the catalase activity significantly fell off. The catalase activity of the Zn²⁺-treated RBCs does not seem to change significantly under the effect of flavonoids even at concentrations of 200 μM.

Discussion

In this study, red blood cells (RBCs) were subjected to *in vitro* extracellular metal ion-induced hemolysis. We investigated the response of RBCs to the generated damage and their (anti)oxidant status. The impact of three relevant flavonoids was examined on the hemolysis and the (anti)oxidant markers of the treated RBCs. These biochemical data were compared to physico-chemical data related on metal binding.

In human cells, H_2O_2 , a by-product of oxygen metabolism, may be present at micromolar concentrations and is relatively harmless as it reacts with biomolecules at relatively low rates. In addition, specific enzymes facilitate its removal. However, metal (*e.g.*, Fe, Cu) ions can catalyze the conversion of H_2O_2 into a highly reactive and damaging hydroxyl radical (HO^\bullet). Fe mostly occurs in its +II and +III oxidation states. The ferrous ions are soluble in biological fluids and generate in the presence of oxygen or hydrogen peroxide (*i.e.*, Fenton reaction) damaging and deleterious reactive oxygen species (ROS). The ferrous ions are unstable in aqueous media and tend to rapidly react with molecular oxygen to form ferric ions and superoxide anion radical [76]. On the other hand, polyphenols can either bind Fe(II), thus preventing this pro-oxidant cation from reacting with hydrogen peroxide or promote the oxidation of the less stable ferrous complexes into more stable ferric ones (*i.e.*, auto-oxidation) that cannot participate any longer to the Fenton reaction. Our findings support previous studies [77] that showed that flavonoids act as valuable antioxidants because of their chelating properties. It was also reported that metal chelates are more prone to scavenge free radicals than the free flavonoids.

Exposure of RBCs to the Fe-induced stress leads to intracellular damages in the form of hemoglobin oxidation and membrane-bound hemichromes production and modification of the membrane components as a consequence of enhanced lipid peroxidation [78]. This damage led to important hemolysis (37% at 400 μM under our experimental conditions). Pre-treatment of RBCs with quercetin, rutin and catechin almost completely prevented the hemolysis caused by Fe^{2+} . This can be explained by several closely related events. These three relevant flavonoids act as efficient oxygen radical scavengers as shown by their potent antiradical properties towards the DDPH $^\bullet$ radical assay (Table 2) [79]. The formation of very stable Fe complexes (Table 1) also prevents ROS formation (*e.g.*, it has been shown that only traces of H_2O_2 are detected in presence of Fe ions and quercetin [80, 81]). In addition, Fe(II) is rapidly oxidised to Fe(III) in the presence or absence of the polyphenol and the oxidation rate constants of the ferrous complexes have been demonstrated to be intimately related to the anti-oxidant properties of the flavonoid compounds [82].

These features are reflected by the oxidant and antioxidant biomarkers. Both oxidant markers—MDA and carbonyl proteins—significantly increased (Fig 6) on supply of the RBCs with Fe^{2+} cations in line with an Fe-induced oxidative stress of the RBCs that ultimately leads to hemolysis. However, pre-treatment of the RBCs with quercetin, rutin or (+)catechin maintained the MDA and carbonyl proteins at initial levels. On the other hand, the antioxidant markers—GSH and catalase activity—(Fig 7) also significantly decreased in the presence of 400 μM of Fe^{2+} ions in agreement with the oxidative stress experienced by the cells. Following a pre-treatment of the RBCs with quercetin, rutin or (+)catechin in the presence of 400 μM of Fe^{2+} , the GSH concentrations were found to be at normal levels even at low flavonoid concentration (100 μM). The catalase activity was also reinstated at initial levels for quercetin and rutin pre-treatments. By contrast, the catalase activity for the RBCs pre-treated with (+)catechin was comparable to that of 400 μM Fe^{2+} alone. Altogether, these data clearly demonstrated that the antioxidant/chelating properties of the catechol-based flavonoids prevented Fe^{2+} toxicity thus allowing regulating the redox homeostasis of the treated RBCs.

Cu(II) stands in contrast with Fe(II). Reaction of free Cu(II) with H₂O₂ can occur via a “free radical” mechanism in which Cu(II) oxidizes H₂O₂ to O₂^{•-} with Cu(I) being formed. Cu(I) which is produced can react with excess of H₂O₂ to form HO[•] (Fenton-like reaction). It was also proposed that Cu(I) (generated from the reaction between Cu(II) and O₂^{•-}) reacted with H₂O₂ to lead to a deleterious and highly reactive Cu(III) intermediate oxidizing species [30]. Cu can also induce oxidative stress and subsequent damages to cells (by alteration of membrane permeability or by affecting chromatin structure, protein synthesis, and various enzyme activities) through redox cycling between Cu(I) and Cu(II), particularly in the presence of H₂O₂ (*i.e.*, Fenton-like reaction), a by-product of oxygen metabolism [83, 84]. Cu was also shown to decrease the glutathione levels [85] as also evidenced in the present study (Fig 7). We demonstrated that Cu²⁺ ions led to less harmful damages (22% of hemolysis) than Fe²⁺ does (37% of hemolysis). Similarly to Fe²⁺ (see above), the higher levels of carbonyl proteins and MDA, used as oxidant markers (Fig 6), confirmed that the RBCs hemolysis mainly resulted from metal-induced oxidative stress. Carbonyl proteins levels seem to be, however, more altered than MDA. By contrast with Fe²⁺, complexation of Cu²⁺ by catecholate based ligands such as particular flavonoids (*i.e.*, those bearing catechol binding sites) do not represent a significant advantage for the antioxidant activity as catechol-mediated reduction to Cu(I) can occur and trigger and exacerbate the copper toxicity [86]. Pre-treatment of the RBCs with the investigated flavonoids indeed clearly demonstrated a prooxidant effect as evidenced by the further increase of the hemolysis to 42%, 30% and 29% for quercetin, (+)catechin and rutin, respectively. We demonstrated that the investigated flavonoids are able to complex Cu²⁺ (Table 1), but led to less stable complexes than with Fe³⁺. The Cu-quercetin complexation was suggested to occur *via* the 4-keto group of the C-ring with additional involvement of the 3OH or 5OH group [74]. Complexation of Cu(II) by catecholate-based ligands was demonstrated to favour metal reduction and lead to more deleterious Cu(I) cations [86]. For instance, it was shown that during the Cu-initiated autoxidation of quercetin, H₂O₂ rapidly accumulates. Furthermore, the main autoxidation products of quercetin was shown to be the solvent adducts on the p-quinone-methide intermediate formed upon two-electron oxidation of quercetin [80]. For 100 μM quercetin, the oxidant markers (Fig 6) were significantly increased while the antioxidant ones (Fig 7) concomitantly decreased in agreement with an altered oxidative environment induced by Cu²⁺ complexation. Increasing quercetin concentration up to 200 μM improved the redox status of the treated RBCs. Even though the same effects were also observed for rutin and (+)catechin, the magnitude of oxidative stress was not significant as for quercetin. This can be related to the nature of the Cu complexes with these two polyphenols. Rutin is substituted on its 3OH position by a rutinose while (+)catechin is a flavan-3-ol. At high concentrations (200 μM), an antioxidant activity was observed for (+)catechin and rutin, while they displayed a prooxidant effect at much lower concentrations (100 μM). It was proposed that for polyphenols exhibiting both antioxidant and prooxidant activity, a redox-cycling pathway may occur at low concentrations when there is not enough polyphenols to scavenge radicals, and at higher concentrations radicals are scavenged at once [87].

A particular behaviour among the investigated metals is occupied by the redox inert metal ion Zn²⁺. Zn²⁺ is an essential component of numerous proteins involved in the defence against oxidative stress [88]. It is the most common non-redox transition metal [89] and its total concentration in human plasma is ranging between 12 and 20 mM [90]. However, its free concentration is several orders of magnitude lower, mainly due to binding by albumin [91, 92]. Cellular Zn homeostasis is tightly regulated because of the regulatory roles of intracellular Zn²⁺. Specialized proteins are responsible for controlling Zn import and export, as well as its intracellular distribution [93]. We have showed that a supply of Zn²⁺ cations at a concentration of 400 μM is responsible of potent hemolysis of RBCs (32%,). At such high concentrations, Zn²⁺ could also affect

transport systems across the RBCs and therefore increase the permeability of the membranes to small molecules and lead to subsequent hemolysis [94]. Flavonoids protection was clearly shown to be less efficient against the Zn^{2+} -induced hemolysis (16% of the initial hemolysis for quercetin and rutin and 55% for catechin at a concentration of 200 μM). This is most likely due to the less stable complexes that can be formed with the three flavonoids considered in this work. Even though Zn^{2+} is an inert metal cation, it induces a significant increase of the oxidant markers such as MDA and carbonyl proteins (Fig 6). Flavonoids protection was efficient at a concentration of 200 μM . The GSH levels were highly decreased by Zn^{2+} ions, but pre-incubation with flavonoids kept these values close to their normal levels (Fig 7). The catalase activity was slightly affected by Zn^{2+} and pre-incubation with flavonoids allowed maintaining the catalase to its normal levels. We thus hypothesized that flavonoids mainly act as antioxidant compounds rather than binding the supplied Zn^{2+} .

Conclusion

In summary, flavonoids possessing a catechol group are more likely to interact with metals especially Fe. Our results showed that at high Fe and Zn concentrations, flavonoids were able to inhibit their hemolytic activity. However, when interacting with Cu a weak antioxidant effect was observed at high metal concentrations, and prooxidant activity was observed at low concentrations. This was corroborated by the data obtained with oxidative stress markers. Flavonoids are then able to exhibit both prooxidant and antioxidant activities depending on their concentration and on the metal concentration.

Supporting Information

S1 Fig. Absorption spectrophotometric titration of quercetin by FeNTA. (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[FeNTA]_0$. Solvent: CH_3OH/H_2O (80/20 by weight); pH = 7.4 (Hepes buffer); $T = 25.0(2^\circ C)$; $l = 1$ cm; (1) $[Quercetin]_0 = 1.90 \times 10^{-5}$ M; (2) $[FeNTA]_0/[Quercetin]_0 = 5.58$. (D) Electrospray mass spectra of quercetin (noted LH_5) ferric complex in the presence of NTA. Solvent: CH_3OH , capillary voltage = 4000 V. $[LH_5FeNTA]_0 = 5 \times 10^{-5}$ M; negative mode; Fragmentor = -200 V. (DOCX)

S2 Fig. Absorption spectrophotometric titration of rutin by FeNTA. (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[FeNTA]_0$. Solvent: CH_3OH/H_2O (80/20 by weight); pH = 7.4 (Hepes buffer); $T = 25.0(2^\circ C)$; $l = 1$ cm. (1) $[Rutin]_0 = 4.94 \times 10^{-5}$ M; (2) $[FeNTA]_0/[Rutin]_0 = 2.02$. (D) Electrospray mass spectra of rutin (noted LH_4) ferric complex in the presence of NTA. Solvent: CH_3OH , capillary voltage = 4000 V. $[LH_4FeNTA]_0 = 5 \times 10^{-5}$ M; positive mode; Fragmentor = +50 V. (DOCX)

S3 Fig. Absorption spectrophotometric titration of quercetin3OH by FeNTA. (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[FeNTA]_0$. Solvent: CH_3OH/H_2O (80/20 by weight); pH = 7.4 (Hepes buffer); $T = 25.0(2^\circ C)$; $l = 1$ cm. (1) $[Quercetin3OH]_0 = 1.51 \times 10^{-5}$ M; (2) $[FeNTA]_0/[Quercetin3OH]_0 = 2.49$. (D) Electrospray mass spectra of quercetin3OH (noted LH) ferric complex in the presence of NTA. Solvent: CH_3OH , capillary voltage = 4000 V. $[LFeNTA]_0 = 5 \times 10^{-5}$ M; positive mode; Fragmentor = +150 V. (DOCX)

S4 Fig. Absorption spectrophotometric titration of quercetin35OH by FeNTA. (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function

of the $[\text{FeNTA}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm; (1) $[\text{Quercetin35OH}]_0 = 4.0 \times 10^{-5}$ M; (2) $[\text{FeNTA}]_0/[\text{Quercetin35OH}]_0 = 1.77$. (DOCX)

S5 Fig. Absorption spectrophotometric titration of quercetin by Cu(II). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[\text{Cu(II)}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm. (1) $[\text{Quercetin}]_0 = 4.98 \times 10^{-5}$ M; (2) $[\text{Cu(II)}]_0/[\text{Quercetin}]_0 = 0.94$. (DOCX)

S6 Fig. Absorption spectrophotometric titration of quercetin by Zn(II). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[\text{Zn(II)}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm. (1) $[\text{Quercetin}]_0 = 4.94 \times 10^{-5}$ M; (2) $[\text{Zn(II)}]_0/[\text{Quercetin}]_0 = 1.95$. (DOCX)

S7 Fig. Absorption spectrophotometric titration of rutin by Cu(II). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[\text{Cu(II)}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm. (1) $[\text{Rutin}]_0 = 4.94 \times 10^{-5}$ M; (2) $[\text{Cu(II)}]_0/[\text{Rutin}]_0 = 1.40$. (D) Electrospray mass spectra of the rutin (noted LH_4) Cu(II) complex. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 w/w), capillary voltage = 4000 V. $[\text{LH}_4]_0 = [\text{Cu}]_0 = 8.7 \times 10^{-5}$ M; negative mode; Fragmentor = -280 V. (DOCX)

S8 Fig. Absorption spectrophotometric titration of rutin by Zn(II). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[\text{Zn(II)}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm. (1) $[\text{Rutin}]_0 = 4.94 \times 10^{-5}$ M; (2) $[\text{Zn(II)}]_0/[\text{Rutin}]_0 = 28.54$. (DOCX)

S9 Fig. Absorption spectrophotometric titration of (+)catechin by Cu(II). (A) Absorption spectra, (B) absorption electronic spectra, and (C) complex formation evolution as a function of the $[\text{Cu(II)}]_0$. Solvent: $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (80/20 by weight); $\text{pH} = 7.4$ (Hepes buffer); $T = 25.0(2^\circ\text{C})$; $l = 1$ cm. (1) $[(+)\text{Catechin}]_0 = 2.55 \times 10^{-4}$ M; (2) $[\text{Cu(II)}]_0/[(+)\text{Catechin}]_0 = 1.50$. (DOCX)

Acknowledgments

The Ministry of Foreign Affairs of the French Government, the Ministry of Higher Education and Research of the Algerian Government and the French Foreign Office (Campus France) are acknowledged for facilitating the scientific collaboration between the University of Tlemcen (University Abou Bekr Belkaid) and the CNRS (UMR 7509).

Author Contributions

Conceptualization: NMS HM ME.

Data curation: SAC.

Formal analysis: SAC NMS ME.

Funding acquisition: NMS ME.

Investigation: SAC FB BB AC.

Methodology: SAC NMS HM ME.

Project administration: NMS ME.

Resources: NMS HM ME.

Software: SAC ME.

Supervision: NMS HM ME.

Validation: SAC NMS HM ME.

Visualization: SAC NMS HM ME.

Writing – original draft: SAC NMS HM ME.

Writing – review & editing: SAC NMS HM ME.

References

1. Correa J, Souza I, Ladeira A, Young M, Aragushi M. Allelopathic potential of *Eupatorium maximiliani* Schrad. leaves. *Allelopathy J.* 2000; 7:225–34.
2. Ignat I, Volf I, Popa VI. A critical review of methods for characterisation of polyphenolic compounds in fruits and vegetables. *Food Chemistry.* 2011; 126(4):1821–35. doi: [10.1016/j.foodchem.2010.12.026](https://doi.org/10.1016/j.foodchem.2010.12.026) PMID: [25213963](https://pubmed.ncbi.nlm.nih.gov/25213963/)
3. Geoghegan F, Wong R, Rabie A. Inhibitory effect of quercetin on periodontal pathogens in vitro. *Phytotherapy Research.* 2010; 24(6):817–20. doi: [10.1002/ptr.3014](https://doi.org/10.1002/ptr.3014) PMID: [19957242](https://pubmed.ncbi.nlm.nih.gov/19957242/)
4. Hassan S, Mathesius U. The role of flavonoids in root–rhizosphere signalling: opportunities and challenges for improving plant–microbe interactions. *Journal of experimental botany.* 2012; 63(9):3429–44. doi: [10.1093/jxb/err430](https://doi.org/10.1093/jxb/err430) PMID: [22213816](https://pubmed.ncbi.nlm.nih.gov/22213816/)
5. Lourenço RMDC, da Silva Melo P, de Almeida ABA. Flavonoids as antifungal agents. *Antifungal Metabolites from Plants:* Springer; 2013. p. 283–300.
6. Salunke B, Kotkar H, Mendki P, Upasani S, Maheshwari V. Efficacy of flavonoids in controlling *Callosobruchus chinensis* (L.)(Coleoptera: Bruchidae), a post-harvest pest of grain legumes. *Crop Protection.* 2005; 24(10):888–93.
7. Moon YJ, Wang X, Morris ME. Dietary flavonoids: effects on xenobiotic and carcinogen metabolism. *Toxicology in vitro.* 2006; 20(2):187–210. doi: [10.1016/j.tiv.2005.06.048](https://doi.org/10.1016/j.tiv.2005.06.048) PMID: [16289744](https://pubmed.ncbi.nlm.nih.gov/16289744/)
8. Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food chemistry.* 2006; 99(1):191–203.
9. Hayder N, Bouhlel I, Skandrani I, Kadri M, Steiman R, Guiraud P, et al. In vitro antioxidant and antige-notoxic potentials of myricetin-3-o-galactoside and myricetin-3-o-rhamnoside from *Myrtus communis*: modulation of expression of genes involved in cell defence system using cDNA microarray. *Toxicology in vitro.* 2008; 22(3):567–81. doi: [10.1016/j.tiv.2007.11.015](https://doi.org/10.1016/j.tiv.2007.11.015) PMID: [18222061](https://pubmed.ncbi.nlm.nih.gov/18222061/)
10. Knekt P, Kumpulainen J, Järvinen R, Rissanen H, Heliövaara M, Reunanen A, et al. Flavonoid intake and risk of chronic diseases. *The American journal of clinical nutrition.* 2002; 76(3):560–8. PMID: [12198000](https://pubmed.ncbi.nlm.nih.gov/12198000/)
11. Chvátalová K, Slaninová I, Březinová L, Slanina J. Influence of dietary phenolic acids on redox status of iron: Ferrous iron autoxidation and ferric iron reduction. *Food Chemistry.* 2008; 106(2):650–60.
12. Pisoschi AM, Pop A. The role of antioxidants in the chemistry of oxidative stress: a review. *European journal of medicinal chemistry.* 2015; 97:55–74. doi: [10.1016/j.ejmech.2015.04.040](https://doi.org/10.1016/j.ejmech.2015.04.040) PMID: [25942353](https://pubmed.ncbi.nlm.nih.gov/25942353/)
13. Agati G, Azzarello E, Pollastri S, Tattini M. Flavonoids as antioxidants in plants: location and functional significance. *Plant Science.* 2012; 196:67–76. doi: [10.1016/j.plantsci.2012.07.014](https://doi.org/10.1016/j.plantsci.2012.07.014) PMID: [23017900](https://pubmed.ncbi.nlm.nih.gov/23017900/)
14. Gil ES, Cout RO. Flavonoid electrochemistry: a review on the electroanalytical applications. *Revista Brasileira de Farmacognosia.* 2013; 23(3):542–58.
15. Pietta P-G. Flavonoids as antioxidants. *Journal of natural products.* 2000; 63(7):1035–42. PMID: [10924197](https://pubmed.ncbi.nlm.nih.gov/10924197/)
16. Robak J, Gryglewski RJ. Flavonoids are scavengers of superoxide anions. *Biochemical pharmacology.* 1988; 37(5):837–41. PMID: [2830882](https://pubmed.ncbi.nlm.nih.gov/2830882/)
17. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacology & therapeutics.* 2001; 90(2):157–77.

18. Chen K, Thomas SR, Keaney JF. Beyond LDL oxidation: ROS in vascular signal transduction. *Free Radical Biology and Medicine*. 2003; 35(2):117–32. PMID: [12853068](#)
19. Baccarin T, Mitjans M, Lemos-Senna E, Vinardell MP. Protection against oxidative damage in human erythrocytes and preliminary photosafety assessment of *Punica granatum* seed oil nanoemulsions entrapping polyphenol-rich ethyl acetate fraction. *Toxicology in Vitro*. 2015; 30(1):421–8.
20. Wilms LC, Kleinjans JC, Moonen EJ, Briedé JJ. Discriminative protection against hydroxyl and superoxide anion radicals by quercetin in human leucocytes in vitro. *Toxicology in vitro*. 2008; 22(2):301–7. doi: [10.1016/j.tiv.2007.09.002](#) PMID: [17959353](#)
21. Procházková D, Boušová I, Wilhelmová N. Antioxidant and prooxidant properties of flavonoids. *Fitoterapia*. 2011; 82(4):513–23. doi: [10.1016/j.fitote.2011.01.018](#) PMID: [21277359](#)
22. Burda S, Oleszek W. Antioxidant and antiradical activities of flavonoids. *Journal of Agricultural and Food Chemistry*. 2001; 49(6):2774–9. PMID: [11409965](#)
23. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *The Journal of nutritional biochemistry*. 2002; 13(10):572–84. PMID: [12550068](#)
24. Bourassa P, Côté R, Hutchandani S, Samson G, Tajmir-Riahi H-A. The effect of milk alpha-casein on the antioxidant activity of tea polyphenols. *Journal of Photochemistry and Photobiology B: Biology*. 2013; 128:43–9. doi: [10.1016/j.jphotobiol.2013.07.021](#) PMID: [24001682](#)
25. Nijveldt RJ, Van Nood E, Van Hoorn DE, Boelens PG, Van Norren K, Van Leeuwen PA. Flavonoids: a review of probable mechanisms of action and potential applications. *The American journal of clinical nutrition*. 2001; 74(4):418–25. PMID: [11566638](#)
26. Kim G-N, Kwon Y-I, Jang H-D. Protective mechanism of quercetin and rutin on 2, 2'-azobis (2-amidopropane) dihydrochloride or Cu 2+-induced oxidative stress in HepG2 cells. *Toxicology in vitro*. 2011; 25(1):138–44. doi: [10.1016/j.tiv.2010.10.005](#) PMID: [20950682](#)
27. Mehta R, Dedina L, O'Brien PJ. Rescuing hepatocytes from iron-catalyzed oxidative stress using vitamins B1 and B6. *Toxicology in Vitro*. 2011; 25(5):1114–22. doi: [10.1016/j.tiv.2011.03.015](#) PMID: [21457772](#)
28. Hippeli S, Elstner EF. Transition metal ion-catalyzed oxygen activation during pathogenic processes. *FEBS letters*. 1999; 443(1):1–7. PMID: [9928941](#)
29. Prousek J. Fenton chemistry in biology and medicine. *Pure and applied chemistry*. 2007; 79(12):2325–38.
30. Pham AN, Xing G, Miller CJ, Waite TD. Fenton-like copper redox chemistry revisited: hydrogen peroxide and superoxide mediation of copper-catalyzed oxidant production. *Journal of catalysis*. 2013; 301:54–64.
31. Robbins MH, Drago RS. Activation of hydrogen peroxide for oxidation by copper (II) complexes. *Journal of Catalysis*. 1997; 170(2):295–303.
32. Leopoldini M, Russo N, Toscano M. The molecular basis of working mechanism of natural polyphenolic antioxidants. *Food Chemistry*. 2011; 125(2):288–306.
33. Psotová J, Lasovsky J, Vicar J. Metal-chelating properties, electrochemical behavior, scavenging and cytoprotective activities of six natural phenolics. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2003; 147(2):147–53. PMID: [15037895](#)
34. Dehghan G, Khoshkam Z. Tin (II)–quercetin complex: Synthesis, spectral characterisation and antioxidant activity. *Food Chemistry*. 2012; 131(2):422–6.
35. Pasban Ziyarat F, Asoodeh A, Sharif Barfeh Z, Pirouzi M, Chamani J. Probing the interaction of lysozyme with ciprofloxacin in the presence of different-sized Ag nano-particles by multispectroscopic techniques and isothermal titration calorimetry. *Journal of Biomolecular Structure and Dynamics*. 2014; 32(4):613–29. doi: [10.1080/07391102.2013.785919](#) PMID: [23659247](#)
36. Mabile L, Piolot A, Boulet L, Fortin L-J, Doyle N, Rodriguez C, et al. Moderate intake of n-3 fatty acids is associated with stable erythrocyte resistance to oxidative stress in hypertriglyceridemic subjects. *The American journal of clinical nutrition*. 2001; 74(4):449–56. PMID: [11566642](#)
37. Kabel AM. Free Radicals and Antioxidants: Role of Enzymes and Nutrition. *World Journal of Nutrition and Health*. 2014; 2(3):35–8.
38. Delmas-Beauvieux M, Peuchant E, Dumon M, Receveur M, Le Bras M, Clerc M. Relationship between red blood cell antioxidant enzymatic system status and lipoperoxidation during the acute phase of malaria. *Clinical biochemistry*. 1995; 28(2):163–9. PMID: [7628075](#)
39. Rodriguez-Mateos A, Vauzour D, Krueger CG, Shanmuganayagam D, Reed J, Calani L, et al. Bioavailability, bioactivity and impact on health of dietary flavonoids and related compounds: an update. *Archives of toxicology*. 2014; 88(10):1803–53. doi: [10.1007/s00204-014-1330-7](#) PMID: [25182418](#)

40. Moosavi-Movahedi AA, Hakimelahi S, Chamani J, Khodarahmi GA, Hassanzadeh F, Luo F-T, et al. Design, synthesis, and anticancer activity of phosphonic acid diphosphate derivative of adenine-containing butenolide and its water-soluble derivatives of paclitaxel with high antitumor activity. *Bioorganic & medicinal chemistry*. 2003; 11(20):4303–13.
41. Davis AL, Cai Y, Davies AP, Lewis J. ¹H and ¹³C NMR assignments of some green tea polyphenols. *Magnetic resonance in Chemistry*. 1996; 34(11):887–90.
42. Han Y, Ding Y, Xie D, Hu D, Li P, Li X, et al. Design, synthesis, and antiviral activity of novel rutin derivatives containing 1, 4-pentadien-3-one moiety. *European journal of medicinal chemistry*. 2015; 92:732–7. doi: [10.1016/j.ejmech.2015.01.017](https://doi.org/10.1016/j.ejmech.2015.01.017) PMID: [25618020](https://pubmed.ncbi.nlm.nih.gov/25618020/)
43. JURD L. Selective Alkylation of Polyphenols. I. The Use of Diphenylmethylene as a Protective Grouping for o-Dihydroxyflavones. *The Journal of Organic Chemistry*. 1962; 27(3):872–5.
44. Shi Z-H, Li N-G, Tang Y-P, Yang J-P, Duan J-A. Metabolism-based synthesis, biologic evaluation and SARs analysis of O-methylated analogs of quercetin as thrombin inhibitors. *European journal of medicinal chemistry*. 2012; 54:210–22. doi: [10.1016/j.ejmech.2012.04.044](https://doi.org/10.1016/j.ejmech.2012.04.044) PMID: [22647223](https://pubmed.ncbi.nlm.nih.gov/22647223/)
45. Méthodes d'Analyses Complexométriques avec les Titriplex EM, Darmstadt.
46. Bastian R, Weberling R, Palilla F. Determination of iron by ultraviolet spectrophotometry. *Analytical Chemistry*. 1956; 28(4):459–62.
47. Raymond K. Tragic consequence with acetonitrile adduct. AMER CHEMICAL SOC 1155 16TH ST, NW, WASHINGTON, DC 20036; 1983. p. 4-
48. Gampp H, Maeder M, Meyer CJ, Zuberbühler AD. Calculation of equilibrium constants from multiwavelength spectroscopic data—II: SPECFIT: two user-friendly programs in basic and standard FORTRAN 77. *Talanta*. 1985; 32(4):257–64. PMID: [18963840](https://pubmed.ncbi.nlm.nih.gov/18963840/)
49. Maeder M, Zuberbuehler AD. Nonlinear least-squares fitting of multivariate absorption data. *Analytical Chemistry*. 1990; 62(20):2220–4.
50. Marquardt DW. An algorithm for least-squares estimation of nonlinear parameters. *Journal of the society for Industrial and Applied Mathematics*. 1963; 11(2):431–41.
51. Goupy P, Dufour C, Loonis M, Dangles O. Quantitative kinetic analysis of hydrogen transfer reactions from dietary polyphenols to the DPPH radical. *Journal of Agricultural and Food Chemistry*. 2003; 51(3):615–22. doi: [10.1021/jf025938i](https://doi.org/10.1021/jf025938i) PMID: [12537431](https://pubmed.ncbi.nlm.nih.gov/12537431/)
52. Tanzadehpanah H, Asoodeh A, Chamani J. An antioxidant peptide derived from Ostrich (*Struthio camelus*) egg white protein hydrolysates. *Food research international*. 2012; 49(1):105–11.
53. Memarpour-Yazdi M, Asoodeh A, Chamani J. A novel antioxidant and antimicrobial peptide from hen egg white lysozyme hydrolysates. *Journal of Functional Foods*. 2012; 4(1):278–86.
54. Ghavami S, Asoodeh A, Klonisch T, Halayko AJ, Kadkhoda K, Krocak TJ, et al. Brevinin-2R1 semi-selectively kills cancer cells by a distinct mechanism, which involves the lysosomal-mitochondrial death pathway. *Journal of cellular and molecular medicine*. 2008; 12(3):1005–22. doi: [10.1111/j.1582-4934.2008.00129.x](https://doi.org/10.1111/j.1582-4934.2008.00129.x) PMID: [18494941](https://pubmed.ncbi.nlm.nih.gov/18494941/)
55. Fiorani M, Accorsi A, Cantoni O. Human red blood cells as a natural flavonoid reservoir. *Free radical research*. 2003; 37(12):1331–8. PMID: [14753757](https://pubmed.ncbi.nlm.nih.gov/14753757/)
56. Ellman GL. Tissue sulfhydryl groups. *Archives of biochemistry and biophysics*. 1959; 82(1):70–7. PMID: [13650640](https://pubmed.ncbi.nlm.nih.gov/13650640/)
57. Aebi H. [13] Catalase in vitro. *Methods in enzymology*. 1984; 105:121–6. PMID: [6727660](https://pubmed.ncbi.nlm.nih.gov/6727660/)
58. Levine RL, Garland D, Oliver CN, Amici A, Climent I, Lenz A-G, et al. [49] Determination of carbonyl content in oxidatively modified proteins. *Methods in enzymology*. 1990; 186:464–78. PMID: [1978225](https://pubmed.ncbi.nlm.nih.gov/1978225/)
59. Draper H, Hadley M. [43] Malondialdehyde determination as index of lipid Peroxidation. *Methods in enzymology*. 1990; 186:421–31. PMID: [2233309](https://pubmed.ncbi.nlm.nih.gov/2233309/)
60. Chimi H, Morel I, Lescoat G, Padeloup N, Cillard P, Cillard J. Inhibition of iron toxicity in rat hepatocyte culture by natural phenolic compounds. *Toxicology in vitro*. 1995; 9(5):695–702. PMID: [20650146](https://pubmed.ncbi.nlm.nih.gov/20650146/)
61. Kakhlon O, Cabantchik ZI. The labile iron pool: characterization, measurement, and participation in cellular processes. *Free Radical Biology and Medicine*. 2002; 33(8):1037–46. PMID: [12374615](https://pubmed.ncbi.nlm.nih.gov/12374615/)
62. Gabricevic M, Anderson DS, Mietzner TA, Crumbliss AL. Kinetics and mechanism of iron (III) complexation by ferric binding protein: the role of phosphate. *Biochemistry*. 2004; 43(19):5811–9. doi: [10.1021/bi036217y](https://doi.org/10.1021/bi036217y) PMID: [15134455](https://pubmed.ncbi.nlm.nih.gov/15134455/)
63. Faller B, Nick H. Kinetics and mechanism of iron (III) removal from citrate by desferrioxamine B and 3-hydroxy-1, 2-dimethyl-4-pyridone. *Journal of the American Chemical Society*. 1994; 116(9):3860–5.
64. Carrër C. Chélation de métaux de transition par des polyphénols du régime alimentaire: Strasbourg 1; 2005.

65. Elhabiri M, Carrër C, Marmolle F, Traboulsi H. Complexation of iron (III) by catecholate-type polyphenols. *Inorganica Chimica Acta*. 2007; 360(1):353–9.
66. Gabricevic M, Crumbliss AL. Kinetics and mechanism of iron (III)-nitrilotriacetate complex reactions with phosphate and acetohydroxamic acid. *Inorganic chemistry*. 2003; 42(13):4098–101. doi: [10.1021/ic026281o](https://doi.org/10.1021/ic026281o) PMID: [12817967](https://pubmed.ncbi.nlm.nih.gov/12817967/)
67. El Hage Chahine J, Pakdaman R. Transferrin, a mechanism for the holoprotein formation from nitrilotriacetato-Fe (III)-transferrin mixed complex. *Journal de chimie physique*. 1996; 93(2):283–99.
68. Nowack B. Environmental chemistry of aminopolycarboxylate chelating agents. *Environmental Science & Technology*. 2002; 36(19):4009–16.
69. Jn Sanchiz, Esparza P, Domínguez S, Brito F, Mederos A. Solution studies of complexes of iron (III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry. *Inorganica chimica acta*. 1999; 291(1):158–65.
70. Palaniandavar M, Velusamy M, Mayilmurugan R. Iron (III) complexes of certain tetradentate phenolate ligands as functional models for catechol dioxygenases. *Journal of Chemical Sciences*. 2006; 118(6):601–10.
71. Das PK, Bhattacharya SG, Banerjee R, Banerjea D. Thermodynamic and kinetic studies on the iron (III)-hydroxamate interaction in acid media. *Journal of coordination chemistry*. 1989; 19(4):311–20.
72. Brandel J, Humbert N, Elhabiri M, Schalk IJ, Mislin GL, Albrecht-Gary A-M. Pyochelin, a siderophore of *Pseudomonas aeruginosa*: physicochemical characterization of the iron (III), copper (II) and zinc (II) complexes. *Dalton transactions*. 2012; 41(9):2820–34. doi: [10.1039/c1dt11804h](https://doi.org/10.1039/c1dt11804h) PMID: [22261733](https://pubmed.ncbi.nlm.nih.gov/22261733/)
73. Schneider HJ. Binding mechanisms in supramolecular complexes. *Angewandte Chemie International Edition*. 2009; 48(22):3924–77.
74. Mira L, Tereza Fernandez M, Santos M, Rocha R, Helena Florêncio M, Jennings KR. Interactions of flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free radical research*. 2002; 36(11):1199–208. PMID: [12592672](https://pubmed.ncbi.nlm.nih.gov/12592672/)
75. Ilbert M, Bonnefoy V. Insight into the evolution of the iron oxidation pathways. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*. 2013; 1827(2):161–75.
76. Jones -Lee A, Lee GF. Role of iron chemistry in controlling the release of pollutants from resuspended sediments. *Remediation Journal*. 2005; 16(1):33–41.
77. Symonowicz M, Kolanek M. Flavonoids and their properties to form chelate complexes. 2012.
78. Rice-Evans C. Iron-mediated oxidative stress and erythrocytes. *Erythroid Cells*: Springer; 1990. p. 429–53.
79. Cesquini M, Torsoni M, Stoppa G, Ogo St-. t-BOOH-induced oxidative damage in sickle red blood cells and the role of flavonoids. *Biomedicine & pharmacotherapy*. 2003; 57(3):124–9.
80. Hajji HE, Nkhili E, Tomao V, Dangles O. Interactions of quercetin with iron and copper ions: complexation and autoxidation. *Free radical research*. 2006; 40(3):303–20. doi: [10.1080/10715760500484351](https://doi.org/10.1080/10715760500484351) PMID: [16484047](https://pubmed.ncbi.nlm.nih.gov/16484047/)
81. Nowak D, Kuźniar A, Kopacz M. Solid complexes of iron (II) and iron (III) with rutin. *Structural Chemistry*. 2010; 21(2):323–30.
82. Perron NR, Wang HC, DeGuire SN, Jenkins M, Lawson M, Brumaghim JL. Kinetics of iron oxidation upon polyphenol binding. *Dalton Transactions*. 2010; 39(41):9982–7. doi: [10.1039/c0dt00752h](https://doi.org/10.1039/c0dt00752h) PMID: [20871896](https://pubmed.ncbi.nlm.nih.gov/20871896/)
83. Iwasaki Y, Hirasawa T, Maruyama Y, Ishii Y, Ito R, Saito K, et al. Effect of interaction between phenolic compounds and copper ion on antioxidant and pro-oxidant activities. *Toxicology in Vitro*. 2011; 25(7):1320–7. doi: [10.1016/j.tiv.2011.04.024](https://doi.org/10.1016/j.tiv.2011.04.024) PMID: [21600975](https://pubmed.ncbi.nlm.nih.gov/21600975/)
84. Liochev SI, Fridovich I. The Haber-Weiss cycle—70 years later: an alternative view. *Redox report*. 2002; 7(1):55–7. doi: [10.1179/135100002125000190](https://doi.org/10.1179/135100002125000190) PMID: [11981456](https://pubmed.ncbi.nlm.nih.gov/11981456/)
85. Speisky H, Gómez M, Burgos-Bravo F, López-Alarcón C, Jullian C, Olea-Azar C, et al. Generation of superoxide radicals by copper–glutathione complexes: redox-consequences associated with their interaction with reduced glutathione. *Bioorganic & medicinal chemistry*. 2009; 17(5):1803–10.
86. Chaturvedi KS, Hung CS, Crowley JR, Stapleton AE, Henderson JP. The siderophore yersiniabactin binds copper to protect pathogens during infection. *Nature chemical biology*. 2012; 8(8):731–6. doi: [10.1038/nchembio.1020](https://doi.org/10.1038/nchembio.1020) PMID: [22772152](https://pubmed.ncbi.nlm.nih.gov/22772152/)
87. Perron NR, García CR, Pinzón JR, Chaur MN, Brumaghim JL. Antioxidant and prooxidant effects of polyphenol compounds on copper-mediated DNA damage. *Journal of inorganic biochemistry*. 2011; 105(5):745–53. doi: [10.1016/j.jinorgbio.2011.02.009](https://doi.org/10.1016/j.jinorgbio.2011.02.009) PMID: [21481816](https://pubmed.ncbi.nlm.nih.gov/21481816/)
88. Jomova K, Valko M. Advances in metal-induced oxidative stress and human disease. *Toxicology*. 2011; 283(2):65–87.

89. Andreini C, Bertini I, Cavallaro G, Holliday GL, Thornton JM. Metal ions in biological catalysis: from enzyme databases to general principles. *JBIC Journal of Biological Inorganic Chemistry*. 2008; 13(8):1205–18. doi: [10.1007/s00775-008-0404-5](https://doi.org/10.1007/s00775-008-0404-5) PMID: [18604568](https://pubmed.ncbi.nlm.nih.gov/18604568/)
90. Davies I, Musa M, Dormandy T. Measurements of plasma zinc Part I In health and disease. *Journal of clinical pathology*. 1968; 21(3):359–63. PMID: [5303355](https://pubmed.ncbi.nlm.nih.gov/5303355/)
91. Haase H, Hebel S, Engelhardt G, Rink L. The biochemical effects of extracellular Zn²⁺ and other metal ions are severely affected by their speciation in cell culture media. *Metallomics*. 2015; 7(1):102–11. doi: [10.1039/c4mt00206g](https://doi.org/10.1039/c4mt00206g) PMID: [25360687](https://pubmed.ncbi.nlm.nih.gov/25360687/)
92. Hosainzadeh A, Gharanfoli M, Saberi MR, Chamani J. Probing the interaction of human serum albumin with bilirubin in the presence of aspirin by multi-spectroscopic, molecular modeling and zeta potential techniques: Insight on binary and ternary systems. *Journal of Biomolecular Structure and Dynamics*. 2012; 29(5):1013–50. doi: [10.1080/073911012010525029](https://doi.org/10.1080/073911012010525029) PMID: [22292958](https://pubmed.ncbi.nlm.nih.gov/22292958/)
93. Nygaard SB, Larsen A, Knuhtsen A, Rungby J, Smidt K. Effects of zinc supplementation and zinc chelation on in vitro β -cell function in INS-1E cells. *BMC research notes*. 2014; 7(1):84.
94. Akahori A, Gabryelak T, Józwiak Z, Gondko R. Zinc induced damage to carp (*Cyprinus carpio* L.) erythrocytes in vitro. *IUBMB Life*. 1999; 47(1):89–98.