



Characterization of a male reproductive transcriptome for *Peromyscus eremicus* (Cactus mouse)

Lauren L. Kordonowy* and Matthew D. MacManes*

Department of Molecular, Cellular, and Biological Sciences, University of New Hampshire, Durham, NH, United States

*These authors contributed equally to this work.

ABSTRACT

Rodents of the genus *Peromyscus* have become increasingly utilized models for investigations into adaptive biology. This genus is particularly powerful for research linking genetics with adaptive physiology or behaviors, and recent research has capitalized on the unique opportunities afforded by the ecological diversity of these rodents. Well characterized genomic and transcriptomic data is intrinsic to explorations of the genetic architecture responsible for ecological adaptations. Therefore, this study characterizes the transcriptome of three male reproductive tissues (testes, epididymis and vas deferens) of *Peromyscus eremicus* (Cactus mouse), a desert specialist. The transcriptome assembly process was optimized in order to produce a high quality and substantially complete annotated transcriptome. This composite transcriptome was generated to characterize the expressed transcripts in the male reproductive tract of *P. eremicus*, which will serve as a crucial resource for future research investigating our hypothesis that the male Cactus mouse possesses an adaptive reproductive phenotype to mitigate water-loss from ejaculate. This study reports genes under positive selection in the male Cactus mouse reproductive transcriptome relative to transcriptomes from *Peromyscus maniculatus* (deer mouse) and *Mus musculus*. Thus, this study expands upon existing genetic research in this species, and we provide a high quality transcriptome to enable further explorations of our proposed hypothesis for male Cactus mouse reproductive adaptations to minimize seminal fluid loss.

Submitted 20 April 2016
Accepted 27 September 2016
Published 27 October 2016

Corresponding author
Lauren L. Kordonowy,
lauren.kordonowy@unh.edu

Academic editor
Ana Pavasovic

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.2617

© Copyright
2016 Kordonowy and MacManes

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Bioinformatics, Evolutionary Studies, Genetics, Genomics, Zoology

Keywords *Peromyscus eremicus*, Transcriptome, Genomics, Bioinformatics, Adaptation, Desert physiology, Cactus mouse, Reproduction

INTRODUCTION

The rapid infusion of novel bioinformatics approaches in the fields of genomics and transcriptomics has enabled the coalescence of the fields of genetics, physiology and ecology into innovative studies for adaptation in evolutionary biology. Indeed, studies on the biology of adaptation had previously been dominated by research painstakingly documenting morphological shifts associated with ecological gradients (e.g., in *Peromyscus*: Carleton, 1989; MacMillen & Garland, 1989). However, the discipline of bioinformatics has breathed new life into the field of adaptation biology. Specifically, while the morphological basis as well as the physiological mechanisms of adaptation have been explored for a variety of

species in extreme environments, the genetic underpinnings of these adaptations have only recently become a larger area of research (Hoekstra et al., 2006; Cheviron & Brumfield, 2011; Lorenzo et al., 2014; Guillen et al., 2015). High-throughput sequencing technology of model and non-model organisms (Ellegren, 2014) enables evolutionary biologists to conduct genome and transcriptome wide analyses and link patterns of gene selection with functional adaptations.

Studies on the genetic basis of adaptation have included a wide variety of taxa. For example, butterflies in the *Heliconius* genus have been a particularly effective study system for determining the genetic basis of pigmentation patterns, and there is evidence of interspecific introgression for genes enabling adaptive mimicry patterns (Hines et al., 2012; *Heliconius Genome Consortium*, 2012). In addition, a population genomic study in three-spine sticklebacks has elucidated many loci responsible for divergent adaptations from marine to freshwater environments (Jones et al., 2012). Furthermore, researchers have developed a list of candidate genes that have evolved in multiple populations of freshwater adapted three-spine sticklebacks (Hohenlohe et al., 2010). Another active area of adaptation genetics research focuses on species residing in extreme environments. High altitude adaptations to hemoglobin variants have been identified in multiple organisms, including humans (Lorenzo et al., 2014), several species of Andean ducks (McCracken et al., 2009a; McCracken et al., 2009b), and deer mice, *Peromyscus maniculatus* (Storz et al., 2010; Natarajan et al., 2015). The genetic pathways responsible for physiological adaptations to desert habitats remain enigmatic; however, considerable progress has been made developing candidate gene sets for future analyses (e.g., Guillen et al., 2015; MacManes & Eisen, 2014; Marra et al., 2012; Marra, Romero & DeWoody, 2014). Functional studies will stem from this foundational research aimed at identifying the genomic underpinnings of adaptations to extreme environments; yet, it is inherently challenging and critically important to demonstrate that specific loci are functionally responsible for adaptations (Storz & Wheat, 2010).

Rodents of the genus *Peromyscus* have been at the forefront of research elucidating the genetic basis for adaptation (reviewed in Bedford & Hoekstra, 2015). This diverse genus has served as an ideal platform for adaptation research spanning from the genetic basis of behavioral adaptations—such as complex burrowing in *Peromyscus polionotus* (Weber & Hoekstra, 2009; Weber, Peterson & Hoekstra, 2013)—to the loci responsible for adaptive morphology—such as coat coloration in *Peromyscus polionotus leucocephalus* (Hoekstra et al., 2006), and including kidney desert adaptations in *Peromyscus eremicus* (MacManes & Eisen, 2014). We are currently using *Peromyscus eremicus* as a model species for investigating the genetic bases of desert adaptations. This paper describes a crucial component of this research aim.

Initial steps toward understanding the genetics of adaptation must include the genomic and transcriptomic characterization of target study species (MacManes & Eisen, 2014). To this end, we assembled and characterized a composite transcriptome for three male reproductive tissues in the desert specialist, *P. eremicus*. This species is an exceptional example of desert adaptation, as individuals may live entirely without water access (Veal & Caire, 2001). MacManes & Eisen (2014) assembled transcriptomes from kidney, hypothalamus, liver, and testes of this species, and they identified several candidate genes potentially

underlying adaptive renal physiology. However, to our knowledge, potential physiological reproductive adaptations to water limitation have not been studied in this species or in other desert rodents. We hypothesize that male *P. eremicus* possess reproductive adaptations to mitigate water loss.

The adaptive kidney physiology in kangaroo rat species (genus: *Dipodomys*), which produce highly concentrated urine via a disproportionately long loop of Henle ([Schmidt-Nielsen et al., 1948](#); [Schmidt-Nielsen & Schmidt-Nielsen, 1952](#); [Vimtrup & Schmidt-Nielsen, 1952](#); [Urity et al., 2012](#)), has been well-established. We propose that there may also be reproductive adaptations to mitigate water loss in desert rodents. Recent findings pertaining to the genetic signatures of adaptive kidney function in desert rodents ([Marra et al., 2012](#); [Marra, Romero & DeWoody, 2014](#); [MacManes & Eisen, 2014](#)), suggest that such hypothesized reproductive adaptations may be detectable at the genetic level (should they exist) using similar comparative transcriptomic methods. We present this hypothesis in light of the mounting body of research for high rates of reproductive protein evolution (reviewed in [Swanson & Vacquier, 2002](#); [Ramm et al., 2014](#)), which we propose indicates that reproductive tissues may possess a significant capacity for evolving in response to strong selective pressures.

[Dewsbury \(1982\)](#) made the assertion that producing ejaculates incurs a cost for male mammals, which produce relatively high sperm counts, and his analysis utilizes *P. maniculatus* as a model. Ejaculation has also been demonstrated to be costly in *Mus musculus domesticus*; [Ramm & Stockley \(2007\)](#) found that mice are able to manipulate the quantity of sperm released in response to varying competition for females. *Peromyscus* and other rodents exhibit rapid evolution of testis-expressed proteins ([Turner, Chuong & Hoekstra, 2008](#)). In addition, the epididymal transcriptome of *M. musculus* shows evidence for positive selection among epididymis-specialized genes that are secreted, which the authors attribute to their putative evolutionary importance ([Dean, Good & Nachman, 2008](#)). Moreover, a recent analysis for sperm genes from multiple mouse strains found that sperm proteins involved in both motility and in sperm-egg interactions show signatures of positive selection, potentially facilitating evolutionary mechanisms for sperm-competition and sexual conflict ([Vicens, Luke & Roldan, 2014](#)). We therefore propose that both the inherently costly nature of producing ejaculate and the rapid evolution of genes in murine testes, epididymis, and sperm would be ideal for desert rodents to evolve ejaculate adaptations to limited water-availability.

In order to develop genomic resources that will allow us to begin to test our hypothesis related to reproductive water conservation, we developed a transcriptome comprising three male specific reproductive tissues in *P. eremicus*. Specifically, we assembled a composite reproductive transcriptome for the epididymis, testes, and vas deferens. These tissues were chosen because they constitute numerous physiological roles in spermatogenesis and in the generation and transportation of seminal fluids. We posit that one or more of these reproductive tissues possess phenotypic characteristics to mitigate water loss from seminal fluids. We also propose that any molecular mechanisms responsible for such an adaptation should be elucidated through future differential gene expression and comparative transcriptomic studies. However, the initial assembly and characterization of a comprehensive male reproductive tissue transcriptome in the current manuscript will be essential

for research studies exploring this hypothesized male *P. eremicus* reproductive desert adaptation. For example, this transcriptome will be instrumental in further experimental studies investigating differential gene expression in these tissue types in response to variable hydration levels or in large-scale comparative transcriptomic studies spanning numerous desert and non-desert rodents.

Here, we characterize a male Cactus mouse tissue-specific transcriptome by presenting a preliminary exploration of transcript abundance and putative homology. We also perform comparative transcriptomic analyses to identify evidence of positive selection in genes potentially related to the hypothesized reproductive desert adaptations. It is beyond the scope of this manuscript to evaluate the functionality of these candidate genes in the context of desert adaptation, much less male specific reproductive adaptations. However, the elucidation of candidate genes in the context of male reproductive tissues will be instrumental for future studies aimed at determining which genes are functionally responsible for the proposed reproductive adaptations to water limitation in male Cactus mouse.

METHODS

Tissue samples, RNA extraction, cDNA library preparation and sequencing

The *Peromyscus eremicus* male used for this study was captive born and descendant from a population from the *Peromyscus* Genetic Stock Center (Columbia, South Carolina). This individual was housed in a facility at the University of New Hampshire designed to mimic desert conditions in the Southwestern United States. Specifically, the temperature increases gradually during the light hours until it peaks at 90° Fahrenheit in the afternoon, and the temperature decreases during hours of darkness to 75°. Humidity levels are 10% during the daylight hours and 25% during darkness. The photoperiodic cycle in this desert chamber is for long days of photo-stimulation, with 16 h of light, and 8 h of darkness. The colony includes males and females which are housed within a single room, providing olfactory cues that stimulate reproductive maturity. The photoperiod and shared housing in this colony result in the attainment of reproductive maturity in both sexes. Males are deemed reproductively mature when they are fully scrotal. The males do not undergo seasonal testicular atrophy, as evidenced by their consistent scrotal condition and their year-round successful reproduction.

A single reproductively mature *P. eremicus* male was sacrificed via isoflurane overdose and decapitation. This was done in accordance with University of New Hampshire Animal Care and Use Committee guidelines (protocol number 130902) and guidelines established by the American Society of Mammalogists (*Sikes et al., 2011*). Testes, epididymis, and vas deferens were immediately harvested (within ten minutes of euthanasia), placed in RNAlater (Ambion Life Technologies) and stored at –80 °C until RNA extraction. We used a standard TRIzol, chloroform protocol for total RNA extraction (Ambion Life Technologies). We evaluated the quantity and quality of the RNA product with a Qubit 2.0 Fluorometer (Invitrogen) and a Tapestation 2200 (Agilent Technologies, Palo Alto, USA).

We used a TURBO DNase kit (Ambion) to eliminate DNA from the samples prior to the library preparation. Libraries were made with a TruSeq Stranded mRNA Sample Prep

LS Kit (Illumina). Each of the three samples was labeled with a unique hexamer adapter for identification after multiplex single lane sequencing. Following library completion, we confirmed the quality and quantity of the DNA product with the Qubit and TapeStation. We submitted the multiplexed sample of the libraries for running on a single lane at the New York Genome Center Sequencing Facility (NY, New York). Paired end sequencing reads of length 125 bp were generated on an Illumina 2500 platform. Reads were parsed by tissue type according to their unique hexamer IDs in preparation for transcriptome assembly.

Reproductive transcriptome assembly

The composite reproductive transcriptome was assembled with reads from the testes, epididymis and vas deferens using the previously developed Oyster River Protocol for *de novo* transcriptome assembly pipeline (MacManes, 2016). Briefly, the reads were error corrected with Rcorrector v1.0.1 (Song & Florea, 2015). We used the *de novo* transcriptome assembler Trinity v2.1.1 (Haas et al., 2013; Grabherr et al., 2011). Within the Trinity platform, we ran Trimmomatic (Bolger, Lohse & Usadel, 2014) to remove the adapters, and we also trimmed at PHRED < 2, as recommended by MacManes (2014).

Next we evaluated transcriptome assembly quality and completeness using BUSCO v1.1b1 and Transrate v1.0.1. BUSCO (Simão et al., 2015) reports the number of complete, fragmented, and missing orthologs in assembled genomes, transcriptomes, or gene sets relative to compiled ortholog databases. We ran BUSCO on the assembly using the ortholog database for vertebrates, which includes 3,023 genes. The assembly was also analyzed by Transrate using the *Mus musculus* peptide database from Ensembl (downloaded 2/24/16) as a reference. The Transrate score provided a metric of *de novo* transcriptome assembly quality, and the software also generated an improved assembly comprised of highly supported contigs (Smith-Unna et al., 2016). Finally, we re-ran BUSCO on the improved assembly generated by Transrate to determine if this assembly had similar metric scores for completeness as the original assembly produced by Trinity. As alternatives to the original Trinity assembly and the optimized Transrate assembly, we proceeded with our optimization determinations by filtering out low abundance contigs from the original Trinity assembly. First we calculated the relative abundance of the transcripts with Kallisto v0.42.4 and Salmon v0.5.1. Kallisto utilizes a pseudo-alignment algorithm to map RNA-seq data reads to targets for transcript abundance quantification (Bray et al., 2015). In contrast, Salmon employs a lightweight quasi-alignment method and a high speed streaming algorithm to quantify transcripts (Patro, Duggal & Kingsford, 2015). After determining transcript abundance in both Kallisto and Salmon, we removed contigs with transcripts per million (TPM) estimates of less than 0.5 and of less than 1.0 in two separate optimization trials (as per MacManes, 2016). Finally, we evaluated these two filtered assemblies with Transrate and BUSCO to determine the relative quality and completeness of both assemblies. We chose the optimal assembly version by comparing Transrate and BUSCO metrics and also through careful consideration of total contig numbers across all filtering and optimizing versions. The chosen assembly was the Transrate optimized TPM > 0.5 filtered assembly, and this assembly was used for all subsequent analyses.

Annotation, transcript abundance, and database searches

We used dammit v0.2.7.1 (Scott, 2016) to annotate the optimized transcriptome assembly (as per MacManes, 2016). Within the dammit platform, we predicted protein coding regions for each tissue with TransDecoder v2.0.1 (Haas et al., 2013), which was used to find open reading frames (ORFs). Furthermore, dammit utilizes multiple database searches for annotating transcriptomes. These database searches include searches in Rfam v12.0 to find non-coding RNAs (Nawrocki et al., 2015), searches for protein domains in Pfam-A v29.0 (Sonnhammer, Eddy & Durbin, 1997; Finn et al., 2016), the execution of a LAST search for known proteins in the UniRef90 database (Suzek et al., 2007; Suzek et al., 2015), ortholog matches in the BUSCO database, and orthology searches in OrthoDB (Kriventseva et al., 2015).

We used the assembly annotated by dammit to re-run Kallisto to determine transcript abundance within each of the three tissue types. We used TPM counts of expression for all three tissues to generate counts of transcripts specific to and shared across tissue types. We also downloaded the ncRNA database for *Mus musculus* from Ensembl (v 2/25/16), and we did a BLASTn (Altschul et al., 1990; Madden, 2002) search for these ncRNAs in our assembly. This database has 16,274 sequences, and we determined the number of transcript ID matches and the number of unique ncRNA sequence matches for our assembly. We also counted how many transcript matches were present in each of the tissues, and we referenced the corresponding Kallisto derived TPM values to determine the number of unique and ubiquitous transcript matches for each tissue.

We searched the annotated assembly for transporter protein matches within the Transporter Classification Database (tcd.org). This database has 13,846 sequences representing proteins in transmembrane molecular transport systems (Saier et al., 2014). We executed a BLASTx (Altschul et al., 1990; Madden, 2002) search to find the number of transcript ID matches and the number of unique transporter protein matches within the assembly. Next we determined how many transcript ID matches were found in each of the three tissues. As previously described above, we also cross-referenced these matches with the Kallisto derived TPM values to find the number of ubiquitous and unique transcript matches by tissue type.

Comparative analysis for genes under positive selection

We performed a three-species comparative analysis to identify genes under positive selection in the male reproductive transcriptome for the *P. eremicus* lineage relative to *M. musculus* and *P. maniculatus*. The *M. musculus* nucleotide and protein sequences were downloaded (version GRCm38) from Ensembl (ensembl.org). The *P. maniculatus* nucleotide and protein files were downloaded (version GCF_000500345.1_Pman_1.9) from NCBI (ncbi.nlm.nih.gov). The comparative analysis was conducted as described below. The corresponding nucleotide and protein files for all three species were modified so that the header names for the sequences in each species file pair were concordant. Next, we found orthologous groups of protein sequences using OrthoFinder v0.6.1 (Emms & Kelly, 2015). The two OrthoFinder command scripts aligned sequences with MAFFT v7.123b (Katoh et al., 2002), built trees with FastTreeMP v1 (Price, Dehal & Arkin, 2010), and generated orthogroups

based on sequence similarity. Our next script selected the single copy orthologs (SCOs) from among these orthogroups for analysis. Then we selected the cds file transcripts for all three species corresponding to the previously identified single copy orthologs. Finally, we ran a script which aligned the sequences with PRANK v150803 (Löytynoja, 2014) and performed the analyses for positive selection for the single copy orthologous sequences with codeml in PAML v4.8 (Yang, 1997; Yang, 2007). Specifically, we performed the *M2a* branch site test (Zhang, Nielsen & Yang, 2005) for positive selection after stipulating *P. eremicus* as the foreground lineage. Genes were deemed under positive selection if the omega values (ω) exceeded 1 within the *M2a* model and if they yielded statistically significant results for the likelihood ratio test (LRT) comparison between the null and alternative models. To perform the LRT, we determined if $2\Delta\ln L$ (two times the difference between the log likelihood values for the alternative—*M2a*—and the null—*M1a*—models) exceeded a chi-square value corresponding to a significant p -value ($p < 0.05$) (Yang, 1998) after applying the Benjamini-Hochburg correction for multiple comparisons (Benjamini & Hochberg, 1995). For each of the *P. eremicus* gene sequences demonstrated to be under positive selection according to the LRT, we selected the *M. musculus* cds sequence from the SCO group corresponding to the *P. eremicus* gene sequence. Then we performed a BLASTn search on the *M. musculus* sequence to find gene matches on NCBI to determine the gene identity for the *P. eremicus* sequence under positive selection.

Of note, the code for performing all of the above analyses can be found at GitHub (https://github.com/macmanes-lab/peer_reproductive/transcriptome). The data files are available on Dryad (doi: [10.5061/dryad.01c3t](https://doi.org/10.5061/dryad.01c3t)).

RESULTS AND DISCUSSION

Reproductive transcriptome assembly

There were 45–94 million paired reads produced for each of the three transcriptome datasets, yielding a total of 415,960,428 reads. The raw reads are available at the European Nucleotide Archive under study accession number [PRJEB13364](https://www.ebi.ac.uk/ena/record/PRJEB13364).

We assembled a *de novo* composite reproductive transcriptome with reads from testes, epididymis and vas deferens. The evaluation of alternative optimized assemblies allowed us to generate a substantially complete transcriptome of high quality. The alternative assemblies had raw Transrate scores ranging from 0.156–0.194 (Table 1). However, the scores for the improved assemblies generated by Transrate, consisting of only highly supported contigs, ranged between 0.285 and 0.349, which is well above the threshold Transrate score of 0.22 for an acceptable assembly. The BUSCO results indicated that the assemblies were highly complete, with complete matches ranging from 73–90% of vertebrate orthologs (Table 2). These BUSCO benchmark values are consistent with the most complete reported assessments for transcriptomes from other vertebrate taxa (busco.ezlab.org). Furthermore, our BUSCO values exceed that of the only available reported male reproductive tissue (from a coelacanth: *Latimeria menadoensis* testes), which was 71% complete (Simão et al., 2015). The assembly version with the highest quality in relation to the Transrate metrics was the Transrate optimized Trinity assembly.

Table 1 Transrate results for the reproductive transcriptome assembly produced by different optimization methods.

Assembly	Transrate score	Optimized score ^a	# Read pairs (fragments)	Contigs (n_seqs)	# Good contigs	% Good contigs
Trinity original	0.1944	0.3492	207,980,214	856,711	657,952	0.77
Filter TPM < 0.5	0.1672	0.3013	207,980,214	147,966	78,424	0.53
Filter TPM < 1.0	0.156	0.2854	207,980,214	80,165	54,140	0.68

Notes.

^aThis is the score of the Transrate optimized assembly in Table 2.

Table 2 BUSCO metrics for the reproductive transcriptome assembly produced by different optimization methods.

Assembly	% Complete	% Duplicated	% Fragmented	% Missing
Trinity original	90	49	3.4	5.5
Transrate optimized	85	44	4.3	9.7
Filter TPM < 0.5	85	38	3.0	11
Transrate TPM < 0.5	73	31	3.9	22
Filter TPM < 1.0	80	28	2.8	16
Transrate TPM < 1.0	74	25	3.4	21

Specifically, the optimized Transrate score was 0.3492, and the percent coverage of the reference assembly was also highest, with 45% of the mouse database represented. This assembly was highly competitive for completeness, as indicated by the fact that it contained 85% of vertebrate single copy orthologs. However, this assembly had an exorbitantly high number of contigs (657,952 contigs), which is nearly an order of magnitude more contigs than the next best performing assembly: the Transrate optimized TPM > 0.5 filtered assembly (78,424 contigs). In consideration of the dramatically more realistic contig number for the Transrate optimized TPM > 0.5 filtered assembly, and in light of its second best performance for Transrate score (0.3013), reasonable Transrate mouse reference assembly coverage (37%), and sufficiently high BUSCO completeness (73% orthologs found), we chose this assembly as our optimized transcriptome. Therefore, we proceeded with this optimized assembly version as our finalized transcriptome assembly for our analyses.

Annotation, transcript abundance and database searches

The reproductive transcriptome assembly annotations were produced by dammit, and they are available on Dryad in a gff3 file format. Furthermore, TransDecoder was used to predict coding regions in the assembly. TransDecoder predicted that 49.5% (38,342) of the transcripts (78,424 total) contained ORFs, of which 63.9% (24,808) had complete ORFs containing a start and stop codon. The predicted protein coding regions generated by TransDecoder are reported in five file types, and they are available on Dryad. Furthermore, the Pfam results yielded 30.7% of transcripts (24,107) matching to the protein family database. In contrast, the LAST search found that 75.9% of transcripts (59,503) matched to the UniRef90 database. We have uploaded the homology search results generated by Pfam and UniRef90 matches onto Dryad. In addition, 1.04% (816) of transcripts matched to the



Figure 1 Venn Diagram of transcript expression differences and similarities between the three reproductive tissues for a single male mouse. The total number of transcripts is 78,424.

Rfam database for ncRNAs, and these results are posted in Dryad. Of note, 80.1% (62,835) of the transcripts were annotated using one or more of the above described methods (the dammit.gff3 file is posted in Dryad), and it is this final annotated assembly that was used for all subsequent analyses (this annotated transcriptome is available on Dryad).

The Kallisto generated TPM counts of expression (available on Dryad) were utilized to determine which transcripts were ubiquitous and specific to the three tissue types, which we have depicted in a Venn diagram format (Fig. 1). The assembly consisted of 78,424 different transcript IDs, of which 64,553 were shared across all three tissues. The number of unique transcripts were as follows: 3,563 in testes, 342 in epididymis, and 502 in vas deferens. The relatively large number of unique transcripts in the testes is consistent with previous findings which describe the testes as the tissue with the highest number of tissue-enriched genes in the human body (Djureinovic et al., 2014; Uhlén et al., 2015). However, because this expression data was generated by a single individual, we want to emphasize that these results have no statistical power. Rather, we view these numerical comparisons of unique and ubiquitous transcript counts as an exploratory evaluation of potential relationships between tissue types. Such comparisons across tissue types for a single individual of this species previously indicated that the kidney had relatively higher numbers of unique transcripts than testes (MacManes & Eisen, 2014). Future research with multiple individuals will be necessary to statistically evaluate the relative rates of transcript expression between these reproductive tissues, as well as their relationship with other non-reproductive tissue types.

In addition, we searched for *Mus musculus* ncRNA sequence matches within our assembly. There were 15,964 transcript matches, which correspond to 2,320 unique




Figure 2 Venn Diagram of transcript matches between the three reproductive tissues to ncRNA sequences in *Mus musculus*. The total number of transcript matches across the tissue types is 15,964.

ncRNA matches, and they are posted on Dryad. The transcript matches by tissue type were found using the Kallisto TPM determinations, and they were as follows, testes: 15,260, epididymis: 15,552, and vas deferens: 15,558. A Venn Diagram depicts unique and shared transcript matches by tissue type (Fig. 2). The majority of transcript matches were ubiquitous to all three tissues (14,724), and there were far fewer tissue specific matches. The testes had more unique transcript matches (185) than the epididymis (26) or the vas deferens (45). These findings are consistent with our results above regarding the relative numbers of total unique transcripts in the assembly by tissue type. However, these counts for relative transcript matches among tissue types were generated with transcripts from a single individual; therefore, the comparative results across tissue type were not statistically evaluated. It is seemingly probable that the diversity of transcripts for regulation should be highest in tissues generating relatively diverse proteins, in this case, the testes, which did have the highest number of unique transcript matches. The role of ncRNA in reproductive tissues throughout multiple developmental stages has recently been reviewed in detail (Hale, Yang & Ross, 2014). In addition, ncRNAs have been found to be highly abundant in murine testes (Sun, Lin & Wu, 2013). Furthermore, sperm from humans and mice contain a significant number of ncRNAs (Krawetz et al., 2011; Kowano et al., 2012). However, we are unaware of any research investigating the involvement of ncRNA in desert adaptations; therefore, we cannot speculate on particular ncRNA matches within our dataset that may have potential desert adaptive roles.

Our search for transporter protein matches within the Transporter Classification Database yielded 7,521 different transcript matches, corresponding to 1,373 unique transporter protein matches, and they are posted on Dryad. The number of transcript




Figure 3 Venn Diagram of transcript matches between the three reproductive tissues to protein sequences in the Transporter Classification Database. The total number of transcript matches across the tissue types is 7,521.

matches was highly similar between the tissue types (testes 7,025; epididymis 7,115; vas deferens: 7,071). We generated a Venn Diagram to display the numbers of shared and unique transcript matches to the transporter protein sequences (Fig. 3). Most transcript matches were present in all three tissues (6,472), and there were relatively few unique matches in the three tissue types. However, the testes had the highest number of unique transcript matches (215) relative to the epididymis (19) and the vas deferens (37). These comparative results across tissue types represent exploratory findings for a single mouse, and the count data have not been statistically tested, but these preliminary results should be investigated in future studies. Furthermore, our BLASTx search of this transporter protein database yielded transcript matches for multiple solute carrier proteins. We are particularly interested in solute carrier proteins because previous research has found candidate genes in this protein family for desert adaptations in kidneys of the kangaroo rat (Marra *et al.*, 2012; Marra, Romero & DeWoody, 2014) and the Cactus mouse (MacManes & Eisen, 2014). In addition, we had multiple matches to aquaporins, which are water channels allowing transport across cellular membranes. One transcript matched specifically to Aquaporin 3, a sperm water channel found in mice and humans, which is essential to maintaining sperm cellular integrity in response to the hypotonic environment within the female reproductive tract (Chen *et al.*, 2011).

Comparative analysis for genes under positive selection

To find evidence for genes undergoing positive selection in the male reproductive transcriptome of *P. eremicus*, we compared this species with two other generalist rodents. We chose

M. musculus as the non-desert adapted outgroup because this species possesses transcriptomic resources which are exceptional in their annotation and completeness. The widely distributed ([Carleton, 1989](#)) habitat generalist deer mouse, *P. maniculatus*, was chosen because it harbors the most complete transcriptomic data available among the *Peromyscus* genus.

There were 3,731 panorthologous groups (single copy orthologs) in our three species comparison. The branch test was successfully implemented for SCOs when all three sequences aligned adequately with PRANK and when codeml produced both *M1a* and *M2a* output files for the LRT comparison ($n = 2,820$ in total). The *M2a* test indicated that 42 genes were evolving under a model of positive selection in the Cactus mouse ([Table 3](#)). Therefore, we investigated whether previous research on either rodent reproductive tissues or on desert specialized rodents documented evidence of positive selection for any of these genes or gene families. Only one of these 42 genes matched an epididymis-specialized secreted gene undergoing positive selection in another rodent (C57/BL6 mice). Namely, *Qsox2* is a match for *Qscn6l1*, a member of the sulfhydryl oxidase/quiescin-6 family, which is purportedly involved in neuroblastoma apoptosis ([Dean, Good & Nachman, 2008](#)). However, we cannot speculate regarding the functionality this gene has in male rodent reproductive tissue, or why it appears to be evolving under a model of positive selection in these two studies. Another of our 42 positively selected genes, *Lrrc46*, may have some similarity to *Lrrc50*, a gene under positive selection in testes of *Peromyscus* ([Turner, Chuong & Hoekstra, 2008](#)). Leucine-rich repeat containing (*Lrrc*) genes have diverse biological roles; therefore, we also will not speculate on any correspondence between these two genes. As expected given our experimental design, there was no concordance between our 42 genes and those found to be under positive selection in a recent study on mouse spermatozoa proteins ([Vicens, Luke & Roldan, 2014](#)).

Our search for gene matches from the current study with other desert rodent research revealed notable similarities. Two solute carrier proteins, *Slc15a3* and *Slc47a1*, were found to be under positive selection in our analysis. This finding bears particular significance because another protein in this family, *Slc2a9*, shows signatures of positive selection in desert rodent kidney transcriptomes in *Dipodomys spectabilis* and *Chaetodipus baileyi* ([Marra, Romero & DeWoody, 2014](#)) and *P. eremicus* ([MacManes & Eisen, 2014](#)). Solute carriers are a large family of cell membrane proteins that are responsible for transporting solutes (reviewed in [Hediger et al., 2004](#); [Hediger et al., 2013](#); [César-Razquin et al., 2015](#)). Furthermore, [Marra, Romero & DeWoody \(2014\)](#), hypothesize that solute carriers are critical for osmoregulation in desert rodents, and they assert that these genes may be under evolutionary pressure in such rodents. In response to the potential relevance of the two solute carriers under positive selection in our study to desert rodent osmoregulation, we generated STRING ([Snel et al., 2000](#); [Szklarczyk et al., 2015](#)) diagrams for their protein-protein interactions ([Fig. 4](#)). These diagrams demonstrate multiple connections to other solute carriers for both proteins, thereby suggesting their potential functional roles.

This analysis of genes undergoing positive selection in *P. eremicus* relative to *P. maniculatus* and *M. musculus* provides candidate genes for desert specialization in the Cactus mouse which can be the target of future studies focused on ascertaining which genes may be functionally responsible for our hypothesized male reproductive desert adaptation.

Table 3 The 42 genes that reached statistical significance ($p < 0.05$) after correcting for multiple hypothesis testing for the *M2a* branch-site test for positive selection in PAML in the male Cactus mouse reproductive transcriptome.

Orthogroup ID	<i>p</i> -value	BLASTn description
OG0010592	1.18E-10	Q6 sulfhydryl oxidase 2 (Qsox2)
OG0010774	1.04E-03	Adenylate kinase 6 (Ak6)
OG0010833	4.90E-05	Plakophilin 2 (Pkp2)
OG0011177	3.15E-03	1-acylglycerol-3-phosphate O-acyltransferase 2 (Agpat2)
OG0011272	1.06E-04	cDNA sequence BC089491
OG0011374	2.83E-03	Hepatoma derived growth factor-like 1 (Hdglf1)
OG0011384	2.05E-03	Chitinase, acidic 1 (Chia1)
OG0011551	7.05E-04	Zinc finger protein 770 (Zfp770)
OG0011784	1.63E-04	Zinc finger, DHHC domain containing 19 (Zdhhc19)
OG0011914	1.33E-09	Fumarylacetoacetate hydrolase domain containing 2A (Fahd2a)
OG0012115	0.00E+00	NUT midline carcinoma, family member 1 (Nutm1)
OG0012232	2.06E-03	Leucine rich repeat neuronal 2 (Lrrn2)
OG0012396	1.19E-06	Leucine rich repeat containing 46 (Lrrc46)
OG0012449	2.79E-04	F-box and leucine-rich repeat protein 14 (Fbxl14)
OG0012511	3.27E-08	SMAD family member 6 (Smad6)
OG0012690	0.00E+00	Solute carrier family 47, member 1 (Slc47a1)
OG0012869	4.59E-02	Bromodomain and WD repeat domain containing 3 (Brwd3)
OG0013171	5.47E-06	ArfGAP with SH3 domain, ankyrin repeat and PH domain 3 (Asap3)
OG0013288	6.09E-03	Phosphodiesterase 3B, cGMP-inhibited (Pde3b)
OG0013304	3.42E-02	Persephin (Pspn)
OG0013342	4.11E-02	Mitochondrial methionyl-tRNA formyltransferase (Mtfmt)
OG0013590	1.77E-02	Matrix metalloproteinase 15 (Mmp15)
OG0013771	1.77E-02	Neuronal pentraxin 2 (Nptx2)
OG0013841	4.34E-03	Transformation related protein 63 regulated like (Tprgl)
OG0014000	1.25E-02	Alkaline phosphatase, liver/bone/kidney (Alpl)
OG0014048	0.00E+00	BPI fold containing family A, member 5 (Bpifa5)
OG0014062	1.42E-02	SPARC related modular calcium binding 2 (Smoc2)
OG0014193	1.05E-06	Carbonic anhydrase 11 (Car11)
OG0014286	9.65E-03	cDNA 4930550C14 gene
OG0014309	7.40E-03	Chymotrypsin-like elastase family, member 1 (Cela1)
OG0014333	2.11E-02	Minichromosome maintenance 9 homologous recombination repair factor (Mcm9)
OG0014414	1.60E-04	Growth arrest specific 6 (Gas6)
OG0014634	2.62E-03	Glutathione peroxidase 2 (Gpx2)
OG0014827	0.00E+00	Secretory carrier membrane protein 5 (Scamp5)
OG0014913	4.34E-04	STIP1 homology and U-Box containing protein 1 (Stub1)
OG0015025	3.23E-08	Pancreatic lipase-related protein 2 (Pnlipr2)
OG0015310	1.06E-03	Claspin (Clspn)
OG0015466	7.62E-05	Excision repair cross-complementing rodent repair deficiency complementation group 6 like (Ercc6l)

(continued on next page)

Table 3 (continued)

Orthogroup ID	p-value	BLASTn description
OG0015627	4.18E-03	Solute carrier family 15, member 3 (Slc15a3)
OG0015662	1.52E-03	Epithelial membrane protein 2 (Emp2)
OG0015704	3.96E-03	Cubilin (intrinsic factor-cobalamin receptor) (Cubn)
OG0015713	4.97E-07	Suppressor APC domain containing 1 (Sapcd1)

However, desert specialization is not the sole difference between *P. eremicus* and *P. maniculatus*, much less between *P. eremicus* and *M. musculus*. *P. maniculatus* is highly promiscuous, while *P. eremicus* is relatively socially monogamous (Wolff, 1989). Indeed, *P. maniculatus* has been the subject of considerable sperm competition research (Dewsbury, 1988; Fisher & Hoekstra, 2010). Differences in reproductive systems, and even potentially in sperm aggregation of *P. maniculatus* (Fisher & Hoekstra, 2010), may manifest themselves as evidence of selection patterns between these two *Peromyscus* species. Therefore, we are not proposing that the 42 genes we found to be under positive selection in the male Cactus mouse are functionally responsible for adaptive desert physiology. Rather, we are proposing that they are interesting candidate genes for future studies investigating the genetic underpinnings of physiological desert adaptations, including our hypothesized male reproductive adaptation, on a functional level. Several of these genes, specifically those in the solute carrier family, seem particularly promising for such work because they are undergoing rapid evolution in multiple desert rodent species.

CONCLUSIONS

Although researchers have determined that renal adaptations are responsible for mitigating water loss in kangaroo rats via the genitourinary tract (Schmidt-Nielsen et al., 1948; Schmidt-Nielsen & Schmidt-Nielsen, 1952; Vimtrup & Schmidt-Nielsen, 1952; Urity et al., 2012), we present the novel hypothesis that there may also be male reproductive adaptations to arid environments that allow desert specialists like the Cactus mouse to conserve water during reproduction. Previous efforts to elucidate the genomic basis of desert adaptations have described candidate genes for adaptive renal physiology in some desert specialized rodents (Marra et al., 2012; Marra, Romero & DeWoody, 2014), including the Cactus mouse (MacManes & Eisen, 2014). In light of these findings, we propose that if the male Cactus mouse possesses an adaptive reproductive phenotype to mitigate water loss via seminal fluids in response to limited-water availability, such an adaptation will be detectable through transcriptomic analyses. The current study generates and characterizes a transcriptome for male reproductive tissues from the Cactus mouse as an initial step towards future efforts to explore this hypothesized reproductive adaptation. This study describes a composite transcriptome from three male reproductive tissues in the desert specialist *Peromyscus eremicus*. Our analyses include quality and completeness assessments of this reproductive assembly, which we generated using reads from testes, epididymis, and vas deferens of a male Cactus mouse. We generate annotations and search relevant databases for ncRNAs and transporter protein sequences. We also describe the degree of ubiquity between transcripts among the three tissues as well as identify transcripts unique to those tissues utilizing




Figure 4 STRING diagrams of protein interactions for two proteins evolving under a model of positive selection in *P. eremicus*: Slc15a3 and Slc47a1.

preliminary (based on a single individual) transcript differences between tissue types. Furthermore, we find genes evolving under a model of positive selection in the *P. eremicus* male reproductive transcriptome relative to *P. maniculatus* and *M. musculus* in order to generate a list of candidate genes for future investigations in desert adaptation genetics. Our future research will investigate the hypothesized male reproductive physiological adaptation to water limitation in Cactus mouse through a differential gene expression study, and the characterization of this reproductive transcriptome will form the foundation of studies along this vein. Moreover, this research contributes transcriptomic materials to a larger body of work in the expanding field of adaptation genetics, which benefits tremendously from enhanced opportunities for comparative analyses.

DRYAD DATA FILE LIST

Final Annotated Reproductive Tissue Transcriptome: *reproductive.annotated.fasta* (127 MB)

Transdecoder (Five Files):

transdecoder.gff3 (49 MB)

transdecoder.pep (27 MB)

transdecoder.cds (62 MB)

transdecoder.mRNA (172 MB)

transdecoder.bed (10 MB)

Pfam Annotation: *reproductive.pfam.gff3* (32 MB)

Rfam Annotation: *reproductive.rfam.gff3* (191 KB)

Dammit Annotation: *reproductive.dammit.gff3* (98 MB)

UniRef90 Annotation: *reproductive.uniref.gff3* (9.5 MB)

Kallisto Results for Annotated Transcriptome (Three Files):

kallisto.testes.tsv (3.4 MB)

kallisto.epi.tsv (3.4 MB)

kallisto.vas.tsv (3.4 MB)

ncRNA Database Matches (three files):

epi.tpm.plus.ncRNA.txt (4.3 MB)

vas.tpm.plus.ncRNA.txt (4.3 MB)

testes.tpm.plus.ncRNA.txt (4.3 MB)

tcdB Database Matches (three files):

epi.tpm.plus.tcdB.txt (946 KB)

vas.tpm.plus.tcdB.txt (946 KB)

testes.tpm.plus.tcdB.txt (946 KB)

Comparative analysis for genes under positive selection (three files):

PAMLresults.txt (354 KB)

significant_orthogroups.txt (6 KB)

42SeqsM2aPosSel.txt (89 KB).

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The work was funded by startup funds from the University of New Hampshire. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
University of New Hampshire.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Lauren L. Kordonowy conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Matthew D. MacManes conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

University of New Hampshire Animal Care and Use Committee: (protocol number 130902).

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

European Nucleotide Archive ([PRJEB13364](#), ID: [318672](#)).

Data Availability

The following information was supplied regarding data availability:

Dryad DOI: [10.5061/dryad.01c3t](#); Github:

https://github.com/macmanes-lab/peer_reproductive-transcriptome.

REFERENCES

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* **215**(3):403–410

DOI [10.1016/S0022-2836\(05\)80360-2](#).

Bedford NL, Hoekstra HE. 2015. The natural history of model organisms: *Peromyscus* mice as a model for studying natural variation. *eLife* **4**:e06813

DOI [10.7554/eLife.06813](#).

- Benjamini Y, Hochberg Y. 1995.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**(1):289–300.
- Bolger AM, Lohse M, Usadel B. 2014.** Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics* **30**(15):2114–2120 DOI [10.1093/bioinformatics/btu170](https://doi.org/10.1093/bioinformatics/btu170).
- Bray N, Pimentel H, Melsted P, Pachter L. 2015.** Near-optimal RNA-seq quantification. ArXiv preprint. [arXiv:1505.02710](https://arxiv.org/abs/1505.02710).
- Carleton M. 1989.** Systematics and evolution. In: Kirkland Jr LG, Layne JN, eds. *Advances in the study of Peromyscus (Rodentia)*. Lubbock: Texas Tech University Press, 7–142.
- César-Razquin A, Snijder B, Frappier-Brinton T, Isserlin R, Gyimesi G, Bai X, Reithmeier RA, Hepworth D, Hediger MA, Edwards AM, Superti-Furga G. 2015.** A call for systematic research on solute carriers. *Cell* **162**(3):478–487 DOI [10.1016/j.cell.2015.07.022](https://doi.org/10.1016/j.cell.2015.07.022).
- Chen Q, Peng H, Lei L, Zhang Y, Kuang H, Cao Y, Shi Q, Ma T, Duan T. 2011.** Aquaporin 3 is a sperm water channel essential for postcopulatory sperm osmoadaptation and migration. *Cell Research* **21**(6):922–933 DOI [10.1038/cr.2010.169](https://doi.org/10.1038/cr.2010.169).
- Chevron ZA, Brumfield RT. 2011.** Genomic insights into adaptation to high-altitude environments. *Heredity* **108**(4):354–361 DOI [10.1038/hdy.2011.85](https://doi.org/10.1038/hdy.2011.85).
- Dean MD, Good JM, Nachman MW. 2008.** Adaptive evolution of proteins secreted during sperm maturation: an analysis of the mouse epididymal transcriptome. *Molecular Biology and Evolution* **25**(2):383–392 DOI [10.1093/molbev/msm265](https://doi.org/10.1093/molbev/msm265).
- Dewsbury DA. 1982.** Ejaculate cost and male choice. *The American Naturalist* **119**(5):601–610.
- Dewsbury DA. 1988.** A test of the role of copulatory plugs in sperm competition in deer mice (*Peromyscus maniculatus*). *Journal of Mammalogy* **69**(4):854–857 DOI [10.2307/1381648](https://doi.org/10.2307/1381648).
- Djureinovic D, Fagerberg L, Hallstrom B, Danielsson A, Lindskog C, Uhlen M, Ponten F. 2014.** The human testes-specific proteome defined by transcriptomics and antibody-based profiling. *Molecular Human Reproduction* **20**(6):476–488 DOI [10.1093/molehr/gau018](https://doi.org/10.1093/molehr/gau018).
- Ellegren H. 2014.** Genome sequencing and population genomics in non-model organisms. *Trends in Ecology and Evolution* **29**(1):1–13 DOI [10.1016/j.tree.2013.09.008](https://doi.org/10.1016/j.tree.2013.09.008).
- Emms DM, Kelly S. 2015.** OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biology* **16**:157 DOI [10.1186/s13059-015-0721-2](https://doi.org/10.1186/s13059-015-0721-2).
- Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016.** The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Research* **44**:D279–D285 DOI [10.1093/nar/gkv1344](https://doi.org/10.1093/nar/gkv1344).
- Fisher HS, Hoekstra HE. 2010.** Competition drives cooperation among closely related sperm of deer mice. *Nature* **463**(7282):801–803 DOI [10.1038/nature08736](https://doi.org/10.1038/nature08736).

- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, Di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29:644–652 DOI 10.1038/nbt.1883.
- Guillen Y, Rius N, Delprat A, Williford A, Muyas F, Puig M, Casillas S, Ramia M, Egea R, Negre B, Mir G, Camps J, Moncunill V, Ruiz-Ruano FJ, Cabrero J, De Lima LG, Dias GB, Ruiz JC, Kapusta A, Garcia-Mas J, Gut M, Gut IG, Torrents D, Camacho JP, Kuhn GCS, Feschotte C, Clark AG, Betran E, Barbadilla A, Ruiz A. 2015. Genomics of ecological adaptation in cactophilic drosophila. *Genome Biology and Evolution* 7(1):349–366 DOI 10.1093/gbe/evu291.
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, Macmanes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T, Dewey CN, Henschel R, Leduc RD, Friedman N, Regev A. 2013. *De novo* transcript sequence reconstruction from RNA-Seq: reference generation and analysis with Trinity. *Nature Protocols* 8(8):1494–1512 DOI 10.1038/nprot.2013.084.
- Hale BJ, Yang CX, Ross JW. 2014. Small RNA regulation of reproductive function. *Molecular Reproduction and Development* 81(2):148–159 DOI 10.1002/mrd.22272.
- Hediger MA, Clemenccon B, Burrier RE, Bruford EA. 2013. The ABCs of membrane transporters in health and disease (SLC series): introduction. *Molecular Aspects of Medicine* 34(2–3):95–107 DOI 10.1016/j.mam.2012.12.009.
- Hediger MA, Romero MF, Peng J-B, Rolfs A, Takanaga H, Bruford EA. 2004. The ABCs of solute carriers: physiological, pathological and therapeutic implications of human membrane transport proteins: introduction. *Pfugers Archiv* 447(5):465–548 DOI 10.1007/s00424-003-1192-y.
- Heliconius Genome Consortium. 2012. Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. *Nature* 487:94–98 DOI 10.1038/nature11041.
- Hines HM, Papa R, Ruiz M, Papanicolaou A, Wang C, Nijhout HF, McMillan WO, Reed RD. 2012. Transcriptome analysis reveals novel patterning and pigmentation genes underlying *Heliconius* butterfly wing pattern variation. *BMC Genomics* 13:288 DOI 10.1186/1471-2164-13-288.
- Hoekstra HE, Hirschmann RJ, Bunday RA, Insel PA, Crossland JP. 2006. A single amino acid mutation contributes to adaptive beach mouse color patterns. *Science* 313:101–104 DOI 10.1126/science.1126121.
- Hohenlohe PA, Bassham S, Etter PD, Stiffler N, Johnson EA, Cresko WA. 2010. Population genomics of parallel adaptation in threespine stickleback using sequenced RAD Tags. *PLoS Genetics* 6(2):e1000862 DOI 10.1371/journal.pgen.1000862.
- Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, Swofford R, Pirun M, Zody MC, White S, Birney E, Searle S, Schmutz J, Grimwood J, Dickson MC, Myers RM, Miller CT, Summers BR, Knecht AK, Brady SD, Zhang H, Pollen AA, Howes T, Amemiya C, Broad Institute Genome Sequencing Platform & Whole Assembly Team, Lander ES, Di Palma F, Lindblad-Toh K, Kingsley DM. 2012. The

- genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**:55–61 DOI [10.1038/nature10944](https://doi.org/10.1038/nature10944).
- Katoh K, Misawa K, Kuma K, Miyata T. 2002.** MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**(14):3059–3066 DOI [10.1093/nar/gkf436](https://doi.org/10.1093/nar/gkf436).
- Kowano M, Kawaji H, Grandejean V, Kiani J, Rassoulzadegan M. 2012.** Novel small noncoding RNAs in mouse spermatozoa, zygotes and early embryos. *PLoS ONE* **7**(9):e44542 DOI [10.1371/journal.pone.0044542](https://doi.org/10.1371/journal.pone.0044542).
- Krawetz SA, Kruger A, Lalancette C, Tagett R, Anton E, Draghici S, Diamond MP. 2011.** A survey of small RNAs in human sperm. *Human Reproduction* **26**(12):3401–3412 DOI [10.1093/humrep/der329](https://doi.org/10.1093/humrep/der329).
- Kriventseva EV, Tegenfeldt R, Petty TJ, Waterhouse RM, Simao FA, Pozdnyakov IA, Ioannidis P, Zdobnov EM. 2015.** OrthoDBv8: update of the hierarchical catalog of orthologs and the underlying free software. *Nucleic Acids Research* **43**:D250–D256 DOI [10.1093/nar/gku1220](https://doi.org/10.1093/nar/gku1220).
- Lorenzo FR, Huff C, Myllymäki M, Olenchock B, Swierczek S, Tashi T, Gordeuk V, Wuren T, Ri-Li G, McClain DA, Khan TM, Koul PA, Guchhait P, Salama ME, Xing J, Semenza GL, Liberzon E, Wilson A, Simonson TS, Jorde LB, Kaelin Jr WG, Koivunen P, Prchal JT. 2014.** A genetic mechanism for Tibetan high-altitude adaptation. *Nature Genetics* **46**(9):951–956 DOI [10.1038/ng.3067](https://doi.org/10.1038/ng.3067).
- Löytynoja A. 2014.** Phylogeny-aware alignment with PRANK. *Methods in Molecular Biology* **1079**:155–170 DOI [10.1007/978-1-62703-646-7_10](https://doi.org/10.1007/978-1-62703-646-7_10).
- MacManes MD. 2014.** On the optimal trimming of high-throughput mRNA sequence data. *Frontiers in Genetics* **5**:13 DOI [10.3389/fgene.2014.00013](https://doi.org/10.3389/fgene.2014.00013).
- MacManes MD. 2016.** Establishing evidenced-based best practice for the de novo assembly and evaluation of transcriptomes from non-model organisms. *bioRxiv* DOI [10.1101/035642](https://doi.org/10.1101/035642).
- MacManes MD, Eisen MB. 2014.** Characterization of the transcriptome, nucleotide sequence polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus*. *PeerJ* **2**:e642 DOI [10.7717/peerj.642](https://doi.org/10.7717/peerj.642).
- MacMillen RE, Garland Jr T. 1989.** Kirkland Jr LG, Layne JN, eds. *Advances in the study of peromyscus (Rodentia)*. Lubbock: Texas Tech University Press, 143–168.
- Madden T. 2002.** The BLAST sequence analysis tool. In: McEntyre J, Ostell J, eds. *The NCBI Handbook*. Bethesda: National Center for Biotechnology Information, Chapter 16. Available at <https://www.ncbi.nlm.nih.gov/books/NBK153387/>.
- Marra NJ, Eo SH, Hale MC, Waser PM, DeWoody JA. 2012.** *A priori* and *a posteriori* approaches for finding genes of evolutionary interest in non-model species: osmoregulatory genes in the kidney transcriptome of the desert rodent *Dipodomys spectabilis* (banner-tailed kangaroo rat). *Comparative Biochemistry and Physiology, Part D: Genomics Proteomics* **7**(4):328–339 DOI [10.1016/j.cbd.2012.07.001](https://doi.org/10.1016/j.cbd.2012.07.001).
- Marra NJ, Romero A, DeWoody A. 2014.** Natural selection and the genetic basis of osmoregulation in heteromyid rodents as revealed by RNA-seq. *Molecular Ecology* **23**(11):2699–2711 DOI [10.1111/mec.12764](https://doi.org/10.1111/mec.12764).

- McCracken KG, Barger CP, Bulgarella M, Johnson KP, Kuhner MK, Moore AV, Peters JL, Trucco J, Valqui TH, Winker K, Wilson RE. 2009a. Signatures of high-altitude adaptation in the major hemoglobin of five species of andean dabbling ducks. *The American Naturalist* 174(5):631–650 DOI 10.1086/606020.
- McCracken KG, Barger CP, Bulgarella M, Johnson KP, Sonsthagen SA, Trucco J, Valqui TH, Wilson RE, Winker K, Sorenson MD. 2009b. Parallel evolution in the major haemoglobin genes of eight species of Andean waterfowl. *Molecular Ecology* 18(19):3992–4005 DOI 10.1111/j.1365-294X.2009.04352.x.
- Natarajan C, Hoffman FG, Lanier HC, Wolf CJ, Cheviron ZA, Spangler ML, Weber RE, Fago A, Storz JF. 2015. Intraspecific polymorphism, interspecific divergence, and origins of function-altering mutation in deer mouse hemoglobin. *Molecular Biology and Evolution* 32(4):978–997 DOI 10.1093/molbev/msu403.
- Nawrocki EP, Burge SW, Bateman A, Daub J, Eberhardt RY, Eddy SR, Floden EW, Gardner PP, Jones TA, Tate J, Finn RD. 2015. Rfam 12.0: updates to the RNA families database. *Nucleic Acids Research* 43:D130–D137 DOI 10.1093/nar/gku1063.
- Patro R, Duggal G, Kingsford C. 2015. Salmon: accurate, versatile and ultrafast quantification from RNA-seq data using lightweight-alignment. *bioRxiv* DOI 10.1101/021592.
- Price M, Dehal P, Arkin A. 2010. FastTree2—approximately maximum-likelihood trees for large alignments. *PLoS ONE* 5(3):e9490 DOI 10.1371/journal.pone.009490.
- Ramm SA, Scharer L, Ehmcke J, Wistuba J. 2014. Sperm competition and the evolution of spermatogenesis. *Molecular Human Reproduction* 20(12):1169–1179 DOI 10.1093/molehr/gau070.
- Ramm SA, Stockley P. 2007. Ejaculate allocation under varying sperm competition risk in the house mouse, *Mus musculus domesticus*. *Behavioral Ecology* 18(2):491–495 DOI 10.1093/beheco/arm003.
- Saier MH, Reddy VS, Tamang DG, Vastermark A. 2014. The transporter classification database. *Nucleic Acids Research* 42:D251–D258 DOI 10.1093/nar/gkt1097.
- Schmidt-Nielsen K, Schmidt-Nielsen B. 1952. Water metabolism of desert mammals 1. *Physiological Reviews* 32(2):135–166.
- Schmidt-Nielsen B, Schmidt-Nielsen K, Brokaw A, Schneiderman H. 1948. Water conservation in desert rodents. *Journal of Cellular Physiology* 32(3):331–360 DOI 10.1002/jcp.1030320306.
- Scott C. 2016. dammit: an open and accessible de novo transcriptome annotator. Available at www.camillescott.org/dammit.
- Sikes RS, Gannon WL, Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American society of mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92(1):235–253 DOI 10.1644/10-MAMM-F-355.1.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31(19):3210–3212 DOI 10.1093/bioinformatics/btv351.

- Smith-Unna R, Bournnell C, Patro R, Hibberd JM, Kelley S. 2016.** TransRate: reference free quality assessment of de-novo transcriptome assemblies. *Genome Research* 26:1134–1144 DOI [10.1101/gr.196469.115](https://doi.org/10.1101/gr.196469.115).
- Snel B, Lehmann G, Bork P, Huynen MA. 2000.** STRING: a web-server to retrieve and display the repeatedly occurring neighbourhood of a gene. *Nucleic Acids Research* 28(18):3442–3444 DOI [10.1093/nar/28.18.3442](https://doi.org/10.1093/nar/28.18.3442).
- Song L, Florea L. 2015.** Rcorrector: efficient and accurate error correction for Illumina RNA-seq reads. *Gigascience* 4:48 DOI [10.1186/s13742-015-0089-y](https://doi.org/10.1186/s13742-015-0089-y).
- Sonnhammer ELL, Eddy SR, Durbin R. 1997.** Pfam: a comprehensive database of protein families based on seed alignments. *Proteins* 28(3):405–420.
- Storz JF, Runck AM, Moriyama H, Weber RE, Fago A. 2010.** Genetic differences in hemoglobin function between highland and lowland deer mice. *Journal of Experimental Biology* 213(15):2565–2574 DOI [10.1242/jeb.042598](https://doi.org/10.1242/jeb.042598).
- Storz JF, Wheat CW. 2010.** Integrating evolutionary and functional approaches to infer adaptation at specific loci. *Evolution* 64(9):2489–2509 DOI [10.1111/j.1558-5646.2010.01044.x](https://doi.org/10.1111/j.1558-5646.2010.01044.x).
- Sun J, Lin Y, Wu J. 2013.** Long non-coding RNA expression profiling of mouse testis during post-natal development. *PLoS ONE* 8(10):e75750 DOI [10.1371/journal.pone.0075750](https://doi.org/10.1371/journal.pone.0075750).
- Suzek BE, Huang H, McGarvey PB, Mazumder R, Wu CH. 2007.** UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics* 23(10):1282–1288 DOI [10.1093/bioinformatics/btm098](https://doi.org/10.1093/bioinformatics/btm098).
- Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH. 2015.** UniRef clusters: a comprehensive and scalable alternative for improving sequence similarity searches. *Bioinformatics* 31(6):926–932 DOI [10.1093/bioinformatics/btu739](https://doi.org/10.1093/bioinformatics/btu739).
- Swanson WJ, Vacquier VD. 2002.** The rapid evolution of reproductive proteins. *Nature Reviews Genetics* 3(2):137–144 DOI [10.1038/nrg733](https://doi.org/10.1038/nrg733).
- Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A, Santos A, Tsafou KP, Kuhn M, Bork P, Jensen LJ, Von Mering C. 2015.** STRINGv10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Research* 43:D447–D452 DOI [10.1093/nar/gku1003](https://doi.org/10.1093/nar/gku1003).
- Turner LM, Chuong EB, Hoekstra HE. 2008.** comparative analysis of testis protein evolution in Rodents. *Genetics* 179(4):2075–2089 DOI [10.1534/genetics.107.085902](https://doi.org/10.1534/genetics.107.085902).
- Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigartyo CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, Von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, Von Heijne G, Nielsen J, Pontén F. 2015.** Tissue-based map of the human proteome. *Science* 347(6220):1260419 DOI [10.1126/science.1260419](https://doi.org/10.1126/science.1260419).
- Urity VB, Issaian T, Braun EJ, Dantzler WH, Pannabecker TL. 2012.** Architecture of kangaroo rat inner medulla: segmentation of descending thin limb of Henle’s loop. *American Journal of Physiology* 302(6):R720–R726 DOI [10.1152/ajpregu.00549.2011](https://doi.org/10.1152/ajpregu.00549.2011).

- Veal R, Caire W. 2001.** *Peromyscus eremicus*. *Mammalian Species* **118**:1–6.
- Vicens A, Luke L, Roldan E.RS. 2014.** Proteins involved in motility and sperm-egg interaction evolve more rapidly in mouse spermatozoa. *PLoS ONE* **9**(3):e91302 DOI [10.1371/journal.pone.0091302](https://doi.org/10.1371/journal.pone.0091302).
- Vimtrup BJ, Schmidt-Nielsen B. 1952.** The histology of the kidney of kangaroo rats. *The Anatomical Record* **114**(4):515–528 DOI [10.1002/ar.1091140402](https://doi.org/10.1002/ar.1091140402).
- Weber JN, Hoekstra HE. 2009.** The evolution of burrowing behavior in deer mice (genus *Peromyscus*). *Animal Behavior* **77**(3):603–609 DOI [10.1016/j.anbehav.2008.10.031](https://doi.org/10.1016/j.anbehav.2008.10.031).
- Weber JN, Peterson BK, Hoekstra HE. 2013.** Discrete genetic modules are responsible for complex burrow evolution in *Peromyscus* mice. *Nature* **493**(7342):402–405 DOI [10.1038/nature11816](https://doi.org/10.1038/nature11816).
- Wolff J. 1989.** Social behavior. In: Kirkland Jr LG, Layne JN, eds. *Advances in the study of Peromyscus (Rodentia)*. Lubbock: Texas Tech University Press, 271–291.
- Yang Z. 1997.** PAML: a program package for phylogenetic analysis by maximum likelihood. *Computer Applications in BioSciences* **13**(5):555–556.
- Yang Z. 1998.** Likelihood ratio tests for detecting positive selection and application to primate lysozyme evolution. *Molecular Biology and Evolution* **15**(5):568–573.
- Yang Z. 2007.** PAML4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution* **24**(8):1586–1591 DOI [10.1093/molbev/msm088](https://doi.org/10.1093/molbev/msm088).
- Zhang J, Nielsen R, Yang Z. 2005.** Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Molecular Biology and Evolution* **22**(12):2472–2479 DOI [10.1093/molbev/msi237](https://doi.org/10.1093/molbev/msi237).