



In this article “Predicting Coronary Heart Disease Using Risk Factor Categories for a Japanese Urban Population, and Comparison with the Framingham Risk Score: The Suita Study” by Kunihiro Nishimura, et al., which appeared in JAT 2014, 21: 784-798 are incorrect.

Table 1 on page 787 and “Population Characteristics” on page 788 are corrected as below:

Original Article

Predicting Coronary Heart Disease Using Risk Factor Categories for a Japanese Urban Population, and Comparison with the Framingham Risk Score: The Suita Study

Kunihiro Nishimura¹, Tomonori Okamura², Makoto Watanabe¹, Michikazu Nakai¹, Misa Takegami¹, Aya Higashiyama³, Yoshihiro Kokubo¹, Akira Okayama⁴ and Yoshihiro Miyamoto¹

Incorrect

Table 1. Population characteristics of the study cohort

	Males (n = 2796)	Females (n = 2725)
Age (years, mean ± SD)	56.1 ± 13.3	54.5 ± 12.9
DM (%)	6	5.8
Current smoker (%)	49.67	11.91
Blood pressure (mmHg, %)		
Optimal (SBP < 120, DBP < 80)	30.74	41.68
Normal (SBP < 130, DBP < 85)	19.31	17.30
High normal (SBP < 140, DBP < 90)	17.98	15.69
Stage I HT (SBP < 160, DBP < 100)	20.39	15.94
Stage II to IV HT (SBP ≥ 160, DBP ≥ 100)	11.59	9.40
Total cholesterol (mg/dl, %)		
< 160	10.12	6.88
160-199	39.75	30.52
200-239	37.41	39.60
240-279	10.98	18.51
≥ 280	1.74	4.49
LDL cholesterol (mg/dl, %)		
> 130	55.54	45.78
130-159	28.19	30.86
> 160	16.26	23.34
HDL cholesterol (mg/dl, %)		
< 35	11.40	3.28
35-44	28.71	16.36
45-49	15.87	12.25
50-59	23.82	29.95
≥ 60	20.20	38.14
Creatinine (mg/dl, mean ± SD)	0.91 ± 0.21	0.69 ± 0.22
eGFR (mean ± SD)	64.7 ± 24.9	90.6 ± 29.3
CKD (≥ Stage 3) (%)	46.2	11.3

LDL, Low-density lipoprotein; HDL, high-density lipoprotein;
eGFR, estimated glomerular filtration rate (ml/min/1.73 m²); CKD, chronic kidney disease; HT, hypertension

Correct**Table 1.** Population characteristics of the study cohort

	Males (n = 2796)	Females (n = 2725)
Age (years, mean ± SD)	56.1 ± 13.3	54.5 ± 12.9
DM (%)	6	5.8
Current smoker (%)	49.67	11.91
Blood pressure (mmHg, %)		
Optimal (SBP < 120, DBP < 80)	30.74	41.68
Normal (SBP < 130, DBP < 85)	19.31	17.30
High normal (SBP < 140, DBP < 90)	17.98	15.69
Stage I HT (SBP < 160, DBP < 100)	20.39	15.94
Stage II to IV HT (SBP ≥ 160, DBP ≥ 100)	11.59	9.40
Total cholesterol (mg/dl, %)		
< 160	10.12	6.88
160-199	39.75	30.52
200-239	37.41	39.60
240-279	10.98	18.51
≥ 280	1.74	4.49
LDL cholesterol (mg/dl, %)		
> 130	55.54	45.78
130-159	28.19	30.86
> 160	16.26	23.34
HDL cholesterol (mg/dl, %)		
< 35	11.40	3.28
35-44	28.71	16.36
45-49	15.87	12.25
50-59	23.82	29.95
≥ 60	20.20	38.14
Creatinine (mg/dl, mean ± SD)	0.91 ± 0.21	0.69 ± 0.22
eGFR (mean ± SD)	73.1 ± 25.7	90.6 ± 29.3
CKD (≥ Stage 3) (%)	27.9	11.3

LDL, Low-density lipoprotein; HDL, high-density lipoprotein;
eGFR, estimated glomerular filtration rate (ml/min/1.73 m²); CKD, chronic kidney disease; HT, hypertension

Incorrect**Results****Population Characteristics**

The number of person-years studied consisted of 75,776 (34,480 for males and 41,296 for females), with a mean follow-up period of 11.8 years. During follow-up period, there were **213** incidents of CHD. The population characteristics are summarized in Table 1. The univariate Cox regression analysis indicated all variables in FRS were statistically significant (data not shown).

Correct**Results****Population Characteristics**

The number of person-years studied consisted of 75,776 (34,480 for males and 41,296 for females), with a mean follow-up period of 11.8 years. During follow-up period, there were **179** incidents of CHD. The population characteristics are summarized in Table 1. The univariate Cox regression analysis indicated all variables in FRS were statistically significant (data not shown).