




Available online at www.sciencedirect.com

ScienceDirect



Journal of Current Ophthalmology 28 (2016) 194–198

<http://www.journals.elsevier.com/journal-of-current-ophthalmology>

Original research

Corneal aberration changes after rigid gas permeable contact lens wear in keratokonic patients

Fereshteh Shokrollahzadeh ^a, Hassan Hashemi ^b, Ebrahim Jafarzadehpur ^{c,*}, Ali Mirzajani ^c, Mehdi Khabazkhoob ^d, Abbasali Yekta ^e, Soheila Asgari ^f

^a Noor Research Center for Ophthalmic Epidemiology, Noor Eye Hospital, Tehran, Iran

^b Noor Ophthalmology Research Center, Noor Eye Hospital, Tehran, Iran

^c Department of Optometry, School of Rehabilitation Science, Iran University of Medical Sciences, Tehran, Iran

^d Department of Medical Surgical Nursing, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^e Department of Optometry, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran

^f Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, International Campus, Tehran, Iran

Received 11 June 2016; accepted 6 August 2016

Available online 30 August 2016

Abstract

Purpose: To determine the short-term effect of rigid gas permeable (RGP) contact lenses on corneal aberrations in keratoconic patients.

Method: Sixteen keratoconic eyes with no history of RGP lens wear were included. They all had corneal aberrometry using Pentacam, and different aberration indices of the anterior and posterior surfaces of the cornea were measured before and 3 months after fitting RGP lenses. The effect of baseline parameters on these changes was tested in univariate and multiple models.

Results: Total aberrations and individual Zernike coefficients did not show statistically significant changes after using RGP lenses. Although not statistically significant, vertical coma decreased in the anterior ($p = 0.073$) and posterior surface ($p = 0.095$). Relationships that remained statistically significant in the multiple model were between baseline central corneal thickness and changes in total higher order aberrations and anterior 4th order astigmatism 0° , and between baseline 2nd order astigmatism 45° and its changes.

Conclusion: In this study, corneal aberrations remained unchanged 3 months after wearing RGP contact lens. Further studies with sufficient samples in different groups of keratoconus severity or baseline aberrations are needed to obtain more accurate results.

Copyright © 2016, Iranian Society of Ophthalmology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Aberrometry; Keratoconus; Rigid gas permeable contact lens; Pentacam

Introduction

The cornea is a major source of aberrations which can cause limitations in visual acuity and quality.^{1–3} In keratoconus patients, decreased visual acuity is one of the main complaints, and in comparison with normal corneas, keratoconic eyes exhibit many corneal irregularities, irregular astigmatism, and

aberrations.^{4–7} Coma, especially vertical coma, is the most common aberration in keratoconic patients.⁸ These aberrations are found and can be evaluated in both the posterior and anterior surfaces of the cornea.⁹ Measuring corneal aberrations changes improves our understanding of the changes in the shape and optical properties of the cornea and keratoconus progression.¹⁰ Studies suggest that vertical coma and total root

Conflict of interest: No conflicting relationship exists for any author.

Financial Support: This project was funded by the Noor Ophthalmology Research Center.

* Corresponding author.

E-mail address: jafarzadehpour.e@iums.ac.ir (E. Jafarzadehpur).

Peer review under responsibility of the Iranian Society of Ophthalmology.

<http://dx.doi.org/10.1016/j.joco.2016.08.002>

2452-2325/Copyright © 2016, Iranian Society of Ophthalmology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

mean square (RMS) aberrations are the best aberrometry indices to detect keratoconus.^{11–13}

Vision improvement in keratoconic patients can be achieved through the use of contact lenses or surgical methods such as corneal ring implantation and corneal grafting. Rigid gas permeable (RGP) lenses, the first lens choice for keratoconic patients, are believed to change corneal astigmatism as a result of the optical characteristics of the tear layer behind the lens.¹⁴ However, studies in this regard are inconclusive; they vary in methodology and follow-up time, and there are still many questions concerning the effects of RGP lenses on different corneal parameters.^{15–18} This study was designed to examine changes in the total RMS and Zernike polynomials in keratoconic patients after 3 months of RGP lens wear.

Method

In this prospective, before–after case series, patients with mild to severe keratoconus referring to the contact lens clinic were included. The study was approved by the Ethics Committee of Iran University of Medical Sciences. All patients had already had complete ophthalmic evaluation by a cornea specialist, based on which the diagnosis of keratoconus was made and confirmed through imaging modalities. Objectives and methods of the study were explained to patients, and they all signed informed consents before participation in the study.

Inclusion criteria were age between 19 and 35 years and being eligible for fitting RGP spherical lenses. Patients with a history of RGP lens use, corneal scarring, infectious or inflammatory ocular diseases, corneal graft surgery, ring implantation or corneal collagen crosslinking, and those referred for fitting mini-scleral and ClearKone lenses were excluded.

Patients had complete ocular examination including the measurement of visual acuity using a Snellen chart (Nidek-34605-6004-LCD Chart, Japan) at 4 m, objective refraction with a retinoscope (Heine Beta 200, Germany) and an auto-refractometer (Nidek, Japan), and subjective refraction with a trial lens set and frame. The appropriate aspheric RGP lens (Iran, Lens Gostar, Tehran, Iran) was prescribed using the diagnostic fitting method.¹⁹ On slit-lamp examination, good tear exchange was verified by observing the fluorescein pattern with mild apical clearance over the corneal cone and slight edge and midperipheral clearance.²⁰ Patients were instructed to wear their lenses 3 h a day during the first week, and add an hour a day per week up to a maximum of 8 h daily. Follow-up calls were made to patients to stay in touch with them and ensure adherence to the study protocol.

To measure aberrations of the anterior and posterior surfaces of the cornea at baseline and at 3 months, we used the Pentacam corneal topographer (Oculus Optikgeräte GmbH, Wetzlar, Germany) which has shown to be highly reliable.²¹ Extracted indices for the anterior and posterior corneal surfaces included the total RMS, 2nd order astigmatism 45°, 3rd order coma 0° and 90°, 3rd order trefoil 0° and 30°, and 4th

order secondary astigmatism 0° and 45° measured over a 6.0 mm aperture.

Statistical analyses were done using the Statistical Package for Social Sciences (SPSS) Version 20.0 (Chicago, IL, USA). Results were compared using repeated measures analysis of covariance (ANCOVA), and the correlation between fellow eyes was accounted for. Changes were considered significant based on a significance level of 5%. Relationships between aberration changes and other variables were also explored using single and multiple linear regression analyses.

Results

A total of 10 patients, 3 women and 7 men, with a mean age of 27.2 ± 4.9 years (range, 20–35) were enrolled, and their 16 keratoconic eyes were evaluated. Based on the indices and classification scheme suggested by Rabinowitz, the number of eyes that could be classified as keratoconus and early keratoconus was 12 and 4, respectively.²² The diameter of the prescribed RGP lenses was 9.60 mm in all cases, and they had a mean base curve radius of 7.70 ± 0.28 mm (range, 7.20–8.10) and a mean power of -1.56 ± 1.26 diopters (D) (range, -3.00 to +1.25). At baseline, patients had a mean maximum keratometry reading of 54.06 ± 5.73 D (range, 43.90–67.10 D), mean central corneal thickness (CCT) of 468.40 ± 56.11 μm (range, 381–547), and mean anterior Q-value of -0.80 ± 0.39 (range, -1.52 to -0.28), as determined with the Pentacam. Table 1 presents the mean values for these variables at 3 months and their changes.

Studied aberrations in the anterior and posterior corneal surface before and 3 months after using RGP lenses are summarized in Table 1. None of the studied aberration indices in the anterior or posterior corneal surfaces showed no significant change.

In the second set of analyses, we examined the effect of baseline values of different parameters on the amount of 3-month changes in the studied aberrations. With univariate regression analyses, baseline CCT significantly impacted changes in total RMS HOA ($\beta = 0.009$, $p = 0.017$), anterior surface 3rd order coma 90° ($\beta = -0.010$, $p = 0.034$), and anterior surface 4th order astigmatism 0° ($\beta = 0.002$, $p = 0.029$). CCT relationships with total RMS HOA and anterior 4th order astigmatism 0° retained their significance in the multiple model as well ($\beta = -0.841$, $p = 0.001$ and $\beta = 0.003$, $p = 0.045$, respectively). Also, changes in anterior spherical aberration ($\beta = -0.525$, $p = 0.005$), 2nd order astigmatism 45° ($\beta = -0.321$, $p = 0.018$), and 4th order astigmatism 45° ($\beta = -0.279$, $p = 0.014$) significantly related with their baseline values; the second relationship retained its significance in the multiple model ($\beta = -0.313$, $p = 0.045$) and is illustrated in Fig. 1. No other significant relation was found between changes in any of the aberration variables and baseline maximum keratometry reading, CCT, or anterior Q-value (all $P > 0.05$).

Table 1

Corneal curvature, thickness, Q-value, and anterior and posterior aberrations in the studied sample at baseline and 3 months after wearing rigid gas permeable contact lenses.

	Baseline	At 3 months	Mean change (95% confidence interval)	p Value ^a
Maximum k-reading (diopter)	54.06 ± 5.73	54.52 ± 6.07	0.46 ± 2.10 (-0.70, 1.62)	0.410
Central corneal thickness (micron)	468.40 ± 56.11	465.40 ± 59.34	-3.00 ± 10.34 (-8.73, 2.73)	0.280
Anterior Q-value	-0.79 ± 0.39	-0.80 ± 0.45	-0.01 ± 0.20 (-0.13, 0.10)	0.785
Posterior Q-value	-0.87 ± 0.66	-0.84 ± 0.76	0.03 ± 0.27 (-0.12, 0.18)	0.658
Total RMS LOA	9.00 ± 4.40	8.69 ± 5.53	-0.31 ± 2.35 (-1.56, 0.94)	0.605
Total RMS HOA	2.54 ± 1.41	2.52 ± 2.04	-0.02 ± 1.10 (-0.60, 0.56)	0.942
Anterior corneal surface aberrations (micron)				
Total RMS	10.95 ± 5.34	10.69 ± 4.92	-0.26 ± 1.42 (-1.04, 0.53)	0.495
2nd order astigmatism 0°	-1.00 ± 2.64	-0.87 ± 2.86	0.13 ± 0.49 (-0.13, 0.39)	0.304
2nd order astigmatism 45°	-0.29 ± 2.60	-0.64 ± 2.11	-0.35 ± 1.43 (-1.11, 0.41)	0.343
3rd order coma 0°	-0.40 ± 1.30	-0.56 ± 1.58	-0.16 ± 0.75 (-0.56, 0.24)	0.414
3rd order coma 90°	-2.09 ± 1.29	-1.61 ± 1.36	0.48 ± 1.00 (-0.05, 1.01)	0.073
3rd order trefoil 0°	0.06 ± 1.12	0.38 ± 1.41	0.32 ± 0.94 (-0.18, 0.82)	0.189
3rd order trefoil 30°	0.02 ± 0.49	-0.33 ± 1.15	-0.35 ± 1.02 (-0.90, 0.19)	0.187
4th order spherical aberration	-0.73 ± 0.95	-0.42 ± 0.72	0.31 ± 0.75 (-0.90, 0.71)	0.119
4th order vertical astigmatism 0°	0.03 ± 0.66	-0.02 ± 0.64	-0.05 ± 0.18 (-0.15, 0.04)	0.273
4th order oblique astigmatism 45°	0.06 ± 0.61	0.17 ± 0.49	0.11 ± 0.28 (-0.04, 0.27)	0.124
Posterior corneal surface aberrations (micron)				
Total RMS	2.71 ± 0.98	2.75 ± 1.17	0.05 ± 0.48 (-0.21, 0.31)	0.693
2nd order astigmatism 0°	0.45 ± 0.27	0.47 ± 0.27	0.01 ± 0.15 (-0.06, 0.09)	0.718
2nd order astigmatism 45°	-0.01 ± 0.39	0.02 ± 0.36	0.03 ± 0.17 (-0.06, 0.12)	0.499
3rd order coma 0°	0.04 ± 0.31	0.05 ± 0.29	0.01 ± 0.17 (-0.08, 0.11)	0.774
3rd order coma 90°	0.50 ± 0.28	0.39 ± 0.36	-0.12 ± 0.26 (-0.25, 0.02)	0.095
3rd order trefoil 0°	0.05 ± 0.24	0.01 ± 0.12	-0.04 ± 0.22 (-0.16, 0.08)	0.472
3rd order trefoil 30°	-0.09 ± 0.17	0.00 ± 0.26	0.09 ± 0.25 (-0.04, 0.22)	0.170
4th order spherical aberration	0.03 ± 0.16	0.01 ± 0.17	-0.02 ± 0.06 (-0.05, 0.01)	0.180
4th order astigmatism 0°	-0.06 ± 0.10	-0.05 ± 0.11	0.01 ± 0.06 (-0.02, 0.04)	0.519
4th order astigmatism 45°	0.00 ± 0.11	-0.01 ± 0.13	-0.01 ± 0.06 (-0.04, 0.02)	0.635

HOA: High order aberration, LOA: Low order aberration, RMS: Root mean square.

^a Repeated measures analysis of covariance.




Fig. 1. Regression line showing the relation between changes in 2nd order oblique astigmatism in microns 3 months after wearing rigid gas permeable contact lenses and its baseline values in the studied sample of keratoconus patients.

Discussion

The impacts of contact lens wear on the human corneal topography and optical performance have been examined through various studies which vary by design, sample size and type, follow-up periods, and measurement tools.

Similarly, studies of RGP lenses on keratoconus patients not only vary by these parameters, but also fitting methods and choice of lens, and results have been inconclusive. Of note, most of the few available studies have focused on changes after suspending contact lens wear or with lenses in place.^{18,23,24} In this study, we examined changes in ocular aberrations after 3 months of RGP lens wear in keratoconus patients using the Pentacam Zernike Analysis software. Determining the role of baseline parameters may provide practitioners with information that can be useful for predicting outcomes.

As presented in Table 1, we observed no significant change in any aberration index of the anterior or posterior cornea. Total RMS is an important parameter for the analysis and understanding the theoretical values of ocular aberrations²⁵ and encompasses all higher and lower order aberrations. In a recent comparative study of apical touch and three point touch fitting methods in 50 keratoconic eyes of 31 patients, total RMS, as measured with the Pentacam, was reported to decrease after 14 days of RGP lens wear.²⁶ In terms of total RMS HOA reduced levels were observed in a study of 40 keratoconic eyes using a Hartman Shack aberrometer before and after fitting RGP lenses.²⁷ In a retrospective study by Negishi et al, changes were examined in 13 eyes of 10 patients with mild keratoconus, and the authors reported significant decreases in the total RMS HOA, third order coefficients, and fourth order Zernike coefficients.¹⁷

In our study, vertical coma slightly reduced 3 months after RGP lens wear; mean change was $0.48 \pm 0.10 \mu\text{m}$ for the anterior corneal surface and $-0.11 \pm 0.26 \mu\text{m}$ for the posterior corneal surface. Statistical analysis showed a borderline significant decrease 3 months after using RGP lenses. The importance of this finding is that a larger sample size may generate results with statistical significance. Also, changes in the vertical coma of the anterior and posterior corneal surfaces had opposite signs. In keratoconic eyes, coma, especially vertical coma, tends to increase as the disease progresses²⁸ which is due to the inferior position of the corneal cone in most patients.^{29–31} Very few studies have evaluated coma changes in relation to RGP lens wear in keratoconus patients. Kosaki et al studied 19 eyes of 15 keratoconic patients and reported a decrease in coma from $0.35 \pm 0.29 \mu\text{m}$ to $0.16 \pm 0.08 \mu\text{m}$ and secondary astigmatism from $0.11 \pm 0.07 \mu\text{m}$ to $0.04 \pm 0.02 \mu\text{m}$ in the anterior surface of the cornea.³² They also examined the effect of RGP lens wear on the posterior surface of the cornea.³² In agreement with our results, the posterior surface was less influenced by RGP lens wear, compared to the anterior corneal surface.

Studies suggest that contact lens wear-related aberrometry changes and direction of coma depend on baseline aberrations and corneal shape.^{33,34} In our study, we found significant relationships between baseline CCT and changes in total RMS HOA, anterior surface 3rd order coma 90° , and anterior surface 4th order astigmatism 0° . Also, changes in anterior spherical aberration, 2nd order astigmatism 45° , and 4th order astigmatism 45° significantly related with their baseline values. In a 2007 comparative study between 22 myopic eyes and 14 keratoconic eyes, the authors concluded that HOA can be enhanced or reduced based on baseline ocular aberrations as an effect of changes in vertical coma. In the mentioned study, HOA decreased from $0.54 \mu\text{m}$ to $0.36 \mu\text{m}$ while coma increased from $-0.185 \mu\text{m}$ to $0.134 \mu\text{m}$, and the orientation of coma changed from inferior to superior cornea in keratoconus patients.¹⁶ These observations call for further studies with sufficient samples in different groups of keratoconus severity or baseline aberrations.

This study has several limitations. The sample size of our study was too small to draw a definitive conclusion. Also, the number of eyes in the different subgroups was not enough to allow for analysis in categories defined by corneal thickness, keratoconus grade, and baseline aberrations.

In conclusion short-term RGP lens wear in keratoconus patients seems to be associated with no significant increase or even a decrease in aberrations. Further long-term studies with larger sample sizes are needed to complete our knowledge in these areas.

References

- Seiler T, Mrochen M, Kaemmerer M. Operative correction of ocular aberrations to improve visual acuity. *J Refract Surg.* 2000;16:S619–S622.
- Rabinowitz YS. Keratoconus. *Surv Ophthalmol.* 1998;42:297–319.
- He JC, Marcos S, Webb R, Burns SA. Measurement of the wave-front aberration of the eye by a fast psychophysical procedure. *JOSA A.* 1998;15:2449–2456.
- Schlegel Z, Lteif Y, Bains HS, Gatinel D. Total, corneal, and internal ocular optical aberrations in patients with keratoconus. *J Refract Surg.* 2009;25:951.
- Li X, Yang H, Rabinowitz YS. Longitudinal study of keratoconus progression. *Exp Eye Res.* 2007;85:502–507.
- Zadnik K, Barr JT, Edrington TB, et al. Baseline findings in the Collaborative Longitudinal Evaluation of keratoconus (CLEK) study. *Invest Ophthalmol Vis Sci.* 1998;39:2537–2546.
- Alió JL, Shabayek MH. Corneal higher order aberrations: a method to grade keratoconus. *J Refract Surg.* 2006;22:539.
- Pantanelli S, MacRae S, Jeong TM, Yoon G. Characterizing the wave aberration in eyes with keratoconus or penetrating keratoplasty using a high-dynamic range wavefront sensor. *Ophthalmology.* 2007;114:2013–2021.
- Nakagawa T, Maeda N, Kosaki R, et al. Higher-order aberrations due to the posterior corneal surface in patients with keratoconus. *Invest Ophthalmol Vis Sci.* 2009;50:2660–2665.
- He JC, Sun P, Held R, Thorn F, Sun X, Gwiazda JE. Wavefront aberrations in eyes of emmetropic and moderately myopic school children and young adults. *Vis Res.* 2002;42:1063–1070.
- Jafri B, Li X, Yang H, Rabinowitz YS. Higher order wavefront aberrations and topography in early and suspected keratoconus. *J Refract Surg.* 2007;23:774–781.
- Gordon-Shaag A, Millodot M, Ifrah R, Shneor E. Aberrations and topography in normal, keratoconus-suspect, and keratoconic eyes. *Optom Vis Sci.* Apr 2012;89:411–418.
- Agarwal A, Agarwal A, Jacob S. *Dr. Agarwal's Textbook on Corneal Topography: Including Pentacam and Anterior Segment OCT.* Jaypee Brothers Publishers; 2010.
- Lawless M, Coster DJ, Phillips AJ, Loane M. Keratoconus: diagnosis and management. *Aust N Z J Ophthalmol.* 1989;17(1):33–60.
- Liu Z, Pflugfelder SC. The effects of long-term contact lens wear on corneal thickness, curvature, and surface regularity. *Ophthalmology.* 2000;107:105–111.
- Choi J, Wee WR, Lee JH, Kim MK. Changes of ocular higher order aberration in on-and off-eye of rigid gas permeable contact lenses. *Optom Vis Sci.* 2007;84:42–51.
- Negishi K, Kumanomido T, Utsumi Y, Tsubota K. Effect of higher-order aberrations on visual function in keratoconic eyes with a rigid gas permeable contact lens. *Am J Ophthalmol.* 2007;144:924–929, e921.
- Jinabhai A, Radhakrishnan H, O'Donnell C. Corneal changes after suspending contact lens wear in early pellucid marginal corneal degeneration and moderate keratoconus. *Eye Contact Lens.* 2011;37:99–105.
- Bennett ES, Weissman BA. *Clinical Contact Lens Practice.* Lippincott Williams & Wilkins; 2005.
- Herranz RM, Zarzuelo GR, de Juan Herráez V. *Contact Lens Correction of Regular and Irregular Astigmatism.* INTECH Open Access Publisher; 2012.
- Doménech Amigot B, Mas Candela D, Ronda Pérez E, Pérez Rodríguez J, Espinosa Tomás J, Illueca Contrí C. *Repeatability and Concordance of the Pentacam System: Comparative Study of Corneal Parameters Measured with Pentacam and Atlas.* 2009.
- Rabinowitz YS. Corneal topography. In: Bennett ES, Weissman BA, eds. *Clinical Contact Lens Practice.* Philadelphia, PA: Lippincott, Williams and Wilkins; 2005:215–232.
- Marsack JD, Parker KE, Pesudovs K, Donnelly 3rd WJ, Applegate RA. Uncorrected wavefront error and visual performance during RGP wear in keratoconus. *Optom Vis Sci.* 2007;84:463–470.
- Jinabhai A, O'Donnell C, Radhakrishnan H. Changes in refraction, ocular aberrations, and corneal structure after suspending rigid gas-permeable contact lens wear in keratoconus. *Cornea.* May 2012;31:500–508.
- Copeland R, Afshari N. *Principles and Practice of Cornea.* New Delhi: Jaypee Brothers Medical Publishers; 2013.
- Romero-Jiménez M, Santodomingo-Rubido J, Flores-Rodríguez P, González-Méijome J-M. Short-term corneal changes with gas-permeable contact lens wear in keratoconus subjects: a comparison of two fitting approaches. *J Optom.* 2015;8:48–55.

27. Baghbani F, Heravian J, Derakhshan A, et al. Evaluation of predictive value of aberrometry visual performance of keratoconus patients before and after rigid gas permeable lenses. *Arak Med Univ J.* 2013;16: 23–29.
28. Schwiegerling J, Greivenkamp JE. Using corneal height maps and polynomial decomposition to determine corneal aberrations. *Optom Vis Sci.* 1997;74:906–916.
29. Applegate RA, Hilmantel G, Howland HC, Tu EY, Starck T, Zayac EJ. Corneal first surface optical aberrations and visual performance. *J Refract Surg.* 2000;16:507–514.
30. Gobbe M, Guillon M. Corneal wavefront aberration measurements to detect keratoconus patients. *Contact Lens Anterior Eye.* 2005;28:57–66.
31. Wilson SE, Lin DT, Klyce SD. Corneal topography of keratoconus. *Cornea.* 1991;10:2–8.
32. Kosaki R, Maeda N, Bessho K, et al. Magnitude and orientation of Zernike terms in patients with keratoconus. *Invest Ophthalmol Vis Sci.* 2007; 48:3062.
33. Lu F, Mao X, Qu J, Xu D, He JC. Monochromatic wavefront aberrations in the human eye with contact lenses. *Optom Vis Sci.* 2003;80: 135–141.
34. Dorronsoro C, Barbero S, Llorente L, Marcos S. On-eye measurement of optical performance of rigid gas permeable contact lenses based on ocular and corneal aberrometry. *Optom Vis Sci.* 2003;80:115–125.