

Serological survey of *Encephalitozoon cuniculi* infection in cats in Japan

Ryusuke TSUKADA¹), Yuki OSAKA¹), Tomomi TAKANO²), Mizuki SASAKI³), Mitsuhiro INOSE⁴) and Hiromi IKADAI¹)*

¹) Laboratory of Veterinary Parasitology, School of Veterinary Medicine, Kitasato University, Towada, Aomori 034-8628, Japan

²) Laboratory of Veterinary Infectious Disease, School of Veterinary Medicine, Kitasato University, Towada, Aomori 034-8628, Japan

³) Department of Parasitology, Asahikawa Medical University, Asahikawa, Hokkaido 078-8510, Japan

⁴) NPO Nagano Animal Welfare Society, Nagano 380-0941, Japan

(Received 28 September 2015/Accepted 8 June 2016/Published online in J-STAGE 20 June 2016)

ABSTRACT. Antibodies to *Encephalitozoon cuniculi* (*E. cuniculi*) were examined by enzyme-linked immunosorbent assay using *E. cuniculi* PTP2 recombinant protein from serum samples that had been collected from a total of 295 cats in Japan. Of these samples, 6.1% (18/295) had antibodies against *E. cuniculi*, which included 6.3% (6/96) of the male cats and 6.0% (12/199) of the female cats. The incidence was slightly higher in feral cats (8.3%, 11/132) compared to domesticated cats (4.3%, 7/163). This suggests the possibility that the cats of our country have become a reservoir of *E. cuniculi*. This study is the first to demonstrate the prevalence of *E. cuniculi* infection in cats in Japan.

KEY WORDS: cat, ELISA, *Encephalitozoon cuniculi*, serological survey

doi: 10.1292/jvms.15-0545; *J. Vet. Med. Sci.* 78(10): 1615–1617, 2016

Encephalitozoon cuniculi (*E. cuniculi*) is an obligate intracellular microsporidian parasite and infects a wide range of mammalian hosts, such as rabbits, rodents, dogs and humans [10, 11, 15]. *E. cuniculi* is an opportunistic pathogen in human patients with acquired immunodeficiency syndrome and other immunocompromised people [3, 8]. Infected animals are arguably the reservoirs of human encephalitozoonosis [3]. Encephalitozoonosis is commonly observed in rabbits, and therefore, clinical assessment and diagnosis is well established in this species [8]. Although infections due to *E. cuniculi* are not common in cats, *E. cuniculi* infections have been reported to occur in cats abroad [2, 8, 14]. One author reported that the potential role of cats in the transmission of the infectious agent cannot be excluded [14]. In Japan, a seroepidemiological survey of *E. cuniculi* infection in domesticated dogs and domesticated rabbits has been performed [7, 16], but the prevalence of *E. cuniculi* infections in cats has not been reported. *E. cuniculi* in subclinically infected cats may represent a reservoir and potential risk for immunocompromised patients [11, 12]. The purpose of this study was to determine the seroprevalence of *E. cuniculi* in cats in Japan to serve as baseline epidemiologic data and a potential source of microsporidial infection in Japan. In addition, we compared the seropositive rate of domesticated cats with feral cats.

Furthermore, we examined four infectious diseases including *Toxoplasma gondii* (*T. gondii*), Feline coronavirus (FCoV), feline leukemia virus (FeLV) and feline immuno-

deficiency virus (FIV) that are major diseases of the cat and compared the seroprevalence in feral cats and domesticated cats.

Serum samples were collected from 295 cats that were brought in for surgical neutering to a private animal hospital in the Nagano Prefecture in Japan from September 2011 to September 2012. On examination, there were no clinical signs. The living environment of the examined cats was investigated by questioning the owners, and cats were classified into two groups: feral cats (i.e., no owner, lives outdoors) or domesticated cats (i.e., lives indoors and sometimes outdoors). The age was unidentified because not managed breeding environment. Serum samples were stored at -30°C prior to serological examination. Animal experimentation protocols (e.g., epidemiology investigation) were approved by the president of Kitasato University in accordance with the judgment of the Institutional Animal Care and Use Committee of Kitasato University (Towada, Japan; approval no. 15-064).

Testing for antibodies against *E. cuniculi* was performed using an enzyme-linked immunosorbent assay (ELISA) with glutathione *S*-transferase fusion polar tube protein 2 as the antigen [16]. Individual cat serum samples diluted to 1:400 and horseradish peroxidase-conjugated goat anti-feline immunoglobulin G antibody (MP Biomedicals, Santa Ana, CA, U.S.A.) diluted to 1:10,000 were used as the first and second antibodies, respectively.

Testing for antibodies against *T. gondii* was performed using a commercial latex agglutination test kit (Toxotest, Eiken Chemical Co., Ltd., Tokyo, Japan) in accordance with the manufacturer's instructions. Tests for antibodies against FCoV were performed using a previously published ELISA protocol with feline infectious peritonitis virus solubilization antigen as the antigen [17]. Tests for FeLV antigen and FIV antibody were performed using a commercial assay kit (SNAP FIV/FeLV Combo; IDEXX Laboratories, West-

*CORRESPONDENCE TO: IKADAI, H., Laboratory of Veterinary Parasitology, School of Veterinary Medicine, Kitasato University, Towada, Aomori 034-8628, Japan. e-mail: ikadai@vmas.kitasato-u.ac.jp

©2016 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

brook, ME, U.S.A.) in accordance with the product manual.

The overall seropositive rate to *E. cuniculi* was 6.1% (18/295), which included 6.3% (6/96) of the male cats and 6.0% (12/199) of the female cats; the incidence in feral cats (8.3%, 11/132) was slightly higher than in domesticated cats (4.3%, 7/163) (Table 1). This was not statistically significant. Moreover, 20.0% (59/295) were seropositive to *T. gondii*, 23.7% (70/295) to FCoV, 2.4% (7/295) to FeLV and 10.5% (31/295) to FIV. *Toxoplasma gondii* and FIV seropositive rates were 31.8% and 16.7%, respectively, in feral cats and 10.4% and 5.5%, respectively, in domesticated cats. The FIV seropositivity was significantly higher in males (15.6%) than in females (8.0%) (Table 2). The association between *E. cuniculi* positive samples and the positive samples of four major infectious diseases was assessed, however, this was not significant.

E. cuniculi infects a wide range of mammalian hosts, including humans [3]. In animals, the main target organs are the central nervous system and the kidney, which can result in a granulomatous encephalitis and nephritis [12]. The life cycle of *E. cuniculi* is simple and direct and, like other microsporidia, involves a proliferative merogonic stage, followed by a sporogonic stage resulting in rupture of the host cell and release of small ($1.5 \times 2.5 \mu\text{m}$), environmentally resistant, infective spores [12]. Spores can survive for many months in humid environments [9]. However, the natural mechanisms of *E. cuniculi* transmission are not fully understood. It is believed that the disease is spread horizontally in breeding with larger numbers of animals by the fecal-oral route, but above all, along the oro-urinal pathway [4]. Vertical, transplacental transmission of the infection may also play a key role in the epidemiology and pathogenesis, especially in carnivores and rodents [6, 13]. However, little is known about the occurrence of *E. cuniculi* in wildlife [12].

In cats, clinical disease is reportedly rare [12]. Infections with *E. cuniculi* occur in subclinically infected cats and may represent a reservoir and potential risk for immunocompromised patients and animals [7, 8].

In our study, serum samples of 295 cats were examined for the presence of antibodies against *E. cuniculi*. Though there were no clinical signs on general inspection, we found 6.1% of cats in Japan that had *E. cuniculi* antibodies (Table 1). A seroprevalence of 24% (17/72) of cats has been reported in Eastern Slovakia [4], and a recent study in Virginia, U.S.A., found a seroprevalence of 6.5% (15/232) [5]. These

Table 1. Seroprevalence of *E. cuniculi* infections in tested serum samples and stratified by living environment and sex

	Total	Living environment		Sex	
		Feral	Domesticated	Male	Female
Samples	295	132	163	96	199
Positive samples	18	11	7	6	12
Positive rate	6.1%	8.3%	4.3%	6.3%	6.0%

differences may reflect the different habitats where samples were obtained and the level of exposure to other sources of infection [12]. Furthermore, it has been reported that there are no sex differences in the prevalence of *E. cuniculi* [1], and the results of our study agree.

In our study, *E. cuniculi* seropositivity was not significantly different in domesticated cats compared to feral cats. This result may reflect the habitat of cats from which samples were obtained and that samples were not obtained from cats that resided completely indoor. On the other hand, it is reported that wildlife species have the potential to be significant reservoirs of infection for both domesticated animals and humans [12]. Furthermore, previous studies have sequenced a mouse strain in cats, indicating that the mouse may be a reservoir for infection in cats [1]. In a previous study, seroprevalence investigations in Japan revealed *E. cuniculi* infection in feral rodents [18]. In addition, we found 8.3% of feral cats in Japan that had *E. cuniculi* antibodies. In the cats of Japan, the mouse can be regarded as one of the important source of *E. cuniculi* infection.

In conclusion, this is the first report of *E. cuniculi* infection in cats in Japan. This suggests the possibility that the cats of our country have become a reservoir of *E. cuniculi*. This result indicates the importance of performing screening examinations of animals with the aim of reducing or halting of the spread of this disease. To further elucidate the route of infection, further studies are needed including the distribution of *E. cuniculi* infection in cats in Japan, other potential hosts or to discover the mouse strain of infection in cats by sequencing.

CONFLICT OF INTEREST. The authors declare no conflicts of interest directly relevant to the content of this article.

Table 2. Seroprevalence of *Toxoplasma gondii* (*T. gondii*), Feline coronavirus (FCoV), feline leukemia virus (FeLV) and feline immunodeficiency virus (FIV) infections in tested serum samples, and stratified by living environment and sex

	Positive samples (positive rate)	Living environment		Sex	
		Feral	Domesticated	Male	Female
<i>T. gondii</i>	59/295 (20%)	42/132 (31.8%)	17/163 (10.4%)	20/96 (20.8%)	39/199 (19.6%)
FCoV	70/295 (23.7%)	28/132 (21.2%)	42/163 (25.8%)	22/96 (23%)	48/199 (24.1%)
FeLV	7/295 (2.4%)	3/132 (2.3%)	4/163 (2.5%)	4/96 (4.2%)	3/199 (1.5%)
FIV	31/295 (10.5%)	22/132 (16.7%)	9/163 (5.5%)	15/96 (15.6%)	16/199 (8.0%)

REFERENCES

1. Benz, P., Maass, G., Csokai, J., Fuchs-Baumgartinger, A., Schwendenwein, I., Tichy, A. and Nell, B. 2011. Detection of *Encephalitozoon cuniculi* in the feline cataractous lens. *Vet. Ophthalmol.* **14** Suppl 1: 37–47. [Medline] [CrossRef]
2. Csokai, J., Fuchs-Baumgartinger, A., Maass, G. and Joachim, A. 2010. Detection of *Encephalitozoon cuniculi* infection (strain II) by PCR in a cat with anterior uveitis. *Wien. Tierarztl. Monatsschr.* **97**: 210–215.
3. Didier, E. S., Visvesvara, G. S., Baker, M. D., Rogers, L. B., Bertucci, D. C., De Groot, M. A. and Vossbrinck, C. R. 1996. A microsporidian isolated from an AIDS patient corresponds to *Encephalitozoon cuniculi* III, originally isolated from domestic dogs. *J. Clin. Microbiol.* **34**: 2835–2837. [Medline]
4. Halánová, M., Cisláková, L., Valencáková, A., Bálent, P., Adam, J. and Trávníček, M. 2003. Serological screening of occurrence of antibodies to *Encephalitozoon cuniculi* in humans and animals in Eastern Slovakia. *Ann. Agric. Environ. Med.* **10**: 117–120. [Medline]
5. Hsu, V., Grant, D. C., Zajac, A. M., Witonsky, S. G. and Lindsay, D. S. 2011. Prevalence of IgG antibodies to *Encephalitozoon cuniculi* and *Toxoplasma gondii* in cats with and without chronic kidney disease from Virginia. *Vet. Parasitol.* **176**: 23–26. [Medline] [CrossRef]
6. Hunt, R. D., King, N. W. and Foster, H. L. 1972. Encephalitozoonosis: evidence for vertical transmission. *J. Infect. Dis.* **126**: 212–214. [Medline] [CrossRef]
7. Igarashi, M., Oohashi, E., Dautu, G., Ueno, A., Kariya, T. and Furuya, K. 2008. High seroprevalence of *Encephalitozoon cuniculi* in pet rabbits in Japan. *J. Vet. Med. Sci.* **70**: 1301–1304. [Medline] [CrossRef]
8. Künzel, F., Peschke, R., Tichy, A. and Joachim, A. 2014. Comparison of an indirect fluorescent antibody test with Western blot for the detection of serum antibodies against *Encephalitozoon cuniculi* in cats. *Parasitol. Res.* **113**: 4457–4462. [Medline] [CrossRef]
9. Li, X., Palmer, R., Trout, J. M. and Fayer, R. 2003. Infectivity of microsporidia spores stored in water at environmental temperatures. *J. Parasitol.* **89**: 185–188. [Medline] [CrossRef]
10. Lindsay, D. S., Goodwin, D. G., Zajac, A. M., Cortés-Vecino, J. A., Gennari, S. M., Rosypal, A. C. and Dubey, J. P. 2009. Serological survey for antibodies to *Encephalitozoon cuniculi* in ownerless dogs from urban areas of Brazil and Colombia. *J. Parasitol.* **95**: 760–763. [Medline] [CrossRef]
11. Mathis, A., Weber, R. and Deplazes, P. 2005. Zoonotic potential of the microsporidia. *Clin. Microbiol. Rev.* **18**: 423–445. [Medline] [CrossRef]
12. Meredith, A. L., Cleaveland, S. C., Brown, J., Mahajan, A. and Shaw, D. J. 2015. Seroprevalence of *Encephalitozoon cuniculi* in wild rodents, foxes and domestic cats in three sites in the United Kingdom. *Transbound. Emerg. Dis.* **62**: 148–156. [Medline] [CrossRef]
13. Mohn, S. F., Nordstoga, K. and Dishington, I. W. 1982. Experimental encephalitozoonosis in the blue fox. Clinical, serological and pathological examinations of vixens after oral and intrauterine inoculation. *Acta Vet. Scand.* **23**: 490–502. [Medline]
14. Rebel-Bauder, B., Leschnik, M., Maderner, A. and Url, A. 2011. Generalized encephalitozoonosis in a young kitten with cerebellar hypoplasia. *J. Comp. Pathol.* **145**: 126–131. [Medline] [CrossRef]
15. Sak, B., Kváč, M., Květoňová, D., Albrecht, T. and Piálek, J. 2011. The first report on natural *Enterocytozoon bieneusi* and *Encephalitozoon* spp. infections in wild East-European House Mice (*Mus musculus musculus*) and West-European House Mice (*M. m. domesticus*) in a hybrid zone across the Czech Republic-Germany border. *Vet. Parasitol.* **178**: 246–250. [Medline] [CrossRef]
16. Sasaki, M., Yamazaki, A., Haraguchi, A., Tatsumi, M., Ishida, K. and Ikadai, H. 2011. Serological survey of *Encephalitozoon cuniculi* infection in Japanese dogs. *J. Parasitol.* **97**: 167–169. [Medline] [CrossRef]
17. Takano, T., Kawakami, C., Yamada, S., Satoh, R. and Hohdatsu, T. 2008. Antibody-dependent enhancement occurs upon reinfection with the identical serotype virus in feline infectious peritonitis virus infection. *J. Vet. Med. Sci.* **70**: 1315–1321. [Medline] [CrossRef]
18. Tsukada, R., Tsuchiyama, A., Sasaki, M., Park, C. H., Fujii, Y., Takesue, M., Hatai, H., Kudo, N. and Ikadai, H. 2013. *Encephalitozoon* infections in Rodentia and Soricomorpha in Japan. *Vet. Parasitol.* **198**: 193–196. [Medline] [CrossRef]