

A New Antifungal Isocoumarin from The Endophytic Fungus *Trichoderma* Sp. 09 of *Myoporum bontioides* A. Gray

Wensheng Li, Jiaxin Xu, Fenqi Li, Li Xu, Chunyuan Li

Department of Applied Chemistry, College of Materials and Energy (Formerly College of Science), South China Agricultural University, Guangzhou, China

Submitted: 16-11-2015

Revised: 28-12-2015

Published: 13-10-2016

ABSTRACT


Background: *Myoporum bontioides* A. Gray is a commonly used medicinal plant in China. Recently, the chemical and bioactive investigations to the endophytic fungi of this plant have led to several new compounds with antimicrobial and cytotoxic activities. To find out more active molecules, the metabolites of an endophytic fungus, *Trichoderma* sp. 09 from the root of *Myoporum bontioides* were investigated. **Materials and Methods:** The metabolites were isolated by column chromatography on silica gel, and their structures were elucidated on the basis of spectroscopic analysis [one-dimensional (1D), two-dimensional (2D)-nuclear magnetic resonance (NMR), Mass spectrometry (MS)], and by comparison with the published data. The dilution method was used for the evaluation of antifungal activity. **Results:** Four metabolites were isolated and identified as: dichlorodiaportinolide (1), dichlorodiaportin (2), diaportinol (3), and diaportin (4). Compounds 1 and 2 showed weak to high antifungal activities against *Colletotrichum musae* (Berk. and M. A. Curtis) Arx and *Rhizoctonia solani* Kühn, as compared with the positive control. **Conclusions:** Compound 1 was a new isocoumarin being worthy of consideration for the development and research of antifungal agents.

Key words: Antifungal activity, isocoumarin, *Myoporum bontioides*, *trichoderma* sp. 09

SUMMARY

- A new isocoumarin named dichlorodiaportinolide, along with dichlorodiaportin, diaportinol, and diaportin were isolated from the endophytic fungus *Trichoderma* sp. 09 of the root of *Myoporum bontioides*.
- Dichlorodiaportinolide and dichlorodiaportin showed weak to high antifungal activities against *musae* and *R. solani* (MIC values from 6.25 to 150 µg/mL).

- Dichlorodiaportinolide and dichlorodiaportin were inactive to *P. italic* and *F. graminearum* (MIC values > 200 µg/mL).



Abbreviations used: IR: Infrared Radiation, HR-ESI-MS: High resolution electrospray ionization mass spectrometry, LCMS-IT-TOF: Liquid chromatography mass spectrometry-ion trap-Time-of-flight, UV: Ultraviolet-visible, HMBC: Heteronuclear multiple bond correlation, NOE: Nuclear Overhauser effect.

Correspondence:

Dr. Chunyuan Li, College of Materials and Energy, South China Agricultural University, Guangzhou, China.

E-mail: chunyuanyi@scau.edu.cn

DOI: 10.4103/0973-1296.192204

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

Myoporum bontioides A. Gray is a small evergreen shrub distributed along the coastal regions of southern Kyushu islands through Okinawa, Taiwan, southern China, and Indochina.^[1] In China, the decoction of *Myoporum bontioides* is used as a folk medicine for antidermatosis, and as antipyretic and antipsychotic.^[2] The previous studies reported that this plant contained sesquiterpenoids, iridoids, monoterpenes, phenylethanoids, and flavonoids etc.^[3-5] Endophytes, residing in the tissues of the living plants, serve as an important source of bioactive compounds with novel structures.^[6] Recently, our chemical and bioactive investigations to the endophytic fungi of *Myoporum bontioides*, have led to several new compounds with antimicrobial^[7] and cytotoxic activities.^[8,9] In the study, the metabolites of an endophytic fungus, *Trichoderma* sp. 09 isolated from the root of *Myoporum bontioides* were investigated. As a result, a novel chlorine-containing isocoumarin named dichlorodiaportinolide (1), along with three known analogues, dichlorodiaportin (2), diaportinol (3),^[10] and diaportin (4) [Figure 1],^[11] were isolated. The isolation, structural elucidation, and the antifungal activity against several plant pathogenic fungi of the metabolites are reported herein.

MATERIALS AND METHODS


General

IR spectra was obtained with a Nicolet 5DX-Fourier transformer infrared spectrophotometer (Thermo Electron Corporation, Madison, USA). All NMR experiments were recorded in a Bruker AVIII 600MHz NMR spectrometer (Bruker BioSpin GmbH company, Rheinstetten, Germany), using deuterated dimethyl sulfoxide, or acetone, or chloroform as solvent and the residual solvent resonance as internal standard. The coupling constants (*J*) were in Hz. HR-ESI-MS, and MS was performed on LCMS-

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Li W, Xu J, Li F, Xu L, Li C. A new antifungal isocoumarin from the endophytic fungus *Trichoderma* Sp. 09 of *Myoporum bontioides* A. gray. Phcog Mag 2016;12:259-61.



IT-TOF (Shimadzu, Japan) mass spectrometer. The optical rotations were measured on a Jasco P-1020 digital polarimeter. Chromatography was carried out on a silica gel column (200-300 mesh; Qingdao haiyang chemicals Co., Ltd., Qingdao, China). All other reagents used were of analytical grade.

Fungus and cell material

The *Trichoderma* sp. 09 strain was isolated from the root of *Myoporium bontioides* A. Gray collected in Leizhou Peninsula, Guangdong Province, China. Its species was identified by comparing with the morphological characteristics with that of *Trichoderma* sp. 09^[12] This fungus is deposited in College of Materials and Energy (Formerly College of Science), South China Agricultural University, Guangzhou, China. A small scrap of agar slice with mycelium was added into a 500 mL Erlenmeyer flask containing 250 mL of GYT medium (1% glucose, 0.1% yeast extract, 0.2% peptone, 0.2% crude sea salt) aseptically, and incubated at 28 °C, 180 rpm for 6 days as seed culture. The fermentation was obtained with 100 Erlenmeyer flasks (the size of each was 1 L) and each containing a rice medium (100 mL water, 100 g rice, 0.3 g crude sea salt). 4 mL seed culture was added into each flask and then cultivated at room temperature for 30 days under static conditions.

Extraction and isolation

[200] mL methanol was added into each of the flasks and extracted for 72 h at room temperature. The solvent was concentrated to 3L with the rotary evaporator in vacuo in a 50°C water bath and extracted three times in sequence with the equal volume of ethyl acetate. The extracts were evaporated to dryness in vacuo and combined. The combined extract was chromatographed repeatedly on silica gel column using gradient elution to obtain (1), (2), (3), and (4), from the petroleum ether/ethyl acetate 80:20, 75:25, 70:30, and 85:15 (v/v) fractions, respectively.

Antifungal activity

The following four phytopathogenic fungi were used for bioassay: *Colletotrichum musae* (Berk. and M. A. Curtis) Arx. (*C. musae*), *Fusarium graminearum* Schw. (*F. graminearum*), *Penicillium italicum* Wehme (P. italicum), and *Rhizoctonia solani* Kühn (*R. solani*). They were obtained from College of Agriculture, South China Agricultural University. The quantitative tests of the antimicrobial activities of the pure compounds were performed by the broth dilution method, as described in the previous report to determine the minimum inhibitory concentration (MIC).^[7,13] The carbendazim was used as the positive control of growth inhibition, and the negative control of solvent influence was detected parallelly. The results were presented as mean values of three measurements.

RESULTS

Dichlorodiaportinolide (1)

White needles; mp 185-187°C; $[\alpha]_D^{25} +17^\circ$ (c 0.04, acetone); IR (KBr) cm^{-1} 3382, 2810, 1760, 1681, 1625, 1511, 1302. UV λ_{max} (CH₃OH) nm 330, 274, 245. ¹H (600 MHz, CD₃COCD₃) and ¹³C-NMR (150MHz, CD₃COCD₃) data, see [Table 1]. HR-ESI-MS m/z : 389.0184 ([M+H]⁺, C₁₆H₁₄Cl₂O₇, calcd.389.0188)

DISCUSSION

Compound 1 was obtained as white needles and it has a molecular formula of C₁₆H₁₄Cl₂O₇ as determined by HR-ESI-MS m/z : 389.0184 ([M+H]⁺, calcd.389.0188), indicating nine degrees of unsaturation. Its UV spectrum with absorption maxima at 245, 274 and 330 nm, and IR spectrum with absorption bands at 1681, 1625, and 1511 cm^{-1} revealed the presence of typical isocoumarin moiety in comparison with known isocoumarins.^[14,15] The ¹H and ¹³C NMR spectral data [Table 1] of 1 were closely comparable to those of 2 and 3 which displayed similar NMR data,^[10] and suggested that a hydroxyl (δ_{H} 11.02), a methoxyl (δ_{H} 3.92, δ_{C} 55.5), and a methine (δ_{H} 3.32, δ_{C} 40.3) were connected to the isocoumarin moiety at C-8 (δ_{C} 165.3), C-6 (δ_{C} 165.6), and C-3 (δ_{C} 150.9), respectively, in the same manner as 2 and 3. The isocoumarin group, along with a carbonyl (δ_{C} 174.3), accounted for 8 of the 9 elements of unsaturation in 1. It suggested that the remaining one unit of unsaturation was due to another monocyclic ring. Moreover, the HMBC correlations [Table 1] from H-9 (δ_{H} 3.32) to C-3 (δ_{C} 150.9), C-4 (δ_{C} 109.0), C-10 (δ_{C} 85.2), and C-11 (δ_{C} 36.6), from H-11 (δ_{H} 2.42) to C-9 (δ_{C} 40.3), C-10 (δ_{C} 85.2), C-12 (δ_{C} 67.5), and C-13 (δ_{C} 174.7), from H-12 (δ_{H} 4.47) to C-10 (δ_{C} 85.2), C-11 (δ_{C} 36.6), and C-13 (δ_{C} 174.7), and from 12-OH (δ_{H} 5.55) to C-11 (δ_{C} 36.6), C-12 (δ_{C} 67.5), and C-13 (δ_{C} 174.7), indicated that an α -hydroxyl- γ -lactone ring was formed by ring closure involving the oxygen atom bridged to C-10 (δ_{C} 85.2) and C-13 (δ_{C} 174.7), and its location at C-9 (δ_{C} 40.3). Thus, the only position left to place the remaining chlorine atoms was at C-14. The HMBC correlations from H-14 (δ_{H} 6.52) to C-9 (δ_{C} 40.3), C-10 (δ_{C} 85.2), and C-11 (δ_{C} 36.6), further demonstrated the connectivity of C-14 and C-10. Therefore, the planar structure of compound 1 was established as shown in [Figure 1].

Table 1: NMR Data for compound 1 (600/150MHz, CD₃COCD₃, δ /ppm, J/Hz)

	¹ H NMR	¹³ C NMR	¹ H- ¹ H COSY	HMBC
1		167.0		
3		150.9		
4	6.70 (1H, s)	109.0		C-3, 5, 4a, 8a, 9
4a		139.1		
5	6.63 (1H, d, 2.4)	101.7	H-7	C-4, 6, 7, 8a
6		165.6		
6-OCH ₃	3.92 (3H, s)	55.5		C-6
7	6.51 (1H, d, 2.4)	100.8	H-5	C-5, 6, 8, 8a
8		165.3		
8a		100.1		
9	3.32 (2H, dd, 15.0, 8.4)	40.3		C-3, 4, 10, 11, 14
10		85.2		
11	a2.42 (1H, dd, 7.2, 15.0) b3.03 (1H, dd, 7.2, 15.0)	36.6	H-11b, 12 H-11a, 12	C-9, 10, 12, 13, 14
8-OH	11.02 (1H, s)			
12	4.74 (1H, m)	67.5	H-11a, 11b	C-10, 11, 13
13		174.7		
14	6.52 (1H, s)	76.3		C-9, 10, 11
12-OH	5.55 (1H, d, 6.0)			C-11, 12, 13

The relative configuration of compound **1** was inferred as 10S* and 12S* from the NOE correlations between H-14 (δ_{H} 6.52) and H-11b (δ_{H} 3.03), between 12-OH (δ_{H} 5.55) and H11b (δ_{H} 3.03), and between 12-OH (δ_{H} 5.55) and H14 (δ_{H} 6.52), as well as no NOE correlations observed between H-14 and H-11a or between 12-OH and H11a in the NOESY spectrum. Compound **1** was named as dichlorodiaportinolide, according to its similar structure with dichlorodiaportin.^[10]

The antifungal activity of the isolated compounds was assessed quantitatively by the broth dilution method to determine the minimum inhibitory concentration (MIC).^[7,13] The results are summarized in [Table 2]. According to the results, compound **1** showed significant antifungal activity against *R. solani* with a MIC value of 6.25 $\mu\text{g mL}^{-1}$ which was the same as the positive control carbendazim and moderate activity against *C. musae* with a MIC value of 25 $\mu\text{g mL}^{-1}$. However, compound **2** only exhibited weak activities toward these two fungi with MIC values of 150 $\mu\text{g mL}^{-1}$. Neither of them was active to *P. italic* and *F. graminearum* (MICs > 200 $\mu\text{g mL}^{-1}$). The activities of compounds **3** and **4** were not detected due to their small amounts. These results suggested that compound **1**, could be valuable for a development of a new fungicide or lead compound for the treatment of some fungal infections.

Financial support and sponsorship

This work was supported by the National Natural Science Foundation of China (21102049), the Natural Science Foundation of Guangdong Province (2015A030313405, 9451064201003751), the Science and Technology Project for public welfare research and capacity building of Guangdong Province (2016A020222019), the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry ([2015]311), and the Innovation Experiment Program for University students of Guangdong Province(201510564196).

Conflicts of interest

There are no conflicts of interest

REFERENCES

- Deng Y, Yang Z, Yu Y, Bi X. Inhibitory activity against plant pathogenic fungi of extracts from *Myoporium bontioides*. A. Gray and identification of active ingredients. Pest Management Science 2008;64:203-7.

Table 2: The antifungal activity of the compounds by the MIC values ($\mu\text{g/mL}$)

	<i>F.graminearum</i>	<i>C.musae</i>	<i>P.italicm</i>	<i>R.solani</i>
1	>200	25	>200	6.25
2	>200	150	>200	150
carbendazim ^a	6.25	6.25	3.125	6.25

^apresented as positive control.

- Huang LL, Li JW, Ni CL, Gu WX, Li CY. Isolation, crystal structure and inhibitory activity against *Magnaporthe Grisea* of (2R,3R)-3,5,7-trihydroxyflavanone 3-acetate from *Myoporium Bontioides* A. Gray. Chin J Struct Chem 2011;30:1298-304.
- Wang QG, Ma CL, Zhai JJ. Furanoeudesmane-B, a new eudesmane sesquiterpenoid from *Myoporium bontioides*. Acta Cryst C 2000;56:e569.
- Kanemoto M, Matsunami K, Otsuka H, Shinzato T, Ishigaki C, Takeda Y. *et al.* Chlorine-containing iridoid and iridoid glucoside, and other glucosides from leaves of *Myoporium bontioides*. Phytochemistry 2008;69:2517-22.
- Li XZ, Li CY, Wu LX, Yang FB, Gu WX. Chemical constituents from leaves of *Myoporium bontioides*. Chin Tradit Herb Drugs 2011;42:2204-7.
- Strobel G, Daisy B. Bioprospecting for microbial endophytes and their natural products. Microbiology and Molecular Biology Reviews 2003;67:491-92.
- Wang JH, Ding WJ, Wang RM, Du YP, Liu HL, Kong XH. *et al.* Identification and bioactivity of compounds from the mangrove endophytic fungus *Alternaria* sp. Marine Drugs 2015;13:4492-504.
- Wang JH, Cox DG, Ding WJ, Huang GH, Lin YC, Li CY. *et al.* Three new resveratrol derivatives from the mangrove endophytic fungus *Alternaria* sp. Marine Drugs 2014;12:2840-50.
- Li CY, Gong B, Cox DG, Li CL, Wang JH, Ding WJ. *et al.* Dichlorodiaportinol A, a new chlorine-containing isocoumarin from an endophytic fungus *Trichoderma* sp. 09 from *Myoporium bontioides* A. Gray and its cytotoxic activity. Pharmacognosy magazine 2014;10:s153-8.
- Larsen TO, Breinholt J. Dichlorodiaportin, diaportinol, and diaportinic acid: three novel isocoumarins from *Penicillium nalgioense*. Journal of Natural Products 1999;62:1182-4.
- Mantle PG. Biosynthesis of diaporthin and orthosporin by *Aspergillus ochraceus*. Phytochemistry 2001;57:165-9.
- Amarean K, Bhagat N, Madhuri S, Srivastava K. RCI Isolation and characterization of *Trichoderma* spp. for antagonistic activity against root rot and foliar pathogens. Indian Journal of Microbiology 2012;52:137-44.
- Feng CL, Ma YM. Isolation and anti-phytopathogenic activity of secondary metabolites from *Alternaria* sp. FL25, an endophytic fungus in *Ficus carica*[J]. Chin J Appl Environ Biol 2010;16:76-8.
- Ichihara A, Hashimoto M, Hirai T, Takeda I, Sasamura Y, Sakamura S. *et al.* Structure, synthesis, and stereochemistry of (+)-orthosporin, a phytotoxic metabolite of *Rhynchosporium orthosporium*. Chem Lett 1989;18:1495-8.
- Yang SX, Gao JM, Zhang Q, Laatsch H. Toxic polyketides produced by *Fusarium* sp., an endophytic fungus isolated from *Melia azedarach*. Bioorganic and Medicinal Chemistry Letters 2011;21:1887-9.



Chunyuan Li

ABOUT AUTHORS

Dr. Chunyuan Li, Obtained his Ph. D. degree in 2006 from School of Chemistry and Chemical Engineering, Sun Yat-sen University, China. Currently, he is positioned as associate professor of College of Materials and Energy, South China Agricultural University, China. Dr. Li is working in the area of Pharmacognosy and Chemistry of Natural Products, mainly in isolation, structure elucidation and bioactivity evaluation of natural products from marine plant endophytic fungi.