

ORIGINAL ARTICLE

Metal stressors consistently modulate bacterial conjugal plasmid uptake potential in a phylogenetically conserved manner

Uli Klümper^{1,4}, Arnaud Dechesne¹, Leise Riber², Kristian K Brandt³, Arda Gülay¹, Søren J Sørensen² and Barth F Smets¹

¹Department of Environmental Engineering, Technical University of Denmark, Lyngby, Denmark;

²Department of Biology, University of Copenhagen, Copenhagen, Denmark and ³Department of Plant and Environmental Sciences, University of Copenhagen, Frederiksberg, Denmark

The environmental stimulants and inhibitors of conjugal plasmid transfer in microbial communities are poorly understood. Specifically, it is not known whether exposure to stressors may cause a community to alter its plasmid uptake ability. We assessed whether metals (Cu, Cd, Ni, Zn) and one metalloid (As), at concentrations causing partial growth inhibition, modulate community permissiveness (that is, uptake ability) against a broad-host-range IncP-type plasmid (pKJK5). Cells were extracted from an agricultural soil as recipient community and a cultivation-minimal filter mating assay was conducted with an exogenous *E. coli* donor strain. The donor hosted a *gfp*-tagged pKJK5 derivative from which conjugation events could be microscopically quantified and transconjugants isolated and phylogenetically described at high resolution via FACS and 16S rRNA amplicon sequencing. Metal stress consistently decreased plasmid transfer frequencies to the community, while the transconjugal pool richness remained unaffected with OTUs belonging to 12 bacterial phyla. The taxonomic composition of the transconjugal pools was distinct from their respective recipient communities and clustered dependent on the stress type and dose. However, for certain OTUs, stress increased or decreased permissiveness by more than 1000-fold and this response was typically correlated across different metals and doses. The response to some stresses was, in addition, phylogenetically conserved. This is the first demonstration that community permissiveness is sensitive to metal(loid) stress in a manner that is both partially consistent across stressors and phylogenetically conserved.

The ISME Journal (2017) 11, 152–165; doi:10.1038/ismej.2016.98; published online 2 August 2016

Introduction

Horizontal gene transfer among bacterial species is recognized as a major evolutionary process (Zhaxybayeva and Doolittle, 2011). Mobile genetic elements can spread across diverse bacterial phyla (Klümper *et al.*, 2015) linking distinct genetic pools (Halary *et al.*, 2009; Norman *et al.*, 2009). A main parameter in assessing the ecology of plasmid transfer is community permissiveness, which refers to the ability of a community to receive a plasmid, both in terms of transfer frequency and transconjugal phylogeny (Musovic *et al.*, 2010).

The spread of plasmid-mediated antibiotic resistance across bacterial communities has recently been identified as a major threat to human health by the WHO (World Health Organization) (WHO, 2014). By carrying sets of genes coding for resistance, conjugal plasmids can facilitate bacterial community adaptation to stress imposed by antibiotics or by other toxicants such as metals and metalloids at elevated concentrations (Sørensen *et al.*, 2005; Heuer and Smalla, 2012).

Here, stress is defined as the exposure of bacteria to a toxicant at a dose that causes a transcriptional response (Cases and de Lorenzo, 2005) as a result of the toxicants incompatibility with normal biological functions (Richardson, 1993). Low exposure levels might act as a stimulant or signal for the transcription of single genes (Pérez-Martín and de Lorenzo, 1996). An elevated dose of the stressor may cause the activation of entire regulons that are part of global stress-response mechanisms. Known stress-responses involve switching to slow growth, biofilm formation or the activation of efflux pumps

Correspondence: BF Smets or U Klümper, Department of Environmental Engineering, Technical University of Denmark, Miljøvej 113, 2840 Lyngby, Denmark.

E-mail: bfm@env.dtu.dk or u.klumper@exeter.ac.uk

⁴Current address: European Centre for Environment and Human Health, University of Exeter, Medical School, Royal Cornwall Hospital, Truro, UK.

Received 4 March 2016; revised 30 May 2016; accepted 7 June 2016; published online 2 August 2016

(Mah and O'Toole, 2001). We hypothesize that, additionally, bacterial plasmid uptake potential or individual permissiveness, might increase as a response to stress. This increased permissiveness could either be the direct or indirect effect of an evolutionary evolved stress-response mechanism.

Single strain experiments have shown modulation in gene uptake as a result of stress exposure. In *Streptococcus pneumoniae*, exposure to antibiotics caused increased competence and increased promiscuity toward foreign DNA (Slager *et al.*, 2014). Heat-shock treatment of Gram-positive Corynebacteria enhanced their receipt of an IncP-type plasmid from a Gram-negative *E. coli* donor, likely due to inactivation of restriction-modification systems (RM) (Schafer *et al.*, 1990). Similarly, in the Gram-negative *Pseudomonas putida*, pre-exposure to SDS increased its ability to receive and maintain plasmids, possibly by repressing RM systems (Arango Pinedo and Smets, 2005). Cells with a turned-off RM system can become hypersusceptible to foreign DNA or plasmid receipt (Schafer *et al.*, 1990; Corvaglia *et al.*, 2010; Roer *et al.*, 2015). Also, in *Bacillus subtilis*, sub-inhibitory concentrations of ethanol activated the conjugative transposition of transposon Tn916 (Seier-Petersen *et al.*, 2013) encoding tetracycline resistance (Franke and Clewell, 1981). In contrast, in *Escherichia coli*, cell-envelope-targeted stress induced the expression of CRISPR-associated (CRISPR-cas) genes which would serve as an intracellular defense against foreign DNA (Perez-Rodriguez *et al.*, 2011), thus decreasing permissiveness. While these mechanisms are described for specific strains under specific stress conditions, there have been no studies to date that evaluate modulation of permissiveness as a general bacterial stress-response mechanism using complex bacterial communities and multiple stressors.

A typical environmental stress for soil microbes are toxic metals accumulating to elevated concentrations due to agronomic practices, industrial activities or atmospheric deposition (Nicholson *et al.*, 2003; Zhao *et al.*, 2015). Metal toxicity is mainly caused by metal ions disrupting iron-sulfur clusters of metallo-enzymes (Macomber and Imlay, 2009; Macomber and Hausinger, 2011; Xu and Imlay, 2012). Metals, at elevated concentrations, have been documented to affect the composition and function of soil bacterial communities (Giller *et al.*, 1998; Gans *et al.*, 2005; Berg *et al.*, 2012), although development of community tolerance is common even when no effects are observed at the levels of activity or community composition (Brandt *et al.*, 2010). Selection for adaptive plasmids within microbial communities has been described as a long-term effect to metal exposure (Campbell *et al.*, 1995; Giller *et al.*, 1998). However, the existence and magnitude of a short-term effect of metal stress on the permissiveness of bacterial communities remain to be explored.

In this study, we combined a cultivation-minimal assay to measure community-wide permissiveness

(Klümper *et al.*, 2015) with normalized stress exposure to five of the most environmentally relevant metal(oid)s (Cu^{2+} , Cd^{2+} , Ni^{2+} , Zn^{2+} , AsO_3^{3-}). This allowed for direct comparison of short-term effects of metal stress on the permissiveness toward the *gfp*-labeled, model broad-host-range IncP-1 ϵ plasmid pKJK5 (Bahl *et al.*, 2007) introduced through an *E. coli* donor strain in a reference soil bacterial community. Plasmid transfer frequencies and the phylogenetic composition of the transconjugal pools were compared with corresponding controls to assess stress-induced modulation of permissiveness of the community and of individual community members.

Materials and methods

Soil sampling

Soil ($16 \pm 1\%$ clay, $15 \pm 1\%$ silt, 43% fine sand and $26 \pm 1\%$ coarse sand, with a $\text{pH}(\text{H}_2\text{O})$ of 7.16 ± 0.13) was sampled from unfertilized control plots ($n = 3$) of the CRUCIAL agricultural field site (December 2013, Taastrup, Denmark) (Poulsen *et al.*, 2013; Lekfeldt *et al.*, 2014). Five hundred grams of soil were collected at three locations at each plot from a depth of 5–15 cm. The nine soil samples were pooled, sieved through a 1 mm^2 mesh filter, homogenized and stored at 4°C for up to one month before conducting the experiments.

Soil bacterial community extraction

Soil bacterial communities were recovered from 30 g sub-samples of homogenized soil by Nycodenz (Nycomed Pharmaceuticals, Zürich, Switzerland)-extraction (Lindahl and Bakken, 1995). Extracted cells were resuspended in sterile 0.9% NaCl solution, filtered, washed, quantified using a Thoma counting chamber (Sigma-Aldrich, St Louis, MO, USA) and adjusted to 10^7 cells per ml to quantify metal-induced stress (measured as growth rate inhibition) or to be used as recipients in the filter mating assay.

[^3H]leucine incorporation inhibition assay

A [^3H]leucine incorporation assay (Supplementary Text 1) was used to estimate the concentrations of individual metal(oid)s inhibiting soil bacterial growth rates by 20% (IC_{20}) and 50% (IC_{50}), respectively (Lekfeldt *et al.*, 2014). The experimental data linking leucine incorporation to metal dose were fitted with a four parameter log-logistic dose-response curve using the *drc* (Analysis of Dose-Response Curves) package for R (Knezevic *et al.*, 2007) (Supplementary Figure 1).

Plasmid and donor strain

Cells of *E. coli* MG1655::*lac^q-pLpp-mCherry-Km^R* (Klümper *et al.*, 2015) carrying the IncP-1 ϵ broad-host range plasmid pKJK5::*gfpmut3b* (Bahl *et al.*, 2007) were used as donors. Plasmid pKJK5 has an

extremely broad transfer range, including both Gram-negative and Gram-positive phyla (Klümper *et al.*, 2015). In addition, pJK5 does not encode metal-related resistance mechanisms (Bahl *et al.*, 2007).

The plasmid was marked with an entranceposon-derived (Bahl *et al.*, 2009) genetic tag that carries a *LacI^q* repressible promoter upstream the conditionally expressed *gfpmut3b* gene, encoding the green fluorescent protein (GFP). The plasmid donor strain was chromosomally tagged with a gene cassette encoding constitutive red fluorescence, expressed by the *mCherry* gene, and constitutive *lacI^q* production. As a result, *gfp* expression is repressed in the donor strain, but on successful conjugal transfer to a soil bacterium, plasmid-encoded *gfp* expression is de-repressed, resulting in green fluorescent cells or microcolonies, which can be retrieved by fluorescence-activated cell sorting (FACS) or detected by fluorescence microscopy, respectively (Sørensen *et al.*, 2005; Klümper *et al.*, 2015). The donor strain was grown overnight in LB medium supplemented with trimethoprim ($30 \mu\text{g ml}^{-1}$), harvested by centrifugation, washed in 0.9% NaCl solution, adjusted in cell density (OD_{600}) and used in filter mating assays as described previously (Klümper *et al.*, 2014a). The viability and plasmid transfer ability of the donor strain under all metal stress conditions were verified in control mating assays with an untagged *E. coli* MG1655 recipient through microscopic observations.

Filter matings

The extracted soil bacterial community was challenged with the exogenous plasmid via solid-surface filter matings (Musovic *et al.*, 2010; Klümper *et al.*, 2014a) to maximize cell-to-cell contact, as physical barriers would limit the contact between freshly introduced plasmid donors and potential recipients in an intact soil matrix (Dechesne *et al.*, 2005). Compared with the original protocol (Klümper *et al.*, 2014a) the initial ratio of donor to recipient cells was increased to 10:1 to maximize recipient cell/microcolony contact with the donor cells during the 48 h incubation period.

Filters were placed on an agar-solidified 10% soil extract medium after Musovic *et al.* (2010). The limited amount of nutrients resulted in monolayer cell coverage of the filter after 48 h as confirmed with confocal laser scanning microscopy (CLSM). With an initial density of $\sim 300\,000$ cells per mm^2 and assuming an average cell surface area of $1 \mu\text{m}^2$, the average number of cell doublings per inoculated cell during incubation was thus estimated at < 5 .

Filter matings were performed on medium without metal amendment (controls) or supplemented with CuSO_4 , NiSO_4 , ZnSO_4 , CdCl_2 or Na_2HAsO_3 at concentrations corresponding to IC_{20} and IC_{50} based on [^3H]leucine incorporation of the recipient community. Filters were incubated for 48 h at 25°C ,

before 72 h storage at 4°C for GFP maturation. Conjugation events were checked by epifluorescence stereomicroscopy and CLSM (Musovic *et al.*, 2010).

Transfer frequency quantification

Conjugation events were visualized by stereomicroscopy and quantified by automated image analysis (Image Pro Plus 7.1; Media Cybernetics, Silver Spring, MD, USA) (Supplementary Text 1) as previously described (Musovic *et al.*, 2010; Klümper *et al.*, 2014b). The number of detected transfer events was scaled up to the total filter area and transfer frequency was calculated by dividing by the number of originally added recipient cells (Supplementary Figure 2).

FACS and sequencing

For each mating condition, cells from triplicate filters were combined in 2 ml 0.9% NaCl solution and detached by vortexing at 3000 r.p.m. for 30 s. Transconjugant cells and recipient community were sorted using FACS (Supplementary Text 1, Supplementary Figure 2). Gating and sorting of transconjugants for each of the combinations were performed based on bacterial size, green fluorescence and exclusion of red fluorescent donor cells as described earlier (Klümper *et al.*, 2015).

Recipient community cells were sorted from the same samples using the same conditions, including both colorless recipient and green fluorescent transconjugal cells and excluding red fluorescent donor cells. In all cases, a minimum of 20 000 *gfp*-expressing transconjugal cells or recipient cells were sorted. Sorted bacterial cells were collected, lysed and subject to tag-encoded 16s rRNA amplicon sequencing (Supplementary Text 1) (Klümper *et al.*, 2015).

Sequence analysis

All sequences documented in this study were deposited in the European Nucleotide Archive under study accession number PRJEB13628.

Sequence analysis was carried out using mothur v.1.32.1 (Schloss *et al.*, 2009) and the MiSeq SOP (Kozich *et al.*, 2013) as accessed on 1 December 2014 from http://www.mothur.org/wiki/MiSeq_SOP. Sequences were classified based on the RDP classifier (Wang *et al.*, 2007).

Diversity was assessed based on observed operational taxonomic units (OTUs) at 97% sequence similarity. Phylogenetic trees were constructed using iTOL (<http://itol.embl.de/>; Letunic and Bork, 2007).

NMDS plots showing the phylogenetic distance between samples were produced based on the theta Yue Clayton dissimilarity index (Yue and Clayton, 2005) using mothur software and clustering of samples was tested for significance by analysis of molecular variance (AMOVA) (Excoffier *et al.*, 1992). Based on NMDS clustering and because the number

of replicates per condition differed, replicates were combined for subsequent statistical analysis.

OTU-level permissiveness analysis

At the community level, precise estimates of initial recipient abundance (R) and number of conjugation events (T) made it possible to measure permissiveness accurately. This was impossible at the OTU level, where only relative abundances of an OTU in the recipient community (after mating incubation) and in the transconjugal pool were measured (Supplementary Figure 2). These values are not solid estimators of the initial recipient abundance or of the number of conjugation events for a specific OTU as they are affected by the growth rate of that OTU relative to other OTUs, the possibility of multiple transfers within an OTU microcolony, and the permissiveness of other recipients. Increases or decreases in recipient OTU cell densities during the mating incubations *per se* (that is, due to a competitive advantage or disadvantage compared with other OTUs) are possible, but equivalent effects would hold for the transconjugant cell densities of that OTU, as the pKJK5 plasmid does not encode for any growth-beneficial phenotype. Hence, an apparent OTU-level permissiveness (AP) can be calculated, based on the relative recipient and transconjugant abundances at the end of the mating incubation:

$$AP_i = \frac{\text{rel. abundance in transconjugal pool of } OTU_i}{\text{rel. abundance in recipient community of } OTU_i} = \frac{T_i}{R_i}$$

Although this apparent permissiveness is not only affected by transfer, but also by the fact that multiple transfers can occur in a microcolony and by recipient and transconjugant growth; these phenomena are expected to occur both under the reference and the metal stress conditions. We therefore use the ratio (δ) of an OTU's apparent permissiveness under stress ($AP_{i \text{ stress}}$) and reference conditions ($AP_{i \text{ ref}}$) to describe stress-induced permissiveness changes in that OTU:

$$\delta_i = \frac{AP_{i \text{ stress}}}{AP_{i \text{ ref}}} = \frac{T_{i \text{ stress}}/R_{i \text{ stress}}}{T_{i \text{ ref}}/R_{i \text{ ref}}}$$

If the stress has no effect on an OTU's permissiveness, δ would be equal to 1 while values above or below 1 would indicate an increase or decrease in permissiveness under the stress condition, respectively.

To analyze whether the variability of δ -values was phylogenetically conserved, we created a maximum likelihood tree of the dominant (average relative abundance in all transconjugal pools >0.05%) transconjugal OTUs based on similarity of their δ -values across stresses. Phylogenetic conservation of permissiveness response for each stress condition was tested using the analysis of trait module of Phylocom v4.2 (Webb *et al.*, 2008).

Exploratory statistics were carried out in R using the package car (Companion to Applied Regression,

version 2.0–25). The package psych (Revelle, 2014) was used to calculate Spearman correlations between parameters and test their significance, after adjusting the P -values for multiple testing (Hommel, 1988).

Results

Metal stress reduces transfer frequencies more than metabolic activity

Inhibitory metal concentrations (IC_{20} or IC_{50}) applied in the mating experiments were determined by dose-response modeling of [3H]leucine incorporation data (Supplementary Figure 1). Molar metal toxicity increased in the order (Table 1): $AsO_3^{3-} < Zn^{2+} < Cd^{2+} < Cu^{2+} < Ni^{2+}$. Matings conducted with metal stress will subsequently be identified with a combination of the elemental symbol and the inhibitory level (for example, As20 see Table 1) and compared with the non-stressed reference condition.

Conjugation events were detected as green fluorescent cells and microcolonies in microscopic images of the filter matings. Red fluorescent donor cells were detected in all matings, albeit with a slight decrease in density under stress conditions (Figure 1). Flow cytometric counts of intact red fluorescent donor cells confirmed that donor cell counts exceeded total recipient counts under each condition to ensure that each recipient microcolony was in contact with multiple donor cells.

Conjugation events were detected at ~ 1 in 15 000 ($6.8 \times 10^{-5} \pm 3.8 \times 10^{-5}$) of the initial soil recipient bacterial cells under reference conditions. This frequency decreased under stress conditions for all metals (Figure 1). Community permissiveness significantly decreased ($P < 0.05$) under conditions of metal stress in a dose-dependent manner (Figure 2). This decrease was metal specific and was more severe than the effect of the metal on community-wide metabolic activity (Table 1, Figure 2). Cu50, Zn50 (both >90% reduction) and Cd50 (no transfer detected) caused the largest permissiveness reduction.

Reduction in community permissiveness could result from a general decrease of transfer across all taxa or from taxon-specific modulation of individual permissiveness. We FACS-sorted at least 20 000 transconjugant cells from each mating, which were

Table 1 Inhibitory concentrations causing 20% (IC_{20}) and 50% (IC_{50}) bacterial growth inhibition as inferred by dose-response modeling of [3H]leucine incorporation data (Supplementary Figure 1)

Metal	IC_{20} (μM)		IC_{50} (μM)	
AsO_3^{3-}	40.5	(As20)	125.2	(As50)
Cd^{2+}	12.6	(Cd20)	63.6	(Cd50)
Ni^{2+}	3.7	(Ni20)	11.5	(Ni50)
Zn^{2+}	24.7	(Zn20)	80.7	(Zn50)
Cu^{2+}	6.9	(Cu20)	28.9	(Cu50)

Treatment names used throughout this paper are indicated in brackets.




Figure 1 Images of plasmid transfer from the *E. coli* MG1655 donor strain to soil bacteria at reference condition (no stress) (left panel) or as affected by metal stress. Type of metal (Ni, As or Cu) and degree of stress are indicated in the top left corner of the other panels; number refers to % inhibition of [³H]leucine incorporation for the Nycodenz-extracted recipient bacterial community. Donor bacteria, chromosomally tagged with *mCherry* appear red, recipient bacteria appear initially colorless until successful plasmid receipt and subsequent *gfp* expression, transconjugants appear green.

subjected to 16S rRNA amplicon sequencing to analyze their phylogenetic composition.

All transconjugal pool sequence libraries showed coverage above 98%, and no subsampling effect was found at a depth 50,000 sequences. Ten subsampled sets of each sample showed no significant dissimilarity ($P > 0.99$) based on weighted UNIFRAC comparisons when compared with the original sequencing library. Thus, further analysis of transconjugal pools was done at a subsampling depth of 50 000 sequences.

Diversity of transconjugal pools is maintained in spite of transfer frequency reduction

We report an extremely diverse phylogenetic range of transconjugants. Across all pools, 206 permissive OTUs, distributed over 12 phyla, were identified (Figure 3a). These included the dominant Proteobacteria (α , β , γ and δ) and Bacteroidetes (Figure 3b), as well as other Gram-negative phyla (Acidobacteria, Nitrospira, Fusobacteria, Planctomycetes, Gemmatimonadetes and Verrucomicrobia), diverse Gram-positive phyla (Firmicutes, Actinobacteria and Chloroflexi) and the candidate phylum Cand. Saccharibacteria.

Plasmid transfer was detected both in abundant and rare taxa of the initial recipient community (Figure 3a). The phylogenetic composition of all transconjugal pools was similar at the phylum level (Figure 3b). The only exception was As50, where the Bacteroidetes phylum was nearly absent from the transconjugal pool, in spite of its abundance in the corresponding recipient community after incubation. Larger shifts were identified in the less abundant phyla with Gram-positive Firmicutes (Figure 3b) and Actinobacteria (Figure 3C, Supplementary Table 2)




Figure 2 Reduction of pKJK5 transfer frequency under metal stress conditions as compared with the reference condition (0% value). Type of metal (As, Cd, Cu, Ni or Zn) and degree of stress are indicated on individual bars; numbers refer to inhibitory concentrations decreasing [³H]leucine incorporation rates of the recipient bacterial community by 20% (IC₂₀, gray bars) and 50% (IC₅₀, black bars). Plasmid transfer frequencies were calculated based on analysis of 90–150 images per condition.

increasing in richness and up to eightfold in abundance under several stress conditions.

The average richness (α -diversity) of the transconjugal pools was 196.0 ± 14.4 OTUs, irrespective of transfer frequency reduction (Figure 4). While $38.5 \pm 2.2\%$ of the recipient community OTUs (identified after 48 h incubation) were permissive to pKJK5 under reference conditions, this increased to 57–96% under metal stress (Figure 4). This increase resulted from a dramatic ($> 50\%$, $P = 0.0053$) decrease in the recipient community richness under metal stress conditions. As 94–97% of the transconjugal sequences under metal stress conditions




Figure 3 Phylogenetic composition of transconjugal pools. **(a)** Phylogenetic tree showing all 206 identified transconjugal OTUs. Colors of the branches mark different phylogenetic groups. The 38 most abundant OTUs (>0.05% relative abundance across all pools), used in subsequent OTU permissiveness analysis are shown in white letters. The blue heatmap-circle at the periphery of the tree represents log-transformed relative OTU abundance in the reference recipient community. **(b)** Phylogenetic composition of the transconjugal pools obtained at various stress conditions. **(c)** Relative abundance of rare phyla (<1% mean relative abundance) in transconjugal pools for the various stress conditions. Experimental treatments (type of metal and degree of stress) are indicated below the bars for **b** and **c**; numbers refer to inhibitory concentrations decreasing [³H]leucine incorporation rates of the recipient bacterial community by 20% (IC₂₀) and 50% (IC₅₀). ‘Ref’ denotes reference condition (unstressed control).




Figure 4 Observed number of unique operational taxonomic units (OTUs; 97% sequence similarity) as a function of reduction in plasmid transfer frequency caused by metal stress. The OTU richness of the transconjugal pools (triangles) and the corresponding sorted soil recipient communities (diamonds) is shown. Experimental treatments (type of metal and degree of stress) are indicated; numbers refer to inhibitory concentrations decreasing [³H]leucine incorporation rates of the recipient bacterial community by 20% (IC₂₀) and 50% (IC₅₀).

belonged to OTUs also found in the reference transconjugal pool, the OTUs permissive to pKJK5 survived better under metal stress than the non-permissive OTUs. However, 86 additional OTUs, rare in abundance, were exclusively detected in transconjugal pools under metal stress conditions.

Transconjugal pools cluster according to stress

The overall OTU richness and phylum level composition of the transconjugal pools remained unchanged under most stress conditions. However, at the OTU level their phylogenetic composition was significantly altered by stress. Some OTUs in the recipient communities were not permissive to plasmid pKJK5 under any of the tested conditions and were excluded prior to ordination (Figure 5). Transconjugal pools nevertheless clustered significantly apart from their respective recipient communities (AMOVA test, $P < 0.001$).

Replicate transconjugal pools and recipient communities consistently grouped together dependent on the type and degree of the metal stress. Three main clusters were observed among the transconjugal pools (Figure 5). Cluster I, comprised of the Cu50, Cd20, Ni20 and Ni50 replicate pools, grouped significantly apart ($P < 0.001$) from the reference transconjugal pools. Replicate transconjugal pools of Cu20 formed a separate Cluster II with a phylogenetic composition significantly different ($P < 0.05$) from pools obtained at any other stress condition. By contrast, the transconjugal pools of As20, As50 and Zn20 did not significantly shift in phylogenetic composition compared with the reference pool ($P > 0.05$), with which they formed Cluster III.




Figure 5 Non-metric 2-dimensional scaling analysis (NMDS) revealing distinct clustering of transconjugal pools (circles) and recipient communities (triangles). Three transconjugal clusters are identified and named with roman numerals. Experimental treatments (type of metal and degree of stress as operationally defined by % inhibition of [³H]leucine incorporation rates) are shown by color and size of symbols. Amplicon pools were subsampled at 50 000 sequences. OTUs from the recipient pool that were not represented in any transconjugal pool were removed prior to subsampling and ordination based on weighted OTU abundance using the Theta Yue Clayton (Yue and Clayton, 2005) algorithm.

Stress-specific responses at OTU level are resolved by transconjugal taxonomy

The relative abundance of transconjugal OTUs varied across metal treatments (Figure 5). Two factors may have caused these variations: a metal-specific shift in the composition of the recipient community or a metal-specific modulation in the permissiveness of some OTUs. We investigated whether the second mechanism, that is, modulation of OTU permissiveness, could be detected. Hence, we analyzed, for each of the 38 dominant transconjugal OTUs (Figure 3), the change in permissiveness (δ) between the stressed conditions and the unstressed reference.

For some OTUs, under certain stress conditions, permissiveness increased over 1000-fold or decreased up to 10 000 fold (Figure 6). The permissiveness of a high number of OTUs (138 of 342 tested combinations) increased or decreased more than 10-fold in response to stress. Under some conditions (As20, Cu20, Cu50, Ni20 and Zn20), permissiveness increased slightly for most of the OTUs (Supplementary Figure 4). In contrast, arsenic at high concentration slightly decreased the permissiveness of the majority of the OTUs.

Individual OTUs tended to respond in a consistent manner to different metal stresses. Of 15 pairwise correlations of the δ -values between the metal stresses (Cd20, Cu20, Cu50, Ni20, Ni50 and Zn20), 12 were significant ($P < 0.05$) (Figure 6). The δ -values for Zn50 correlated poorly with the others, except for Ni20, mainly caused by few Rhizobiales OTUs with a highly decreased permissiveness. The metalloid arsenic at As50 did not correlate well with the metals; here all OTUs belonging to the Bacteroidetes phylum displayed a strong decrease in permissiveness (Figure 6). Despite a general consistency across most




Figure 6 Pairwise correlation analysis of the modulation of the permissiveness ($\log(\delta)$) across metal treatments for all 206 bacterial OTUs to explore whether two stress conditions resulted in similar response. Each circle symbol in the diagrams corresponds to one OTU plotted according to its $\log(\delta)$ under the two compared conditions. Lines show linear regression. Star(s) in the top left corner indicate the significance of the Spearman correlation ($*P < 0.05$, $**P < 0.005$) after correction for multiple testing (Hommel, 1988). Experimental treatments (type of metal and degree of stress) are indicated by numbers referring to inhibitory concentrations decreasing [^3H]leucine incorporation rates of the recipient bacterial community by 20% (IC_{20}) and 50% (IC_{50}).

metal stresses, almost none of the 38 most abundant OTUs responded similarly to all stress conditions (Figure 7). Only one OTU, a member of the alphaproteobacterial Rhodobacterales family, responded to every applied stress with a significant decrease in plasmid uptake (always >100 fold and up to 5000 fold) (Figure 7).

Phylogenetically similar OTUs responded similarly to metal stress. All Bacteroidetes OTUs responded similarly to the metal exposure: for all stresses except As50, the plasmid uptake ability of the OTUs belonging to the Bacteroidetes phylum increased by >10 -fold (Figure 7). The only Gram-positive OTU among the 38 most abundant transconjugant OTUs, a member of the Firmicutes phylum, responded in a similar way. For most of the transconjugal OTUs in the phylum Proteobacteria, the stress response was more variable. Four of these OTUs, similar to Bacteroidetes, became increasingly permissive under stress conditions. Especially an OTU classified as Xanthomonadales responded with an increased permissiveness to most stresses and showed the highest relative increase in permissiveness (>1000 -fold for Cu50). Most Alphaproteobacteria, such as the

Rhizobiales, displayed little permissiveness modulation under stress conditions.

The distribution of δ -values across the community's phylogenetic tree was significantly correlated with OTU phylogeny ($P < 0.05$) for two (Zn20, Ni20) of the nine stress conditions and additionally showed a tendency for phylogenetic conservation ($P = 0.074\text{--}0.258$) for six of the remaining seven conditions (Supplementary Table 1).

Discussion

Broad-host range plasmids introduced to microbial communities can spread among a wide variety of Gram-negative and Gram-positive bacterial species (De Gelder *et al.*, 2005; Musovic *et al.*, 2014; Shintani *et al.*, 2014; Klümper *et al.*, 2015). Here we demonstrate pKJK5 transfer to an extremely diverse fraction of a soil bacterial community both under reference and metal stressed conditions. The transconjugant pool included abundant, as well as rare species-level OTUs and covered 12 different phyla, thereby expanding the realized transfer range of IncP-type




Figure 7 Maximum likelihood tree clustering of the 38 most abundant OTUs in the transconjugal pools (relative average abundance > 0.05%) based on similarity in their changes in permissiveness under different stress conditions. The heatmap shows the relative change in permissiveness (δ value) of each OTU: an increased plasmid receipt response is shown in red, and a decreased one in green. The average relative abundance of an OTU across all transconjugal pools is shown in violet (right). Their taxonomy is indicated by the color of the font (for Proteobacteria, the class is shown in brackets). Experimental treatments (type of metal and degree of stress) are indicated above the heatmap with numbers referring to inhibitory concentrations decreasing [^3H]leucine incorporation rates of the recipient bacterial community by 20% (IC_{20}) and 50% (IC_{50}).

plasmids in soil beyond our previous observations with two more phyla, Nitrospira and Cand. Saccharibacteria (Klümper *et al.*, 2015). About 86 transconjugant OTUs were only detected under specific stress conditions, which might stem from a stimulation of permissiveness in these OTUs associated with metal stress exposure. Because of the low abundance of these rare OTUs (<0.01% rel. abundance) this possibility requires confirmation.

More importantly, we show, for the first time, that metal stress can modulate – increase or decrease – the permissiveness of different members of a soil bacterial community toward an IncP plasmid. Elevated levels of plasmids and plasmid-encoded genes have been reported in soil communities subject to long-term anthropogenic selective pressure through agricultural use (Agersø *et al.*, 2006; Heuer *et al.*, 2011; You *et al.*, 2012), and exposure to long-term metal stress was shown to increase plasmid mobilization capacity of a soil community (Top *et al.*, 1995). However, these long-term effects offer no direct evidence for increased permissiveness toward plasmids because they may also be caused by indirect selection of specific beneficial phenotypes. By employing a plasmid (Bahl *et al.*, 2009) that does not encode a beneficial phenotype when exposed to the chosen stressors and examining transfer directly, we aim at effectively uncoupling plasmid transfer from selective forces.

While earlier studies revealed that the plasmid uptake of individual strains can differ under stress conditions (Arango Pinedo and Smets, 2005; Slager *et al.*, 2014), this is the first assessment of OTU permissiveness in a community context. Taking advantage of our high-resolution approach, we demonstrated that stress-induced modulation of permissiveness is indeed unique for each OTU.

The composition and density of the transconjugal pools is affected by (1) transconjugant growth and (2) *de novo* transconjugant formation by plasmid transfer to recipient cells. Both processes can be affected by heavy metal exposure. While we employ a cultivation-minimal assay with a high-initial bacterial density on the filters, growth may occur on the filters. However, transconjugant cells are expected to grow at the same rate as recipient cells of the same OTU, as pJK5 plasmid carriage does not selectively affect growth. An artefactual underestimation of *T/R* due to recipient growth is, therefore, unlikely as similar growth is expected for the respective transconjugant. An overestimation of *T/R*, due to transfer occurring before die-off of heavy-metal-sensitive recipient OTUs, is unlikely as transfer can only be detected in transconjugant cells that maintain sufficient metabolic activity to ensure GFP production. Hence, *T-to-R* ratios can be compared across treatment conditions, to evaluate permissiveness differences across OTUs.

Artificial permissiveness changes could be inferred if the abundance of non-participatory (those with zero permissiveness) recipient OTUs would dramatically

increase or decrease due to selective growth or death. However, 94–97% of all sequences within each recipient pool belonged to OTUs also found in the corresponding transconjugal pools, and an effect of changes in non-participatory OTU cell abundance was excluded by calculating alternative δ -values based on relative abundances after excluding all non-participatory OTUs for each treatment. These corrected δ -values are virtually identical to those calculated for the whole recipient community (Supplementary Figure 3) due to the low abundance of non-participatory OTUs. In addition, such effects would manifest in a consistent increase or decrease in all δ -values across all OTUs within one treatment. For all treatments, δ -values were distributed with a median insignificantly different from 1, the no-effect level on permissiveness if all OTUs would participate in plasmid transfer (Supplementary Figure 4). Thus, changes in the recipient pool composition did not affect the measured δ -values to a significant degree.

Our experimental conditions resemble the agronomic practice of manure application to soil. Pig manures (especially from piglets) often contain elevated levels of Cu and Zn (Nicholson *et al.*, 2003) used as feed additives (Jondreville *et al.*, 2003). These are introduced to soil alongside a high load of fecal bacteria hosting various antibiotic and metal resistance plasmids (Smalla *et al.*, 2000; Zhu *et al.*, 2013). We similarly introduce an enterobacterial exogenous donor strain to a soil bacterial community while simultaneously challenging it with metal stress. However, we remove the physical barriers that would limit the contact between freshly introduced plasmid donors and potential recipients in an intact soil matrix (Dechesne *et al.*, 2005), thus assessing the full potential of the soil bacterial permissiveness.

Under reference conditions, ~1 out of 15 000 soil bacterial cells had the potential to receive and, at least transiently, host plasmid pJK5, consistent with earlier permissiveness measurements of soil bacterial communities for IncP-1 α , IncP-1 ϵ and IncPromA broad-host range plasmids (Musovic *et al.*, 2014; Klümper *et al.*, 2015). Despite stress being commonly associated with increasing bacterial evolvability and promoting gene transfer (Gillings and Stokes, 2012), we here show that community-level plasmid acquisition decreased consistently and significantly (27–100%, $P < 0.05$) with metal-stress independent of the metal, and even exceeded the degree of growth inhibition. This reduction in permissiveness might be the consequence of a decrease in metabolic status, the result of a specific response of plasmid transfer, plasmid replication or marker gene expression to metal exposure or a combination of these. Higher stressor doses were furthermore associated with a further decrease in transfer frequency. Plasmid transfer frequency reduction, previously documented for Zn stress (De Rore *et al.*, 1994), might be common across soil bacterial communities for a multitude of different metals.

In spite of the observed absolute decrease in community permissiveness, several OTUs had a markedly increased permissiveness under stress conditions. For example, several OTUs belonging to the Bacteroidetes phylum had a more than 10-fold increased proportion of recipient cells carrying plasmid pKJK5 after stress treatment. Thus, gene transfer from the enterobacterial host across phyla to the soil indigenous community can increase due to metal stress. This short-term increase might play a crucial role for the dissemination of plasmid-encoded antibiotic resistance genes from manure as their original fecal hosts are quickly lost due to competitive exclusion (Estrada *et al.*, 2004). In this way, the soil bacterial community can serve as a long-term reservoir for plasmids as elevated retention of plasmids can be observed under long-term metal stress, even for plasmids not coding for metal resistance (Smets *et al.*, 2003).

Although studied for decades, our current knowledge of metal-bacteria interactions is insufficient to pinpoint the mechanisms that link metal exposure to the observed modulation in permissiveness. Several generic and specific mechanisms of metal stress have been described. The cationic metals used in this study (Cd, Cu, Ni, Zn), as well as the anion arsenite share the toxic mechanism of disrupting iron–sulfur clusters of metallo-enzymes (Hughes, 2002; Macomber and Imlay, 2009; Macomber and Hausinger, 2011; Xu and Imlay, 2012). Metal cations are excreted from bacterial cells as a stress response using efflux systems of similar types (P-type, RND, CDF) (Nies, 1999), while arsenite has its own type of efflux system (A-type) (Nies, 1999). All bacteria have a certain tolerance level to metal stress. Thus, for a given exposure, a gradient of stress levels ranging from sub-toxic to toxic or even lethal conditions can exist within a community (Rensing *et al.*, 2002). Due to the multitude of mechanisms of metal toxicity and resistance present in different bacteria, it is safe to assume that although introduced at the same community growth inhibition doses, each element caused diverse specific short-term stress-responses in individual community members.

Taking advantage of the high-resolution identification of transconjugal OTUs, we demonstrated that stress-induced modifications of permissiveness are indeed unique for each OTU. However, the modified permissiveness of an OTU was generally consistent across most metal stresses. In contrast, the bacterial response to the metalloid arsenic was not as strongly correlated with the four metals. One potential explanation could stem from a connection between the regulation of efflux pumps, which are similar across metal ions but different for arsenite (Nies, 1999), and that of permissiveness.

Furthermore, phylogenetically similar OTUs tended to act similarly in their short-term response to different stresses. The stress-induced regulation of permissiveness thus seems to be taxon-dependent,

potentially due to evolutionary conservation, and might well be connected to the regulation of the defense mechanisms against foreign DNA, such as CRISPR-cas or RM systems in the specific phylogenetic groups. While the different regulatory control systems for CRISPR-cas systems are not well-understood (Mojica and Díez-Villaseñor, 2010), specific stress responses can lead to induced expression of CRISPR-cas genes, decreasing the plasmid receipt potential (Perez-Rodriguez *et al.*, 2011).

The expression of RM genes is not constant, but depends on environmental conditions (Bayliss *et al.*, 2006), and diverse stressors can play a role in their regulation (Schafer *et al.*, 1994). The observed phylogenetic conservation holds not only true across the highly correlated heavy metal stresses, but also for arsenic stress, thus further indicating that permissiveness toward plasmids might be directly coupled to different stress responses.

In pure cultures, the ability to uptake foreign DNA changes as a response to stress for specific stressors (Arango Pinedo and Smets, 2005; Pérez-Mendoza and de la Cruz, 2009). Due to the phylogenetic conservation of the stress response, results from single strain experiments might have predictive value based on their phylogeny and for diverse stresses of the same type. However, based on the highly diverse responses across different community members neither single strain nor community averaged analyses suffice to predict the effect of stress on plasmid receipt across complex communities. Our results show that understanding the effect of stress on the ecology of plasmid transfer can only be achieved by examination at both community and individual strain level.

We demonstrated here that the stress-response mechanisms affecting the permissiveness among phylogenetically related bacterial groups are similar and consistent for a variety of different heavy metal stresses. Extrapolation of our results to phylogenetically related groups of bacteria and other stressors might therefore be valid. To further understand the stress-induced modulation of permissiveness, the exact cellular responses remain to be elucidated.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank J Magid for access to the CRUCIAL field plot, LK Jensen for technical assistance in the laboratory and SM Milani for assistance in FACS sorting. This work was funded by the Villum Kann Rasmussen Foundation Center of Excellence CREAM (Center for Environmental and Agricultural Microbiology). UK is currently supported through an MRC/BBSRC grant (MR/N007174/1).

References

- Agersø Y, Wulff G, Vaclavik E, Halling-Sørensen B, Jensen LB. (2006). Effect of tetracycline residues in pig manure slurry on tetracycline-resistant bacteria and resistance gene *tet(M)* in soil microcosms. *Environ Int* **32**: 876–882.
- Arango Pinedo C, Smets BF. (2005). Conjugal TOL transfer from *Pseudomonas putida* to *Pseudomonas aeruginosa*: Effects of restriction proficiency, toxicant exposure, cell density ratios, and conjugation detection method on observed transfer efficiencies. *Appl Environ Microbiol* **71**: 51–57.
- Bahl MI, Hansen LH, Goesmann A, Sørensen SJ. (2007). The multiple antibiotic resistance IncP-1 plasmid pKJK5 isolated from a soil environment is phylogenetically divergent from members of the previously established α , β and δ sub-groups. *Plasmid* **58**: 31–43.
- Bahl MI, Oregaard G, Sørensen SJ, Hansen LH. (2009). Construction and use of flow cytometry optimized plasmid-sensor strains. *Methods Mol Biol* **532**: 257–268.
- Bayliss CD, Callaghan MJ, Moxon ER. (2006). High allelic diversity in the methyltransferase gene of a phase variable type III restriction-modification system has implications for the fitness of haemophilus influenzae. *Nucleic Acids Res* **34**: 4046–4059.
- Berg J, Brandt KK, Al-Soud WA, Holm PE, Hansen LH, Sørensen SJ et al. (2012). Selection for Cu-tolerant bacterial communities with altered composition, but unaltered richness, via long-term Cu exposure. *Appl Environ Microbiol* **78**: 7438–7446.
- Brandt KK, Frandsen RJN, Holm PE, Nybrø O. (2010). Development of pollution-induced community tolerance is linked to structural and functional resilience of a soil bacterial community following a five-year field exposure to copper. *Soil Biol Biochem* **42**: 748–757.
- Campbell JIA, Jacobsen CS, Sørensen J. (1995). Species variation and plasmid incidence among fluorescent *Pseudomonas* strains isolated from agricultural and industrial soils. *FEMS Microbiol Ecol* **18**: 51–62.
- Cases I, de Lorenzo V. (2005). Promoters in the environment: transcriptional regulation in its natural context. *Nat Rev Microbiol* **3**: 105–118.
- Corvaglia AR, François P, Hernandez D, Perron K, Linder P, Schrenzel J. (2010). A type III-like restriction endonuclease functions as a major barrier to horizontal gene transfer in clinical *Staphylococcus aureus* strains. *Proc Natl Acad Sci USA* **107**: 11954–11958.
- Dechesne A, Pallud C, Bertolla F, Grundmann GL. (2005). Impact of the microscale distribution of a *Pseudomonas* strain introduced into soil on potential contacts with indigenous bacteria. *Appl Environ Microbiol* **71**: 8123–8131.
- De Gelder L, Vandecasteele FPJ, Brown CJ, Forney LJ, Top EM. (2005). Plasmid donor affects host range of promiscuous IncP-1 β plasmid pB10 in an activated-sludge microbial community. *Appl Environ Microbiol* **71**: 5309–5317.
- De Rore H, Top E, Houwen F, Verstraete W. (1994). Evolution of heavy metal resistant transconjugants in a soil environment with a concomitant selective pressure. *FEMS Microbiol Ecol* **14**: 263–274.
- Estrada IB, Aller A, Aller F, Gómez X, Morán A. (2004). The survival of *Escherichia coli*, faecal coliforms and enterobacteriaceae in general in soil treated with sludge from wastewater treatment plants. *Bioresour Technol* **93**: 191–198.
- Excoffier L, Smouse PE, Quattro JM. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics* **131**: 479–491.
- Franke AE, Clewell DB. (1981). Evidence for a chromosome-borne resistance transposon (Tn916) in *Streptococcus faecalis* that is capable of ‘conjugal’ transfer in the absence of a conjugative plasmid. *J Bacteriol* **145**: 494–502.
- Gans J, Wolinsky M, Dunbar J. (2005). Computational improvements reveal great bacterial diversity and high metal toxicity in soil. *Science* **309**: 1387–1390.
- Giller KE, Witter E, Mcgrath SP. (1998). Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: a review. *Soil Biol Biochem* **30**: 1389–1414.
- Gillings MR, Stokes HW. (2012). Are humans increasing bacterial evolvability? *Trends Ecol Evol* **27**: 346–352.
- Halary S, Leigh JW, Cheaib B, Lopez P, Baptiste E. (2009). Network analyses structure genetic diversity in independent genetic worlds. *Proc Natl Acad Sci USA* **107**: 127–132.
- Heuer H, Schmitt H, Smalla K. (2011). Antibiotic resistance gene spread due to manure application on agricultural fields. *Curr Opin Microbiol* **14**: 236–243.
- Heuer H, Smalla K. (2012). Plasmids foster diversification and adaptation of bacterial populations in soil. *FEMS Microbiol Rev* **36**: 1083–1104.
- Hommel G. (1988). A stagewise rejective multiple procedure based on a modified Bonferroni. *Biometrika* **75**: 383–386.
- Hughes MF. (2002). Arsenic toxicity and potential mechanisms of action. *Toxicol Lett* **133**: 1–16.
- Jondreville C, Revy P, Dourmad J. (2003). Dietary means to better control the environmental impact of copper and zinc by pigs from weaning to slaughter. *Livest Prod Sci* **84**: 147–156.
- Klümper U, Dechesne A, Smets BF. (2014a). Protocol for evaluating the permissiveness of bacterial communities toward conjugal plasmids by quantification and isolation of transconjugants. In: McGenity TJ, Timmis KN, Nogales B (eds), *Hydrocarbon and Lipid Microbiology Protocols, Springer Protocols Handbook*. Springer-Verlag: Berlin, Heidelberg, pp 1–14.
- Klümper U, Droumpali A, Dechesne A, Smets BF. (2014b). Novel assay to measure the plasmid mobilizing potential of mixed microbial communities. *Front Microbiol* **5**: 730.
- Klümper U, Riber L, Dechesne A, Sannazzarro A, Hansen LH, Sørensen SJ et al. (2015). Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community. *ISME J* **9**: 934–945.
- Knezevic SZ, Streibig JC, Ritz C. (2007). Utilizing R Software package for dose-response studies: The concept and data analysis. *Weed Technol* **21**: 840–848.
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Appl Environ Microbiol* **79**: 5112–5120.
- Lekfeldt JDS, Magid J, Holm PE, Nybrø O, Brandt KK. (2014). Evaluation of the leucine incorporation technique for detection of pollution-induced community tolerance to copper in a long-term agricultural field

- trial with urban waste fertilizers. *Environ Pollut* **194**: 78–85.
- Letunic I, Bork P. (2007). Interactive Tree Of Life (iTOL): An online tool for phylogenetic tree display and annotation. *Bioinformatics* **23**: 127–128.
- Lindahl V, Bakken LR. (1995). Evaluation of methods for extraction of bacteria from soil. *FEMS Microbiol Ecol* **16**: 135–142.
- Macomber L, Hausinger RP. (2011). Mechanisms of nickel toxicity in microorganisms. *Metallomics* **3**: 1153.
- Macomber L, Imlay JA. (2009). The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci USA* **106**: 8344–8349.
- Mah TFC, O'Toole GA. (2001). Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* **9**: 34–39.
- Mojica FJM, Díez-Villaseñor C. (2010). The on-off switch of CRISPR immunity against phages in *Escherichia coli*: MicroCommentary. *Mol Microbiol* **77**: 1341–1345.
- Musovic S, Dechesne A, Sørensen J, Smets BF. (2010). Novel assay to assess permissiveness of a soil microbial community toward receipt of mobile genetic elements. *Appl Environ Microbiol* **76**: 4813–4818.
- Musovic S, Klümper U, Dechesne A, Magid J, Smets BF. (2014). Long-term manure exposure increases soil bacterial community potential for plasmid uptake. *Environ Microbiol Rep* **6**: 125–130.
- Nicholson FA, Smith SR, Alloway BJ, Carlton-Smith C, Chambers BJ. (2003). An inventory of heavy metals inputs to agricultural soils in England and Wales. *Sci Total Environ* **311**: 205–219.
- Nies DH. (1999). Microbial heavy-metal resistance. *Appl Microbiol Biotechnol* **51**: 730–750.
- Norman A, Hansen LH, Sørensen SJ. (2009). Conjugal plasmids: vessels of the communal gene pool. *Philos Trans R Soc Lond B Biol Sci* **364**: 2275–2289.
- Pérez-Martín J, de Lorenzo V. (1996). In vitro activities of an N-terminal truncated form of *XylR*, a sigma 54-dependent transcriptional activator of *Pseudomonas putida*. *J Mol Biol* **258**: 575–587.
- Pérez-Mendoza D, de la Cruz F. (2009). *Escherichia coli* genes affecting recipient ability in plasmid conjugation: are there any? *BMC Genomics* **10**: 71.
- Perez-Rodriguez R, Haitjema C, Huang Q, Nam KH, Bernardis S, Ke A et al. (2011). Envelope stress is a trigger of CRISPR RNA-mediated DNA silencing in *Escherichia coli*. *Mol Microbiol* **79**: 584–599.
- Poulsen PHB, Al-Soud WA, Bergmark L, Magid J, Hansen LH, Sørensen SJ. (2013). Effects of fertilization with urban and agricultural organic wastes in a field trial - Prokaryotic diversity investigated by pyrosequencing. *Soil Biol Biochem* **57**: 784–793.
- Rensing C, Newby DT, Pepper IL. (2002). The role of selective pressure and selfish DNA in horizontal gene transfer and soil microbial community adaptation. *Soil Biol Biochem* **34**: 285–296.
- Revelle W. (2014). *Psych: Procedures for Personality and Psychological Research*, Northwestern University, Evanston, IL, USA, <http://CRAN.R-project.org/package=psych> Version = 1.6.6.
- Richardson M. (1993). *Epitoxicology Monitoring*. Richardson ML (ed). VCH Verlagsgesellschaft: New York, NY, USA, pp 335–343.
- Roer L, Aarestrup FM, Hasman H. (2015). The EcoKI type I restriction-modification system in *Escherichia coli* affects but is not an absolute barrier for conjugation. *J Bacteriol* **197**: 337–342.
- Schafer A, Kalinowski J, Simon R, Seep-Feldhaus AH, Puhler A. (1990). High-frequency conjugal plasmid transfer from gram-negative *Escherichia coli* to various gram-positive coryneform bacteria. *J Bacteriol* **172**: 1663–1666.
- Schafer A, Kalinowski J, Puhler A. (1994). Increased fertility of *Corynebacterium glutamicum* recipients in intergeneric matings with *Escherichia coli* after stress exposure. *Appl Environ Microbiol* **60**: 756–759.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB et al. (2009). Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**: 7537–7541.
- Seier-Petersen MA, Jasni A, Aarestrup FM, Vigre H, Mullany P, Roberts AP et al. (2013). Effect of subinhibitory concentrations of four commonly used biocides on the conjugative transfer of Tn916 in *Bacillus subtilis*. *J Antimicrob Chemother* **69**: 343–348.
- Shintani M, Matsui K, Inoue J, Hosoyama A, Ohji S, Yamazoe A et al. (2014). Single-cell analyses revealed transfer ranges of IncP-1, IncP-7, and IncP-9 plasmids in a soil bacterial community. *Appl Environ Microbiol* **80**: 138–145.
- Slager J, Kjos M, Attaiech L, Veening JW. (2014). Antibiotic-induced replication stress triggers bacterial competence by increasing gene dosage near the origin. *Cell* **157**: 395–406.
- Smalla K, Heuer H, Gotz A, Niemeyer D, Krogerrecklenfort E, Tietze E. (2000). Exogenous isolation of antibiotic resistance plasmids from piggy manure slurries reveals a high prevalence and diversity of IncQ-like plasmids. *Appl Environ Microbiol* **66**: 4854–4862.
- Smets BF, Morrow JB, Arango Pinedo C. (2003). Plasmid introduction in metal-stressed, subsurface-derived microcosms: plasmid fate and community response. *Appl Environ Microbiol* **69**: 4087–4097.
- Sørensen SJ, Bailey M, Hansen LH, Kroer N, Wuertz S. (2005). Studying plasmid horizontal transfer in situ: a critical review. *Nat Rev Microbiol* **3**: 700–710.
- Top EM, Rore H, Collard J-M, Gellens V, Slobodkina G, Verstraete W et al. (1995). Retromobilization of heavy metal resistance genes in unpolluted and heavy metal polluted soil. *FEMS Microbiol Ecol* **18**: 191–203.
- Wang Q, Garrity GM, Tiedje JM, Cole JR. (2007). Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73**: 5261–5267.
- Webb CO, Ackerly DD, Kembel SW. (2008). Phylocom: software for the analysis of phylogenetic community structure and trait evolution. *Bioinformatics* **24**: 2098–2100.
- World Health Organization (WHO). (2014). *Antimicrobial Resistance Global Report on Surveillance*. Summary: Geneva, Switzerland.
- Xu FF, Imlay JA. (2012). Silver(I), mercury(II), cadmium (II), and zinc(II) target exposed enzymic iron-sulfur clusters when they toxify *Escherichia coli*. *Appl Environ Microbiol* **78**: 3614–3621.
- You Y, Hilpert M, Ward MJ. (2012). Detection of a common and persistent *tet(L)*-carrying plasmid in chicken-waste-impacted farm soil. *Appl Environ Microbiol* **78**: 3203–3213.

- Yue JC, Clayton MK. (2005). A similarity measure based on species proportions. *Commun StatTheory Methods* **34**: 2123–2131.
- Zhao FJ, Ma Y, Zhu YG, Tang Z, McGrath SP. (2015). Soil contamination in China: Current status and mitigation strategies. *Environ Sci Technol* **49**: 750–759.

- Zhaxybayeva O, Doolittle WF. (2011). Lateral gene transfer. *Curr Biol* **21**: R242–R246.
- Zhu Y-G, Johnson TA, Su J-Q, Qiao M, Guo G-X, Stedtfeld RD *et al.* (2013). Diverse and abundant antibiotic resistance genes in Chinese swine farms. *Proc Natl Acad Sci USA* **110**: 3435–3440.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)