

Inhibitory phosphorylation of Cdk1 mediates prolonged prophase I arrest in female germ cells and is essential for female reproductive lifespan

Deepak Adhikari^{1,4,*}, Kiran Busayavalasa^{1,*}, Jingjing Zhang¹, Mengwen Hu¹, Sanjiv Risal¹, Mustafa Bilal Bayazit¹, Meenakshi Singh¹, M Kasim Diril^{2,5}, Philipp Kaldis^{2,3}, Kui Liu¹

¹Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg SE-405 30, Sweden; ²Institute of Molecular and Cell Biology (IMCB), A*STAR (Agency for Science, Technology and Research), 61 Biopolis Drive, Proteos #3-09, Singapore 138673, Republic of Singapore; ³Department of Biochemistry, National University of Singapore (NUS), Singapore 117597, Republic of Singapore

A unique feature of female germ cell development in mammals is their remarkably long arrest at the prophase of meiosis I, which lasts up to 50 years in humans. Both dormant and growing oocytes are arrested at prophase I and completely lack the ability to resume meiosis. Here, we show that the prolonged meiotic arrest of female germ cells is largely achieved via the inhibitory phosphorylation of Cdk1 (cyclin-dependent kinase 1). In two mouse models where we have introduced mutant Cdk1^{T14AY15F} which cannot be inhibited by phosphorylation (Cdk1AF) in small meiotically incompetent oocytes, the prophase I arrest is interrupted, leading to a premature loss of female germ cells. We show that in growing oocytes, Cdk1AF leads to premature resumption of meiosis with condensed chromosomes and germinal vesicle breakdown followed by oocyte death, whereas in dormant oocytes, Cdk1AF leads to oocyte death directly, and both situations damage the ovarian reserve that maintains the female reproductive lifespan, which should be around 1 year in mice. Furthermore, interruption of the inhibitory phosphorylation of Cdk1 results in DNA damage, which is accompanied by induction of the Chk2 (checkpoint kinase 2)-p53/p63-dependent cell death pathway, which eventually causes global oocyte death. Together, our data demonstrate that the phosphorylation-mediated suppression of Cdk1 activity is one of the crucial factors that maintain the lengthy prophase arrest in mammalian female germ cells, which is essential for preserving the germ cell pool and reproductive lifespan in female mammals.

Keywords: meiotic prophase I; Cdk1 inhibitory phosphorylation; immature oocyte; female fertility lifespan

Cell Research (2016) 26:1212-1225. doi:10.1038/cr.2016.119; published online 21 October 2016

Introduction

In mammals, female germ cells enter meiosis in the embryonic gonad, but they are subsequently arrested at

the dictyate stage of prophase I in the form of dormant oocytes that are enclosed within primordial follicles [1]. Such meiosis arrest can last for a very long time, which can be, for example, up to 50 years in humans and more than 1 year in mice. In addition, before the oocytes can resume meiotic maturation, they must undergo a prolonged growth period from small primary oocytes to fully grown oocytes. During such growth, which usually takes weeks to months or even longer, the oocytes remain arrested at prophase I [2-4]. In other words, throughout the lengthy dormancy and growth of oocytes, the oocytes are incompetent for meiosis resumption [5].

The pool of primordial follicles with dormant oocytes provides the ovarian reserve that maintains the female reproductive lifespan [4]. Only a limited number of dor-

*These two authors contributed equally to this work.

Correspondence: Philipp Kaldis^a, Kui Liu^b

^aE-mail: kaldis@imcb.a-star.edu.sg

^bE-mail: kui.liu@gu.se

⁴Current address: Department of Anatomy and Developmental Biology, Monash Biomedicine Discovery Institute, Monash University, Melbourne, Victoria 3800, Australia

⁵Current address: Izmir Biomedicine and Genome Institute, Dokuz Eylul University, Izmir 35340, Turkey

Received 7 May 2016; revised 8 August 2016; accepted 18 August 2016; published online 21 October 2016

mant oocytes are recruited into the growth phase, and of these only some will eventually reach the fully grown stage and gain the ability to resume meiosis (illustrated in Figure 1A). Importantly, the maintenance of genomic integrity in the meiotically incompetent oocytes is crucial for generating future healthy progeny. The prophase I arrest of the small oocytes could thus be an essential mechanism for maintaining the genomic integrity of female germ cells.

Although the prophase I arrest in dormant and growing oocytes appears to be similar to that of fully grown germinal vesicle (GV)-stage oocytes, their cell cycle behaviors are fundamentally different. For example, in mice, meiotic resumption can occur in fully grown oocytes either upon stimulation by luteinizing hormone or upon release from the follicular environment *in vitro*. However, in growing oocytes, such as oocytes smaller than 60 μm or oocytes from the ovaries of postnatal day (PD) 15 or younger mice, release from the follicular environment *in vitro* does not lead to the resumption of meiosis [6, 7].

Based on published data, we hypothesize that the molecular regulation of meiotic arrest in dormant and growing oocytes is distinct from mechanisms maintaining the GV-stage of fully grown oocytes. For example, high cyclic AMP (cAMP) levels maintain the meiotic arrest in fully grown oocytes, but they do not maintain the meiotic arrest of smaller oocytes because even if the cAMP levels are decreased in growing oocytes, these oocytes do not resume meiosis [8, 9]. In addition, prophase I arrest in fully grown mouse oocytes is maintained by the constant degradation of cyclin B1 by the anaphase-promoting complex/cyclosome (APC/C)-Cdh1, which keeps the activity of cyclin-dependent kinase 1 (Cdk1) low [10]. However, the prevention of cyclin B1 degradation in growing *Cdh1*-null oocytes, where APC/C activity is suppressed, is not sufficient to release the oocytes from prophase I arrest, and these growing *Cdh1*-null oocytes still do not resume meiosis *in vivo* (i.e., inside the follicles) [11]. Moreover, increasing the amount of Cdk1 protein by microinjection of *Cdk1* mRNA into growing mouse oocytes does not lead to the resumption of meiosis [12]. Similarly, more rapid GV breakdown (GVBD) occurs in fully grown mouse oocytes only when the activity of Cdk1 is increased by knocking out the lysine-specific demethylase 1 [13] or by downregulating Wee1B that can phosphorylate and inhibit Cdk1 [14], but GVBD is not seen in growing oocytes in these mutant models. These results raise the important question of whether Cdk1 activity is regulated in dormant and growing oocytes, and if so, what the physiological significance of Cdk1 regulation is in these immature oocytes.

By generating two knock-in mouse models to express a Cdk1 variant that is non-inhibitable by phosphorylation (Cdk1AF) in dormant and growing oocytes, we were able to demonstrate that the simple phosphorylation-mediated suppression of Cdk1 activity plays an important role in maintaining the seemingly complicated lengthy dictyate arrest of mammalian female germ cells. This mechanism also prevents the accumulation of DNA damage and subsequent death of female germ cells, which is essential for maintaining the normal female reproductive lifespan.

Results

Expression of Cdk1AF in dormant mouse oocytes causes DNA damage and rapid oocyte death

To first determine the effects of premature Cdk1 activation on dormant oocytes, we generated a mouse model in which two point mutations lead to substitutions of T14 and Y15 in Cdk1 with alanine 14 and phenylalanine 15 (Figure 1B and Supplementary information, Figure S1). Because these two amino acids are non-phosphorylatable, Cdk1 with these two mutations becomes non-inhibitable by phosphorylation (Cdk1AF) [15, 16] (Figure 1B). We introduced the T14A and Y15F mutations into one *Cdk1* allele to generate the *Cdk1*^{T14AY15F} (in short, *Cdk1AF*) allele, but with a *loxP*-flanked *STOP* (S) sequence in front of the *Cdk1AF* allele to prevent its expression in the absence of Cre recombinase. Wild-type Cdk1 is expressed by the other allele (Supplementary information, Figure S1). The mice carrying one floxed *STOP-Cdk1AF* (*SAF*) allele and one wild-type *Cdk1* allele are referred to as *Cdk1*^{+/SAF} mice. The *Cdk1*^{+/SAF} mice displayed normal ovarian development and fertility (see below) and were used as controls.

By crossing *Cdk1*^{+/SAF} mice with *growth differentiation factor 9 (Gdf9)-Cre* mice in which the Cre recombinase is efficiently expressed in dormant oocytes of primordial follicles [17, 18] (Figure 1C), we specifically deleted the *STOP* sequence and induced the expression of the *Cdk1AF* in one allele of the *Cdk1* gene (referred to as *OoCdk1*^{+/AF}; *Gdf9-Cre*) in dormant and growing oocytes (illustrated in Figure 1C), while the other allele carried the wild-type *Cdk1* gene (*OoCdk1*⁺).

To maintain the normal length of female reproductive life, the majority of primordial follicles must be maintained in a dormant state throughout the reproductive lifespan. In comparison with the healthy ovaries from control *Cdk1*^{+/SAF} mice at PD8 (Figure 2A and 2C) and PD16 (Figure 2E), the mutant *OoCdk1*^{+/AF}; *Gdf9-Cre* ovaries contained only a few follicles at PD8 (Figure 2B and 2D, arrows, and Supplementary information, Figure S2G), and they were depleted of almost all oocytes by




Figure 2 Expression of *Cdk1AF* in dormant oocytes causes DNA damage and oocyte depletion in *OoCdk1*^{+/AF}; *Gdf9-Cre* mice. (A–F) Hematoxylin-stained sections of mouse ovary at PD8 and PD16. By PD8, only a few follicles were left in *OoCdk1*^{+/AF}; *Gdf9-Cre* ovaries (B arrows), and the oocytes of surviving primordial follicles remained arrested at GV stage (D, arrows). The follicular structures had mostly disappeared in the mutant ovaries by PD16 (F). As controls, normal ovaries of PD8 (A and C) and PD16 (E) *Cdk1*^{+/SAF} mice containing healthy follicles are shown. The experiments were repeated more than three times each, and for each time and each age ovaries from one mouse of each genotype were used. (G) Comparison of the average cumulative number of pups per *OoCdk1*^{+/AF}; *Gdf9-Cre* female (dotted blue line) and per *Cdk1*^{+/SAF} female (solid black line), indicating that the *OoCdk1*^{+/AF}; *Gdf9-Cre* females were infertile. The numbers of females used are shown as *n*. (H and I) Prominent γ -H2AX staining was observed in dormant oocytes enclosed in primordial follicles of PD6 *OoCdk1*^{+/AF}; *Gdf9-Cre* ovaries (I, arrows) but not in the control oocytes (H, arrowheads). DDX4 was used to label the oocytes and the same sections were counterstained with hematoxylin to visualize ovarian histology. The experiments were repeated more than three times each.


Cdk1 kinase activity can cause DNA damage [19]. One of the first cellular responses to DNA double-strand breaks is the phosphorylation of histone H2AX [20]. Immunostaining for phosphorylated histone H2AX at Ser139 (known as γ -H2AX) in PD6 *OoCdk1*^{+/AF}; *Gdf9-Cre* ovarian sections revealed substantially increased DNA damage in the dormant oocytes (Figure 2I, arrows). In contrast, almost no γ -H2AX foci were seen in control *Cdk1*^{+/SAF} oocytes (Figure 2H, arrowheads). These results demonstrate that the direct suppression of *Cdk1* kinase activity by its phosphorylation is essential for protecting genomic integrity of dormant oocytes arrested at prophase I, thereby maintaining the female reproductive lifespan.

Premature meiotic resumption in growing oocytes expressing *Cdk1AF*

The rapid depletion of all of the oocytes in *OoCdk1*^{+/AF}; *Gdf9-Cre* ovaries makes it impossible to obtain mutant oocytes to study the underlying mechanisms of oocyte depletion caused by *Cdk1AF*. Therefore, we crossed the

Cdk1^{+/SAF} mice with *Zona pellucida 3 (Zp3)-Cre* mice [21] to introduce the expression of *Cdk1AF* starting in the growing oocytes but not in the dormant oocytes (Figure 1A and 1C). The resulting mice are referred to as *OoCdk1*^{+/AF}; *Zp3-Cre* mice. We have previously validated that the *Zp3* promoter-mediated Cre recombinase expression is oocyte-specific and is expressed only in growing oocytes but not in dormant oocytes [22]. We used small growing oocytes from PD14 ovaries from *OoCdk1*^{+/AF}; *Zp3-Cre* mice in the subsequent experiments.

We first isolated growing oocytes from the ovaries of PD14 *OoCdk1*^{+/AF}; *Zp3-Cre* and control *Cdk1*^{+/SAF} mice and measured their *Cdk1* kinase activity. As shown in Figure 3A, *Cdk1* kinase activity was significantly elevated in the PD14 *OoCdk1*^{+/AF}; *Zp3-Cre* oocytes, and as expected it was accompanied by a decrease in phosphorylation at Y15 of *Cdk1* due to the expression of a *Cdk1AF* allele in *OoCdk1*^{+/AF}; *Zp3-Cre* oocytes (Figure 3A), and the stability of *Cdk1* protein in growing oocytes seems to be positively affected by its dephosphorylation (Figure 3A). We found that the *OoCdk1*^{+/AF}; *Zp3-Cre* mice



were completely infertile (Figure 3B). To study how the expression of *Cdk1AF* in oocytes leads to infertility, we compared the follicular development in *OoCdk1^{+/-AF}; Zp3-Cre* mice with that in *Cdk1^{+/-SAF}* mice. No apparent morphological difference was found in PD8 ovaries of *Cdk1^{+/-SAF}* and *OoCdk1^{+/-AF}; Zp3-Cre* mice, where ovaries of both genotypes had mostly primordial and some growing follicles (Supplementary information, Figure S2A and S2B) and contained comparable numbers of total follicles (Supplementary information, Figure S2H). By PD14, however, the *OoCdk1^{+/-AF}; Zp3-Cre* ovaries (Supplementary information, Figure S2D) appeared smaller than the *Cdk1^{+/-SAF}* ovaries (Supplementary information, Figure S2C) and contained less number of follicles (Supplementary information, Figure S2H). By PD24, most of the growing follicles in *OoCdk1^{+/-AF}; Zp3-Cre* ovaries were undergoing atresia and contained degenerating oocytes (Supplementary information, Figure S2F, arrows) as compared with the healthy follicles in the *Cdk1^{+/-SAF}* ovaries (Supplementary information, Figure S2E, arrows). Quantification results showed that PD24 *OoCdk1^{+/-AF}; Zp3-Cre* ovaries contained significantly less number of follicles than in the *Cdk1^{+/-SAF}* ovaries (Supplementary information, Figure S2H). Compared with the healthy control *Cdk1^{+/-SAF}* ovaries at 6 weeks (Figure 3C) and 3

months (Figure 3E) of age, the mutant *OoCdk1^{+/-AF}; Zp3-Cre* ovaries contained only a few follicles at 6 weeks of age (Figure 3D, arrows and Supplementary information, Figure S2H), and they were depleted of almost all oocytes by 3 months of age (Figure 3F and Supplementary information, Figure S2H).

In the control mouse ovary, all analyzed oocytes within small or large follicles remained arrested at the GV stage at any given age, as seen in the ovarian section of PD14 control *Cdk1^{+/-SAF}* mice (Figure 3G, arrowhead). However, we found that the young PD14 *OoCdk1^{+/-AF}; Zp3-Cre* oocytes, where *Cdk1AF* was expressed, had already undergone GVBD within the follicles *in vivo* and their chromosomes were condensed (Figure 3H, arrow).

Quantification results showed that within the primary, secondary, and preantral follicles in PD14 *OoCdk1^{+/-AF}; Zp3-Cre* ovaries, 1.5% ($n = 220$), 46.5% ($n = 97$), and 64.4% ($n = 78$) of the oocytes had already resumed meiosis, respectively (Figure 3I). This was in sharp contrast to the control PD14 *Cdk1^{+/-SAF}* ovaries in which all oocytes were at the meiotically arrested GV stage (Figure 3I).

Using chromosome spreads, we observed condensed chromosomes in PD14 *OoCdk1^{+/-AF}; Zp3-Cre* oocytes that had already undergone GVBD (Figure 3K and 3M, arrows). In sharp contrast, the chromosomes remained

Figure 3 Premature *in vivo* resumption of meiosis in small growing oocytes in *OoCdk1^{+/-AF}; Zp3-Cre* mice. **(A)** Elevated *Cdk1* kinase activity and total *Cdk1* level but decreased inhibitory phosphorylation of *Cdk1* (p-*Cdk1*, Y15) were observed in *OoCdk1^{+/-AF}; Zp3-Cre* oocytes, suggesting that *Cdk1AF* was successfully introduced into the growing oocytes in the mutant mice. β -Actin was used as the loading control. For the *Cdk1* kinase assay, 10 oocytes per reaction were used. For Western blots (WB), lysate from 200 oocytes was loaded in each lane. The experiments were repeated more than three times each. **(B)** Comparison of the average cumulative number of pups per *OoCdk1^{+/-AF}; Zp3-Cre* female (red dotted line) and per *Cdk1^{+/-SAF}* female (solid blue line). All *OoCdk1^{+/-AF}; Zp3-Cre* females were infertile. The numbers of females are shown as n . **(C-F)** Hematoxylin-stained sections from mouse ovary at 6 weeks and 3 months. By 6 weeks of age, only a few follicles were observed in *OoCdk1^{+/-AF}; Zp3-Cre* ovaries **(D, arrows)**, and healthy follicular structures had mostly disappeared in 3-month-old *OoCdk1^{+/-AF}; Zp3-Cre* ovaries **(F)**. Control *Cdk1^{+/-SAF}* ovaries **(C and E)** contained healthy oocytes and follicles. CL, corpus luteum. The experiments were repeated more than three times each, and for each time and each age; ovaries from one mouse of each genotype were used. **(G, H)** Premature *in vivo* GVBD and chromosome condensation in small primary oocytes of *OoCdk1^{+/-AF}; Zp3-Cre* mice. Ovaries from PD14 *OoCdk1^{+/-AF}; Zp3-Cre* and *Cdk1^{+/-SAF}* females were embedded in paraffin, and serial sections of 8- μ m thickness were prepared and stained with hematoxylin. **(G)** A wild-type primary oocyte in a secondary follicle showing the GV **(G, arrowhead)** from a PD14 *Cdk1^{+/-SAF}* ovary. **(H)** A representative growing primary oocyte from a PD14 *OoCdk1^{+/-AF}; Zp3-Cre* ovary where GVBD and chromosome condensation had already occurred **(H, arrow)**. The experiments were repeated more than three times each, and for each time and each age ovaries from one mouse of each genotype were used. **(I)** Quantification of the percentages of oocytes that resumed meiosis *in vivo* in PD14 ovaries within primary, secondary, and preantral follicles. Numbers of each type of follicles included (n) are shown. **(J-Q)** Condensed chromosomes with cohesin localization at the chromosome axis in PD14 *OoCdk1^{+/-AF}; Zp3-Cre* oocytes. Staining with DAPI for DNA **(L, M, blue)**, SMC3 for cohesin **(N, O, green)**, CREST for kinetochores **(P, Q, red)**, and merge **(J, K)** in *OoCdk1^{+/-AF}; Zp3-Cre* and *Cdk1^{+/-SAF}* oocytes. The experiments were repeated more than three times each, and the representative images from one experiment are shown. **(R, S)** Increased GVBD in mutant *OoCdk1^{+/-AF}; Zp3-Cre* oocytes upon release from the follicular environment. In wild-type PD14 *Cdk1^{+/-SAF}* oocytes, GV was maintained even if the oocytes were released from the follicles by enzymatic digestion **(R, arrowheads)**. However, an increased rate of GVBD was seen in PD14 *OoCdk1^{+/-AF}; Zp3-Cre* oocytes upon release from the follicles **(S, arrows)** by enzymatic digestion of ovaries from PD14 mice. Representative Hoechst-stained oocyte nuclei are shown to the right. Quantification of GV and GVBD rates of oocytes immediately after their isolation from follicles in PD14 *Cdk1^{+/-SAF}* and *OoCdk1^{+/-AF}; Zp3-Cre* mice **(T)** and after further *in vitro* culture **(U)**. The numbers of oocytes analyzed (n) are shown.

decondensed in the control oocytes (Figure 3J and 3L, arrowheads). This indicates that meiosis had already resumed in the small oocytes of PD14 *OoCdk1*^{+/*AF*}; *Zp3-Cre* ovaries. The homologous chromosome arms are held together by cohesin during meiosis I until late metaphase I when cohesin is cleaved by separase [23]. Simultaneous staining of SMC3 (structural maintenance of chromosome 3) (Figure 3O, arrows), and CREST (Figure 3Q, arrows) showed that SMC3 was localized along the chromosome axis, indicating that the chromosomes were at prometaphase I in the *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes. These results demonstrate that these growing oocytes expressing Cdk1AF are not arrested at the diplotene stage of prophase I.

Earlier studies have shown that oocytes isolated from prepubertal mouse ovaries or oocytes that are smaller than 60 μm in diameter are meiotically incompetent because they fail to resume meiosis spontaneously when denuded and cultured *in vitro* [6, 7]. This was observed in oocytes isolated from PD14 control *Cdk1*^{+/*SAF*} ovaries where all oocytes were arrested at the GV stage and contained decondensed DNA (Figure 3R, arrowheads, and Figure 3T control oocytes, $n = 320$). We also isolated oocytes from PD14 mutant *OoCdk1*^{+/*AF*}; *Zp3-Cre* ovaries and quantified the proportion of oocytes that had resumed meiosis as evidenced by GVBD. We observed premature *in vivo* resumption of meiosis in PD14 mutant *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes as small as 25 μm in diameter. The results showed that 48% of the oocytes isolated from PD14 mutant *OoCdk1*^{+/*AF*}; *Zp3-Cre* ovaries, which were all smaller than 60 μm in diameter, had undergone premature GVBD with condensed DNA (Figure 3S, arrows, and quantification in Figure 3T, $n = 450$).

Notably, when the isolated growing *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes were cultured *in vitro* for 16 h, the proportion of mutant oocytes undergoing GVBD increased to 68% (Figure 3U, $n = 450$), whereas none of the control *Cdk1*^{+/*SAF*} oocytes underwent GVBD in the *in vitro* culture (Figure 3U, $n = 340$). Therefore, the Cdk1AF increases the rate of meiosis resumption in otherwise meiotically incompetent mouse oocytes. These results indicate that the inhibition of Cdk1 kinase activity by its inhibitory phosphorylation in growing mouse oocytes is important for sustaining arrest in the middle of prophase I.

Accumulation of DNA damage in growing OoCdk1^{+/*AF*}; *Zp3-Cre* oocytes

Increased Cdk1 activity led to premature resumption of meiosis in the *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes (Figure 3G–3U), which was followed by oocyte depletion (Figure 3F), and immunostaining for γ -H2AX in PD14 ovarian sections revealed increased DNA damage in the *OoC-*

dk1^{+/*AF*}; *Zp3-Cre* oocytes (Figure 4B, arrows). In contrast, γ -H2AX foci were mostly absent in control *Cdk1*^{+/*SAF*} oocytes (Figure 4A, arrowheads).

We next isolated oocytes from PD14 *OoCdk1*^{+/*AF*}; *Zp3-Cre* and control *Cdk1*^{+/*SAF*} ovaries and immunostained for γ -H2AX. The DNA in *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes showed intense γ -H2AX staining (Figure 4D, arrow), whereas the γ -H2AX signal was mostly absent in the DNA of control *Cdk1*^{+/*SAF*} oocytes (Figure 4C, arrowhead). Quantification of the γ -H2AX foci-positive oocytes indicated that 94% ($n = 85$) of the *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes were positive for nuclear γ -H2AX foci compared with only 4% ($n = 80$) of the control *Cdk1*^{+/*SAF*} oocytes (Figure 4E). Western blot analysis also confirmed the increased level of γ -H2AX in *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes (Figure 4F). The level of 53BP1, another marker of DNA double-strand break [24] was also increased in *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes (Figure 4F). These results indicate that the elevated Cdk1 kinase activity leads to DNA damage in the growing *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes and that this is apparently the cause of the accelerated oocyte death.

The Cdk1-ATM cascade activates Chk2 in OoCdk1^{+/*AF*}; *Zp3-Cre* oocytes

In somatic cells, DNA double-strand breaks cause the activation of ATM, the master regulator of the DNA damage checkpoint. ATM is the major kinase involved in the phosphorylation of γ -H2AX during the cellular response to double-strand breaks [25, 26]. However, the induction of DNA damage in fully grown mouse oocytes does not lead to their death or prevent the oocytes from maturing because in fully grown oocytes the DNA damage-response pathway is not activated effectively [27, 28].

To test whether the DNA damage-response pathway in growing *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes is triggered, we analyzed the levels of p-ATM (S1981) in relation to γ -H2AX in these oocytes. We found that both p-ATM (Figure 4J, arrow) and γ -H2AX (Figure 4K, arrow) foci were present in the nuclei of GVBD oocytes in PD14 *OoCdk1*^{+/*AF*}; *Zp3-Cre* ovaries, whereas staining for p-ATM (Figure 4G, arrowhead) and γ -H2AX foci (Figure 4H, arrowhead) were absent in control *Cdk1*^{+/*SAF*} oocytes. In addition, we also observed DNA damage and increased p-ATM signal in analyzed GV stage *OoCdk1*^{+/*AF*}; *Zp3-Cre* oocytes (Supplementary information, Figure S3B, arrows). Thus, these results suggest that elevated Cdk1 activation in growing oocytes triggers DNA damage, and unlike in fully grown oocytes, the DNA damage in growing oocytes lead to oocyte death.




Figure 4 Expression of Cdk1AF leads to DNA damage and ATM activation in growing *OoCdk1^{+/AF}; Zp3-Cre* oocytes. **(A)** Absence of γ -H2AX staining in the oocytes of control PD14 *Cdk1^{+/SAF}* ovarian sections (arrowheads). **(B)** Prominent γ -H2AX staining in the oocytes of PD14 *OoCdk1^{+/AF}; Zp3-Cre* ovaries (arrows). DDX4 was used to label the oocytes and the same sections were counterstained with hematoxylin to visualize ovarian histology. The experiments were repeated more than three times each, and representative images are shown. **(C)** Absence of γ -H2AX staining in isolated oocytes of control PD14 *Cdk1^{+/SAF}* mice (arrowhead). **(D)** Strong γ -H2AX staining in isolated oocytes from PD14 *OoCdk1^{+/AF}; Zp3-Cre* ovaries (arrow). **(E)** Quantification of γ -H2AX-positive oocytes that were isolated from PD14 *Cdk1^{+/SAF}* and *OoCdk1^{+/AF}; Zp3-Cre* ovaries. Numbers of each genotype of oocytes checked (*n*) are shown. **(F)** Western blot for γ -H2AX and 53BP1 levels in oocytes isolated from PD14 *Cdk1^{+/SAF}* and *OoCdk1^{+/AF}; Zp3-Cre* ovaries. β -actin was used as the loading control. Lysate from 200 oocytes was loaded in each lane. The experiments were repeated more than three times. **(G-L)** High level of p-ATM (S1981) accompanied by increased γ -H2AX staining in PD14 *OoCdk1^{+/AF}; Zp3-Cre* oocytes (**J-L**, arrows), in comparison with PD14 control *Cdk1^{+/SAF}* oocytes where there was almost no signal for p-ATM (S1981) or γ -H2AX (**G-I**, arrowheads). The experiments were repeated more than three times each, and representative images are shown.

Ablation of *Chk2* prevents the death of *OoCdk1^{+/AF}; Zp3-Cre* oocytes

Chk2 is a downstream effector of ATM kinase that responds to DNA damage and it is a component of the DNA damage-sensing checkpoint [29-31]. ATM activates *Chk2* by phosphorylating it at T68 [32]. We found that although p-*Chk2* (T68) and total *Chk2* levels were almost undetectable in control PD14 *Cdk1^{+/SAF}* oocytes, the p-*Chk2* (T68) and total *Chk2* levels were elevated in *OoCdk1^{+/AF}; Zp3-Cre* oocytes (Figure 5A), showing that the DNA damage in *OoCdk1^{+/AF}; Zp3-Cre* oocytes leads to the activation of the *Chk2* pathway. We also found that the level of active caspase 3 was elevated in *OoCdk1^{+/AF}; Zp3-Cre* oocytes as compared with the control PD14 *Cdk1^{+/SAF}* oocytes (Figure 5B).

To provide *in vivo* evidence that Cdk1AF activates the *Chk2* pathway and leads to the death of *OoCdk1^{+/AF}; Zp3-Cre* oocytes, we generated *OoCdk1^{+/AF}; Zp3-Cre* oocytes in a *Chk2*-null background (referred to as *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* mice). We found that although the ovaries in *OoCdk1^{+/AF}; Zp3-Cre* mice were depleted of almost all follicles by 2 months of age (Figure 5C and Supplementary information, Figure S3C), *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* ovaries of the same age had many healthy-looking follicles (Figure 5D, arrows and Supplementary information, Figure S3C). These results suggest that the loss of *Chk2* largely prevented the death of oocytes caused by the Cdk1AF in *OoCdk1^{+/AF}; Zp3-Cre* oocytes.

We further stained for γ -H2AX in ovarian sections from 2-month-old *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* mice and found that the surviving oocytes still exhibited DNA damage (Figure 5E, arrows). As a control, γ -H2AX foci were mostly absent in the oocytes in 2-month-old control




Figure 5 Cdk1AF triggers the Chk2-p63/p53 and cell death pathways in *OoCdk1^{+/AF}; Zp3-Cre* oocytes. **(A)** Western blot for p-Chk2 (T68) and Chk2 in oocytes isolated from PD14 *Cdk1^{+/SAF}* and *OoCdk1^{+/AF}; Zp3-Cre* ovaries. **(B)** Western blot for active caspase 3 in oocytes isolated from PD14 *Cdk1^{+/SAF}* and *OoCdk1^{+/AF}; Zp3-Cre* ovaries. β -actin was used as the loading control. Lysate from 200 oocytes was loaded in each lane. The experiments were repeated more than three times each. **(C, D)** Hematoxylin-stained sections of mouse ovary at 2 month of age. Rescue of follicle depletion by the concomitant deletion of *Chk2* in *OoCdk1^{+/AF}; Zp3-Cre* ovaries. By 2 months of age, although healthy follicular structures and oocytes had mostly disappeared in mutant *OoCdk1^{+/AF}; Zp3-Cre* ovaries **(C)**, concomitant deletion of *Chk2* largely prevented the death of follicles in double-mutant *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* ovaries, and many surviving oocytes could be observed **(D)**, arrows). The experiments were repeated more than three times each, and for each time and each age ovaries from one mouse of each genotype were used. **(E)** Surviving oocytes in PD60 double-mutant *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* ovaries stained positive for γ -H2AX (arrows) indicating that DNA damage is still present. **(F)** Control *Cdk1^{+/SAF}* ovaries with γ -H2AX-negative oocytes. The experiments were repeated more than three times, and representative images are shown. **(G)** Western blot for p63 and p53 levels in oocytes isolated from PD14 *Cdk1^{+/SAF}*, *OoCdk1^{+/AF}; Zp3-Cre*, and *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* ovaries. β -actin was used as the loading control, and lysate from 200 oocytes was loaded in each lane. The experiments were repeated more than three times.

Cdk1^{+/SAF} ovarian sections (Figure 5F, arrowheads). Thus, the prevention of oocyte death was mostly due to the lack of Chk2 or its downstream effectors but not due to the repair of DNA because the DNA damage was already present in *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* oocytes.

We checked whether the activated Chk2 in mutant *OoCdk1^{+/AF}; Zp3-Cre* oocytes (Figure 5A) triggered the accumulation of p63 and p53 which may mediate the depletion of oocytes. Indeed, the levels of p63 and p53 in *OoCdk1^{+/AF}; Zp3-Cre* oocytes were much higher than in the control *Cdk1^{+/SAF}* oocytes (Figure 5G). Moreover, the levels of p63 and p53 in *OoCdk1^{+/AF}; Zp3-Cre; Chk2^{-/-}* oocytes returned to the level in *Cdk1^{+/SAF}* oocytes (Figure 5G). Thus, DNA damage in *OoCdk1^{+/AF}; Zp3-Cre* oocyte leads to the Chk2-dependent activation of p53 and p63, which in turn leads to oocyte elimination. These results

show that the phosphorylation-mediated suppression of Cdk1 in small oocytes is essential in preventing loss of oocytes prematurely.

Discussion

The mechanisms sustaining the prolonged meiotic arrest in female germ cells are of great importance and far from clear. In the present study, we have demonstrated that the maintenance of meiotic arrest in mammalian female germ cells relies on the constant suppression of Cdk1 activity, which is largely achieved by inhibitory phosphorylation of Cdk1. This constant suppression of Cdk1 activity is not only needed for maintaining the arrested meiotic cell cycle, but it is also essential for safeguarding the integrity of the DNA in the dormant and

growing oocytes. These molecular mechanisms are essential for preserving the germ cell pool and maintaining the female reproductive lifespan.

There are at least three processes known to be required for Cdk1 activity: the availability of cyclin, dephosphorylation of T14 and Y15, and phosphorylation of the T-loop. In fully grown mouse oocytes, the activity of Cdk1 remains suppressed by the maintenance of low levels of cyclin B1 and cyclin B2 [10, 33] or through Wee1B/Myt1-mediated phosphorylation of Cdk1 [14]. A previous study has shown that growing oocytes lacking *Cdh1* accumulate higher levels of cyclin B1 due to low APC/C activation, but higher levels of cyclin B1 are not sufficient to drive smaller oocytes into premature resumption of meiosis *in vivo* within the follicles [11]. Even so, it is still likely that Cdk1-activating cyclins in growing oocytes have not reached high enough levels to activate Cdk1. This is because that although the growing, meiotically incompetent oocytes already contain higher concentration of cyclin B1 protein than Cdk1, the level of cyclin B1 further increases several folds by the time the oocytes are fully grown [34].

Similarly, downregulation of Wee1B in mouse oocytes causes more rapid GVBD in fully grown oocytes, but GVBD is not seen in growing oocytes, which implies that Wee1B is not important for regulating Cdk1 phosphorylation in growing oocytes [14]. However, the lack of GVBD in growing oocytes by knockdown of Wee1B in this transgenic mouse model might be due to the fact that Wee1B was only partially depleted [14]. In addition, another kinase in oocytes, Myt1, might still be able to maintain meiotic arrest in the absence of Wee1B [14]. It is also possible that the growing oocytes might not sufficiently activate Cdc25 for the removal of Cdk1 inhibitory phosphorylation. Therefore, whether Cdk1 is regulated in smaller oocytes is far from understood. However, our direct genetic approach revealed that the phosphorylation-mediated suppression of Cdk1 activity is one of the crucial factors for the prolonged prophase I arrest in dormant and growing oocytes.

When fully grown oocytes resume meiosis after Cdk1 activation, it does not lead to DNA damage. Even the induction of DNA damage in fully grown mouse oocytes does not lead to their death; instead, these oocytes still progress through meiosis. This is because DNA damage in fully grown oocytes does not effectively activate the DNA damage-response pathway [27, 28]. In contrast, as we have shown in this study, premature activation of Cdk1 in meiotically incompetent dormant and growing oocytes cannot be tolerated and it triggers massive DNA damage and oocyte death. This is because as we showed the activated Cdk1 in smaller oocytes leads to active

response to DNA damage through the ATM-Chk2-p53/p63-mediated pathway and activation of this pathway quickly causes oocyte loss. However, the mechanism for increased level of Chk2 protein observed in growing mutant oocytes remains unclear. It is possible that the degradation of Chk2 might be affected in the mutant oocytes as ubiquitin-mediated Chk2 degradation has been recently reported to regulate Chk2 protein level in other cell lines [35].

Interestingly, we found that genetic ablation of *Chk2* largely prevented the death of oocytes, which is in line with a previous study showing that the death of mouse oocytes due to ionizing radiation-induced DNA double-strand breaks is prevented by concomitant ablation of *Chk2* [29]. Nevertheless, a possibility for a contribution of ATR (ataxia telangiectasia and Rad3-related)-Chk1 pathway [36] for death of the mutant oocytes cannot be ruled out. Our results thus show that a direct phosphorylation-mediated suppression of Cdk1 activity is essential for safeguarding the DNA integrity of meiotically arrested female germ cells.

Studies in both yeast and mammals have revealed that Cdk1 is involved in the regulation of DNA double-strand break processing, DNA repair by homologous recombination, and DNA damage checkpoint activation, through which the DNA-repair capacity is coordinated with progression through the cell cycle [37]. In human MCF-7 cell lines, inhibition of Wee1 increases Cdk1 activity and results in the inactivation of DNA repair [38]. In yeast, the detection of DNA damage in G2 activates a checkpoint that prevents entry into mitosis through the maintenance of the phosphorylation of Y15-Cdc2p (the single Cdk in yeast); however, inactivation of Wee1 and Cdc25 abolishes the G2 checkpoint control [39]. Similarly, Cdk1 hyperactivation in the *Drosophila* imaginal wing disc also causes genome instability and chromosomal abnormalities [40]. These findings suggest that overactivated Cdk1 activity leads to genome instability, indicating that Cdk1 activity needs to be properly suppressed in order to maintain genome stability.

Due to the multiple downstream substrates of Cdk1, elevated Cdk1 activity might cause DNA damage through various routes. For example, Cdk1 has been shown to be involved in the regulation of DNA damage-related molecules such as polo-like kinase 1 (PLK1) in U2OS and HeLa cells [41], making it attractive to hypothesize that Cdk1 activation in quiescent and growing oocytes might enhance PLK1 activity and affect the DNA damage-repair response. Previous studies showed that Wee1 inhibition (and thus elevated Cdk1 activity) in mitotic cells leads to dysregulated endonuclease activity and might cause DNA damage [42]. More in-depth stud-

ies are needed to elucidate the mechanisms behind the DNA damage in dormant and growing oocytes caused by active Cdk1. Understanding the response to DNA damage by mouse oocytes in a developmental stage specific manner may have important implications in helping identify additional targets for fertility preservation therapies in cancer patients who are at the verge of losing ovarian reserve.

The rapid loss of growing follicles in *OoCdk1^{+SAF}; Zp3-Cre* ovaries seems to indirectly increase the activation and subsequent depletion of primordial follicles in the adult ovaries. It is known that the larger follicles secrete inhibitory signals such as Anti-Müllerian hormone to prevent primordial follicle recruitment [43], and it is likely that in the absence of growing follicles in *OoCdk1^{+SAF}; Zp3-Cre* ovaries, the activation of the primordial follicle pool is accelerated (Supplementary information, Figure S4). In summary, our data demonstrate that the Cdk1 inhibitory phosphorylation sustains the prolonged period of meiotic arrest in female germ cells and protects the small dormant and growing oocytes from accumulating DNA damage and their death. Therefore, the phosphorylation-mediated direct suppression of Cdk1 kinase activity is essential for preserving the oocyte pool for maintaining the reproductive lifespan of female mammals.

Materials and methods

Generation of *Cdk1^{AF}* conditional expression knock-in mice

Mouse genomic DNA harboring the *Cdk1* locus was retrieved from the clone pBeloBACII 305J21 (ResGen) and inserted into the pBlight-TK vector as described previously [44]. Using recombineering methods [45], a floxed neomycin-selection cassette adjacent to a strong splice acceptor site (derived from exon 2 of the mouse *Engrailed* gene) with multiple stop codons was introduced to the 5' end of exon 3 of *Cdk1*, which had been modified by site-directed mutagenesis to introduce the point mutations into the *Cdk1* coding sequence to generate the T14A and Y15F substitutions. The resultant targeting vector (PKB945) was linearized by *NotI* digestion and electroporated into ES cells. After positive and negative selection with Geneticin and ganciclovir, respectively, genomic DNA from resistant ES cell colonies was used to screen for homologous recombination by Southern hybridization. Correctly targeted ES cell clone #2681 was used for generation of the *Cdk1^{AF}* conditional expression knock-in mouse strain. The presence of the point mutations in the knock-in mice was verified by sequencing of the DNA from tail snips (see Supplementary information, Figure S1). *Cdk1^{+SAF}* mice carry one allele of wild-type *Cdk1* and one allele of *Cdk1^{SAF}*.

Cdk1^{+SAF} mice of mixed background (129S1/SvImJ and C57BL/6) were backcrossed to C57BL/6J mice for four generations and then crossed with transgenic mice carrying *Gdf-9* promoter-mediated Cre recombinase in a C57BL/6J background¹⁸ to induce the expression of the *Cdk1^{AF}* allele in the oocytes of pri-

mordial follicles and further developed follicles (*OoCdk1^{+AF};Gdf-Cre* mice). The *Cdk1^{+SAF}* mice were also crossed with transgenic mice carrying *Zp3* promoter-mediated Cre recombinase in a C57BL/6J background [21] to obtain mutant female mice expressing the *Cdk1^{AF}* allele only in their growing oocytes but not in the primordial oocytes (*OoCdk1^{+AF}; Zp3-Cre* mice). Littermates that expressed a wild-type *Cdk1* allele and a *Cdk1^{SAF}* allele, but not the *Cre* transgene, were used as controls (*Cdk1^{+SAF}* mice). *OoCdk1^{+AF}; Zp3-Cre*; *Chk2^{-/-}* mice were obtained by multiple rounds of crossings among *Cdk1^{+SAF}; Zp3-Cre*, and *Chk2^{-/-}* mice [46].

All mice were housed under controlled environmental conditions with free access to water and food. Illumination was on between 0600 hours and 1800 hours. Experimental protocols were approved by the regional ethical committee of the University of Gothenburg, Sweden, and by the Institutional Animal Care and Use Committee at the Biological Research Centre mouse facility at Biopolis, Singapore.

Reagents, antibodies, and immunological detection methods

Rabbit polyclonal antibodies to γ -H2AX (phospho S139), DDX4, and Chk2 were purchased from Abcam (Cambridge, UK). Rabbit polyclonal antibodies to phospho-Chk2 (T68) and 53BP1 were purchased from Novus Biologicals (Littleton, CO, USA). Mouse monoclonal antibodies against Cdk1, p63, and p-ATM (S1981) were purchased from Santa Cruz Biotechnology (Dallas, TX, USA). Rabbit polyclonal antibodies against phospho-Cdk1 (Y15) and mouse monoclonal antibodies against p53 were obtained from Cell Signaling Technologies (Beverly, MA, USA). Mouse monoclonal antibodies against β -actin, M2, and M16 media, dibutyryl cyclic AMP (dbcAMP), 4',6-diamidino-2-phenylindole (DAPI), and mineral oil were purchased from Sigma (St. Louis, MO, USA). Human anti-CREST serum was from Antibodies (Davis, CA, USA). Western blots were performed according to the instructions of the suppliers of the different antibodies and visualized using the ECL Prime Western Blotting Detection System (GE Healthcare, UK).

Isolation of oocytes from postnatal mouse ovaries

Mice were killed and their ovaries were dissected free of fat and connective tissue using a stereo microscope (Carl Zeiss, Germany). The ovaries were then minced with a pair of dissection scissors before being incubated in 0.05% collagenase dissolved in Dulbecco's modified Eagle's medium-F12 (DMEM/F12; Invitrogen) supplemented with 4 mg/ml bovine serum albumin (BSA), 100 U/ml penicillin, and 100 μ g/ml streptomycin with frequent agitation and pipetting. After the tissues had mostly been digested by collagenase, usually within 45-60 min, EDTA was added to this mixture to a final concentration of 40 mM, and the mixture was incubated at 37 °C with frequent pipetting for another 15-20 min until clusters of granulosa cells or other cells were completely dispersed. The mixture of cells and oocytes was then washed once with PBS by centrifugation, and pellets were resuspended in 5 ml of the above-mentioned serum-free DMEM/F12 medium and poured into 6-cm culture dishes. Oocytes were picked by mouth pipetting. For Western blots, oocytes were washed in HBSS, and lysed in a buffer containing 50 mM Tris HCl (pH 8.0), 120 mM NaCl, 20 mM NaF, 20 mM β -glycerophosphate, 1 mM EDTA, 6 mM EGTA (pH 8.0), 1% NP-40, 1 mM DTT, 5 mM benzamidine, 1 mM PMSF, 250 μ M sodium orthovanadate, 10 μ g/ml aprotinin,

10 µg/ml leupeptin, and 1 µg/ml pepstatin and lysate from 200 growing oocytes was loaded in each lane.

For the isolation of fully grown oocytes, 3- to 4-week-old female mice were injected with 7.5 IU pregnant mare serum gonadotropin intraperitoneally, and 42–44 h later the mice were killed and their ovaries were collected in 100 µg/ml dbcAMP-containing M2 medium. Fully grown GV-stage oocytes surrounded by cumulus cells were released by puncturing the ovaries in M2 medium supplemented with dbcAMP to maintain meiotic arrest during the *in vitro* operation. Oocytes were freed from the attached cumulus cells by repetitive pipetting through a narrow-bore glass pipette.

Oocyte immunofluorescence and confocal microscopy

Oocytes obtained by collagenase-EDTA digestion were fixed in 4% paraformaldehyde (PFA). After the fixed oocytes were permeabilized with 0.5% Triton X-100 at room temperature (RT) for 20 min, they were blocked with 1% BSA-supplemented PBS for 1 h and incubated at 4 °C with anti-γ-H2AX (phospho S139) antibody (1:500 dilution) and anti-p-ATM (S1981) antibody (1:500 dilution) overnight followed by an incubation with Alexa Fluor 488 goat anti-rabbit IgG and Alexa Fluor 555 goat anti-mouse IgG (Invitrogen Molecular Probes, Carlsbad, CA, USA) for 1 h at RT. DNA was stained with DAPI (1 µg/ml in PBS). Finally, the oocytes were mounted on glass slides with DABCO-containing mounting medium and examined by laser-scanning confocal microscopy (Zeiss LSM700 Inverted).

Chromosome spreads

For chromosome spreads, the oocytes isolated from PD14 *Cdk1^{+/SAF}* and *OoCdk1^{+/AF}*; *Zp3-Cre* ovaries were first freed of the zona pellucida with acidic Tyrode's solution (Sigma-Aldrich). After a brief recovery in M2 medium, the oocytes were transferred onto glass slides and fixed in a solution of 1% PFA in distilled H₂O (pH 9.2) containing 0.15% Triton X-100 and 3 mM dithiothreitol. The slides were left to dry and then blocked with 1% BSA in PBS for 1 h at RT. The spreads were incubated with primary antibodies against CREST and SMC3 (Abcam) overnight at 4 °C using a dilution of 1:100. After three washes, the slides were incubated with the corresponding secondary antibodies (Invitrogen) at 1:500 dilution for 1 h at RT. DNA on the slides was stained with DAPI, and the slides were mounted for observation with laser-scanning confocal microscopy (Zeiss LSM700 Inverted).

Histone H1 kinase assay

Cdk1 activity was measured using histone H1 kinase assays. This assay was performed using 10 oocytes in a 10 µl reaction volume based on a published protocol [23]. In brief, the reaction was carried out in a buffer containing 25 mM HEPES (pH 7.4), 15 mM MgCl₂, 20 mM EGTA, 50 mM NaCl, 0.05% NP-40, 0.1 mM ATP, 100 mM PMSF, 1 mM DTT, 1 µg histone H1 (Roche, Mannheim, Germany), and 3 µCi of (γ-³²P) ATP (Perkin-Elmer, Boston, MA, USA) and incubated at 37 °C for 30 min. Samples were boiled in SDS sample buffer and then separated on a 15% SDS-polyacrylamide gel. The radioactive signal was detected by exposing the gels on an FLA 3000 phosphorimager (Fuji, Tokyo, Japan).

Histological analysis of ovarian tissues and immunofluorescence staining

Histological analysis of ovarian tissue was performed as pre-

viously described [17]. Briefly, ovaries were fixed in 4% PFA, dehydrated, and embedded in paraffin. Paraffin-embedded ovaries were serially sectioned at 8 µm thickness and stained with hematoxylin for morphological observation. To quantify total number of follicles per ovary, every fifth section was counted throughout the entire ovary, beginning with the first section, and the final count was multiplied by 5 as a correction factor. To calculate the ovarian follicles at different stages of development, all the follicles were counted in all sections of an ovary based on the well-accepted standards established by Pedersen and Peters [47]. Oocytes that had resumed meiosis were identified by observation of the condensed chromosomes.

To detect the DNA damage in ovarian sections by immunofluorescence staining, the sections were treated with 0.01% sodium citrate buffer (pH 6.0) for antigen retrieval. After blocking with 1.5% goat serum in PBS, the sections were incubated with anti-γ-H2AX (phospho S139) and DDX4 primary antibodies overnight at 4 °C followed by an incubation with Alexa Fluor 488 goat anti-rabbit IgG and Alexa Fluor 555 goat anti-mouse IgG (Invitrogen Molecular Probes) for 1 h at RT. The sections were then stained with DAPI for 5 min. The fluorescent images were taken with a Zeiss Axio Scope A1 upright microscope. After imaging for γ-H2AX, DDX4, and DAPI, the same sections were stained with hematoxylin to clearly visualize the histology.

Statistical analysis

All experiments were repeated at least three times. For comparisons between groups, the differences were calculated with Student's *t*-test, and a difference was considered to be significant if *p* < 0.05.

Acknowledgments

This study was supported by grants to KL from the Jane and Dan Olssons Foundation, the LUA/ALF-medel Västra Götalandsregionen, AFA Insurance, the Swedish Research Council, and the Swedish Cancer Foundation, Sweden. PK and MKD were supported by the Biomedical Research Council of A*STAR (Agency for Science, Technology and Research), Singapore.

Author Contributions

DA, PK, and KL designed the study. DA, KB, JZ, MH, SR, MBB, MS, and MKD performed the experiments. DA, PK, and KL wrote the manuscript.

Competing Financial Interests

The authors declare no competing financial interests.

References

- 1 Nagaoka SI, Hassold TJ, Hunt PA. Human aneuploidy: mechanisms and new insights into an age-old problem. *Nat Rev Genet* 2012; **13**:493-504.
- 2 Zheng W, Zhang H, Gorre N, Risal S, Shen Y, Liu K. Two classes of ovarian primordial follicles exhibit distinct developmental dynamics and physiological functions. *Hum Mol Genet* 2014; **23**:920-928.
- 3 McGee EA, Hsueh AJ. Initial and cyclic recruitment of ovarian follicles. *Endocr Rev* 2000; **21**:200-214.

- 4 Adhikari D, Liu K. Molecular mechanisms underlying the activation of mammalian primordial follicles. *Endocr Rev* 2009; **30**:438-464.
- 5 Eppig JJ, Viveiros MM, Bivens CM, De La Fuente R. Regulation of mammalian oocyte maturation. In: Leung PC, Adashi EY, eds. *The Ovary*. Academic Press: Amsterdam 2004:113-129.
- 6 Sorensen RA, Wassarman PM. Relationship between growth and meiotic maturation of the mouse oocyte. *Dev Biol* 1976; **50**:531-536.
- 7 Hirao Y, Miyano T, Kato S. Acquisition of maturational competence in *in vitro* grown mouse oocytes. *J Exp Zool* 1993; **267**:543-547.
- 8 Nishimura T, Fujii W, Kano K, Sugiura K, Naito K. Analyses of the involvement of PKA regulation mechanism in meiotic incompetence of porcine growing oocytes. *Biol Reprod* 2012; **87**:53.
- 9 Goren S, Piontkewitz Y, Dekel N. Meiotic arrest in incompetent rat oocytes is not regulated by cAMP. *Dev Biol* 1994; **166**:11-17.
- 10 Reis A, Chang HY, Levasseur M, Jones KT. APCcdh1 activity in mouse oocytes prevents entry into the first meiotic division. *Nat Cell Biol* 2006; **8**:539-540.
- 11 Holt JE, Tran SM, Stewart JL, et al. The APC/C activator FZR1 coordinates the timing of meiotic resumption during prophase I arrest in mammalian oocytes. *Development* 2011; **138**:905-913.
- 12 de Vantery C, Stutz A, Vassalli JD, Schorderet-Slatkine S. Acquisition of meiotic competence in growing mouse oocytes is controlled at both translational and posttranslational levels. *Dev Biol* 1997; **187**:43-54.
- 13 Kim J, Singh AK, Takata Y, et al. LSD1 is essential for oocyte meiotic progression by regulating CDC25B expression in mice. *Nat Commun* 2015; **6**:10116.
- 14 Han SJ, Chen R, Paronetto MP, Conti M. Wee1B is an oocyte-specific kinase involved in the control of meiotic arrest in the mouse. *Curr Biol* 2005; **15**:1670-1676.
- 15 Norbury C, Blow J, Nurse P. Regulatory phosphorylation of the p34cdc2 protein kinase in vertebrates. *EMBO J* 1991; **10**:3321-3329.
- 16 Krek W, Nigg EA. Mutations of p34cdc2 phosphorylation sites induce premature mitotic events in HeLa cells: evidence for a double block to p34cdc2 kinase activation in vertebrates. *EMBO J* 1991; **10**:3331-3341.
- 17 Reddy P, Liu L, Adhikari D, et al. Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool. *Science* 2008; **319**:611-613.
- 18 Lan ZJ, Xu X, Cooney AJ. Differential oocyte-specific expression of Cre recombinase activity in GDF-9-iCre, Zp3cre, and Msx2Cre transgenic mice. *Biol Reprod* 2004; **71**:1469-1474.
- 19 Heijink A, Blomen V, Bisteau X, et al. A haploid genetic screen identifies the G1/S regulatory machinery as a determinant of Wee1 inhibitor sensitivity. *Proc Natl Acad Sci USA* 2015; **112**.
- 20 Rogakou EP, Boon C, Redon C, Bonner WM. Megabase chromatin domains involved in DNA double-strand breaks *in vivo*. *J Cell Biol* 1999; **146**:905-916.
- 21 de Vries WN, Binns LT, Fancher KS, et al. Expression of Cre recombinase in mouse oocytes: a means to study maternal effect genes. *Genesis* 2000; **26**:110-112.
- 22 Adhikari D, Diril MK, Busayavalasa K, et al. Mastl is required for timely activation of APC/C in meiosis I and Cdk1 reactivation in meiosis II. *J Cell Biol* 2014; **206**:843-853.
- 23 Kudo NR, Wassmann K, Anger M, et al. Resolution of chiasmata in oocytes requires separase-mediated proteolysis. *Cell* 2006; **126**:135-146.
- 24 Panier S, Boulton SJ. Double-strand break repair: 53BP1 comes into focus. *Nat Rev Mol Cell Biol* 2014; **15**:7-18.
- 25 Bakkenist CJ, Kastan MB. DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature* 2003; **421**:499-506.
- 26 Shiloh Y, Ziv Y. The ATM protein kinase: regulating the cellular response to genotoxic stress, and more. *Nat Rev Mol Cell Biol* 2013; **14**:197-210.
- 27 Ma JY, Ou Yang YC, Wang ZW, et al. The effects of DNA double-strand breaks on mouse oocyte meiotic maturation. *Cell Cycle* 2013; **12**:1233-1241.
- 28 Marangos P, Carroll J. Oocytes progress beyond prophase in the presence of DNA damage. *Curr Biol* 2012; **22**:989-994.
- 29 Bolcun-Filas E, Rinaldi VD, White ME, Schimenti JC. Reversal of female infertility by Chk2 ablation reveals the oocyte DNA damage checkpoint pathway. *Science* 2014; **343**:533-536.
- 30 Hirao A, Cheung A, Duncan G, et al. Chk2 is a tumor suppressor that regulates apoptosis in both an ataxia telangiectasia mutated (ATM)-dependent and an ATM-independent manner. *Mol Cell Biol* 2002; **22**:6521-6532.
- 31 Matsuoka S, Huang M, Elledge SJ. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 1998; **282**:1893-1897.
- 32 Ahn JY, Schwarz JK, Piwnicka-Worms H, Canman CE. Threonine 68 phosphorylation by ataxia telangiectasia mutated is required for efficient activation of Chk2 in response to ionizing radiation. *Cancer Res* 2000; **60**:5934-5936.
- 33 Gui L, Homer H. Hec1-dependent cyclin B2 stabilization regulates the G2-M transition and early prometaphase in mouse oocytes. *Dev Cell* 2013; **25**:43-54.
- 34 Kanatsu-Shinohara M, Schultz RM, Kopf GS. Acquisition of meiotic competence in mouse oocytes: absolute amounts of p34(cdc2), cyclin B1, cdc25C, and wee1 in meiotically incompetent and competent oocytes. *Biol Reprod* 2000; **63**:1610-1616.
- 35 Garcia-Limones C, Lara-Chica M, Jimenez-Jimenez C, et al. CHK2 stability is regulated by the E3 ubiquitin ligase SIAH2. *Oncogene* 2016.
- 36 Chen Y, Poon RY. The multiple checkpoint functions of CHK1 and CHK2 in maintenance of genome stability. *Front Biosci* 2008; **13**:5016-5029.
- 37 Trovesi C, Manfrini N, Falcettoni M, Longhese MP. Regulation of the DNA damage response by cyclin-dependent kinases. *J Mol Biol* 2013; **425**:4756-4766.
- 38 Krajewska M, Heijink AM, Bisselink YJ, et al. Forced activation of Cdk1 via wee1 inhibition impairs homologous recombination. *Oncogene* 2013; **32**:3001-3008.
- 39 Raleigh JM, O'Connell MJ. The G(2) DNA damage checkpoint targets both Wee1 and Cdc25. *J Cell Sci* 2000; **113**:1727-1736.

- 40 Ayeni JO, Varadarajan R, Mukherjee O, *et al.* Dual phosphorylation of cdk1 coordinates cell proliferation with key developmental processes in *Drosophila*. *Genetics* 2014; **196**:197-210.
- 41 Benada J, Burdova K, Lidak T, von Morgen P, Macurek L. Polo-like kinase 1 inhibits DNA damage response during mitosis. *Cell Cycle* 2015; **14**:219-231.
- 42 Dominguez-Kelly R, Martin Y, Koundrioukoff S, *et al.* Wee1 controls genomic stability during replication by regulating the Mus81-Eme1 endonuclease. *J Cell Biol* 2011; **194**:567-579.
- 43 Durlinger AL, Kramer P, Karels B, *et al.* Control of primordial follicle recruitment by anti-Mullerian hormone in the mouse ovary. *Endocrinology* 1999; **140**:5789-5796.
- 44 Diril MK, Ratnacaram CK, Padmakumar VC, *et al.* Cyclin-dependent kinase 1 (Cdk1) is essential for cell division and suppression of DNA re-replication but not for liver regeneration. *Proc Natl Acad Sci USA* 2012; **109**:3826-3831.
- 45 Lee EC, Yu D, Martinez de Velasco J, *et al.* A highly efficient *Escherichia coli*-based chromosome engineering system adapted for recombinogenic targeting and subcloning of BAC DNA. *Genomics* 2001; **73**:56-65.
- 46 Hirao A, Kong YY, Matsuoka S, *et al.* DNA damage-induced activation of p53 by the checkpoint kinase Chk2. *Science* 2000; **287**:1824-1827.
- 47 Pedersen T, Peters H. Proposal for a classification of oocytes and follicles in the mouse ovary. *J Reprod Fertil* 1968; **17**:555-557.

(Supplementary information is linked to the online version of the paper on the *Cell Research* website.)



This license allows readers to copy, distribute and transmit the Contribution as long as it attributed back to the author. Readers are permitted to alter, transform or build upon the Contribution as long as the resulting work is then distributed under this is a similar license. Readers are not permitted to use the Contribution for commercial purposes. Please read the full license for further details at - <http://creativecommons.org/licenses/by-nc-sa/4.0/>

© The Author(s) 2016