



Published in final edited form as:

*Arterioscler Thromb Vasc Biol.* 2016 October ; 36(10): e82–e86. doi:10.1161/ATVBAHA.116.308261.

## Vascular Smooth Muscle Cells

**Mark W. Majesky**

Center for Developmental Biology & Regenerative Medicine, Seattle Children's Research Institute, and Departments of Pediatrics and Pathology, University of Washington, Seattle, WA 98101

### Keywords

Smooth muscle; aneurysm; TGF $\beta$ ; remodeling; atherosclerosis

### Introduction

Decades of work have shown that vascular smooth muscle cell (SMC) phenotypes are controlled by cues received from the local environment<sup>1-3</sup>. When nestled into a cage of cross-linked collagen and elastin of its own making<sup>4</sup>, medial SMCs exhibit a fully differentiated phenotype conferred by the transcriptional activity of myocardin and serum response factor (SRF) and reinforced by the stabilizing roles of miR-143/145<sup>2,3,5-7</sup>. The central MADS (MCM1, Agamous, Deficiens, SRF) box of SRF is an ancient DNA-binding regulatory platform<sup>8,9</sup> that formed a partnership with the basic and polyglutamine domains of myocardin family proteins early in metazoan evolution<sup>10-12</sup>. This evolutionary innovation allowed for the formation of an extremely versatile type of muscle cell now found in the walls of almost all lumen-containing organs and tissues in adults, and as a transitional form in the embryonic heart.

### Vascular SMCs Adopt Macrophage-Like Phenotypes

Recent studies, many of which have been published or discussed in the pages of *ATVB*, extend the range of phenotypes exhibited by vascular SMCs to those previously thought to be found only in cells of myeloid origin<sup>13-17</sup>. These papers use modern genetic tools to build upon observations made over the past 30 years on SMC responses to injury and cholesterol loading<sup>18-21</sup>. These early studies identified and characterized the SMC “foam cell”, but investigators lacked modern transgenic methods to assess the frequency of SMC foam cell formation in atherogenesis. The surprising result described in recent reports is not that SMC to macrophage-like cell transitions occur in atherosclerosis, but rather the evidently large extent to which a SMC origin can account for cells that express common macrophage markers in the plaque<sup>14-17,22</sup>. The high frequency of SMC-derived plaque macrophages may begin to explain how human atherosclerotic plaques appear as clonal

---

Address correspondence to: Mark W. Majesky, Ph.D., Center for Developmental Biology & Regenerative Medicine, Room 565, M/S C9S-5, Seattle Children's Research Institute, Seattle, WA 98101, mwm84@uw.edu, Phone: (206) 884-3661, Fax: (206) 884-1407.

Disclosures  
None

lesions<sup>23</sup> since the two main cell types in the lesion are now shown to have a common origin.

One question that arises from these observations is why do SMCs so readily express macrophage-like phenotypes?<sup>17,24,25</sup> Is there an inherent sloppiness in SMC regulatory networks with the purposeless outcome that some macrophage marker genes become expressed as SMCs cope with environmental stress? Or is it possible that this property of SMCs has been maintained by selection during vertebrate evolution? Insight into this issue comes from the important work of Vengrenyuk et al who showed that in addition to upregulation of the macrophage markers CD68, Lgals3, Abca1 and Abcg1, cholesterol loading of SMCs also increased expression of Ccl2, Ccl7, and TLR4, which initiates proinflammatory responses to endogenous and microbial ligands<sup>16</sup>. It is possible, therefore, that the dysfunctional macrophage phenotype adopted by SMC foam cells<sup>13-16</sup> may, at least in part, represent an attempt at initiating a wound healing response by injured SMCs. This notion is supported by evidence from multiple groups that disruption of myocardin expression or function in SMCs leads to increased expression of proinflammatory cytokines, chemokines and adhesion molecules<sup>26-28</sup>. The fact that SMCs can also adopt other mesenchymal phenotypes, such as chondrogenic transformation accompanied by down-regulation of Wnt16<sup>29</sup> or produced by Cox2 inhibitors<sup>30</sup>, suggests that permissiveness of SRF MADS box binding interactions with many different activator and repressor cofactor proteins may underlie these phenotype transitions<sup>8,31-33</sup>.

## Control of SMC Phenotypes by TGF $\beta$ and Actin Polymerization

Vascular SMC phenotypes are also controlled by TGF $\beta$  signaling and the state of actin polymerization<sup>34-37</sup>. One factor that has been shown to function in a pathway for actin polymerization and SMC differentiation *in vitro* is the diaphanous-related formin mDia<sup>38,39</sup>. To determine whether mDia has a similar role in promoting the differentiated phenotype of SMCs *in vivo*, Weise-Cross et al expressed a dominant-negative form of mDia from the SM22 $\alpha$  promoter (SM22 $\alpha$ -DNmDia) in transgenic mice<sup>40</sup>. Animals expressing SM22 $\alpha$ -DNmDia exhibit defects in SMC investment in developing arteries, and reduced SMC migration leading to inhibition of neointimal formation after vascular injury in adult mice. In experiments using Myh11creERT2, the authors found that inducible expression of DNmDia in adult arteries reduced the levels of expression of SMC differentiation marker proteins *in vivo*. To identify additional genes regulated by actin polymerization in SMCs, Turczynska et al stimulated SMCs *in vitro* with the F-actin promoting agent jasplakinolide and identified upregulated genes by microarray analysis. Two genes that were identified in this screen were dystrophin and synaptopodin, both of which promote the differentiated phenotype of vascular SMCs *in vitro* and whose expression is downregulated after arterial injury *in vivo*<sup>41</sup>.

In normal healthy arteries, SMCs are organized within medial layers into electrically coupled aggregates. Balint et al showed that SMCs are actually self-organized within normal media via TGF $\beta$ - and p38-dependent assembly of novel adherens junction structures composed of hybrid nanotracks of N-cadherin and cadherin-11<sup>42</sup>. Blocking either N-cadherin or cadherin-11 inhibited formation of SMC collectives and reduced expression of a



ActA2, SM22 $\alpha$ , calponin-1, and smoothelin along with nuclear accumulation of phospho-Smad2 to levels greater than healthy aortas<sup>53</sup>. Elevated expression of these SMC differentiation markers was maintained in aortic medial explant-derived cell cultures along with increased expression of myocardin. Further analysis showed that SMCs obtained from Marfan patients displayed more robust actin stress fibers, increased rhoA-GTP levels, nuclear accumulation of MRTF-A, greater numbers of FAs at the cell surface, and increased levels of nuclear pSmad2 and pSmad3 compared to similar SMC cultures from healthy aortas. Pharmacological inhibition of TGF $\beta$  signaling in Marfan SMCs with the T $\beta$ R1 inhibitor LY364947 (LY) strongly reduced the high pSmad2/pSmad3 levels, SMC differentiation marker expression, and mRNA levels for myocardin and SRF<sup>53</sup>. Atomic force microscopy was used to show that aortic SMCs from Marfan patients were stiffer than counterparts from healthy aortas, and similar results were found for the secreted ECM from these SMC cultures. The authors suggest that upregulated TGF $\beta$  signaling in Marfan SMC cultures may result from reduced TGF $\beta$  binding capacity of defective fibrillin-1 microfibrils in the ECM and the consequent increased levels of free, biologically-active TGF $\beta$  in the medium. Although this is a logical conclusion, it is important to keep in mind that the increased stiffness of aortic cells and matrix in Marfan patients may be a response to aortic wall dilation, rather than a cause of it, and the enhanced expression of a SMC differentiated phenotype may be a feature of lesion progression, rather than lesion initiation. Clearly, it is not a simple matter to tease out cause and effect relations in complex systems in general, and in the aortic wall specifically. More work is required to understand how aortic dilation progresses during aneurysm formation.

## Vascular SMCs and Adaptive Wall Remodeling

Artery wall remodeling responses to intraluminal injuries (catheters, stents, downstream ligation) involves changes at virtually all levels of regulation of SMC gene expression and function. Once identified, some of these regulatory changes offer the potential of a short path to therapeutics. One example of this is found in the report by Wang et al exploring the regulatory role of microRNAs (miR's) in a humanized animal model in which balloon catheter-injured human internal mammary arteries were transplanted into nude rats and subjected to miR profiling<sup>54</sup>. miR-21 was identified as a significantly upregulated target. Experiments employing intravenous fluorescein-tagged locked nucleic acid antagomirs (anti-miR21) showed inhibition of neointimal thickening was obtained along with significant off-target effects on liver, heart, lung and kidney. The authors then tested anti-miR21-coated stents and, compared to bare metal stents, anti-miR21 stents inhibited in-stent restenosis without the off-target effects observed with intravenous injection<sup>54</sup>. miR-21 has a number of targets whose inhibition effectively promotes growth factor signaling pathways<sup>55,56</sup>. Continuing the theme, Xu et al reported that miR-15b/16 inhibits neointimal formation in a rat carotid balloon injury model by promoting the SMC differentiated phenotype, attenuating arterial SMC migration, and inhibiting SMC proliferation<sup>57</sup>. In part, these effects are mediated by the ability of miR-15b/16 to downregulate the expression of the oncoprotein YAP (yes-associated protein). This group previously reported that TEAD1, a DNA-binding partner of YAP, acts to repress SMC differentiation by competing with myocardin for binding to SRF and thereby downregulating expression of the SMC differentiated

phenotype<sup>58</sup>. A role for X-box-binding protein-1 (XBP1)-mediated increases in miR-1274B in neointima formation after femoral artery injury was reported by Zeng et al<sup>59</sup>. In these experiments, miR-1274B was found to target calponin h1 (CNN1) mRNA for inhibition thereby contributing to the PDGFR $\beta$ -mediated increases in SMC migration and proliferation.

Additional novel regulators of vascular SMC migration were identified by Williams et al<sup>60</sup>. These investigators showed that Wnt2 mRNA and protein levels were specifically increased in migrating aortic SMCs<sup>60</sup>. Wnt2 increased Wnt1-inducible signaling pathway protein-1 (WISP1) via  $\beta$ catenin/TCF-dependent pathways in cultured SMCs. Silencing of WISP1 reduced SMC migration *in vitro* and carotid artery ligation in WISP1 $-/-$  mice led to greatly reduced neointima formation compared to WISP1 wild type mice.<sup>60</sup> Vascular SMC migration is also associated with production of a proteoglycan and hyaluronan (HA)-rich provisional ECM.<sup>61</sup> Kiene et al reported an important role for hyaluronic acid synthase (HAS)-3 during neointimal formation in mice.<sup>62</sup> When examined 28d after carotid ligation injury in HAS3-deficient mice, HA content of the artery wall was greatly reduced and formation of a neointima was strongly inhibited compared to wild type animals.<sup>62</sup> Transcriptome analysis in HAS3-deficient carotid arteries suggested that genes involved in cell migration including focal adhesion proteins, SMC integrins, and PDGF-BB signal pathway mediators were downregulated. The authors conclude that HAS3-mediated hyaluronan production supports important steps in SMC migration and neointimal formation *in vivo*.

## Vascular SMCs, Inflammation and Atherosclerosis

Throughout the long history of research on the pathogenesis of atherosclerosis, the close association between intimal SMCs and inflammatory cells has been the subject of intense investigation.<sup>63,64</sup> As discussed earlier in this article, recent studies shed new light on the relation between vascular SMC phenotype transitions and expression of proinflammatory genes.<sup>16,26,27</sup> This relation can also be extended to senescent vascular SMCs that are present within atherosclerotic plaques. Rather than being inert bystanders in plaque progression, senescent human SMCs exhibit a proinflammatory senescence-associated secretory phenotype characterized by release of multiple cytokines and chemokines and driven by autocrine-acting interleukin-1 $\alpha$ .<sup>65</sup> Senescent SMCs release MMP9, secrete less collagen, promote mononuclear cell chemotaxis, and stimulate adjacent endothelial cells to express a proadhesive and proinflammatory phenotype. Thus senescent SMCs, possibly through the same down-regulation of myocardin-SRF regulated SMC gene expression<sup>27</sup>, can contribute to plaque progression by assuming a proinflammatory and cap-destabilizing phenotype *in vivo*. As discussed above, transition of SMCs to macrophage-like foam cells in a hyperlipidemic environment is associated with co-expression of a proinflammatory phenotype.<sup>16</sup> She et al reported that the conversion of SMCs to foam cells may be accelerated by expression of neural/glial antigen-2 (NG2) proteoglycan on the surface of synthetic SMCs.<sup>66</sup> Despite the fact that ablation of NG2 proteoglycan in mice leads to hyperlipidemia and obesity, both risk factors for the progression of atherosclerosis, when tested on an ApoE-null background loss of NG2 expression unexpectedly results in reduced plaque development.<sup>66</sup> These results suggest that local acquisition of a macrophage like

proinflammatory phenotype by plaque SMCs plays a more predominant role in plaque progression than systemic hyperlipidemia or obesity. Finally, an emphasis on the role of local factors in the control of macrophage-like phenotype transition of plaque SMCs emphasizes the role of SMC-derived ECM proteins in plaque progression. Schwanekamp et al examined the role of periostin, a secreted ECM protein, on the development of atherosclerosis in an ApoE<sup>-/-</sup> model.<sup>67</sup> These authors found that periostin-deficient, ApoE-null mice developed reduced plaque areas, fewer plaque macrophages, disorganized plaque ECM, thinner fibrotic caps, and increased MMP-2 and MMP-13 levels than ApoE-null control mice.<sup>67</sup> It would be interesting to determine if the reduced number of plaque macrophages observed in the absence of periostin was the result of reduced plaque SMC transition to a macrophage-like phenotype now that the necessary mouse genetic tools are available to test this possibility.

## Summary

The studies reviewed above suggest that the diverse origins of vascular SMCs in the embryo<sup>47</sup> is mirrored by the diverse fates that these cells can exhibit in response to injury or disease.<sup>13-16, 29-31, 68-70</sup> An understanding of these phenotype transitions will most likely ultimately lead back to control of the SMC CARG box by myocardin and SRF together with the constellation of corepressors and coactivators that have been described to interact with this SMC differentiation platform.<sup>32,71,72</sup> Whether these phenotype transitions to macrophage-like, or osteoblast-like cell types are reversible *in vivo*, and what role resident SMC progenitor cells play in the homeostasis of the artery wall<sup>73-77</sup> are among the important questions going forward.

## Sources of Funding

Research in the authors' laboratory is supported by the National Institutes of Health grants RO1HL123650, RO1HL121877, RO1HL133723, the Loie Power Robinson Stem Cell and Regenerative Medicine Fund, and the Seattle Children's Research Institute, Seattle, WA.

## References

1. Chamley-Campbell J, Campbell G, Ross R. The smooth muscle cell in culture. *Physiol Rev.* 1979; 59:1–61. pmid:108688. [PubMed: 108688]
2. Majesky MW, Dong XR, Regan JN, Høglund VJ. Vascular smooth muscle progenitor cells: building and repairing blood vessels. *Circ Res.* 2011; 108:365–377. doi:10.1161/CIRCRESAHA.110.223800. [PubMed: 21293008]
3. Gomez D, Swiatlowska P, Owens GK. Epigenetic control of smooth muscle cell identity and lineage memory. *Arterioscler Thromb Vasc Biol.* 2015; 35:2508–2516. doi:10.1161/ATVBAHA.115.305044. [PubMed: 26449751]
4. Wagenseil JE, Mecham RP. Vascular extracellular matrix and arterial mechanics. *Physiol Rev.* 2009; 89:957–989. doi:10.1152/physrev.00041.2008. [PubMed: 19584318]
5. Wang DZ, Chang P, Wang Z, Sutherland L, Richardson J, Small E, Krieg P, Olson E. Activation of cardiac gene expression by myocardin, a transcriptional cofactor for serum response factor. *Cell.* 2001; 105:851–862. doi: 10.1016/S0092-8674(01)00404-4. [PubMed: 11439182]
6. Cordes K, Sheehy N, White M, Berry E, Morton S, Muth A, Lee T, Miano J, Ivey K, Srivastava D. miR-143 and miR-145 regulate smooth muscle cell fate and plasticity. *Nature.* 2009; 460:705–710. doi:10.1038/nature08195. [PubMed: 19578358]

7. Albinsson S, Suarez Y, Skoura A, Offermanns S, Miano JM, Sessa WC. MicroRNAs are necessary for vascular smooth muscle growth, differentiation, and function. *Arterioscler Thromb Vasc Biol.* 2010; 30:1118–1126. doi:10.1161/ATVBAHA.109.200873. [PubMed: 20378849]
8. Pellegrini L, Tan S, Richmond TJ. Structure of serum response factor core bound to DNA. *Nature.* 1995; 376:490–498. doi:10.1038/376490a0. [PubMed: 7637780]
9. Gramzow L, Ritz MS, Theissen G. On the origin of MADS-domain transcription factors. *Trends in Genet.* 2010; 26:149–153. doi:10.1016/j.tig.2010.01.004. [PubMed: 20219261]
10. Han Z, Li X, Wu J, Olson EN. A myocardin-related transcription factor regulates activity of serum response factor in *Drosophila*. *Proc Natl Acad Sci USA.* 2004; 101:12567–12572. doi:10.1073/pnas.0405085101. [PubMed: 15314239]
11. Wang DZ, Olson EN. Control of smooth muscle development by the myocardin family of transcriptional coactivators. *Curr Opin Genet Dev.* 2004; 14:558–566. doi: 10.1016/j.gde.2004.08.003. [PubMed: 15380248]
12. Miano JM. Myocardin in biology and disease. *J Biomed Res.* 2015; 29:3–19. doi:10.7555/JBR.29.20140151. [PubMed: 25745471]
13. Rong JX, Shapiro M, Trogan E, Fisher EA. Transdifferentiation of mouse aortic smooth muscle cells to a macrophage-like state after cholesterol loading. *Proc Natl Acad Sci USA.* 2003; 100:13531–13526. doi: 10.1073/pnas.1735526100. [PubMed: 14581613]
14. Feil S, Fehrenbacher B, Lukowski R, Essmann F, Schulze-Osthoff K, Schaller M, Feil R. Transdifferentiation of vascular smooth muscle cells to macrophage-like cells during atherogenesis. *Circ Res.* 2014; 115:662–667. doi:10.1161/CIRCRESAHA.115.304634. [PubMed: 25070003]
15. Shankman LS, Gomez D, Cherepanova OA, Salmon M, Alencar GF, Haskins RM, Swiatlowska P, Newman AA, Greene ES, Straub AC, Isakson B, Randolph GJ, Owens GK. KLF4-dependent phenotypic modulation of smooth muscle cells has a key role in atherosclerotic plaque pathogenesis. *Nat Med.* 2015; 21:628–637. doi:10.1038/nm.3866. [PubMed: 25985364]
16. Vengrenyuk Y, Nishi H, Long X, Ouimet M, Savji N, Martinez FO, Cassella CP, Moore KJ, Ramsey SA, Miano JM, Fisher EA. Cholesterol loading reprograms the microRNA-143/145-myocardin axis to convert aortic smooth muscle cells to a dysfunctional macrophage-like phenotype. *Arterioscler Thromb Vasc Biol.* 2015; 35:535–546. doi:10.1161/ATVBAHA.114.304029. [PubMed: 25573853]
17. Huff MW, Pickering JG. Can a vascular smooth muscle-derived foam-cell really change its spots? *Arterioscler Thromb Vasc Biol.* 2015; 35:492–495. doi:10.1161/ATVBAHA.115.305225. [PubMed: 25717175]
18. Cooke PH, Smith SC. Smooth muscle cells: the source of foam cells in atherosclerotic white carneau pigeons. *Exp Mol Pathol.* 1968; 8:171–189. pmid:5647247. [PubMed: 5647247]
19. Bierman EL, Albers JJ. Lipoprotein uptake and degradation by cultured human arterial smooth muscle cells. *Adv Exp Med Biol.* 1976; 67:437–452. pmid: 179303. [PubMed: 179303]
20. Campbell JH, Popadyne L, Nestel PJ, Campbell GR. Lipid accumulation in arterial smooth muscle cells. Influence of phenotype. *Arteriosclerosis.* 1983; 47:279–283. pmid: 6882501.
21. Faggiotto A, Ross R, Harker L. Studies of hypercholesterolemia in the nonhuman primate. I. Changes that lead to fatty streak formation. *Arteriosclerosis.* 1984; 4:323–340. pmid: 6466191. [PubMed: 6466191]
22. Allahverdian S, Chehroudi AC, McManus BM, Abraham T, Francis GA. Contribution of intimal smooth muscle cells to cholesterol accumulation and macrophage-like cells in human atherosclerosis. *Circulation.* 2014; 129:1551–1559. Doi:10.1161/CIRCULATIONAHA.113.005015. [PubMed: 24481950]
23. Benditt EP, Benditt JM. Evidence for a monoclonal origin of human atherosclerotic plaques. *Proc Natl Acad Sci USA.* 1973; 70:1753–1756. pmid: 4515934. [PubMed: 4515934]
24. Fisher EA, Miano JM. Don't judge books by their covers: vascular smooth muscle cells in arterial pathologies. *Circulation.* 2014; 129:1545–1547. doi:10.1161/CIRCULATIONAHA.114.009075. [PubMed: 24733539]
25. Chellan B, Reardon CA, Getz GS, Hofmann Bowman MA. Enzymatically modified low-density lipoprotein promotes foam cell formation in smooth muscle cells via macropinocytosis and

- enhances receptor-mediated uptake of oxidized low-density lipoprotein. *Arterioscler Thromb Vasc Biol.* 2016; 36:1101–1113. doi:10.1161/ATVBAHA.116.307306. [PubMed: 27079883]
26. Pidkova MA, Cherepanova OA, Yoshida T, Alexander MR, Deaton RA, Thomas JA, Leitinger N, Owens GK. Oxidized phospholipids induce phenotypic switching of vascular smooth muscle cells in vivo and in vitro. *Circ Res.* 2007; 101:792–801. doi:10.1161/CIRCRESAHA.107.152736. [PubMed: 17704209]
27. Ackers-Johnson M, Talasila A, Sage AP, Long X, Bot I, Morrell NW, Bennett MR, Miano JM, Sinha S. Myocardin regulates vascular smooth muscle cell inflammatory activation and disease. *Arterioscler Thromb Vasc Biol.* 2015; 35:817–828. doi:10.1161/ATVBAHA.114.305218. [PubMed: 25614278]
28. Kang I, Barth JL, Sproul EP, Yoon DW, Workman GA, Braun KR, Argraves WS, Wight TN. Expression of V3 versican by rat arterial smooth muscle cells promotes differentiated and anti-inflammatory phenotypes. *J Biol Chem.* 2015; 290:21629–21641. doi:10.1074/jbc.M115.657486. [PubMed: 26152723]
29. Beazley KE, Nurminsky D, Lima F, Gandhi C, Nurminskaya MV. Wnt16 attenuates TGF $\beta$ -induced chondrogenic transformation in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 2015; 35:573–579. doi:10.1161/ATVBAHA.114.304393. [PubMed: 25614285]
30. Gao C, Fu Y, Li Y, Zhang X, Zhang L, Yu F, Xu SS, Xu Q, Zhu Y, Guan Y, Wang X, Kong W. Microsomal prostaglandin E synthase-1-derived PGE2 inhibits vascular smooth muscle cell calcification. *Arterioscler Thromb Vasc Biol.* 2016; 36:108–121. doi:10.1161/ATVBAHA.115.306642. [PubMed: 26543101]
31. Krohn JB, Hutcheson JD, Martinez-Martinez E, Irvin WS, Bouten CV, Bertazzo S, Bendeck MP, Aikawa E. Discoidin domain receptor-1 regulates calcific extracellular vesicle release in vascular smooth muscle cell fibrocalcific response via transforming growth factor- $\beta$  signaling. *Arterioscler Thromb Vasc Biol.* 2016; 36:525–533. doi: 10.1161/ATVBAHA.115.307009. [PubMed: 26800565]
32. Miano JM. Serum response factor: toggling between disparate programs of gene expression. *J Mol Cell Cardiol.* 2003; 35:577–593. pmid:12788374. [PubMed: 12788374]
33. Zhou J, Zhang M, Fang H, El-Mounayri O, Rodenberg JM, Imbalzano AN, Herring BP. The SWI/SNF chromatin remodeling complex regulates myocardin-induced smooth muscle-specific gene expression. *Arterioscler Thromb Vasc Biol.* 2009; 29:921–928. doi: 10.1161/ATVBAHA.109.187229. [PubMed: 19342595]
34. Lu J, Landerholm TE, Wei JS, Dong XR, Wu SP, Liu X, Nagata K, Inagaki M, Majesky MW. Coronary smooth muscle differentiation from proepicardial cells requires rhoA-mediated actin reorganization and p160 rho-kinase activity. *Dev Biol.* 2001; 240:404–418. doi:10.1006/dbio.2001.0403. [PubMed: 11784072]
35. Mack CP, Somlyo AV, Hautmann M, Somlyo AP, Owens GK. Smooth muscle differentiation marker gene expression is regulated by rhoA-mediated actin polymerization. *J Biol Chem.* 2001; 276:341–347. doi:10.1074/jbc.M005505200. [PubMed: 11035001]
36. Du KL, Chen M, Li J, Lepore JJ, Mericko P, Parmacek MS. Megakaryoblastic leukemia factor-1 transduces cytoskeletal signals and induces smooth muscle cell differentiation from undifferentiated embryonic stem cells. *J Biol Chem.* 2004; 279:17578–17586. doi:10.1074/jbc.M400961200. [PubMed: 14970199]
37. Staus DP, Weise-Cross L, Mangum KD, Medlin MD, Mangiante L, Taylor JM, Mack CP. Nuclear RhoA signaling regulates MRTF-dependent SMC-specific transcription. *Am J Physiol Heart Circ Physiol.* 2014; 307:H379–H390. doi: 10.1152/ajpheart.01002.2013. [PubMed: 24906914]
38. Copeland JW, Treisman R. The diaphanous-related forming mDia1 controls serum response factor activity through its effects on actin polymerization. *Mol Biol Cell.* 2002; 13:4088–4099. doi: 10.1091/mbc.02-06-0092. [PubMed: 12429848]
39. Staus DP, Baker AL, Medlin MD, Taylor JM, Mack CP. Formin homology domain-containing protein 1 regulates smooth muscle cell phenotypes. *Arterioscler Thromb Vasc Biol.* 2011; 31:360–367. doi:10.1161/ATVBAHA.110.21.212993. [PubMed: 21106951]
40. Weise-Cross L, Taylor JM, Mack CP. Inhibition of diaphanous formin signaling in vivo impairs cardiovascular development and alters smooth muscle cell phenotype. *Arterioscler Thromb Vasc Biol.* 2015; 35:2374–2383. doi: 10.1161/ATVBAHA.115.305879. [PubMed: 26381868]



41. Turczynska KM, Sward K, Hien TT, Wohlfahrt J, Mattisson IY, Ekman M, Nilsson J, Sjogren J, Murugesan V, Hultgardh-Nilsson A, Cidada P, Hellstrand P, Perez-Garcia MT, Albinsson S. Regulation of smooth muscle dystrophin and synaptopodin 2 expression by actin polymerization and vascular injury. *Arterioscler Thromb Vasc Biol.* 2015; 35:1489–1497. doi:10.1161/ATVBAHA.114.305065. [PubMed: 25857312]
42. Balint B, Yin H, Chakrabarti S, Chu MW, Sims SM, Pickering JG. Collectivization of vascular smooth muscle cells via TGF $\beta$ -cadherin-11-dependent adhesive switching. *Arterioscler Thromb Vasc Biol.* 2015; 35:1254–1264. doi: 10.1161/ATVBAHA.115.305310. [PubMed: 25767275]
43. Thiery JP. Cell adhesion in development: a complex signaling network. *Curr Opin Genet Dev.* 2003; 13:365–371. doi: 10.1016/S0959-437X(03)00088-1. [PubMed: 12888009]
44. Fernandez I, Martin-Garrido A, Zhou DW, Clempus RE, Seidel-Rogol B, Valdivia A, Lessegue B, Garcia AJ, Griendling KK, San Martin A. Hic-5 mediates TGF $\beta$ -induced adhesion in vascular smooth muscle cells by a Nox4-dependent mechanism. *Arterioscler Thromb Vasc Biol.* 2015; 35:1198–1206. doi:10.1161/ATVBAHA.114.305185. [PubMed: 25814672]
45. Massague J. TGF $\beta$  signaling in context. *Nat Rev Mol Cell Biol.* 2012; 13:616–630. doi:10.1038/nrm3434. [PubMed: 22992590]
46. Hu JH, Wei H, Jaffe M, Airhart N, Du L, Angelov SN, Yan J, Allen JK, Kang I, Wight TN, Fox K, Smith A, Enstrom R, Dichek DA. Postnatal deletion of the type II transforming growth factor- $\beta$  receptor in smooth muscle cells causes severe aortopathy in mice. *Arterioscler Thromb Vasc Biol.* 2015; 35:2647–2656. doi: 10.1161/ATVBAHA.115.306573. [PubMed: 26494233]
47. Majesky MW. Developmental basis of vascular smooth muscle diversity. *Arterioscler Thromb Vasc Biol.* 2007; 27:1248–1258. doi:10.1161/ATVBAHA.107.141069. [PubMed: 17379839]
48. Li W, Li Q, Jiao Y, Qin R, Zhou J, Ferruzzi J, Kim RW, Geirsson A, Dietz HC, Offermanns S, Humphrey JD, Tellides G. Tgf $\beta$ R2 disruption in postnatal smooth muscle impairs aortic wall homeostasis. *J Clin Invest.* 2014; 124:755–767. doi:10.1172/JCI69942. [PubMed: 24401272]
49. Chen X, Rateri DL, Howatt DA, Balakrishnan A, Moorleggen JJ, Cassis LA, Daugherty A. TGF $\beta$  neutralization enhances AngII-induced aortic rupture and aneurysm in both thoracic and abdominal regions. *PLoS One.* 2016; 11:e0153811. doi: 10.1371/journal.pone.0153811. [PubMed: 27104863]
50. Siu SC, Silversides CK. Bicuspid aortic valve disease. *J Am Coll Cardiol.* 2010; 55:2789–2800. doi:10.1016/j.jacc.2009.12.068. [PubMed: 20579534]
51. Paloschi V, Gadin JR, Khan S, Bjorck HM, Du L, Maleki S, Roy J, Lindeman JH, Mohamed SA, Tsuda T, Franco-Cereceda A, Eriksson P. Aneurysm development in patients with a bicuspid aortic valve is not associated with transforming growth factor- $\beta$  activation. *Arterioscler Thromb Vasc Biol.* 2015; 35:973–980. doi:10.1161/ATVBAHA.114.304996. [PubMed: 25745062]
52. Gallo EM, Loch DC, Habashi JP, Calderon JF, Chen Y, Bedja D, van Erp C, Gerber EE, Parker SJ, Sauls K, Judge DP, Cooke SK, Lindsay ME, Rouf R, Myers L, Rhys CM, Kent KC, Norris RA, Huso DL, Dietz HC. Angiotensin II-dependent TGF $\beta$  signaling contributes to Loeys-Dietz syndrome vascular pathogenesis. *J Clin Invest.* 2014; 124:448–460. doi:10.1172/JCI69666. [PubMed: 24355923]
53. Crosas-Molist E, Meirelles T, Lopez-Luque J, Serra-Peinado C, Selva J, Caja L, Gorbenko Del Blanco D, Uriate JJ, Bertran E, Mendizabal Y, Hernandez V, Garcia-Calero C, Busnadiago O, Condom E, Toral D, Castella M, Forteza A, Navajas D, Sarri E, Rodriguez-Pascual F, Dietz HC, Fabregat I, Egea G. Vascular smooth muscle cell phenotypic changes in patients with Marfan syndrome. *Arterioscler Thromb Vasc Biol.* 2015; 35:960–972. doi:10.1161/ATVBAHA.114.304412. [PubMed: 25593132]
54. Wang D, Deuse T, Stubbendorff M, Chernogubova E, Erben RG, Eken SM, Jin H, Li Y, Busch A, Heeger CH, Behnisch B, Reichenspurner H, Robbins RC, Spin JM, Tsao PS, Schrepfer S, Maegdefessel L. Local microRNA modulation using a novel anti-miR-21-eluting stent effectively prevents experimental in-stent restenosis. *Arterioscler Thromb Vasc Biol.* 2015; 35:1945–1953. doi:10.1161/ATVBAHA.115.305597. [PubMed: 26183619]
55. Thum T, Gross C, Fiedler J, Fischer T, Kissler S, Bussen M, Galuppo P, Just S, Rottbauer W, Frantz S, Castoldi M, Soutschek J, Koteliansky V, Rosenwald A, Basson MA, Licht JD, Pena JT, Rouhanifard SH, Muckenthaler MU, Tuschl T, Martin GR, Bauersachs J, Engelhardt S.

- MicroRNA-21 contributes to myocardial disease by stimulating MAP kinase signaling in fibroblasts. *Nature*. 2008; 456:980–984. doi:10.1038/nature07511. [PubMed: 19043405]
56. Hartmann D, Thum T. MicroRNAs and vascular (dys)function. *Vascul Pharmacol*. 2011; 55:92–105. doi: 10.1016/j.vph.2011.07.005. [PubMed: 21802526]
57. Xu F, Ahmed AS, Kang X, Hu G, Liu F, Zhang W, Zhou J. MicroRNA-15b/16 attenuates vascular neointima formation by promoting the contractile phenotype through targeting YAP. *Arterioscler Thromb Vasc Biol*. 2015; 35:2145–2152. doi: 10.1161/ATVBAHA.115.305748. [PubMed: 26293467]
58. Liu F, Wang X, Hu G, Wang Y, Zhou J. The transcription factor TEAD1 represses smooth muscle-specific gene expression by abolishing myocardin function. *J Biol Chem*. 2013; 289:3308–3316. doi: 10.1074/jbc.M113.515817. [PubMed: 24344135]
59. Zeng L, Li Y, Yang J, Wang G, Margariti A, Xiao Q, Zampetaki A, Yin X, Mayr M, Mori K, Wang W, Hu Y, Xu Q. XBP1-deficiency abrogates neointimal lesion of injured vessels via cross talk with the PDGF signaling. *Arterioscler Thromb Vasc Biol*. 2015; 35:2134–2144. doi:10.1161/ATVBAHA.115.305420. [PubMed: 26315405]
60. Williams H, Mill CA, Monk BA, Hulin-Curtis S, Johnson JL, George SJ. Wnt2 and WISP-1/CCN4 induce intimal thickening via promotion of smooth muscle cell migration. *Arterioscler Thromb Vasc Biol*. 2016; 36:1416–1424. doi: 10.1161/ATVBAHA.116.307626.
61. Wight TN. Arterial remodeling in vascular disease: a key role for hyaluronan and versican. *Front Biosci*. 2008; 13:4933–4937. pmid: 18508558. [PubMed: 18508558]
62. Kiene LS, Homann S, Suvorava T, Rabausch B, Muller J, Kojda G, Kretschmer I, Twarock S, Dai G, Deenen R, Hartwig S, Lehr S, Kohrer K, Savani RC, Grandoch M, Fischer JW. Deletion of hyaluronan synthase 3 inhibits neointimal hyperplasia in mice. *Arterioscler Thromb Vasc Biol*. 2016; 36:e9–16. doi: 10.1161/ATVBAHA.115.306607. [PubMed: 26586662]
63. Libby P, Tabas I, Fredman G, Fisher EA. Inflammation and its resolution as determinants of acute coronary syndromes. *Circ Res*. 2014; 114:1867–1879. doi: 10.1161/CIRCRESAHA.114.302699. [PubMed: 24902971]
64. Hansson GK, Libby P, Tabas I. Inflammation and plaque vulnerability. *J Intern Med*. 2015; 278:483–493. doi:10.1111/joim.12406. [PubMed: 26260307]
65. Gardner SE, Humphry M, Bennett MR, Clarke MC. Senescent vascular smooth muscle cells drive inflammation through an interleukin-1 $\alpha$ -dependent senescence-associated secretory phenotype. *Arterioscler Thromb Vasc Biol*. 2015; 35:1963–1974. doi: 10.1161/ATVBAHA.115.305896. [PubMed: 26139463]
66. She ZG, Chang Y, Pang HB, Han W, Chen HZ, Smith JW, Stallcup WB. NG2 proteoglycan ablation reduces foam cell formation and atherogenesis via decreased low-density lipoprotein retention by synthetic smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2016; 36:49–59. doi: 10.1161/ATVBAHA.115.306074. [PubMed: 26543095]
67. Schwanekamp JA, Lorts A, Vagnozzi RJ, Vanhoutte D, Molkentin JD. Deletion of periostin protects against atherosclerosis in mice by altering inflammation and extracellular matrix remodeling. *Arterioscler Thromb Vasc Biol*. 2016; 36:60–68. doi: 10.1161/ATVBAHA.115.306397. [PubMed: 26564821]
68. Briot A, Jaroszewicz A, Warren CM, Lu J, Touma M, Rudat C, Hofmann JJ, Airik R, Weinmaster G, Lyons K, Wang Y, Kispert A, Pellegrini M, Iruela-Arispe ML. Repression of Sox9 by Jag1 is continuously required to suppress the default chondrogenic fate of vascular smooth muscle cells. *Dev Cell*. 2014; 31:707–721. doi: 10.1016/j.devcel.2014.11.023. [PubMed: 25535917]
69. Speer MY, Yang HY, Brabb T, Leaf E, Look A, Lin WL, Frutkin A, Dichek DA, Giachelli CM. Smooth muscle cells give rise to osteochondrogenic precursors and chondrocytes in calcifying arteries. *Circ Res*. 2009; 104:733–741. doi: 10.1161/CIRCRESAHA.108.183053. [PubMed: 19197075]
70. Deng L, Huang L, Sun Y, Heath JM, Wu H, Chen Y. Inhibition of FOXO1/3 promotes vascular calcification. *Arterioscler Thromb Vasc Biol*. 2015; 35:175–183. doi: 10.1161/ATVBAHA.114.304786. [PubMed: 25378413]
71. Horita H, Wysoczynski CL, Walker LA, Moulton KS, Li M, Ostriker A, Tucker R, McKinsey TA, Churchill ME, Nemenoff RA, Weiser-Evans MC. Nuclear PTEN functions as an essential

- regulator of SRF-dependent transcription to control smooth muscle differentiation. *Nat Commun.* 2016; 7:10830. doi: 10.1038/ncomms10830. [PubMed: 26940659]
72. Shi N, Chen SY. Smooth muscle cell differentiation: model systems, regulatory mechanisms, and vascular diseases. *J Cell Physiol.* 2016; 231:777–787. doi:10.1002/jcp.25208. [PubMed: 26425843]
73. Hu Y, Zhang Z, Torsney E, Afzal AR, Davison F, Metzler B, Xu Q. Abundant progenitor cells in the adventitia contribute to atherosclerosis of vein grafts in ApoE-deficient mice. *J Clin Invest.* 2004; 113:1258–1265. doi: 10.1172/JCI19628. [PubMed: 15124016]
74. Passman JN, Dong XR, Wu SP, Maguire CT, Hogan KA, Bautch VL, Majesky MW. A sonic hedgehog signaling domain in the arterial adventitia supports resident Sca1<sup>+</sup> smooth muscle progenitor cells. *Proc Natl Acad Sci USA.* 2008; 105:9349–9354. doi: 10.1073/pnas.0711382105. [PubMed: 18591670]
75. Majesky MW, Dong XR, Høglund VJ, Mahoney WM Jr, Daum G. The adventitia: a dynamic interface containing resident progenitor cells. *Arterioscler Thromb Vasc Biol.* 2011; 31:1530–1539. Doi:10.1161/ATVBAHA.110.221549. [PubMed: 21677296]
76. Shikatani EA, Chandy M, Besla R, Li CC, Momen A, El-Mounayri O, Robbins CS, Husain M. c-Myb regulates proliferation and differentiation of adventitial Sca1<sup>+</sup> vascular smooth muscle cell progenitors by transactivation of myocardin. *Arterioscler Thromb Vasc Biol.* 2016; 36:1367–1376. doi:10.1161/ATVBAHA.115.307116. [PubMed: 27174098]
77. Sun Y, Wang K, Ye P, Wu J, Ren L, Zhang A, Huang X, Deng P, Wu C, Yue Z, Chen Z, Ding X, Chen J, Xia J. MicroRNA-155 promotes the directional migration of resident smooth muscle progenitor cells by regulating monocyte chemoattractant protein 1 in transplant arteriosclerosis. *Arterioscler Thromb Vasc Biol.* 2016; 36:1230–1239. doi:10.1161/ATVBAHA.115.306691. [PubMed: 27079875]