

# Can Creutzfeldt-Jakob disease unravel the mysteries of Alzheimer?

Gabor G. Kovacs

Institute of Neurology, Medical University of Vienna, Vienna, Austria

**ABSTRACT.** Recent studies on iatrogenic Creutzfeldt-Jakob disease (CJD) raised concerns that one of the hallmark lesions of Alzheimer disease (AD), amyloid- $\beta$  ( $A\beta$ ), may be transmitted from human-to-human. The neuropathology of AD-related lesions is complex. Therefore, many aspects need to be considered in deciding on this issue. Observations of recent studies can be summarized as follows: 1) The frequency of iatrogenic CJD cases with parenchymal and vascular  $A\beta$  deposits is statistically higher than expected; 2) The morphology and distribution of  $A\beta$  deposition may show distinct features; 3) The pituitary and the dura mater themselves may serve as potential sources of  $A\beta$  seeds; 4) Cadaveric dura mater from 2 examined cases shows  $A\beta$  deposition; and 5) There is a lack of evidence that the clinical phenotype of AD appears following the application of cadaveric pituitary hormone or dura mater transplantation. These studies support the notion that neurodegenerative diseases have common features regarding propagation of disease-associated proteins as seeds. However, until further evidence emerges, prions of transmissible spongiform encephalopathies are the only neurodegenerative disease-related proteins proven to propagate clinicopathological phenotypes.

**KEYWORDS.** Alzheimer disease, Amyloid- $\beta$ , dura mater, iatrogenic Creutzfeldt-Jakob disease, pituitary, prion

## INTRODUCTION

Creutzfeldt-Jakob disease (CJD) is a human prion disease characterized by progressive loss of neurons together with spongiform change of the neuropil and deposition of

disease-associated prion protein ( $PrP^{Sc}$ ), the abnormal conformer of the physiological cellular prion protein. Although very rare, CJD has been in the center of scientific interest due to the fact that it is transmissible even between humans. However, the majority of

---

Correspondence to: Gabor G. Kovacs, MD, PhD; Medical University of Vienna, Institute of Neurology, AKH 4J, Währinger Gürtel 18-20, A-1090 Vienna, Austria; Email: gabor.kovacs@meduniwien.ac.at

Received August 2, 2016; Accepted August 20, 2016.

© 2016 Gabor G. Kovacs.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.

Extra View to: Kovacs GG, Lutz MI, Ricken G, Ströbel T, Höftberger R, Preusser M, Regelsberger G, Hönigschnabl S, Reiner A, Fischer P, et al. Dura mater is a potential source of  $A\beta$  seeds. *Acta Neuropathol* 2016; 131(6):911–923; <http://dx.doi.org/10.1007/s00401-016-1565-x>

CJD cases are sporadic. Less frequently, mutations in the prion protein gene (*PRNP*) cause prion disease. Very rarely medical interventions, such as application of cadaveric pituitary hormones or transplantation of cadaveric dura mater, lead to the development of iatrogenic CJD.<sup>1</sup>

Other neurodegenerative diseases, such as Alzheimer disease (AD) and Parkinson disease, are more frequent than CJD. Like PrP, most of the neurodegenerative conditions are characterized also by the deposition of physicochemically modified disease-related proteins, such as amyloid- $\beta$  ( $A\beta$ ),  $\alpha$ -synuclein, tau or TAR-DNA-binding protein 43 kDa (TDP-43).<sup>2</sup> The concept of protein-based classification of neurodegenerative diseases is widely accepted.<sup>2</sup> It seems that misfolding of endogenous proteins can lead to proteinaceous seeds, which can serve as self-propagating agents associated with the progression of the specific disease.<sup>3</sup> Experimental models and studies on humans revealed that cell-to-cell transmission of disease-associated proteins may lead to sequential dissemination of pathological protein aggregates<sup>4</sup> reminiscent of that reported by Fraser in 1982 for the scrapie agent in the retino-tectal pathway.<sup>5</sup>

$A\beta$  and tau deposition is characteristic for AD but can be found in a wide range of disorders; furthermore, genetic predisposing factors, such as the  $\epsilon 4$  allele of Apolipoprotein E (*APOE*), influence disease risk. The presence of  $A\beta$  and tau pathology is not unusual in sporadic and genetic CJD brains.<sup>6-8</sup> Accordingly, is the increased attention on reports on the detection of  $A\beta$  deposition in iatrogenic CJD warranted?

### *Neuropathology of Alzheimer Disease*

AD is characterized by the intraneuronal deposition of pathological tau in the form of neurofibrillary tangles (NFTs) and neuropil threads and extracellular accumulation of  $A\beta$  in the form of  $A\beta$  plaques.<sup>9</sup> Both are needed to be detected simultaneously to allow definite diagnosis of AD.<sup>10</sup> Neuronal tau pathology can be detected from early adulthood, particularly

in the brainstem.<sup>11</sup> Based on imaging and cerebrospinal fluid biomarkers of  $A\beta$ , the concept of preclinical AD has been proposed.<sup>12</sup> Indeed,  $A\beta$  plaques can be detected in cognitively normal individuals from 30 y of age,<sup>11</sup> particularly associated with the  $\epsilon 4$  allele of the *APOE* gene.<sup>13</sup> In young patients with widespread  $A\beta$  pathology, mutations in the amyloid precursor protein gene (*APP*) and presenilin (*PSEN*) 1 and 2 must be also excluded.<sup>14</sup> Furthermore,  $A\beta$  can deposit in the cerebral blood vessels in the form of cerebral amyloid angiopathy (CAA), which can associate with AD but can also be seen independent of AD.<sup>15</sup>

Development of protein depositions composed of pathological tau and  $A\beta$  has 2 aspects. First, the deposit itself undergoes "maturation;" second, the anatomical distribution shows sequential involvement of different regions. The intracellular development of tau pathology starts from fine granular deposition of hyperphosphorylated tau in the form of pretangles, which become ubiquitinated and argyrophilic when showing the classic morphology of NFTs.<sup>16</sup> Extracellular  $A\beta$  deposits also show different morphologies.<sup>9</sup> Fine granular perineuronal and diffuse deposits are the earliest lesions. These are not detectable with thioflavin S and Congo red staining. Furthermore, focal deposits can be distinguished, which may be associated or not to a neuritic corona. A subset of focal deposits may be detected only by immunohistochemistry for  $A\beta$  while further ones are additionally congophilic and thioflavine S-positive. Dystrophic neurites are important components of a subset of plaques. Although rare tau positive dots and thin neurites appear already in some diffuse deposits, degenerating neurites are visible in the corona of focal deposits called mature senile plaques.<sup>17</sup> These plaques can be grouped based on the presence or lack of chromogranin A, synaptic markers, amyloid precursor protein, ubiquitin, and tau immunoreactivity in the dystrophic component.<sup>9,18-20</sup> Tau-positive dystrophic neurites mostly appear to be associated with plaques only in so far as NFTs appear in those regions where the plaque forms.<sup>20</sup> The whole spectrum of these tau and  $A\beta$  immunoreactivities are seen in typical AD cases.

The anatomical distribution of tau and A $\beta$  deposits show distinct patterns in early stages (tau) and phases (A $\beta$ ). Later these pathologies converge and affect the same anatomical regions. Neuronal tau pathology first involves specific brainstem nuclei and later proceed to involve limbic and neocortical areas.<sup>11,21</sup> A $\beta$  plaques appear first in neocortical areas and later the hippocampus, subcortical nuclei, brainstem and cerebellum become sequentially involved.<sup>22</sup>

Based on these observations, it is difficult to determine whether human-to-human transmission of AD-related pathology occurred in association with the iatrogenic transmission of CJD. Thus, in addition to employing statistical approaches on the frequency of AD-related pathology, the cadaveric tissue must be examined, distinctive morphological features of brain pathology must be sought and genetic or other predisposing factors have to be excluded, to allow for a more definitive conclusion.

### *A $\beta$ and Tau in Sporadic and Genetic CJD*

Tau and A $\beta$  deposition is not rare in sporadic and genetic CJD. One aspect needs to be emphasized: phospho-tau-positive small neuritic dots (1–3  $\mu$ m; smaller than the tau-positive dystrophic neurites around A $\beta$  plaques in AD) are seen in nearly all cases of sporadic and genetic CJD.<sup>6,23</sup> These might be triggered by PrP itself,<sup>23</sup> or simply reflect the degeneration of neurons with consequently damaged neurites.<sup>6</sup> In the following sections on CJD, the term tau pathology will be used to refer to all morphological types of neuronal or glial tau immunoreactivities except these tau-positive neuritic dots.

Tau-positive NFTs and A $\beta$  plaques in CJD is indicative of concomitant AD; the frequency is not high.<sup>6,7</sup> The morphology of A $\beta$  plaques is similar to AD or that seen in cognitively normal individuals.<sup>9,13, 23</sup> The anatomical involvement follows the stages and phases seen in AD<sup>21,22</sup> irrespective of the distribution of PrP deposits. This argues against cross-seeding of disease-associated PrP and A $\beta$  in human brains. CAA is not seen frequently in sporadic CJD. In certain forms of genetic CJD (e.g., E200K mutation), even in younger patients (<55 years), A $\beta$  deposition similar to that seen in AD has been reported.<sup>8,24</sup> Importantly, neuronal and glial tau pathology without A $\beta$  deposition is more frequent in sporadic CJD than A $\beta$  deposition alone or with NFTs.<sup>6</sup> In addition, widespread tau pathologies (neuronal and glial) have been reported in genetic but also in sporadic CJD cases.<sup>6,8</sup>

### *A $\beta$ and Tau in Iatrogenic CJD*

There are 4 published studies reporting the examination of AD-related pathologies in iatrogenic CJD cases (summarized in Table 1). The first study reported an increased frequency of parenchymal and vascular A $\beta$  in relatively young iatrogenic CJD cases treated with cadaveric growth hormone (GH) as compared to genetic and sporadic CJD cases.<sup>25</sup> This study excluded possible genetic predisposing factors and emphasized the lack of tau positive NFTs. Although the study demonstrated a higher load of parenchymal A $\beta$  deposition and CAA in iatrogenic CJD, it did not comment on potentially distinctive morphological features of A $\beta$  pathology in the iatrogenic CJD cases. Finally, the study did not have the opportunity to examine cadaveric samples for the presence of A $\beta$ ; however, it did confirm previous observations that the pituitary itself can exhibit A $\beta$  deposits.<sup>25,26</sup>

The second study focused on iatrogenic CJD cases following dura mater transplantation.<sup>27</sup> It compared the frequency of A $\beta$  deposition in iatrogenic CJD and sporadic CJD and reported that CAA and brain parenchymal A $\beta$  plaques are more frequent in iatrogenic than sporadic CJD. No conspicuous tau pathology could be observed in iatrogenic CJD cases.<sup>27</sup> This study provided further evidence that A $\beta$  deposits can be potentially transmitted from human-to-human.

However, there were still missing pieces to the puzzle. Indeed, there was a lack of literature data whether A $\beta$  can deposit in the dura mater at all; hence, to be able to act as a

TABLE 1. Comparison of studies on iatrogenic Creutzfeldt-Jakob disease (iCJD) and A $\beta$  deposition. AD: Alzheimer disease; GH: growth hormone; NA: not addressed in the study; Nr: number. \* Indicates that these cases were included in Study 2 also.

Study	Study 1	Study 2	Study 3	Study 4
Type of iCJD	Pituitary-derived GH	Dura mater	Dura mater	Dura mater
Number of iCJD cases examined	8	7	2*	16
Nr. of iCJD with parenchymal A $\beta$ deposits	4 + 2 focal	5	2*	13
Distinct morphology or distribution of parenchymal A $\beta$	NA	NA	Yes	Subpial accumulation emphasized. Morphology of parenchymal plaques not commented.
Nr. of iCJD with CAA	3 + 1 focal	5	2*	11
Cadaveric tissue examined	No	No	Yes	No
Non-cadaveric tissue examined	Yes: pituitary (55 samples)	No	Yes: dura mater (84 samples)	No
AD related tau pathology	No	No	No	Yes - details not provided
Genetic aspects	APOE + AD genes	NA	APOE + AD genes	APOE
Statistical difference when compared to sCJD	Yes	Yes	NA	Yes
Age of iCJD cases with parenchymal A $\beta$	4th decade to 51	28–63	28, 33	35–81
Age of iCJD cases with CAA	5th decade to 51	28–63	28, 33	35–81
Age of cases < 40 y with parenchymal A $\beta$ deposits	36 (focal)	28, 33	28, 33	35, 39
Age of cases < 40 y with CAA	–	28, 33	28, 33	35, 39
Clinical phenotype	CJD	CJD	CJD	CJD
Reference	25	27	28	29

source of transmissible protein seeds. Moreover, it was still not known whether 1) the cadaveric tissue used in these iatrogenic CJD cases exhibit A $\beta$  depositions, and 2) there are any distinctive features of brain pathology in these cases. To answer these questions, we performed an additional study on 2 cases.<sup>28</sup> We used 4 anti-A $\beta$  antibodies (clones 6F/3D, 4G8, anti-A $\beta$ <sub>1–40</sub>, and anti-A $\beta$ <sub>1–42</sub>) to provide unequivocal results and not misinterpret non-specific stainings. Our observations on 84 dura mater samples obtained in a longitudinal community-based aging study provided evidence for the first time that the dura mater may harbor A $\beta$  deposits in the form of CAA or amorphous aggregates. Furthermore, this study is the only one, which compared cadaveric and host dura mater samples and demonstrated the presence of A $\beta$  only in the cadaveric dura mater sample in addition to the brain tissue of the host.<sup>28</sup> Importantly, these cases are the

youngest reported with iatrogenic CJD and A $\beta$  deposition, and we excluded genetic predisposing factors (*APOE*, *APPP*, *PSEN 1*, 2). Meticulous analysis of A $\beta$  morphologies suggested peculiar features. First of all, in contrast to disease-associated PrP, which was found widespread in the brain, A $\beta$  deposits were accentuated close to the operation site. In these cases the subpial cortical area near to the traumatic lesion and the surgical intervention showed tissue damage. Strikingly, the morphological spectrum of A $\beta$  depositions in these 2 cases was much more limited and different as compared to that seen in AD, other forms of CJD, traumatic brain injury, and cognitively normal individuals.<sup>28</sup> Instead of diffuse plaques, we observed focal deposits and congophilic cored plaques with and without a neuritic corona. Dystrophic neurites around the plaques accumulated ubiquitin but not hyperphosphorylated tau. We observed clusters

away from the lesion site or columnar alignment of A $\beta$  plaques close to the lesion.<sup>28</sup> We also emphasized that NFTs were completely lacking even in subcortical areas where it may be seen in young individuals.<sup>11</sup> However, we noted the tau immunoreactive small neuritic profiles as seen in all CJD forms.<sup>28</sup>

A fourth study has been recently reported from Japan.<sup>29</sup> They did not find a higher frequency in iatrogenic CJD as compared to sporadic CJD; however, their study included elderly patients (up to 81 years). The youngest iatrogenic CJD cases did show more A $\beta$  than the youngest sporadic CJD cases. The study demonstrated significant association of cadaveric dura mater grafting with subpial A $\beta$  deposition and meningeal amyloid angiopathy using the anti-A $\beta$  antibody 4G8.<sup>29</sup> They discussed the possibility that A $\beta$  pathology propagated from the superficial portions of the brain in the patients with iatrogenic CJD.<sup>29</sup> The immunohistochemical studies were performed only on sections of the frontal, parietal, temporal, and occipital lobes and it was not clear how the A $\beta$  deposits related to the site of the surgical intervention. This study did not provide details on distinctive morphological features of parenchymal A $\beta$  deposits. However, the cohort included 11 cases (from 16 examined) above the age of 60 and some carrying  $\epsilon$ 4 alleles of the *APOE* gene, which could potentially limit any interpretation of this aspect. It is also not surprising that the study documented NFTs following Braak stages,<sup>29</sup> which might also be compatible with findings in the general population of this age group.<sup>11</sup>

In summary, not all aspects needed for exact comparisons have been presented in these studies (Table 1). The question why not all iatrogenic CJD cases show A $\beta$  deposition raises the possibility that cadaver samples with or without A $\beta$  had been used or there are yet unidentified susceptibility/protective factors. Frontzek et al. suggested that the longer latency might be associated with A $\beta$  deposition,<sup>27</sup> which seems to be supported also by the observation that the incubation period between dura mater grafting and death significantly correlated with the degree of subpial A $\beta$  deposition and CAA.<sup>29</sup> The confirmation of A $\beta$  in cadaveric dura mater could

be complemented by studies on the samples of cadaveric pituitary (see also Irwin et al.<sup>26</sup>), or the derived GH used for therapy in iatrogenic CJD cases. Furthermore, potential distinguishing morphological features of human cadaveric GH-related iatrogenic CJD merits further studies. Finally, the exact mechanism by which the external A $\beta$  deposits propagate in the brain tissue and in particular how they induce CAA in humans could not be addressed.

In spite these missing pieces we can conclude, however, that: 1) The frequency of cases with parenchymal and vascular A $\beta$  deposits is high in iatrogenic CJD with cadaveric pituitary-derived GH and cadaveric dura mater transplantation (statistical level), even after excluding an effect of genetic predisposing factors; 2) The morphology and distribution of A $\beta$  deposition shows distinct features at least in the 2 cases where this was commented on in details; 3) The pituitary and the dura mater themselves may harbor A $\beta$  deposits and thus may serve as potential sources of A $\beta$  seeds; 4) The cadaveric dura mater shows A $\beta$  deposition at least in the 2 cases where this was examined; and finally that 5) None of the studies reported iatrogenic CJD cases with the typical AD-type clinical phenotype.

## CONCLUSIONS

When talking about an infectious nature of a protein different levels can be defined (see recent reviews on A $\beta$ ).<sup>3,17,30</sup> Experimental observations suggest that A $\beta$  may propagate (as suggested to be called as 'propagon')<sup>17</sup> its conformation on a molecular level and has the ability to spread within and, in a limited manner, across tissues.<sup>3,17,30</sup> The observations on human iatrogenic CJD with A $\beta$  deposition suggest that A $\beta$ , in addition to prions of CJD, may act as an infectious seed between individuals. However, there are still differences between prions and A $\beta$  seeds. The complete clinicopathological phenotype of AD has not been proven to be transmitted by A $\beta$ , in contrast to prions in CJD. Thus only prions of CJD are 'phenotype propagons'.<sup>28</sup> Although it could be argued that perhaps A $\beta$  could be also classified

as such if more time given. Another explanation could be that classical AD requires an altered microenvironment, including the presence of tau appearing first at a distant location from A $\beta$ . In the rare situation when an A $\beta$  seed is placed in the brain (i.e. “external” source), the microenvironment might not be conducive for the development of the classical AD phenotype. Again, in the 2 young cases reported by our group,<sup>28</sup> even the brainstem nuclei, which are thought to be the earliest to develop neuronal tau pathology during AD,<sup>11</sup> did not show any tau immunoreactivity.<sup>28</sup> Experimental studies have shown that injection of A $\beta$ 42 fibrils into the brains of P301L mutant tau transgenic mice caused prominent increases in the numbers of NFTs.<sup>31</sup> On one hand this can be interpreted that A $\beta$  is needed to accelerate tau pathology, but on the other hand this could mean also that the microenvironment (mutant tau transgenic mice) was favorable for A $\beta$  to achieve this.

It is conceivable that the term ‘prion-like’ could be used to refer to some of the pathogenic aspects resembling prion diseases where it was first described. However, in addition to the studies reviewed here, further observations on humans could not provide unequivocal evidence that the clinical phenotype of AD is transmissible between humans through blood transfusion or cadaver human growth hormone.<sup>26,32</sup> Therefore, until further evidence emerges on the transmissibility of clinicopathological phenotypes between humans, the term ‘prion disease’ is still most compatible with disorders described also as transmissible spongiform encephalopathies characterized by the deposition of the abnormal form of PrP.

#### **DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST**

No potential conflicts of interest were disclosed.

#### **ACKNOWLEDGMENTS**

I am grateful to Drs. Linda K. Kwong and Kurt R. Brunden, Center for Neurodegenerative

Disease Research, University of Pennsylvania, Philadelphia, PA, USA, for the helpful discussions.

#### **REFERENCES**

- [1] Brown P, Brandel JP, Sato T, Nakamura Y, MacKenzie J, Will RG, Ladogana A, Pocchiari M, Leschek EW, Schonberger LB. Iatrogenic Creutzfeldt-Jakob disease, final assessment. *Emerg Infect Dis* 2012; 18:901-7; PMID:22607808; <http://dx.doi.org/10.3201/eid1806.120116>
- [2] Kovacs GG. molecular pathological classification of neurodegenerative diseases: turning towards precision medicine. *Int J Mol Sci* 2016; 17:189; <http://dx.doi.org/10.3390/ijms17020189>
- [3] Jucker M, Walker LC. Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* 2013; 501:45-51; PMID:24005412; <http://dx.doi.org/10.1038/nature12481>
- [4] Brettschneider J, Del Tredici K, Lee VM, Trojanowski JQ. Spreading of pathology in neurodegenerative diseases: a focus on human studies. *Nat Rev Neurosci* 2015; 16:109-20; PMID:25588378; <http://dx.doi.org/10.1038/nrn3887>
- [5] Fraser H. Neuronal spread of scrapie agent and targeting of lesions within the retino-tectal pathway. *Nature* 1982; 295:149-50; PMID:6173756; <http://dx.doi.org/10.1038/295149a0>
- [6] Kovacs GG, Rahimi J, Strobel T, Lutz MI, Regelsberger G, Streichenberger N, Perret-Liaudet A, Hoftberger R, Liberski PP, Budka H, et al. Tau pathology in Creutzfeldt-Jakob disease revisited. *Brain Pathol* 2016; PMID:27377321
- [7] Tousseyn T, Bajsarowicz K, Sanchez H, Gheyara A, Oehler A, Geschwind M, DeArmond B, DeArmond SJ. Prion Disease Induces Alzheimer Disease-Like Neuropathologic Changes. *J Neuropathol Exp Neurol* 2015; 74:873-88; PMID:26226132; <http://dx.doi.org/10.1097/NEN.0000000000000228>
- [8] Kovacs GG, Seguin J, Quadrio I, Hoftberger R, Kapas I, Streichenberger N, Biacabe AG, Meyronet D, Sciot R, Vandenberghe R, et al. Genetic Creutzfeldt-Jakob disease associated with the E200K mutation: characterization of a complex proteinopathy. *Acta Neuropathol* 2011; 121:39-57; PMID:20593190; <http://dx.doi.org/10.1007/s00401-010-0713-y>
- [9] Duyckaerts C, Delatour B, Potier MC. Classification and basic pathology of Alzheimer disease. *Acta Neuropathol* 2009; 118:5-36; PMID:19381658; <http://dx.doi.org/10.1007/s00401-009-0532-1>
- [10] Montine TJ, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, Mirra SS, et al. National Institute

- on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: a practical approach. *Acta Neuropathol* 2012; 123:1-11; PMID:22101365; <http://dx.doi.org/10.1007/s00401-011-0910-3>
- [11] Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. *J Neuropathol Exp Neurol* 2011; 70:960-9; PMID:22002422; <http://dx.doi.org/10.1097/NEN.0b013e318232a379>
- [12] Dubois B, Hampel H, Feldman HH, Scheltens P, Aisen P, Andrieu S, Bakardjian H, Benali H, Bertram L, Blennow K, et al. Preclinical Alzheimer's disease: Definition, natural history, and diagnostic criteria. *Alzheimers Dement* 2016; 12:292-323; PMID:27012484; <http://dx.doi.org/10.1016/j.jalz.2016.02.002>
- [13] Pletnikova O, Rudow GL, Hyde TM, Kleinman JE, Ali SZ, Bharadwaj R, Gangadeen S, Crain BJ, Fowler DR, Rubio AI, et al. Alzheimer Lesions in the Autopsied Brains of People 30 to 50 Years of Age. *Cogn Behav Neurol* 2015; 28:144-52; PMID:26413742; <http://dx.doi.org/10.1097/WNN.0000000000000071>
- [14] Cacace R, Slegers K, Van Broeckhoven C. Molecular genetics of early-onset Alzheimer's disease revisited. *Alzheimers Dement* 2016; 12:733-48; PMID:27016693; <http://dx.doi.org/10.1016/j.jalz.2016.01.012>
- [15] Thal DR, Griffin WS, de Vos RA, Ghebremedhin E. Cerebral amyloid angiopathy and its relationship to Alzheimer's disease. *Acta Neuropathol* 2008; 115:599-609; PMID:18369648; <http://dx.doi.org/10.1007/s00401-008-0366-2>
- [16] Bancher C, Brunner C, Lassmann H, Budka H, Jellinger K, Wiche G, Seitelberger F, Grundke-Iqbal I, Iqbal K, Wisniewski HM. Accumulation of abnormally phosphorylated tau precedes the formation of neurofibrillary tangles in Alzheimer's disease. *Brain Res* 1989; 477:90-9; PMID:2495152; [http://dx.doi.org/10.1016/0006-8993\(89\)91396-6](http://dx.doi.org/10.1016/0006-8993(89)91396-6)
- [17] Eisele YS, Duyckaerts C. Propagation of Abeta pathology: hypotheses, discoveries, and yet unresolved questions from experimental and human brain studies. *Acta Neuropathol* 2016; 131:5-25; PMID:26715565; <http://dx.doi.org/10.1007/s00401-015-1516-y>
- [18] Thal DR, Capetillo-Zarate E, Del Tredici K, Braak H. The development of amyloid beta protein deposits in the aged brain. *Sci Aging Knowledge Environ* 2006; 2006:re1; PMID:16525193; <http://dx.doi.org/10.1126/sageke.2006.6.re1>
- [19] Wang D, Munoz DG. Qualitative and quantitative differences in senile plaque dystrophic neurites of Alzheimer's disease and normal aged brain. *J Neuropathol Exp Neurol* 1995; 54:548-56; PMID:7602328; <http://dx.doi.org/10.1097/00005072-199507000-00009>
- [20] Dickson DW. The pathogenesis of senile plaques. *J Neuropathol Exp Neurol* 1997; 56:321-39; PMID:9100663; <http://dx.doi.org/10.1097/00005072-199704000-00001>
- [21] Braak H, Alafuzoff I, Arzberger T, Kretschmar H, Del Tredici K. Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* 2006; 112:389-404; PMID:16906426; <http://dx.doi.org/10.1007/s00401-006-0127-z>
- [22] Thal DR, Rub U, Orantes M, Braak H. Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* 2002; 58:1791-800; PMID:12084879; <http://dx.doi.org/10.1212/WNL.58.12.1791>
- [23] Reiniger L, Lukic A, Linehan J, Rudge P, Collinge J, Mead S, Brandner S. Tau, prions and Abeta: the triad of neurodegeneration. *Acta Neuropathol* 2011; 121:5-20; PMID:20473510; <http://dx.doi.org/10.1007/s00401-010-0691-0>
- [24] Ghoshal N, Cali I, Perrin RJ, Josephson SA, Sun N, Gambetti P, Morris JC. Codistribution of amyloid beta plaques and spongiform degeneration in familial Creutzfeldt-Jakob disease with the E200K-129M haplotype. *Arch Neurol* 2009; 66:1240-6; PMID:19822779; <http://dx.doi.org/10.1001/archneurol.2009.224>
- [25] Jaunmuktane Z, Mead S, Ellis M, Wadsworth JD, Nicoll AJ, Kenny J, Launchbury F, Linehan J, Richard-Loendt A, Walker AS, et al. Evidence for human transmission of amyloid-beta pathology and cerebral amyloid angiopathy. *Nature* 2015; 525:247-50; PMID:26354483; <http://dx.doi.org/10.1038/nature15369>
- [26] Irwin DJ, Abrams JY, Schonberger LB, Leschek EW, Mills JL, Lee VM, Trojanowski JQ. Evaluation of potential infectivity of Alzheimer and Parkinson disease proteins in recipients of cadaver-derived human growth hormone. *JAMA Neurol* 2013; 70:462-8; PMID:23380910; <http://dx.doi.org/10.1001/jamaneurol.2013.1933>
- [27] Frontzek K, Lutz MI, Aguzzi A, Kovacs GG, Budka H. Amyloid-beta pathology and cerebral amyloid angiopathy are frequent in iatrogenic Creutzfeldt-Jakob disease after dural grafting. *Swiss Med Wkly* 2016; 146:w14287; PMID:26812492
- [28] Kovacs GG, Lutz MI, Ricken G, Strobel T, Hofberger R, Preusser M, Regelsberger G, Honigschnabl S, Reiner A, Fischer P, et al. Dura mater is a potential source of Abeta seeds. *Acta Neuropathol* 2016; 131:911-23; PMID:27016065; <http://dx.doi.org/10.1007/s00401-016-1565-x>

- [29] Hamaguchi T, Taniguchi Y, Sakai K, Kitamoto T, Takao M, Murayama S, Iwasaki Y, Yoshida M, Shimizu H, Kakita A, et al. Significant association of cadaveric dura mater grafting with subpial Abeta deposition and meningeal amyloid angiopathy. *Acta Neuropathol* 2016; 132:313-5; PMID:27314593; <http://dx.doi.org/10.1007/s00401-016-1588-3>
- [30] Morales R, Callegari K, Soto C. Prion-like features of misfolded Abeta and tau aggregates. *Virus Res* 2015; 207:106-12; PMID:25575736; <http://dx.doi.org/10.1016/j.virusres.2014.12.031>
- [31] Gotz J, Chen F, van Dorpe J, Nitsch RM. Formation of neurofibrillary tangles in P3011 tau transgenic mice induced by Abeta 42 fibrils. *Science* 2001; 293:1491-5; PMID:11520988; <http://dx.doi.org/10.1126/science.1062097>
- [32] Edgren G, Hjalgrim H, Rostgaard K, Lambert P, Wikman A, Norda R, Titlestad KE, Erikstrup C, Ullum H, Melbye M, et al. Transmission of Neurodegenerative Disorders Through Blood Transfusion: A Cohort Study. *Ann Intern Med* 2016; 165:316-324