

Improving and accelerating the differentiation and functional maturation of human stem cell-derived neurons: role of extracellular calcium and GABA


Paul J. Kemp¹, David J. Rushton², Polina L. Yarova¹, Christian Schnell¹, Charlene Geater³, Jane M. Hancock⁴, Annalena Wieland¹, Alis Hughes¹, Luned Badder¹, Emma Cope¹, Daniela Riccardi¹, Andrew D. Randall⁴, Jonathan T. Brown⁴, Nicholas D. Allen¹ and Vsevolod Telezhkin¹

¹School of Biosciences, Cardiff University, Cardiff, UK

²Board of Governors Regenerative Medicine Institute, Cedars–Sinai Medical Center, Los Angeles, CA, USA

³Department of Psychiatry and Human Behavior, University of California at Irvine, Irvine, CA, USA

⁴Institute of Biomedical and Clinical Sciences, University of Exeter Medical School, Hatherly Laboratory, Exeter EX4 4PS, UK



Paul J. Kemp completed his DPhil in Physiology in the Department of Human Anatomy at the University of Oxford in 1990. Since then he has held academic positions in the University of Dundee, the University of Southern California and the University of Leeds. Since 2004, he has been Professor of Physiology at Cardiff University School of Biosciences where his group works on gas sensing by ion channels, novel asthma therapeutics (with Daniela Riccardi) and fate determination of human pluripotent stem cells (with Nicholas D Allen). **Vsevolod Telezhkin** was trained in the Bogomoletz Institute in Kiev where he obtained his PhD in 2003. Following several successful post-doctoral positions at the Universities of Vermont, Bath and Cardiff, he worked at University College London with Professor David Brown, FRS, with whom he published a series of important papers on regulation of M-channels by lipids. Currently, he is an associate at Cardiff University where his main interest is in the role of M-channels in pluripotent stem cell differentiation and neuronal maturation.



This review was presented at the symposium “Are stem cell-derived neural cells physiologically credible?”, which took place at Physiology 2015 in Cardiff, UK, 6–8 July 2015.

Abstract Neurons differentiated from pluripotent stem cells using established neural culture conditions often exhibit functional deficits. Recently, we have developed enhanced media which both synchronize the neurogenesis of pluripotent stem cell-derived neural progenitors and accelerate their functional maturation; together these media are termed SynptoJuice. This pair of media are pro-synaptogenic and generate authentic, mature synaptic networks of connected forebrain neurons from a variety of induced pluripotent and embryonic stem cell lines. Such enhanced rate and extent of synchronized maturation of pluripotent stem cell-derived neural progenitor cells generates neurons which are characterized by a relatively hyperpolarized resting membrane potential, higher spontaneous and induced action potential activity, enhanced synaptic activity, more complete development of a mature inhibitory GABA_A receptor phenotype and faster production of electrical network activity when compared to standard differentiation media. This entire process – from pre-patterned neural progenitor to active neuron – takes 3 weeks or less, making it an ideal platform for drug discovery and disease modelling in the fields of human neurodegenerative and neuropsychiatric disorders, such as Huntington's disease, Parkinson's disease, Alzheimer's disease and Schizophrenia.

(Received 22 February 2016; accepted after revision 23 August 2016; first published online 12 September 2016)

Corresponding author P. J Kemp and V. Telezhkin: School of Biosciences, The Sir Martin Evans Building, Cardiff University, Museum Avenue, Cardiff CF10 3AX, UK. Emails: kemp@cf.ac.uk; telezhkinv@cf.ac.uk

Abstract figure legend Rationale and basic protocol for accelerating neuronal maturation using SynptoJuice. Neural precursor cells (NPCs) that have been generated from either human embryonic stem (ES) cells or human induced pluripotent stem (iPS) cells are plated at day 0 in SynptoJuice A. This first medium has been designed to synchronize neurogenesis, by forcing NPCs to exit the cell cycle, whilst increasing GABA-induced Ca²⁺ influx. At day 7, and thereafter, SynptoJuice A is replaced by SynptoJuice B, which is designed to promote synaptogenesis via WNT and TrkB signalling. At Day 21, the resulting neurons demonstrate punctate co-registration of synaptic markers (synaptophysin and PSD95 are shown) and synchronized electrical activity (an exemplar spontaneous train of action potentials is shown).

Introduction

Since the original isolation and characterization of human embryonic stem cells (ESCs) in 1998 (Thomson *et al.* 1998), followed by the discovery some 10 years later of human somatic cell reprogramming into induced pluripotent stem cells (iPSCs) (Takahashi *et al.* 2007), pluripotent stem cells have been regularly hailed as a panacea for understanding and, potentially, treating an ever increasing list of human disorders, providing the raw material for cell replacement therapies, disease modelling and drug screening. Review of clinical trials shows that pluripotent stem cells have been employed with variable success as cell replacements in a variety of disorders, either directly or via a population of artificial support matrices, including tracheal replacement (Jungebluth *et al.* 2011), retinal dysfunction (Zarbin, 2016), spinal cord injury (Lukovic *et al.* 2014) and several neurodegenerative diseases (Levy *et al.* 2016). Unlike replacement therapies, which require the use of either the pluripotent stem cells themselves or pre-patterned/pre-specified cell lineage-specific precursors, disease modelling and drug screening frequently have the explicit requirement for a population of terminally differentiated cells of the type affected in the disease in question. This requirement has led to the rapid and spirited development of protocols

aimed at directing differentiation of pluripotent stem cells to correctly specified cells. Such developments have been particularly competitive and often controversial in the field of neuroscience where the scope for mixed populations of neurons is high but often undesirable when the exact cellular composition *in vitro* may not yet be fully defined.

Independently of specification of a particular neuronal cell type, it is absolutely imperative that differentiated neurons demonstrate appropriate cell biological characteristics and functional activities. In the field of neuronal differentiation, the efficiency and efficacy of the majority of protocols is assessed using an array of established cell-specific markers employing a combination of immunocytochemistry (ICC), reverse transcription polymerase chain reaction (RT-PCR) and, more recently, RNA sequencing (RNAseq) and proteomic analyses. However, except in a handful of exceptional studies (Johnson *et al.* 2007; Song *et al.* 2013; Livesey *et al.* 2014; Telias *et al.* 2014; Bardy *et al.* 2015), integration of functional assays of neuronal function is rare and, where reported, is normally restricted to single examples of induced action potentials and/or miniature synaptic potentials, with no further analyses. This relative dearth of functional information regarding appropriate neuronal activity is troublesome when such protocols are used to develop meaningful disease models and high-throughput

drug screens. In other words, it is unclear whether particular protocols consistently generate functionally active neurons whose activity has not been robustly characterized or, more worryingly, that the protocols do not produce neurons with robust electrophysiological activity characteristic of the neurons being studied. Indeed, we and others have found that published neuronal differentiation protocols generate neurons which express the appropriate marker molecules but which often fail to generate authentic neuronal activity with any consistency across the population. With this in mind, recent work in several laboratories, including our own, has focused on improving basic neuronal function during the *in vitro* differentiation process using small molecule regulators of specific cell biological cascades and manoeuvres aimed at mimicking physiological and environmental cues which are known to be important during neuronal differentiation *in vivo* in order to improve function and to stimulate synaptogenesis and consequent neuronal network activity *in vitro*.

Regulation of physiological P_{O_2} and of extracellular free ionized calcium and their potential roles in neuronal differentiation

Neuronal specification and differentiation *in vivo* occur prenatally in an environment which is markedly different from that postnatally (Mohyeldin *et al.* 2010). Of particular significance is the partial pressure of oxygen (P_{O_2}), which rapidly changes from around 20–30 to 150–160 mmHg following the first breath of the emerging neonate (Ward, 2008). Such low P_{O_2} , or relative hypoxia, has been successfully employed to improve the efficiency of the generation of iPSCs (Bilican *et al.* 2012; Hawkins *et al.* 2013) and a handful of studies have investigated the effect of such relative hypoxia on subsequent neuralization, specification and differentiation of neurons (e.g. Studer *et al.* 2000; Bilican *et al.* 2014).

Less well appreciated is the parallel switch in extracellular free ionized Ca^{2+} concentration ($[Ca^{2+}]_o$) which occurs in the perinatal period. Thus, human fetal plasma $[Ca^{2+}]_o$ is ~ 1.6 – 1.7 mM (Kovacs & Kronenberg, 1997) and this is rapidly reduced within 24 h postnatally to the mean adult level of 1.1–1.3 mM (Brown, 1991); the relative hypercalcaemia of the developing fetus compared to maternal levels is retained independently of any changes in the maternal $[Ca^{2+}]_o$ (Kovacs *et al.* 1998). In health, the relatively low postnatal $[Ca^{2+}]_o$ persists and is strictly regulated within a very narrow range by the action of parathyroid hormone on the kidney, intestine and bone, a homeostatic process which is critically dependent upon the activation by Ca_o^{2+} of the extracellular Ca^{2+} -sensing receptor (CaSR) expressed in the parathyroid gland (Brown *et al.* 1993; Riccardi & Kemp, 2012). The striking transition from high prenatal

to low postnatal $[Ca^{2+}]_o$ is mediated by the CaSR (Kovacs *et al.* 1998).

Many commercially available tissue culture media contain $[Ca^{2+}]_o$ broadly embracing the mean value of postnatal $[Ca^{2+}]_o$ in humans of 1.1–1.3 mM (Brown, 1991). With this knowledge in mind, it is puzzling that the majority of neuronal specification and differentiation protocols employ these standard culture media which contain relatively low $[Ca^{2+}]_o$, or hypocalcaemic, conditions for the developing fetus. Indeed, where studied, the effect of increasing $[Ca^{2+}]_o$ from adult to fetal values is permissive for neurite outgrowth of prenatal primary neurons isolated from the sympathetic cervical ganglion *in vitro* (Vizard *et al.* 2008).

Role of astrocytes and astrocyte-conditioned medium in neuronal differentiation

It has long been recognized that co-culture of primary neurons with astrocytes promotes neurogenesis and enhances synaptogenesis both directly (Pfrieger & Barres, 1997; Ullian *et al.* 2001, 2004; Hama *et al.* 2004) and indirectly via secreted factors (Chang *et al.* 2003; Ullian *et al.* 2004; Chou *et al.* 2008). Furthermore, maturation of human stem cell-derived neurons can be enhanced by co-culture with astrocytes (Johnson *et al.* 2007) or astrocyte conditioned medium (ACM) (Rushton *et al.* 2013; Tang *et al.* 2013). Indeed, we have shown that ACM has the effect of accelerating the functional maturation rate of iPSC-derived neurons during differentiation by hyperpolarizing the resting membrane potential, thereby increasing the spontaneous activity of the developing neurons, often generating neurons with complex, biphasic patterns of excitability (Rushton *et al.* 2013). Such relative hyperpolarization, which is seen during differentiation of human iPSCs to neurons *in vitro*, appears to be physiologically relevant as evidenced by the fact that it can also be observed *in vivo*, where we have preliminary evidence to show that developmentally regulated expression of Kv7.2/7.3 (M-type) currents appears to underlie a gradual hyperpolarization of striatal neurons during the second and third trimesters of mouse gestation (Telezhkin *et al.* 2014).

Previous observations have reported that there are increases in Ca^{2+} channel expression and/or function early in stem cell-derived and native NPC differentiation (Arnhold *et al.* 2000; Mazzanti & Haydon, 2003; D'Ascenzo *et al.* 2006). This led us to investigate voltage-gated Ca^{2+} entry, and the potential role of Ca^{2+} channels, in the ACM-evoked enhancement of human iPSC neuronal differentiation (Rushton *et al.* 2013). Differentiation in ACM evoked a robust enhancement of depolarization-evoked Ca^{2+} entry, principally via L-type, N-type and R-type Ca^{2+} channels, with little contribution of P/Q-type channels (Aga toxin-sensitive) (Fig. 1A).

Consistent with the selective upregulation of Ca^{2+} influx through voltage-gated Ca^{2+} channels, chronic blockade of L-type (nifedipine), N-type (conotoxin), or R-type (SNX-482) Ca^{2+} channels during the differentiation process resulted in complete abolishment of the ACM-evoked enhancements of spontaneous activity and significantly depolarized resting membrane potential (Fig. 1B). Furthermore, similar diminutions were seen when ionotropic GABA_A receptors were inhibited using bicuculline, the first suggestion that a major contributor to Ca^{2+} influx during differentiation might be provided by excitatory GABA_A synaptic input in these developing neuronal networks (Fig. 1B). However, it should be noted that no direct comparison of $[\text{Ca}^{2+}]_i$ in the two conditions was attempted in these studies.

Roles for GABA_A receptors and extracellular Ca^{2+} in neuronal differentiation

The fact that the robust enhancement of function supported by ACM was characterized by increased Ca^{2+} influx through specific Ca^{2+} channels, and that it could be attenuated by their inhibition, suggested that the mechanism of the ACM effect might involve chronic augmentation of intracellular free ionized calcium concentration ($[\text{Ca}^{2+}]_i$) via GABA_A -dependent excitation. In order to test this possibility, we attempted to mimic, at least early in the differentiation process, the permissive effects of ACM by chronically raising the extracellular GABA concentration, and the $[\text{Ca}^{2+}]_o$ to a level similar to that measured in fetal plasma, namely




Figure 1. Potential mechanisms of ACM-evoked enhancement of neuronal function

A, Fura-2-derived relative intracellular Ca^{2+} concentration measurements for iPSC-derived NPCs after 1 week of differentiation in either standard differentiation medium (SDM, upper trace) or (ACM, lower trace). Ca^{2+} influx was evoked by depolarization using 50 mM KCl in the absence and presence of 10 μM nifedipine (L-type Ca^{2+} channel blockade), 100 nM conotoxin (N-type Ca^{2+} channel blockade), 100 nM agatoxin (P/Q Ca^{2+} channel blockade) or 100 nM SNX482 (R-type Ca^{2+} channel inhibition). B, bar graphs comparing the proportion of neurons firing spontaneous action potentials (sAPs, upper bars) and mean resting membrane potentials (V_m , lower bars) following 3 weeks of differentiation in ACM alone, or ACM with 10 μM bicuculline, 2 μM nifedipine, 100 nM conotoxin or 100 nM SNX482. C, comparison of the proportion of neurons firing sAPs (upper bars) and mean V_m (lower bars) following 3 weeks of differentiation in 1.2 or 1.8 mM $[\text{Ca}^{2+}]_o$ in the presence of 10 μM bicuculline, 2 μM nifedipine, 100 nM conotoxin or 100 nM agatoxin. D, comparison of the proportion of cells firing sAPs (upper bars) and mean V_m (lower bars) following 1 and 2 weeks of differentiation in SDM with and without 300 μM GABA. Different from control, *** $P < 0.001$. Adapted from Rushton *et al.* (2013).

1.8 mM (Kovacs & Kronenberg, 1997). These manoeuvres allowed the study of the efficiency and efficacy of differentiation of iPSC-derived neural progenitor cells (NPCs) to neurons under conditions where the $[Ca^{2+}]_i$ was predicted to have been increased tonically (Rushton *et al.* 2013), although it must be noted that no direct comparison of $[Ca^{2+}]_i$ in the two conditions was attempted in these studies.

Thus, differentiating the NPCs for 3 weeks (which is 35 days from the pluripotent stem cell stage), in medium containing 1.8 mM Ca^{2+}_o resulted in an increase in spontaneous activity and a significant hyperpolarization. These effects were attenuated by co-incubation for the entire differentiation with inhibitors of GABA_A receptors or blockers of L-type, N-type or R-type Ca^{2+} channels; consistent with the Ca^{2+} imaging data presented in Fig. 1A, blockade of P/Q channels had no effect on the ability of raised $[Ca^{2+}]_o$ to augment spontaneous activity and to hyperpolarize the resting membrane potential (Fig. 1C). Note that the potent Ca^{2+} channel opener BayK 8644 was also able to enhance spontaneous activity and to hyperpolarize the neurons in standard $[Ca^{2+}]_o$ (Rushton *et al.* 2013).

Chronic incubation of differentiating iPSCs with GABA evoked a significant increase in spontaneous activity and hyperpolarization at week 1. However, this augmentation in function was not further enhanced at week 2 (Fig. 1D), suggesting that GABA might only be useful in promoting neuronal maturation early in the differentiation programme. The idea that GABA might provide the stimulus for Ca^{2+}_o influx early in the neuronal differentiation/maturation process, perhaps both *in vivo* and *in vitro*, is consistent with knowledge that ionotropic GABA_A receptors are actually excitatory in pre-natal neurons. Thus, GABA binding elicits Cl^- exit and consequent depolarization (Ganguly *et al.* 2001; Ben-Ari, 2002; Tozuka *et al.* 2005). The postnatal transition from excitatory to inhibitory is a result of the developmentally modulated Cl^- equilibrium potential, which is itself controlled by the differential expression of $Na^+-K^+-2Cl^-$ cotransporters (NKCC1) and K^+-Cl^- cotransporters (KCC2) (Ganguly *et al.* 2001). Consequently, the GABA_A phenotype (excitatory or inhibitory) of differentiating stem cells is a reliable functional readout for immature *versus* mature neurons. Human stem cell-derived neurons demonstrate robust GABA_A currents and GABA-evoked Ca^{2+}_o influx even at early stages of neuronal differentiation (Joannides *et al.* 2007; HD iPSC Consortium, 2012) and GABA-dependent excitation has been shown to promote neuronal differentiation in several systems, including adult hippocampal stem/progenitors (Tozuka *et al.* 2005) and neuroectodermal stem cells (Jelita *et al.* 2004). Therefore, it seemed likely that GABA might promote neuronal differentiation of iPSCs. Interestingly, although there is transition from excitatory to

inhibitory GABA phenotype during early differentiation, ACM does not significantly alter the time course of this event (Rushton *et al.* 2013), suggesting that regulated GABA_A, NKCC1 and KCC2 expression do not contribute significantly to ACM-induced neuronal maturation *per se* (Rushton *et al.* 2013). The GABA transition has also been well characterized during differentiation of human stem cells to cortical neurons (Livesey *et al.* 2014); this study also showed time-dependent alteration in the expression of AMPA receptor composition. Whether this process is sensitive to $[Ca^{2+}]_o$ is not known (Livesey *et al.* 2014).

Integrating GABA and raised $[Ca^{2+}]_o$ into a pro-synaptogenic differentiation protocol

Having determined that maintaining $[Ca^{2+}]_o$ at 1.8 mM throughout the 3 week differentiation period and that GABA stimulation early in the process were both able to support enhanced neuronal function, as determined by observation of the frequency of spontaneous electrical activity and relative hyperpolarization, we began the rational design of a pair of media which would augment and/or accelerate functional maturation of iPSC-derived neurons. Medium 1 contained the high $[Ca^{2+}]_o$ together with GABA (used during the first week following plate-down of NPCs) whilst medium 2 contained just the high $[Ca^{2+}]_o$ (used from 1 week onwards). With the addition of brain-derived neurotrophic factor (BDNF) to both, these media were termed base media, and to them various small molecules were added. For these particular experiments, pre-patterning to NPCs took 16 days, meaning the entire process, from pluripotent stem cell stage to mature neurons took 35–37 days.

One of the major problems associated with neuronal differentiation from stem cells is the lack of synchronicity in the process. This means that neurons can be 'born' from NPCs at almost any time during the differentiation and this results in distinct lack of neuronal functional homogeneity. To address this issue, we employed inhibitors of both Notch signalling and the cell cycle checkpoint cyclin-dependent kinases CDK4 and CDK6 in order to promote cell cycle exit (Telezhkin *et al.* 2016). Blockade of Notch signalling with the γ -secretase inhibitor *N*-[*N*-(3,5-difluorophenacetyl-L-alanyl)]-(*S*)-phenylglycine *t*-butyl ester (DAPT) or CDK4/6 kinase activity with the selective inhibitor PD0332991 (Toogood *et al.* 2005) significantly reduced the proportion of NPCs actively proliferating at 3 and 7 days post-plate-down. In combination, these small molecules evoked synchronous, sustained cell cycle exit (Fig. 2A and B) and enhanced neural differentiation, as shown by an increase in the proportion of cells expressing the neuronal marker, β -III-tubulin, and a marked reduction in the neural progenitor marker nestin at 7 days (Telezhkin *et al.* 2016).

These data suggested that these molecules should be added to the base media.

It is well known that GABA_A receptor activation leads to Ca²⁺-dependent cAMP response element-binding protein (CREB) and extracellular signal-regulated kinase (ERK) phosphorylation, which positively regulates neurogenic gene expression (Merz *et al.* 2011; Wiegert & Bading, 2011). Furthermore, functional expression of neuronal voltage-gated Na⁺ and K⁺ channels during differentiation can also be affected by agents that elevate cAMP (Aglah *et al.* 2008; Ravni *et al.* 2008). Thus, medium 1, which already contained high [Ca²⁺]_o and GABA to drive the GABA_A-dependent increase in [Ca²⁺]_i, was supplemented with forskolin to raise intracellular [cAMP]_i, leading to CREB phosphorylation (Telezhkin *et al.* 2016).

As GABA-dependent CREB phosphorylation has a transient developmental role in neurogenesis (Jagasia *et al.* 2009), GABA and forskolin were removed from medium 2, to encourage maturation during the later phase differentiation. Similarly, since the pro-neurogenic effects of Notch inhibition are only needed early in neurogenesis, and sustained γ -secretase inhibition can be deleterious (Shen, 2014), DAPT was also removed from medium 2.

A directed screen of growth factors and small molecules was then performed in order to identify compounds which enhance neuronal excitability. Little or no enhancement in excitability was observed with BDNF, fibroblast growth factor 2 (FGF2), insulin-like growth factor 1 (IGF1) or activin A. However, the glycogen synthase kinase 3 β (GSK3 β) antagonist CHIR99021 did increase action potential firing, which prompted us to incorporate it into both media for further analysis. CHIR99021 has been previously demonstrated to promote post-mitotic nociceptor neural differentiation, although this was in combination with DAPT (Chambers *et al.* 2012).

Functional consequences of neuronal differentiation in the enhanced media

With the components of the two media complete, plated NPCs derived from several iPSC sources, derived using both integrating and non-integrating reprogramming vectors, were differentiated for 1 week in medium 1, and 2 more weeks in medium 2. Together, these two media are now termed SynaptoJuice (www.synaptojuice.co.uk). The potential benefits of using these media were determined




Figure 2. Synchronous cell cycle exit and differentiation are promoted by γ -secretase and cyclin dependent kinase 4/6 inhibition

Exemplar fluorescence micrographs (upper) and mean cell counts (lower) of MKI67 immunoreactivity (green) and cell nuclear staining with DAPI (blue) of neural progenitor cells (NPCs) 3 (A) and 7 days (B) post-plating in base medium with no additions (Control), or base medium supplemented with 2 μ M PD0332991 or 10 μ M DAPT, or both. Different from control, *** P < 0.001, ** P < 0.01, * P < 0.05. Scale bars: 100 μ m. Adapted from Telezhkin *et al.* (2016).

by comparing their effects on electrical activity and neurotransmitter responses with those seen in base medium and standard differentiation medium (Telezhkin *et al.* 2016). Differentiation in the enhanced media yielded the highest proportion of neurons demonstrating spontaneous electrical activity (Fig. 3A and B), which was consistent with the data showing that such neurons displayed the most hyperpolarized membrane potentials and suggested that there was a more rapid transition from inactive to active state via evocation of relative hyperpolarization (Fig. 3C). Such a notion is again entirely consistent with our preliminary observations that M-currents are developmentally regulated both *in vivo* and *in vitro* (Telezhkin *et al.* 2014).

In terms of induced electrical activity, differentiation in enhanced media supported the highest proportion of neurons exhibiting action potential trains compared

to those differentiated in base or standard media (Fig. 4; Telezhkin *et al.* 2016); and the current injection *vs.* spike frequency plots showed that the enhanced media supported the greatest excitability (Telezhkin *et al.* 2016). Differentiation in the enhanced media generated neurons with significantly shorter half-widths and larger overshoots, afterhyperpolarizations, spike amplitudes and depolarization rates (Telezhkin *et al.* 2016). Surprisingly, neither voltage-activated Na⁺ nor K⁺ current densities were affected by the differentiation protocol. However, differentiation in the enhanced media generated neurons which displayed significantly altered Na⁺ current activation/inactivation profiles, resulting in larger availability windows and a higher proportion of neurons with V_m values falling within those windows than did either base or standard media resulting in the probability of a neuron firing a spontaneous action




Figure 3. Enhanced differentiation media (SynptoJuice) augment the generation of neurons which exhibit spontaneous action potentials

A, typical current-clamp recordings ($I = 0$ pA) illustrating the three separate types of activity which were observed spontaneously in the neurons: spontaneous action potentials (sAP-full, upper); incomplete initiation of spontaneous action potentials (sAP-incomplete initiation, middle); and no spontaneous action potentials (sAP-None, lower). Note that each sAP type is also the modal activity for neurons differentiated in SynptoJuice, base medium and standard differentiation medium, respectively. B, proportions of neurons which demonstrated each of the individual sAP characteristics when NPCs were differentiated into neurons for 3 weeks in the enhanced media (SynptoJuice), base medium and standard differentiation medium (i.e. 35 days from pluripotent stem cells). C, mean resting membrane potentials (V_m) of neurons differentiated for 3 weeks in SynptoJuice, base medium and standard differentiation medium. *** $P < 0.001$, **** $P < 0.0001$. Time bar applies to all traces. Adapted from Telezhkin *et al.* (2016).

potential being increased (Telezhkin *et al.* 2016). These data suggested that the enhanced media improved functional maturation by facilitating neuronal hyperpolarization (to support higher spontaneous activity), and by increasing Na⁺ current availability (to enhance regenerative action potential train activity).

To examine whether any set of media was able to support functional synaptogenesis, spontaneous miniature synaptic currents (minis), evoked postsynaptic currents, neurotransmitter-evoked changes in intracellular calcium concentration ($[Ca^{2+}]_i$) and network activity were determined. Cells differentiated in standard medium for 3 weeks never exhibited miniature synaptic currents. Differentiation in base media only rarely supported the generation of GABAergic minis. In contrast, up to 75% of neurons differentiated in the enhanced media exhibited large GABAergic and modest glutamatergic

minis. All cells differentiated in any of the media exhibited GABA-evoked currents, whereas there was a distinct enhancement in the proportion of cells exhibiting, and the magnitude of, NMDA-evoked currents in the enhanced media. These observations were supported by the demonstration of punctate, presumably synaptic, staining of NR1, NR2A and NR2B NMDA receptor subunits, co-registration of the pre- and postsynaptic markers, synaptophysin and PSD95 and the appearance of dendritic spines exclusively in the neurons differentiated in the enhanced media (Telezhkin *et al.* 2016).

As the enhanced media appeared to improve the extent of functional neuronal maturation, weekly imaging of the neurotransmitter responses was performed in order to determine the time course of this enhancement. From a very low level at the time of plating the NPCs, most of the cells from all three protocols responded




Figure 4. Enhanced differentiation media (SynptoJuice) augment the generation of neurons exhibiting induced action potential train

A, exemplar current-clamp recordings ($I = 0$ pA) illustrating the five separate types of induced activity which were observed in the neurons: train of induced action potentials (iAP-Train); single action potential with incomplete initiation of action potential trains (iAP-Incomplete initiation); single action potential (iAP-Single); incomplete initiation of single action potential (iAP-Incomplete initiation of single); and no induced action potential (iAP-None, lower). Time bar applies to all traces. Adapted from Telezhkin *et al.* (2016). *B*, proportions of neurons which demonstrated each of the individual iAP characteristics when NPCs were differentiated into neurons for 3 weeks in the enhanced media (SynptoJuice), base medium and standard differentiation medium (i.e. for 35 days from pluripotent stem cells).

to a depolarizing challenge at 1 week of differentiation and thereafter (Telezhkin *et al.* 2016). However, from 2 weeks onwards, the magnitude of the responses was significantly lower for neurons differentiated using either standard or base media and this persisted until 3 weeks in standard medium (Telezhkin *et al.* 2016). This was consistent with augmented expression of voltage-activated Ca^{2+} channels being supported by the enhanced media, which has been previously documented to occur with ACM (Rushton *et al.* 2013). The proportion of neurons responding, and the magnitude of the Ca^{2+} response, to glutamate/glycine increased during differentiation, but the enhanced media encouraged a more rapid augmentation which was delayed by a week in base medium and was never enhanced in standard differentiation medium (Fig. 5A–C). Likewise, the proportion of cells responding, and the magnitude of the response, to GABA in low $[\text{Cl}^-]_o$ remained significantly smaller for standard and base media (Fig. 5A–C). Furthermore, since the response to GABA in

normal $[\text{Cl}^-]_o$ is potentially composed of both excitatory GABA_A and facilitatory GABA_B (Karls & Mynlieff, 2015) receptor responses, it seems reasonable to assume that our enhanced media, which largely abrogate this response to GABA in normal $[\text{Cl}^-]_o$ (Fig. 5C), are promoting the transition from immature to mature GABA_A phenotype and, possibly, causing a reduction in GABA_B expression. As little is known about the ontogeny of GABA_B -dependent Ca^{2+} signalling in developing neurons, but it is known that GABA_A responses transition from excitatory to inhibitory, loss of GABA_B might also be an indication of maturity in pluripotent stem cell-derived neurons. This, and the precise timing of the functional switch in GABA_A phenotype, which is promoted by the enhanced differentiation media, are currently under investigation.

Finally, evidence that the enhanced media supported the formation of excitatory synaptic networks whose activity was being tonically dampened by inhibitory GABAergic tone was provided by experiments using 24-well




Figure 5. Enhanced differentiation media (SynptoJuice) improve time-dependent maturation of post-synaptic neurotransmitter responses and supports regulatable network activity

A, proportion of neurons displaying Ca^{2+} influx in response to 60 mM KCl (KCl), 300 μM glutamate/30 μM glycine (Glu/Gly) and 300 μM GABA in 7.5 mM chloride (Low Cl^- /GABA) following differentiation for 3 weeks of NPCs (i.e. 35 days from pluripotent stem cells) from NPCs in SynptoJuice, base medium, and standard differentiation medium. **B**, mean magnitude of Ca^{2+} influx, expressed as change in fluorescence from baseline (Δ Fluorescence) of the same neurons used for the analysis shown in panel A. **C**, exemplar fura-2-derived intracellular Ca^{2+} recordings from individual neurons of the cohort shown in panels A and B. Note the response to 300 μM GABA alone (GABA) was large in neurons in standard differentiation medium, small in base medium and often not detectable in enhanced differentiation media. **D**, exemplar recording of spontaneous network spike rate before (left) and following (right) the addition of 5 μM gabazine to one of 24 wells in a multi-well microelectrode array. Note the split time axis for addition of drug. Single unit activity is shown as an insert. Neurons were differentiated from NPCs for 3 weeks in enhanced differentiation medium (i.e. 35 days from pluripotent stem cells). Adapted from Telezhkin *et al.* (2016).

microelectrode array (MEA) plates which showed that large and maintained increases in activity could be evoked by the addition of gabazine, a GABA_A receptor inhibitor (Fig. 5D). Furthermore, glutamate was able to stimulate activity whilst tetrodotoxin completely abolished activity in all wells (Telezhkin *et al.* 2016).

Comparison with alternative differentiation media and protocols

Thus, our recent published studies support the notion that forced cell cycle exit is able to synchronize neuronal differentiation. That simple manoeuvre, coupled to simulating fetal conditions by increasing $[Ca^{2+}]_o$ throughout whilst transiently generating a GABA excitatory drive and tonically applying known pro-synaptogenic small molecules (CHIR 99021 and forskolin), results in rapid functional maturation of neurons with highly homogeneous response profiles. Although there has been sustained activity for over a decade aimed at producing neurons from a variety of stem cell sources, it is only recently that there has been brisk and highly competitive progress towards generating demonstrably functional neurons exhibiting characteristically appropriate synaptic properties *in vitro*. There have been several other carefully performed recent studies which have demonstrated maturation by a variety of functional endpoints (Song *et al.* 2013; Livesey *et al.* 2014; Telias *et al.* 2014; Bardy *et al.* 2015). The study by Song and colleagues analysed the electrophysiological time course of pluripotent stem cell-derived neuronal differentiation (Song *et al.* 2013). Although their data are in broad agreement with ours, the homogeneity of the neurons with respect to repetitive action potential train generation is more similar at 3 weeks to those generated by our standard differentiation medium. Even after a further week, the Song protocol still only supported repetitive firing in under 50% of the neurons, likely to be due to the lack of synchronization of the differentiation programme that our enhanced media afford early in neurogenesis. Furthermore, their induced action potentials have lower spike amplitudes, slow depolarization rates and long half-widths, again more reminiscent of those we see in standard differentiation medium. The Livesey protocol (Livesey *et al.* 2014) used several important readouts of neuronal maturation, including GABA_A phenotype (Ganguly *et al.* 2001). They observed GABA_A response transition from excitatory to inhibitory during the 7 week differentiation, which was due to progressive reduction in chloride equilibrium potential (E_{Cl}) as the expression of NKCC1 and KCC2 was reduced and enhanced, respectively (Rivera *et al.* 1999; Yamada *et al.* 2004); no data were available between 1 and 5 weeks. With our enhanced media, inhibition of GABAergic signalling with gabazine resulted in a dramatic increase in neuro-

nal network activity, which suggests that even by 3 weeks synaptic GABA was inhibitory across the network. In the Telias protocol (Telias *et al.* 2014) neurons differentiated from human ESCs demonstrated significant maturation properties by 3 weeks, including a high proportion of neurons with train activity. However, input resistance was high and individual action potential amplitudes were very low compared to neurons differentiated in our enhanced media. Their use of DAPT was encouraging and might have been expected partially to synchronize the neuronal maturation. However, there were no data presented regarding potential synchronization and it might be more prudent to use PD0332991 with DAPT, as we have now shown (Telezhkin *et al.* 2016). Finally, and of particular note, there is the exceptionally elegant study of Bardy and colleagues who, using an evidence-based approach, systematically manipulated the composition of their medium in order to optimize neuronal functional readouts (Bardy *et al.* 2015). This medium supported high levels of synaptic function in both primary neurons and differentiated stem cells. Now termed BrainPhys (Stem Cell Technologies), this medium contains, in contrast to SynptoJuice, adult $[Ca^{2+}]_o$. Compared to our enhanced media, activity appears to take longer to develop, but because the authors have shown that it supports neuronal activity for many weeks longer than SynptoJuice has been assayed so far, it could well be a perfect candidate for the 'third' medium for use after our enhanced media, in order to support long-term ageing of neurons, something that will be important for successful modelling of late-onset neurodegenerative diseases.

References

- Aglah C, Gordon T & Posse de Chaves EI (2008). cAMP promotes neurite outgrowth and extension through protein kinase A but independently of Erk activation in cultured rat motoneurons. *Neuropharmacology* **55**, 8–17.
- Arnhold S, Andressen C, Angelov DN, Vajna R, Volsen SG, Hescheler J & Addicks K (2000). Embryonic stem-cell derived neurones express a maturation dependent pattern of voltage-gated calcium channels and calcium-binding proteins. *Int J Dev Neurosci* **18**, 201–212.
- Bardy C, van den Hurk M, Eames T, Marchand C, Hernandez RV, Kellogg M, Gorris M, Galet B, Palomares V, Brown J, Bang AG, Mertens J, Bohnke L, Boyer L, Simon S & Gage FH (2015). Neuronal medium that supports basic synaptic functions and activity of human neurons *in vitro*. *Proc Natl Acad Sci USA* **112**, E2725–E2734.
- Ben-Ari Y (2002). Excitatory actions of GABA during development: the nature of the nurture. *Nat Rev Neurosci* **3**, 728–739.
- Bilican B, Livesey MR, Haghi G, Qiu J, Burr K, Siller R, Hardingham GE, Wyllie DJ & Chandran S (2014). Physiological normoxia and absence of EGF is required for the long-term propagation of anterior neural precursors from human pluripotent cells. *PLoS One* **9**, e85932.

- Bilican B, Serio A, Barmada SJ, Nishimura AL, Sullivan GJ, Carrasco M, Phatnani HP, Puddifoot CA, Story D, Fletcher J, Park IH, Friedman BA, Daley GQ, Wyllie DJ, Hardingham GE, Wilmut I, Finkbeiner S, Maniatis T, Shaw CE & Chandran S (2012). Mutant induced pluripotent stem cell lines recapitulate aspects of TDP-43 proteinopathies and reveal cell-specific vulnerability. *Proc Natl Acad Sci USA* **109**, 5803–5808.
- Brown EM (1991). Extracellular Ca^{2+} sensing, regulation of parathyroid cell function, and role of Ca^{2+} and other ions as extracellular (first) messengers. *Physiol Rev* **71**, 371–411.
- Brown EM, Gamba G, Riccardi D, Lombardi M, Butters R, Kifor O, Sun A, Hediger MA, Lytton J & Hebert SC (1993). Cloning and characterization of an extracellular Ca^{2+} -sensing receptor from bovine parathyroid. *Nature* **366**, 575–580.
- Chambers SM, Qi Y, Mica Y, Lee G, Zhang XJ, Niu L, Bilslund J, Cao L, Stevens E, Whiting P, Shi SH & Studer L (2012). Combined small-molecule inhibition accelerates developmental timing and converts human pluripotent stem cells into nociceptors. *Nat Biotechnol* **30**, 715–720.
- Chang MY, Son H, Lee YS & Lee SH (2003). Neurons and astrocytes secrete factors that cause stem cells to differentiate into neurons and astrocytes, respectively. *Mol Cell Neurosci* **23**, 414–426.
- Chou SY, Weng JY, Lai HL, Liao F, Sun SH, Tu PH, Dickson DW & Chern Y (2008). Expanded-polyglutamine huntingtin protein suppresses the secretion and production of a chemokine (CCL5/RANTES) by astrocytes. *J Neurosci* **28**, 3277–3290.
- D'Ascenzo M, Piacentini R, Casalbore P, Budoni M, Pallini R, Azzena GB & Grassi C (2006). Role of L-type Ca^{2+} channels in neural stem/progenitor cell differentiation. *Eur J Neurosci* **23**, 935–944.
- Ganguly K, Schinder AF, Wong ST & Poo M (2001). GABA itself promotes the developmental switch of neuronal GABAergic responses from excitation to inhibition. *Cell* **105**, 521–532.
- Hama H, Hara C, Yamaguchi K & Miyawaki A (2004). PKC signaling mediates global enhancement of excitatory synaptogenesis in neurons triggered by local contact with astrocytes. *Neuron* **41**, 405–415.
- Hawkins KE, Sharp TV & McKay TR (2013). The role of hypoxia in stem cell potency and differentiation. *Regen Med* **8**, 771–782.
- HD iPSC Consortium (2012). Induced pluripotent stem cells from patients with Huntington's disease show CAG-repeat-expansion-associated phenotypes. *Cell Stem Cell* **11**, 264–278.
- Jagasia R, Steib K, Englberger E, Herold S, Faus-Kessler T, Saxe M, Gage FH, Song H & Lie DC (2009). GABA-cAMP response element-binding protein signaling regulates maturation and survival of newly generated neurons in the adult hippocampus. *J Neurosci* **29**, 7966–7977.
- Jelitai M, Anderova M, Marko K, Kekesi K, Koncz P, Sykova E & Madarasz E (2004). Role of γ -aminobutyric acid in early neuronal development: studies with an embryonic neuroectodermal stem cell clone. *J Neurosci Res* **76**, 801–811.
- Joannides AJ, Webber DJ, Raineteau O, Kelly C, Irvine KA, Watts C, Rosser AE, Kemp PJ, Blakemore WF, Compston A, Caldwell MA, Allen ND & Chandran S (2007). Environmental signals regulate lineage choice and temporal maturation of neural stem cells from human embryonic stem cells. *Brain* **130**, 1263–1275.
- Johnson MA, Weick JP, Pearce RA & Zhang SC (2007). Functional neural development from human embryonic stem cells: accelerated synaptic activity via astrocyte coculture. *J Neurosci* **27**, 3069–3077.
- Jungebluth P, Alici E, Baiguera S, Le Blanc K, Blomberg P, Bozoky B, Crowley C, Einarsson O, Grinnemo KH, Gudbjartsson T, Le Guyader S, Henriksson G, Hermanson O, Juto JE, Leidner B, Lilja T, Liska J, Luedde T, Lundin V, Moll G, Nilsson B, Roderburg C, Stromblad S, Sutlu T, Teixeira AI, Watz E, Seifalian A & Macchiarini P (2011). Tracheobronchial transplantation with a stem-cell-seeded bioartificial nanocomposite: a proof-of-concept study. *Lancet* **378**, 1997–2004.
- Karls A & Mynlieff M (2015). GABA_B receptors couple to $\text{G}\alpha_q$ to mediate increases in voltage-dependent calcium current during development. *J Neurochem* **135**, 88–100.
- Kovacs CS, Ho-Pao CL, Hunzelman JL, Lanske B, Fox J, Seidman JG, Seidman CE & Kronenberg HM (1998). Regulation of murine fetal-placental calcium metabolism by the calcium-sensing receptor. *J Clin Invest* **101**, 2812–2820.
- Kovacs CS & Kronenberg HM (1997). Maternal-fetal calcium and bone metabolism during pregnancy, puerperium, and lactation. *Endocr Rev* **18**, 832–872.
- Levy M, Boulis N, Rao M & Svendsen CN (2016). Regenerative cellular therapies for neurologic diseases. *Brain Res* **1638**, 88–96.
- Livesey MR, Bilican B, Qiu J, Rzechorzek NM, Haghi G, Burr K, Hardingham GE, Chandran S & Wyllie DJ (2014). Maturation of AMPAR composition and the GABA_AR reversal potential in hPSC-derived cortical neurons. *J Neurosci* **34**, 4070–4075.
- Lukovic D, Stojkovic M, Moreno-Manzano V, Bhattacharya SS & Erceg S (2014). Perspectives and future directions of human pluripotent stem cell-based therapies: lessons from Geron's clinical trial for spinal cord injury. *Stem Cells Dev* **23**, 1–4.
- Mazzanti M & Haydon PG (2003). Astrocytes selectively enhance N-type calcium current in hippocampal neurons. *Glia* **41**, 128–136.
- Merz K, Herold S & Lie DC (2011). CREB in adult neurogenesis—master and partner in the development of adult-born neurons? *Eur J Neurosci* **33**, 1078–1086.
- Mohyeldin A, Garzon-Muvdi T & Quinones-Hinojosa A (2010). Oxygen in stem cell biology: a critical component of the stem cell niche. *Cell Stem Cell* **7**, 150–161.
- Pfrieger FW & Barres BA (1997). Synaptic efficacy enhanced by glial cells in vitro. *Science* **277**, 1684–1687.
- Ravni A, Vaudry D, Gerdin MJ, Eiden MV, Falluel-Morel A, Gonzalez BJ, Vaudry H & Eiden LE (2008). A cAMP-dependent, protein kinase A-independent signaling pathway mediating neuritogenesis through Egr1 in PC12 cells. *Mol Pharmacol* **73**, 1688–1708.

- Riccardi D & Kemp PJ (2012). The calcium-sensing receptor beyond extracellular calcium homeostasis: conception, development, adult physiology, and disease. *Annu Rev Physiol* **74**, 271–297.
- Rivera C, Voipio J, Payne JA, Ruusuvoori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M & Kaila K (1999). The K^+/Cl^- co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* **397**, 251–255.
- Rushton DJ, Mattis VB, Svendsen CN, Allen ND & Kemp PJ (2013). Stimulation of GABA-induced Ca^{2+} influx enhances maturation of human induced pluripotent stem cell-derived neurons. *PLoS One* **8**, e81031.
- Shen J (2014). Function and dysfunction of presenilin. *Neurodegener Dis* **13**, 61–63.
- Song M, Mohamad O, Chen D & Yu SP (2013). Coordinated development of voltage-gated Na^+ and K^+ currents regulates functional maturation of forebrain neurons derived from human induced pluripotent stem cells. *Stem Cells Dev* **22**, 1551–1563.
- Studer L, Csete M, Lee SH, Kabbani N, Walikonis J, Wold B & McKay R (2000). Enhanced proliferation, survival, and dopaminergic differentiation of CNS precursors in lowered oxygen. *J Neurosci* **20**, 7377–7383.
- Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K & Yamanaka S (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* **131**, 861–872.
- Tang X, Zhou L, Wagner AM, Marchetto MC, Muotri AR, Gage FH & Chen G (2013). Astroglial cells regulate the developmental timeline of human neurons differentiated from induced pluripotent stem cells. *Stem Cell Res* **11**, 743–757.
- Telezhkin V, Schnell C, Yarova P, Yung S, Cope E, Hughes A, Thompson BA, Sanders P, Geater C, Hancock JM, Joy S, Badder L, Connor-Robson N, Comella A, Straccia M, Bombau G, Brown JT, Canals JM, Randall AD, Allen ND & Kemp PJ (2016). Forced cell cycle exit and modulation of $GABA_A$, CREB, and GSK3 β signaling promote functional maturation of induced pluripotent stem cell-derived neurons. *Am J Physiol Cell Physiol* **310**, C520–C541.
- Telezhkin V, Thompson BA, Pardo M, Barriga GG, Brown DA, Canals JM, Allen ND & Kemp PJ (2014). $Kv7.2/7.3$ channels are enhanced during striatal development and promote neuronal functional maturation of iPS cell-derived neurons. *Biophys J* **106** (Suppl 1), 142a.
- Telias M, Segal M & Ben-Yosef D (2014). Electrical maturation of neurons derived from human embryonic stem cells. *F1000Res* **3**, 196.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS & Jones JM (1998). Embryonic stem cell lines derived from human blastocysts. *Science* **282**, 1145–1147.
- Toogood PL, Harvey PJ, Repine JT, Sheehan DJ, VanderWel SN, Zhou H, Keller PR, McNamara DJ, Sherry D, Zhu T, Brodfuehrer J, Choi C, Barvian MR & Fry DW (2005). Discovery of a potent and selective inhibitor of cyclin-dependent kinase 4/6. *J Med Chem* **48**, 2388–2406.
- Tozuka Y, Fukuda S, Namba T, Seki T & Hisatsune T (2005). GABAergic excitation promotes neuronal differentiation in adult hippocampal progenitor cells. *Neuron* **47**, 803–815.
- Ullian EM, Christopherson KS & Barres BA (2004). Role for glia in synaptogenesis. *Glia* **47**, 209–216.
- Ullian EM, Sapperstein SK, Christopherson KS & Barres BA (2001). Control of synapse number by glia. *Science* **291**, 657–661.
- Vizard TN, O’Keefe GW, Gutierrez H, Kos CH, Riccardi D & Davies AM (2008). Regulation of axonal and dendritic growth by the extracellular calcium-sensing receptor. *Nat Neurosci* **11**, 285–291.
- Ward JP (2008). Oxygen sensors in context. *Biochim Biophys Acta* **1777**, 1–14.
- Wiegert JS & Bading H (2011). Activity-dependent calcium signaling and ERK-MAP kinases in neurons: A link to structural plasticity of the nucleus and gene transcription regulation. *Cell Calcium* **49**, 296–305.
- Yamada J, Okabe A, Toyoda H, Kilb W, Luhmann HJ & Fukuda A (2004). Cl^- uptake promoting depolarizing GABA actions in immature rat neocortical neurons is mediated by NKCC1. *J Physiol* **557**, 829–841.
- Zarbin M (2016). Cell-based therapy for degenerative retinal disease. *Trends Mol Med* **22**, 115–134.

Additional information

Competing interests

None declared.

Author contributions

All authors have approved the final version of the manuscript and agree to be accountable for all aspects of the work. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

Funding

The authors would like to thank the CHDI Foundation, an FP7 collaborative grant (Repair-HD), and the NC3Rs (Crack-it challenge) for funding parts of this work.