



Published in final edited form as:

Psychosom Med. 2016 ; 78(9): 1114–1119. doi:10.1097/PSY.0000000000000421.

Mechanisms underlying the association between early-life adversity and physical health: Charting a course for the future

Nicole R. Bush, Ph.D.,

University of California San Francisco, Departments of Psychiatry and Pediatrics, Center for Health and Community, Division of Developmental Medicine

Richard D. Lane, M.D., Ph.D., and

University of Arizona, Departments of Psychiatry, Psychology and Neuroscience

Katie A. McLaughlin, Ph.D.

University of Washington, Department of Psychology

Abstract

Early life adversities (ELA) are associated with subsequent pervasive alterations across a wide range of neurobiological systems and psychosocial factors that contribute to accelerated onset of health problems and diseases. In this article, we provide an integrated perspective on recent developments in research on ELA, based on the articles published in this Special Issue of *Psychosomatic Medicine*. We focus on: (1) the distinction between specific versus general aspects of ELA in terms of the nature of exposure (e.g., physical and sexual abuse, emotional abuse or neglect, relative socioeconomic deprivation), biological and behavioral correlates of ELA, and differences across diseases; (2) the importance of timing in the critical phases of exposure to ELA; and (3) adaptive versus dysfunctional responses to ELA and their consequences for biological and behavioral risk factors for adverse health outcomes. This article concludes with outlining important new targets for research in this area, including the neurobiology of affect as mechanism linking ELA to adverse health outcomes, and the need for large-scale longitudinal investigations of multi-system processes relevant to ELA in diverse samples, starting prenatally, continuing to late adolescence, and with long-term follow-up assessments that enable evaluation of incident disease outcomes.

The articles in the special issue demonstrate that increases in physical health problems emerge early in development among children exposed to adversity. These include global indices of health problems and health impairment (1, 2), asthma (3, 4), obesity (5), and pain conditions (6). Papers in this issue also provide evidence that a number of neurobiological mechanisms underlie the associations of adversity experienced in utero and in childhood with physical health problems across the life-course, including inflammatory (7), cardiometabolic (8), epigenetic (9, 10); gene expression (10), and cellular aging pathways (11). It is intriguing that, in contrast to influential theories (e.g. 12), two papers failed to find

Corresponding Author: Nicole R. Bush, UCSF, Center for Health and Community, 3333 California Street, Suite 465, San Francisco, CA 94118, Nicole.bush@ucsf.edu, Phone (415) 476-7655.

Conflicts of Interest: The authors have no conflicts of interest to report.

evidence for cardiovascular mechanisms linking early-life adversity with child and adult physical health outcomes (1, 13). In addition to these neurobiological mechanisms, papers in the special issue highlight the importance of psychosocial mechanisms in the association between early adversity and health. Most notably, multiple papers document a mechanistic role of psychological distress and mental health problems in the link between adversity and health. Psychopathology, particularly anxiety and depression, mediated the association between early-life adversity and postpartum weight retention (3), trajectories of health across childhood and adolescence (2), adult cardiometabolic risk (8), and risk of asthma and allergy in one's offspring (4). Social relationships also partially explained adversity-health associations (3). Not all examinations of social mediators produced positive findings, however. For example, negativity in the parent-child relationship did not mediate the association between family SES and health impairment in young children (1). Finally, the special issue highlights the significance of chronic pain as an outcome associated with early-life adversity beginning early in the life-course (6) and the role of central sensitization as a mechanism explaining this association (14).

Together, these findings reveal pervasive alterations across a variety of neurobiological systems and psychosocial factors among individuals exposed to adverse early environments that, in turn, contribute to accelerated onset of health problems and disease. Yet, the pathways explored here are by no means exhaustive, some papers represent fields in their infancy in terms of bridging social and biological sciences (e.g. social epigenetics), and some papers failed to find support for mechanisms widely argued to play a role in the link between adversity and health despite strong study designs (e.g., (13)). A multitude of fundamental questions about the pathways linking adverse early environments to the onset of disease remain unanswered. We highlight some of the most pressing issues for the field with the goal of stimulating innovative research on these topics.

Are the adversity exposures and mechanisms linking adversity to health universal or specific?

Determining the degree of specificity in terms of exposures, mechanisms, and outcomes in the associations between early-life adversity and health is a critical issue for future research. First, the term adversity is used to refer to a broad range of experiences including child abuse and neglect, parental psychopathology, and poverty. Many studies, including some in this issue, combine these diverse experiences into a cumulative risk index and associate that index with health outcomes and underlying mechanisms. This approach has been advocated to address challenges in measurement, reporting, and statistical modeling of adversity (15). However, this approach has been critiqued for failing to distinguish between diverse experiences that may have differing associations with neurobiological and psychological development that contribute to psychopathology following adversity (16, 17). For example, in some cumulative indices, an individual who experienced a singular experience of sexual abuse by a stranger would have the same adversity score as a child who was chronically abused by a parent for a decade or who had a parent with depression. Emerging evidence suggests that this type of approach clouds specificity in the associations of adversity with emotion, cognition, and at least some of the neurobiological pathways thought to play a role

in the onset of disease, including regulation of stress response systems (18). Indeed, Chen and colleagues demonstrate that although two distinct dimensions of socio-economic status (SES), prestige (parent education) and resources (assets), are each associated with asthma-related clinical outcomes, the mechanisms underlying these associations are distinct such that prestige is associated with better control behaviors within the home and lower exposure to smoke whereas resources are associated with a more advantageous profile of immune regulation (7). The degree to which specificity across different forms of adversity exists in other neurobiological pathways involved in the adversity-health association remains a critical question for the field. Moreover, it is often not feasible to ask the types of detailed questions required for distinguishing adverse experiences, particularly in epidemiological samples. One paper in this issue, however, provides an excellent example of combining detailed assessment with a population-based approach. Baldwin and colleagues demonstrate in a large representative twin birth cohort that childhood bullying, especially when chronic, predicts adolescent overweight, adjusting for a broad range of potential confounds including other adversities such as maltreatment and genetic and fetal liability for overweight (5). Such sophisticated prospective study designs are required to evaluate the degree to which the mechanisms linking heterogeneous forms of adversity with physical health are general versus specific across experiences.

In addition to exposure types, greater attention to specificity in examining mechanistic pathways is needed. Loucks and colleagues' investigation of epigenetic mechanisms linking adversity with adult BMI (9) revealed that childhood SES was associated with adulthood DNA methylation in adipose tissue, but not in blood leukocytes—providing important data for the tissue-specificity debate within the field of social epigenetics and pointing, more broadly, to the importance of investigating specific disease-relevant systems and tissues when examining other neurobiological pathways. Again, the work of Chen and colleagues (7) provides an illustrative example by demonstrating that different dimensions of SES had distinct associations with particular markers of immune regulation in response to specific environmental and corticosteroid triggers but not others. This type of approach has the potential to identify the specific neurobiological mechanisms most relevant for specific exposures, which may ultimately generate innovative targets for early intervention.

Greater specificity would also be useful in the domain of emotional mechanisms, which reflect neurobiological and cognitive processes but are often measured with self-report instruments in the adversity-health literature. Given that many forms of adversity influence emotional development, more research is needed on the *affective* mechanisms linking early adversity and health using novel neuroimaging methodologies. This issue is particularly salient for the field of psychosomatic medicine given that in the Great Debate of 2001, the primary question under consideration was whether emotional processes or stress have direct physiological effects on physical health and whether interventions to improve emotional functioning also improve medical outcomes (19). Atypical patterns of emotional processing are a key mechanism linking early-life adversity with psychopathology (20) but have been investigated less frequently in relation to physical health outcomes. Although several studies in this issue demonstrate the critical role of mental health in linking adversity to physical health, greater specificity in the measurement of affective mechanisms is needed. Given that many forms of early adversity are associated with deficits in emotional awareness and ability

to report on feelings and emotions (21, 22), there is a particular need for future studies to go beyond simple self-reports of affective symptoms in investigating emotional mechanisms linking early adversity with physical health. Advances in functional neuroimaging have provided useful tools for measuring emotional reactivity and regulation, as well other emotional processes. Yet, despite extensive work in the area of adversity and mental health, surprisingly little research has incorporated affective neuroscience measures into the study of mechanisms linking adversity with physical health. This represents an important next step for the field.

A final issue with regard to specificity involves health outcomes. In particular, mental and physical health outcomes are rarely considered simultaneously, precluding progress in identifying mechanisms that are shared across disparate health outcomes and those that are domain-specific. This is surprising, in light of research demonstrating associations between atypical patterns of regulation of physiological stress response systems and both physical and mental health problems (23). A recent epidemiological study examining the associations of a wide range of early-life adversities with both mental and physical health outcomes provides evidence for shared pathways of social support and behavioral factors but also for specificity in the links between particular experiences and specific health outcomes (24). Evaluating the degree to which specific neurobiological, affective, and psychosocial pathways contribute to diverse health outcomes is a critical next step for the field, as innovations in intervention require greater knowledge about shared pathways that could be targeted to prevent a host of adverse health outcomes as well as unique pathways that have relevance to particular outcomes.

Are there sensitive periods for exposure, and does it depend on the system of interest?

A fundamental principle regarding the effects of experience on neurobiological development is that timing matters. Although neural plasticity persists across the life-course, extensive evidence documents that plasticity is heightened during developmental windows in which the nervous system is particularly responsive to certain inputs from the environment. These windows are referred to as critical or sensitive periods (25). Although sensitive periods have been identified with regard to the impact of experience on the development of sensory systems, the degree to which such periods exist within the peripheral nervous system and in other regulatory systems is largely unknown. A recent study, however, documented that psychosocial deprivation early in life exerted pronounced effects on the development of the hypothalamic-pituitary-adrenal (HPA) axis, but only during a sensitive period during the first two years of life (26). Determining whether such periods exist in other systems is critical not only for understanding when and how adversity will have the most potent influences on neurobiological development but also for targeting interventions during periods of heightened plasticity. Relatedly, multiple cascading sensitive periods that unfold over development have been observed in some domains (e.g., language development) (see 27 for discussion) but have yet to be examined in most pathways underlying adversity-health associations. Identifying periods of particular susceptibility to the social environment represents a critical area for future research.

Research on sensitive periods and health is relatively consistent in documenting that exposures occurring very early in development may have more pronounced effects on development than those that occur later. Building upon extant animal literature, Thomfohr-Madsen and colleagues (4) highlight the possibility that exposures during a mother's own childhood can affect her child's risk for developing airway disease, through intergenerational transmission of the effects of adversity. An additional paper in this issue found exposure to maternal depression during the prenatal period—but not during preconception—is most relevant for offspring cortisol regulation (10). The burgeoning fields considering prenatal programming, intergenerational transmission of effects of the social environment, and sensitive periods of development are poised to provide important insights about the etiology of disease opportunities for successful intervention.

Does altered development following adversity reflect adaptation or dysfunction?

It is important to consider that humans are designed to adapt to a broad range of environments to promote survival and ensure reproductive success. Many outcomes of adversity described as dysfunction or dysregulation actually reflect adaptations that are advantageous in an adverse environment, at least in the short-term, despite having pernicious long-term consequences (23, 28). For example, metabolic changes that allow a neonate to respond adaptively to under-nutrition in utero may contribute to enhanced risk for obesity and cardiovascular disease in a calorie-rich environment later in life (29). In addition, some argue that environmentally-shaped enhanced physiological reactivity is not only a risk factor for poor health in risky environments but can promote health and wellbeing in more advantageous environments (see 30 or, 31 for reviews), or that accelerated development provoked by adversity exposure is adaptive evolutionarily by increasing chances for survival and reproduction despite links with earlier onset of disease (32). Complexities in interpreting variation in biological functioning are further exemplified by evidence that associations between adversity and physiological indicators of stress regulation are sometimes non-linear and vary by race/ethnicity (see for example 33). The associations of adversity with early-onset pain conditions (6) and heightened central sensitization, a potential mechanism underlying enhanced pain (14) provide additional examples of adaptations that may confer benefits in an adverse environment in the short-term but result in morbidity and disability in the long-term. In our quest to understand mechanisms, it is critical to attend to the potential benefits conferred by neurobiological and psychosocial adaptations to adverse environments, use caution with labels such as “maladaptive” and “dysfunctional,” and integrate theories and evidence from the multitude of relevant perspectives on developmental variation following adversity.

Where do we go from here?

The papers in this special issue provide a broad illustration of the immense progress in studying mechanisms for the effects of early-life adversity on health across the life-course. Yet, much remains to be accomplished in charting these pathways. Despite increasingly large literature in this area, more large-scale research is needed that contains rich prospective

measurements of early exposures, biological and behavioral mechanisms, brain structure and function as well as multiple aspects of physical health. Many papers studied precursors to disease (e.g., adiposity, cardiometabolic measures, immune processes) or endophenotypes thought to predict variability in health and disease outcomes (e.g., DNA methylation, telomere length) rather than disease outcomes. Additional studies are needed to determine the extent to which the biological embedding is enduring over time and whether such effects mediate broader health consequences.

Another significant challenge for this body of research is to incorporate rigorous methods that can identify causal effects while still capturing the complexity of adverse social experiences (34). To date, experimental evidence for the health effects of adversity have come from animal studies or from naturalistic experiments in humans (e.g., famines, natural disasters, sudden economic shifts). Although their sample size was small, Schnepfer and colleagues used random assignment of primates to parenting environments, increasing confidence in the causal nature of the effects on adult telomere length (11). Experimental manipulation of adversity exposure involves ethical complexities for both animal and human research, and a promising future direction for the field is to examine mechanisms of adversity in the context of randomized preventive interventions. Developmental processes that are altered by intervention and that confer protection against negative health outcomes can be inferred to be causal risk factors for those outcomes (35). Intervention designs are powerful tools that have yet to be widely used for shedding light on mechanisms linking early-life adversity to health. Given the consistent evidence in this issue documenting mental health symptoms as a mechanism linking adversity with physical health outcomes (2–4, 8), intervention designs could be used to determine whether reducing psychological distress and/or treating psychopathology in populations exposed to adversity also confers physical health benefits.

The special issue specifically emphasized mechanisms, but consideration of moderators at multiple levels of influence that confer resilience to physical health problems following adversity exposure is equally important. For example, Hagan and colleagues demonstrate that individual differences in autonomic reactivity moderate the association of SES with child physical health impairment and suggest that better parent-child relationships protect children in low-SES families from poor physical health (1). Identifying modifiable factors that buffer children from the health consequences of adversity may stimulate innovation in preventing the health consequences of adversity by identifying novel targets for early intervention.

Leveraging basic research to develop more effective interventions for children exposed to adversity is the penultimate goal of the research presented in the issue. Although eradicating early-life adversity would likely produce substantial improvements in population health, this is unlikely to occur without substantial policy change (36). In the meantime, identifying mechanisms and protective factors is a particularly useful contribution of basic research in this area because it can contribute to greater knowledge of whether, and how, the negative health consequences of early-life adversity can be reversed or prevented. Relatively little is known currently about how to most effectively target the neurobiological mechanisms highlighted in this issue, although the effects of psychosocial interventions on

neurobiological systems are being studied increasingly and the early evidence for such effects is promising (e.g. 37). The articles in this special issue stimulate progress in this domain because they provide additional foundation for understanding the mechanisms linking early-life adversity to health. These research findings can be used to generate novel strategies for preventing disease across the life-course in individuals who are disproportionately vulnerable to adverse health outcomes.

Acknowledgments

Sources of Funding:

This work was supported by grants from the National Institute of Mental Health (R01-MH103291 and R01-MH106482 to McLaughlin), the National Heart, Lung and Blood Institute (R01 HL116511 to Bush; R01-HL103692 to Lane), the Robert Wood Johnson Foundation Health and Society Scholars Program Health Disparities Working Group (to Bush), a Jacobs Foundation Early Career Research Fellowship (to McLaughlin), and a Rising Star Award grant from AIM for Mental Health, a program of One Mind Institute (IMHRO) (to McLaughlin).

We would also like to thank each of the authors that contributed research to the special issue.

References

1. Hagan MJ, Roubinov DS, Adler NE, Boyce WT, Bush NR. Socioeconomic adversity, negativity in the parent child-relationship, and physiological reactivity: An examination of pathways and interactive processes affecting young children's physical health. *Psychosom Med.* 2016; doi: 10.1097/PSY.0000000000000379
2. Whalen DJ, Belden AC, Tillman R, Barch DM, Luby JL. Early adversity, psychopathology and latent class profiles of global physical health from preschool through early adolescence. *Psychosomatic Medicine.*
3. Luecken LJ, Jewell SL, MacKinnon DP. Prediction of postpartum weight in low-income mexican-origin women from childhood experiences of abuse and family conflict. *Psychosom Med.* 2016; doi: 10.1097/PSY.0000000000000391
4. Tomfohr LM, Bayrampour H, Tough S. Maternal history of childhood abuse and risk of asthma and allergy in 2-year-old children. *Psychosomatic Medicine.*
5. Baldwin JR, Arseneault L, Odgers C, Belsky DW, Matthews T, Ambler A, Caspi A, Moffitt TE, Danese A. Childhood bullying victimization predicts overweight in young adulthood: A cohort study. *Psychosomatic Medicine.* 2016 In press.
6. McLaughlin KA, Basu A, Walsh K, Slopen N, Sumner JA, Koenen KC, Keyes KM. Childhood exposure to violence and chronic physical conditions in a national sample of us adolescents. *Psychosomatic Medicine.* 2016; doi: 10.1097/psy.0000000000000366
7. Chen E, Shalowitz MU, Story RE, Ehrlich KB, Levine CS, Hayen R, Leigh AKK, Miller GE. Dimensions of socioeconomic status and childhood asthma outcomes: Evidence for distinct behavioral and biological associations. *Psychosomatic Medicine.* In Press.
8. Winning A, Glymour M, McCormick M, Gilsanz P, Kubzansky L. Childhood psychological distress as a mediator in the relationship between early-life social disadvantage and adult cardiometabolic risk: Evidence from the 1958 british birth cohort. *Psychosomatic Medicine.*
9. Loucks EB, Yen-Tsung Huang Y, Agha G, Eaton CB, Gilman SE, Buka SL, Kelsey KT. Epigenetic mediators between childhood socioeconomic disadvantage and mid-life body mass index: The new england family study. *Psychosom Med.* In press.
10. Stroud LR, Papandonatos GD, Parade SH, Salisbury AL, Phipps M, Lester B, Padbury JF, Marsit CJ. Prenatal major depressive disorder, placental glucocorticoid and serotonergic signaling pathways, and neonatal stress response. *Psychosomatic Medicine.*
11. Schnepfer LM, Brooks-Gunn J, Notterman DA, Suomi SJ. Early life experiences and telomere length in adult rhesus monkeys: An exploratory study. *Psychosomatic Medicine.* In press.

12. Lovallo WR, Gerin W. Psychophysiological reactivity: Mechanisms and pathways to cardiovascular disease. *Psychosomatic Medicine*. 2003; 65:36–45. [PubMed: 12554814]
13. de Rooij SR, Jones A, Phillips DI, Osmond C, Karemaker JM, Roseboom TJ, Painter RC. Prenatal undernutrition and autonomic function in adulthood. *Psychosom Med*. 2016; doi: 10.1097/PSY.0000000000000393
14. You DS, Meagher MW. Childhood adversity and pain sensitization. *Psychosomatic Medicine*.
15. Evans GW, Li D, Whipple SS. Cumulative risk and child development. *Psychological bulletin*. 2013; 139:1342–96. [PubMed: 23566018]
16. McLaughlin KA, Sheridan MA. Beyond cumulative risk: A dimensional approach to childhood adversity. *Current Directions in Psychological Science*. 2016; 25:239–45. [PubMed: 27773969]
17. McLaughlin KA, Sheridan MA, Lambert HK. Childhood adversity and neural development: Deprivation and threat as distinct dimensions of early experience. *Neuroscience and biobehavioral reviews*. 2014; 47:578–91. [PubMed: 25454359]
18. Busso DS, McLaughlin KA, Sheridan MA. Dimensions of adversity, physiological reactivity, and externalizing psychopathology in adolescence: Deprivation and threat. *Psychosom Med*. 2016; doi: 10.1097/PSY.0000000000000369
19. Freedland KE, Miller GE, Sheps DS. The great debate, revisited. *Psychosomatic Medicine*. 2006; 68:179–84. [PubMed: 16554380]
20. McLaughlin KA. Future directions in childhood adversity and youth psychopathology. *J Clin Child Adolesc Psychol*. 2016; 45:361–82. [PubMed: 26849071]
21. Colvert E, Rutter M, Kreppner J, Beckett C, Castle J, Groothues C, Hawkins A, Stevens S, Sonuga-Barke EJ. Do theory of mind and executive function deficits underlie the adverse outcomes associated with profound early deprivation?: Findings from the english and romanian adoptees study. *Journal of abnormal child psychology*. 2008; 36:1057–68. [PubMed: 18427975]
22. Lane RD, Weihs KL, Herring A, Hishaw A, Smith R. Affective agnosia: Expansion of the alexithymia construct and a new opportunity to integrate and extend freud’s legacy. *Neuroscience and biobehavioral reviews*. 2015; 55:594–611. [PubMed: 26054794]
23. McEwen BS. Brain on stress: How the social environment gets under the skin. *Proc Natl Acad Sci U S A*. 2012; 109(Suppl 2):17180–5. [PubMed: 23045648]
24. Sheikh MA, Abelsen B, Olsen JA. Clarifying associations between childhood adversity, social support, behavioral factors, and mental health, health, and well-being in adulthood: A population-based study. *Front Psychol*. 2016; 7:727. [PubMed: 27252668]
25. Hensch TK. Critical period plasticity in local cortical circuits. *Nat Rev Neurosci*. 2005; 6:877–88. [PubMed: 16261181]
26. McLaughlin KA, Sheridan MA, Tibu F, Fox NA, Zeanah CH, Nelson CA 3rd. Causal effects of the early caregiving environment on development of stress response systems in children. *Proc Natl Acad Sci U S A*. 2015; 112:5637–42. [PubMed: 25902515]
27. Werker JF, Hensch TK. Critical periods in speech perception: New directions. *Annu Rev Psychol*. 2015; 66:173–96. [PubMed: 25251488]
28. Belsky J, Schlomer GL, Ellis BJ. Beyond cumulative risk: Distinguishing harshness and unpredictability as determinants of parenting and early life history strategy. *Developmental psychology*. 2012; 48:662–73. [PubMed: 21744948]
29. Godfrey KM, Gluckman PD, Hanson MA. Developmental origins of metabolic disease: Life course and intergenerational perspectives. *Trends Endocrinol Metab*. 2010; 21:199–205. [PubMed: 20080045]
30. Bush, NR.; Boyce, WT. Differential sensitivity to context: Implications for developmental psychopathology. In: Cicchetti, D., editor. *Developmental psychopathology*. Hoboken, NJ: John Wiley and Sons, Inc; 2015.
31. Ellis BJ, Boyce WT, Belsky J, Bakermans-Kranenburg MJ, van Ijzendoorn MH. Differential susceptibility to the environment: An evolutionary-neurodevelopmental theory. *Dev Psychopathol*. 2011; 23:7–28. [PubMed: 21262036]
32. Ellis, BJ.; Del Giudice, M.; Shirtcliff, EA. The adaptive calibration model of stress responsivity: Concepts, findings, and implications for developmental psychopathology. In: Beauchaine, TP.; Hinshaw, SP., editors. *Child and adolescent psychopathology*. 3. New York: Wiley; in press

33. Bush NR, Obradovic J, Adler N, Boyce WT. Kindergarten stressors and cumulative adrenocortical activation: The “first straws” of allostatic load? *Dev Psychopathol.* 2011; 23:1089–106. [PubMed: 22018083]
34. Adler N, Bush NR, Pantell MS. Rigor, vigor, and the study of health disparities. *Proc Natl Acad Sci U S A.* 2012; 109(Suppl 2):17154–9. [PubMed: 23045672]
35. Cicchetti D, Hinshaw SP. Editorial: Prevention and intervention science: Contributions to developmental theory. *Dev Psychopathol.* 2002; 14:667–71. [PubMed: 12549698]
36. Shonkoff JP. Leveraging the biology of adversity to address the roots of disparities in health and development. *Proc Natl Acad Sci U S A.* 2012; 109(Suppl 2):17302–7. [PubMed: 23045654]
37. Slopen N, McLaughlin KA, Shonkoff JP. Interventions to improve cortisol regulation in children: A systematic review. *Pediatrics.* 2014; 133:312–26. [PubMed: 24420810]