

Overexpression of a Pathogenesis-Related Protein 10 Enhances Biotic and Abiotic Stress Tolerance in Rice

Jingni Wu^{1,2}, Sang Gon Kim³, Kyu Young Kang², Ju-Gon Kim⁴, Sang-Ryeol Park⁵, Ravi Gupta⁶, Yong Hwan Kim⁷, Yiming Wang^{1*}, and Sun Tae Kim^{6*}

¹Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, Cologne 50829, Germany

²Division of Applied Life Science, Gyeongsang National University, Jinju 52828, Korea

³National Institute of Crop Science, Rural Development Administration, Suwon 16429, Korea

⁴College of Agriculture and Life Sciences, Seoul National University, Pyeongchang 25354, Korea

⁵National Institute of Agricultural Science, Rural Development Administration, Jeonju 54875, Korea

⁶Department of Plant Bioscience, Life and Industry Convergence Research Institute, Pusan National University, Miryang 50463, Korea

⁷College of Life and Resource Science, Dankook University, Cheonan 31116, Korea

(Received on June 9, 2016; Revised on July 31, 2016; Accepted on August 4, 2016)

Pathogenesis-related proteins play multiple roles in plant development and biotic and abiotic stress tolerance. Here, we characterize a rice defense related gene named “jasmonic acid inducible pathogenesis-related class 10” (*JIOsPR10*) to gain an insight into its functional properties. Semi-quantitative RT-PCR analysis showed up-regulation of *JIOsPR10* under salt and drought stress conditions. Constitutive over-expression *JIOsPR10* in rice promoted shoot and root development in transgenic plants, however, their productivity was unaltered. Further experiments exhibited that the transgenic plants showed reduced susceptibility to rice blast fungus, and enhanced salt and drought stress tolerance as compared to the wild type. A comparative proteomic profiling of wild type and transgenic plants showed that overexpression of *JIOsPR10* led to

the differential modulation of several proteins mainly related with oxidative stresses, carbohydrate metabolism, and plant defense. Taken together, our findings suggest that *JIOsPR10* plays important roles in biotic and abiotic stresses tolerance probably by activation of stress related proteins.

Keywords : abiotic stress, *JIOsPR10*, *Magnaporthe oryzae*, proteomics, rice

*Co-corresponding authors.

Y Wang

Phone) +49-221-5062-337, FAX) +49-221-5062-353

E-mail) ywang@mpipz.mpg.de

ST Kim

Phone) +82-55-350-5505, FAX) +82-55-350-5509

E-mail) stkim71@pusan.ac.kr

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Plants are continuously exposed to various biotic and abiotic stresses that results in significant decrease in their productivity. During the course of evolution, plants have developed multiple pathways that allow them to sense and adapt these environmental stresses (Gupta et al., 2015). Plant's responses to stress include activation of ion channels, production of reactive oxygen species (ROS) scavenging enzymes, accumulation of hormones, and expression of stress tolerance genes (Fraire-Velázquez et al., 2011). During multiple stresses, plants lead to a complex interaction and crosstalk with multiple signals, and decide the final output of stress-tolerance related genes (Rizhsky et al., 2004). Therefore, plenty of genes are likely to function together and play multiple roles against combined environmental stresses.

Among identified proteins, the pathogenesis-related (PR) family proteins are a highly conserved protein fam-

ily which is involved in plant immune responses (Stintzi et al., 1993). Overexpression of PR proteins has ability to enhance plants tolerance to pathogen infections. For instance, overexpression of PR-5 protein in rice enhances tolerance to *Rhizoctonia solani* (Datta et al., 1999), while overexpression of pepper *PR-1* in tobacco enhances host tolerance to *Phytophthora nicotianae*, *Ralstonia solanacearum*, and *Pseudomonas syringae* (Sarowar et al., 2005). Those PR proteins have also been reported to have multiple roles in adaption to abiotic stresses, like express pepper PR-1 protein in tobacco can enhance heavy metal tolerance (Sarowar et al., 2005). Moreover, transgenic tobacco overexpressing a *Trichoderma harzianum* induced endochitinase enhances the tolerance to both biotic (*P. syringae*), and abiotic (salt and heavy metals) stresses (de las Mercedes Dana et al., 2006). Therefore, study and utilization of PR proteins, which functions in both biotic and abiotic tolerance, may provide new challenges in agriculture.

The PR proteins have been classified into 17 classes based on their amino acid sequence, serological relationship, and biological activities (van Loon et al., 1994; van Loon and van Strien, 1999). Among them, the PR-10 family proteins are involved in multiple anti-pathogen processes, and are generally localized in the intracellular spaces (van Loon and van Strien, 1999). PR-10 family proteins have conserved N-terminal sequences, but very divergent in the C-terminus (Fernandes et al., 2013). The PR10 family proteins consist of three α -helices and seven antiparallel β -strands. Those structure elements enclose a large hydrophobic cavity which is most likely related with their functional relevance (Fernandes et al., 2013). Several studies have shown that PR-10 proteins contain RNase activities (Chen et al., 2007; Kim et al., 2008b; Krishnaswamy et al., 2011; Xie et al., 2010), and cysteine protease inhibitor activity (Andrade et al., 2010).

Previous studies have revealed that PR-10 family proteins are involved in anti-biotic stresses, such as anti-fungus (Chen et al., 2007; Xie et al., 2010), anti-bacteria (Flores et al., 2002; Xie et al., 2010), anti-virus (Park et al., 2004), and anti-nematode (Andrade et al., 2010). To date, four different PR10 family genes have been characterized in rice: *PR10a* (McGee et al., 2001), *PR10b* (McGee et al., 2001), jasmonic acid inducible pathogenesis-related class 10 (*JIOsPR10*) (Jwa et al., 2001), and *RSOsPR10* (Hashimoto et al., 2004). All those PR10 family genes are induced by *Magnaporthe oryzae* infection and jasmonic acid treatment (Hashimoto et al., 2004; Jwa et al., 2001; McGee et al., 2001), suggesting that those PR10 family genes may be functional redundant in rice. Moreover, phytohormone salicylic acid could activate the transcriptional expression of leaf accumulated *PR10a* and

JIOsPR10, but not the root specific *RSOsPR10* (Hashimoto et al., 2004), indicating that *PR10* gene regulation in root and shoot may have different mechanisms. Moreover, rice *PR10* genes are also induced by various abiotic stresses, such as cold, salt and drought (Hashimoto et al., 2004; Kim et al., 2008b), suggesting that PR10 protein may have multiple function in tolerance to both biotic and abiotic stresses. In this study, transgenic rice constitutively overexpressing *JIOsPR10* was constructed. The transgenic plants showed a promotion in root and shoot development, and enhance tolerance to rice blast fungus, drought and salt stresses. Proteomics analysis reveals that plant defense related proteins as well as ROS related proteins were also highly accumulated in overexpression plant. These results implied that *JIOsPR10* play a role in multiple stress tolerance, which may important for future agricultural applicants.

Materials and Methods

Plant material, growth conditions, and stress treatment.

Rice seeds (*Oryzae sativa* cv. Dongjin) were sterilized with 70% ethanol for 10 min, then in 3% sodium hypochlorite for 30 min, followed by washing with sterilized water for three times. Sterilized seeds were imbibed in water at 4°C in dark for 3 days, and allowed to germinate in standard rice soil (Monsanto, Seoul, Korea) in greenhouse or on Murashige & Skoog (MS) phytagel medium at 28°C growth chamber (light/dark time, 16/8 h). For drought and salt stress treatments, 10-day-old seedlings were transferred into container containing either 20% polyethylene glycol (PEG) 4000 or 150 mM NaCl solution. For drought stress to 5- to 6-leaf stages, pots were kept on dried papers to remove water from the pots, and plants were re-watered after 5 days. For salt stress to 4- to 5-leaf stage plants, rice in growth pots were transferred to a container filled with 150 mM NaCl. NaCl solution was changed each two days to maintain the concentration of salt stress.

Fungal spore preparation and infection. *M. oryzae* conidia (KJ301) was cultured on rice-bran agar medium (25 g/l rice bran, 1 g/l sucrose, and 20 g/l agar). Aerial mycelia were removed under fluorescent light to induce synchronous conidia as described previously (Kim et al., 2013). Conidial suspension (1×10^6 conidia/ml) was inoculated to rice leaves using an air sprayer. Inoculated plants were kept in a humidity chamber at 28°C. Leaf phenotypes were confirmed at 72-h post-inoculation, and the infection area were measured by Assess 2.0 software (The American Phytopathological Society Press, St. Paul, MN, USA).

Vector construction, plant transformation, and genotyping. Full length of *JIOsPR10* coding sequence was cloned into pDONR201 vector and sequenced. Then, the *JIOsPR10* was constructed into pMJ101 vector containing a rice cytochrome c promoter (*OsCcl* pro) and basta selection marker (BAR) by using Gateway system. The construct was transformed into *Agrobacterium* LBA4404, and introduced into rice embryonic calli by selection on MS medium contain 50 μ M phosphinothricin. Regenerated plants were transferred to soil pots and grown in greenhouse. Genotyping of these plants were carried out by PCR with specific primer sets (Supplementary Table 1).

Southern blot analysis. Genomic DNA was extracted from 10-day-old seedlings of wild type (WT) and *JIOsPR10* transgenic lines. The genomic DNA was digested with *Eco*RI, electrophoresed on a 0.8% agarose gel and then blotted onto a nylon membrane (NEN Life Science, Boston, MA, USA). BAR gene in the binary vector was amplified by PCR and used as probe. BAR fragment was labeled with [α - 32 P]-dCTP by using Prime-A-Gene Labeling System (Promega, Madison, WI, USA), and hybridized with the blotted membrane. The membrane was exposed to X-ray film (Fuji Film, Tokyo, Japan) for 3 days at -80° C.

RNA extraction and RT-PCR. Total RNA was extracted from the leaves of WT and transgenic lines using TRIzol reagent (ThermoFisher, Darmstadt, Germany). Extracted RNA was treated with DNase and precipitated with 2.5 vol of ethanol and 1/10 vol of 4 M sodium acetate. Five micrograms of RNA was used for cDNA synthesis by Superscript II reverse transcriptase (Invitrogen, Carlsbad,

CA, USA). Gene specific primers were designed by Primer Designer 4 (Sci Ed Central, Cary, NC, USA), and summarized in Supplementary Table 1. The equal amount of cDNA was optimized by *OsActin2* (Os10g36650) primer sets (He et al., 2009).

Measurement of water loss. The water loss from detached leaves of WT and transgenic lines was measured by monitoring the fresh weight loss at the indicated time points (Duan et al., 2012). For each time point, at least twenty independent leaves from each genotype were used for the measurement of fresh and dried weights.

Protein extraction, proteomics analysis, and protein identification. Total leaf proteins were extracted from three independent biological samples and subjected to PEG fractionation as described previously (Kim et al., 2001; Wang et al., 2014). Briefly, leaf powder was homogenized in the Mg/NP-40 extraction buffer (0.5 M Tris-HCl [pH 8.3], 2% [v/v] NP-40, 20 mM MgCl₂, and 2% [v/v] β -mercaptoethanol) and centrifuged to obtain clear supernatant. Supernatant layer was then precipitated by 15% PEG solution and stand by on ice for 30 min. Supernatant was collected after centrifugation at 12,000 rpm for 15 min, and precipitated with 4 volume of methanol contain 0.1 M ammonium acetate. The 2-DE gels were performed as described previously (Kim et al., 2008a). Briefly, 250 μ g of proteins of each samples were dissolved in the rehydration buffer 9 M (w/v) CO(NH₂)₂, 2% (w/v) 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate hydrate (CHAPS), 0.002% (w/v) bromophenol blue, 20 mM dithiothreitol (DTT), and 0.5% (v/v) pharmalyte (pH 5–8) and were loaded on 18 cm IPG




Fig. 1. Transcriptional regulations of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) under drought (A) and salt (B) stresses. Total RNAs were isolated from 10-day-old seedlings of wild type plant at indicated time points after treatment of 150 mM NaCl (A) or 20% polyethylene glycol (PEG) 4000 (B). Rice actin1 was set as internal control. Error bars represent standard error of three biological replicates of samples.

strips (pH 4–7) by rehydration loading for 12 h. Second dimensional separation was carried out on 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and the gels were stained with Coomassie brilliant blue G-250 (CBB). The stained gels were scanned using transmissive scanner (Powerlook III; UMAX, Fort Worth, TX, USA), and images were analyzed with ImageMaster 2D Platinum software 6.0 (Amersham Biosciences AB, Uppsala, Sweden). Protein spots were excised from CBB-stained gels, digested with trypsin, and extracted with method describes previously (Kim et al., 2008a). Extracted peptide mixture was re-dissolved in dissolve solution (distilled water:acetonitrile:trifluoacetic acid = 93:5:2), sonicated for 5 min, and spin down at 14,000 rpm for 2 min. The peptide mixture was dropped on MALDI plate in matrix solution α -cyano-4-hydroxycinnamic acid (Sigma, St. Louis, MO, USA) in acetone (40 mg/ml). The MALDI plate was then washed with 0.1% (v/v) TFA. The gel spots were analyzed using a Voyager-DE STR MALDI-TOF mass spectrometer (PerSeptiveBiosystems, Framingham, MA, USA). Database searches were performed using Mascot (<http://www.matrixscience.com>) and ProteinProspector (<http://prospector.ucsf.edu/prospector/mshome.htm>) as described previously (Kim et al., 2008a).

Western blot analysis. Equal amount of total leaf proteins were separated on 12% SDS-PAGE and transblotted to a PVDF membrane using semidry electrophoretic apparatus (Hoefer, Holliston, MA, USA). The transferred PVDF membranes were blocked for 4 h at room temperature in $1\times$ TTBS buffer (50 mM Tris-HCl, pH 8.2, 0.1% v/v Tween 20, and 150 mM NaCl) containing 7% skim milk. Membranes were then incubated with primary antibodies of DUF26-like protein, Beta-1,3-glucanase and Cu/Zu-SOD, which were generated as described previously (Kim et al., 2014), at 1,000-fold dilution for 1 h at room temperature. Treated membranes were then incubated with a secondary antibody (anti-rabbit IgG conjugated with horseradish peroxidase; 5,000-fold dilution) for 2 h. Membranes were washed with $1\times$ TTBS (Tris-buffered saline, 0.1% Tween 20) for 15 min, 3 times. Washed membranes were then subjected to protein detection by ECL (Perkin Elmer Life Sciences, Boston, MA, USA) using LAS4000 image system (Fujifilm, Tokyo, Japan).

Results

JIOsPR10 is induced under drought and salt stresses.

Previous report showed that *JIOsPR10* protein was induced by drought and salt stress in rice (Kim et al., 2008b). To understand whether *JIOsPR10* transcripts

were induced in response to salt and drought stress, rice seedlings of WT were exposed to drought (20% PEG) and salt (150 mM NaCl) solutions and semi-quantitative RT-PCR analysis was carried out. *JIOsPR10* transcripts were induced after 60 min and 30 min upon exposure to drought and salt stress, respectively, and increased in a time dependent manner (Fig. 1).

Molecular Characterization of transgenic rice over-expressing *JIOsPR10*.

The role of *JIOsPR10* in plant defense is well established (Kim et al., 2004). Our results also showed induction of *JIOsPR10* transcripts by drought and salt stress, which prompted us to analyze the biological functions of this gene in both biotic and abiotic stress tolerance. The *JIOsPR10* cDNA was constructed under constitutive promoter (Fig. 2A), and introduced




Fig. 2. Characterization of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) overexpression plants. (A) Schematic diagram of *JIOsPR10* under rice cytochrome c promoter with basta selection marker was used for transformation. RB, right border; OsCc1 pro, rice cytochrome c promoter; TpinII, pinII terminator; BAR, basta resistance marker; Tnos, nos terminator; LB, left border. (B) Genotyping of overexpression plants by PCR with Basta selection marker specific primers. M, DNA marker; EV, empty vector plasmid set as positive control; WT, wild type. (C) Southern blot analysis of *JIOsPR10* overexpression plants with BAR probe. (D) RT-PCR analysis of *JIOsPR10* expression in WT and overexpression plants. Number indicate the relative expression level of *JIOsPR10* compare to *actin2*. (E) Western blot analysis of *JIOsPR10* in WT and overexpression plants with *JIOsPR10* specific antibody. Equal loading of protein samples was detected by PonceauS staining.

into rice via *Agrobacterium*-mediated transformation. Two independent transgenic lines, Ox-1 and Ox-3, were produced and confirmed by PCR after Basta selection. A

500 bp PCR band was detected in transgenic plant, so as the constructed vector plasmid which set as positive control (Fig. 2B). Southern blot analysis indicated that both




Fig. 3. Overexpression of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) promotes root and shoot growth. Shoot (A) and root (B) phenotypes of 7-day-old seedlings. (C) Average shoot and root length of wild type (WT) and *JIOsPR10* overexpression plants. (D) Seed weight of WT and *JIOsPR10* overexpression plants. Error bars represent standard error of 20 individual plants. * $P < 0.01$, ** $P < 0.001$.




Fig. 4. Overexpression plant of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) enhances drought and salt stress tolerance. (A) Phenotype of wild type (WT) and *JIOsPR10* overexpression plants re-watered after 5 days under drought condition. (B) Leaf water loss of WT and overexpression plants. Error bars represent standard error of samples. (C) Phenotype of WT and *JIOsPR10* overexpression under salt stress (150 mM NaCl). (D) Close-up view of leaf phenotype under salt stress. At least 20 independent plants were measured for each experiment.

transgenic lines have single-copy T-DNA insertion (Fig. 2C). Transcriptional expression of *JIOsPR10* in transgenic lines was then confirmed by semi-quantitative PCR, the results of which showed 4.9- and 2.6-fold increase of *JIOsPR10* transcripts in Ox-1 and Ox-3 plant than WT plant (Fig. 2D). Subsequently, the protein level of *JIOsPR10* was also detected by Western blot analysis which was 25 folds higher in each transgenic line, compared to the WT (Fig. 2E).

Overexpression of *JIOsPR10* promotes rice development. At first, the growth phenotype of transgenic plants was analyzed. Compared to the WT, transgenic lines showed a growth promotion at early developmental stages (Fig. 3A, B). Root and shoot lengths were 39% and 18% higher in Ox-1, and 34% and 29% higher in Ox-3 in comparison with WT at the 7-day-old seedling stage (Fig. 3C). In addition, seed weight and total productivity was also measured which did not show any significant differences in transgenic lines as compared to the WT (Fig. 3D).

***JIOsPR10* overexpression lines increase drought and salt stress tolerance.** To analyze whether *JIOsPR10* overexpression enhances abiotic stress tolerance in transgenic rice, the Ox-1 and Ox-3 lines were exposed to drought and salt stresses (Fig. 4). Over-expression lines (Ox-1 and Ox-3) have similar phenotypes in drought and salt stress-

es. For drought stress treatment, 4- to 5-week-old plants in growth pot were dried on paper in greenhouse, and re-watered after 5 days. Ox-1 and Ox-3 plants showed successful growth recovery, but not WT (Fig. 4A). The leaf water loss was measured in WT, Ox-1, and Ox-3 lines at different treated time points. In WT, a water loss to nearly 30% of leaf fresh weight was detected under drought stress at 72-h post treatment (hpt) (Fig. 4B). Comparably, the water loss rate in Ox-1 and Ox-3 lines was less than 10% at 72 hpt (Fig. 4B). Salt stress treatment was also performed on *JIOsPR10*-overexpression plants. The growth pots were incubated in 150 mM NaCl solution and phenotype were measured after 24 h. Overexpression lines showed enhanced salt stress tolerance as compared with the WT (Fig. 4C), and a significant leaf curling was detected after 24 hpt in WT, but not in Ox-1 and Ox-3 plants (Fig. 4D).

Differential protein expression profile in WT and *JIOsPR10* overexpression plant. Our results also showed that both Ox-1 and Ox-3 lines showed very similar response to both drought and salt stresses, indicating the higher accumulation of *JIOsPR10* in Ox-3 lines is sufficient for stress tolerance. Therefore, we selected the Ox-3 line for proteomic analysis. Total protein profiles of both WT and *JIOsPR10* overexpression plants were analyzed by 2-DE analysis. The 2-DE gels were stained by CBB, and the differentially expressed protein spots were detect-




Fig. 5. Proteomic analysis of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) overexpression plants showing a different protein expression profile as compared to the wild type (WT). (A) Gel map of 2-DE analysis. Same amount of total rice leaf proteins were separated on 2-DE gels, and stained with Coomassie brilliant blue. Arrows indicate the differentially expressed protein spots. (B) On-gel intensity of each protein spots in WT and overexpression plants. Error bars represent standard error of three biological replicates of experiments. *pI*, isoelectric point.

Table 1. Identification of differentially modulated protein spots in wild type and *JIOsPR10* overexpression plants using MALDI-TOF mass spectrometer

Spot No.	Protein name	Accession No.	Score	Database	MP	SC (%)	Ex Mr (kDa)	Th Mr (kDa)	Ex pI	Th pI	Biological function
1	Thaumatococin-like protein	ACA50508	479	Prospector	3	21.0	18.0	18.1	4.4	6.5	Defense
2	DUF26-like protein	AAW34053	475	Prospector	6	38.4	24.4	27.3	4.7	5.0	Defense
3	Beta-1,3-glucanase	BAA77783	422	Prospector	4	15.9	32.2	34.2	4.6	4.6	Carbohydrate metabolic process
4	Peroxidase	CAA46916	86	Mascot	7	29.0	37.1	32.9	5.2	5.8	Response to oxidative stress
5	Enolase	ABB46862	374	Prospector	13	32.0	48.0	48.0	5.5	5.4	Glycolytic process
6	Nucleoside diphosphate kinase	AAT08712	396	Prospector	4	27.0	18.5	17.7	5.4	9.1	Energy
7	Formate dehydrogenase	NP_001057666	112	Mascot	8	32.0	36.6	41.2	6.4	6.9	Response to oxidative stress
8	Cu/Zn-superoxide dismutase	BAB21760	228	Prospector	7	42.0	41.4	21.3	6.6	5.8	Response to oxidative stress
9	Xylanase inhibitor protein 1	NP_001057599	65	Mascot	5	24.0	44.6	27.4	6.9	6.4	Carbohydrate metabolic process
10	Cytoplasmic malate dehydrogenase	AAG13573	372	Prospector	8	21.0	32.9	35.6	6.5	5.7	Tricarboxylic acid cycle

JIOsPR10, jasmonic inducible pathogenesis-related class 10; MP, matched peptides; SC, sequence coverage; Ex Mr, experimental molecular mass; Th Mr, theoretical molecular mass; Ex pI, experimental isoelectric point; Th pI, theoretical isoelectric point.

ed (Fig. 5A, indicated by arrows). Five protein spots (spot 1, 2, 3, 4, and 8) were up-regulated in *JIOsPR10* overexpression plants (Fig. 5B), including thaumatin-like protein (TLP), DUF26-like protein, beta-1,3-glucanase, peroxidase, and Cu/Zn-SOD (Table 1). Five protein spots, which encoding enolase (spot 5), nucleoside diphosphate kinase (NDK, spot 6), formate dehydrogenase (spot 7), xylanase inhibitor protein 1 (spot 9), and cytoplasmic malate dehydrogenase (spot 10), were down-regulated in Ox-3 line than in WT (Fig. 5B, Table 1). Western blot analysis using gene specific antibody of DUF26-like protein, beta-1,3-glucanase, and Cu/Zn-SOD was employed to confirm the differential expression of identified proteins in Ox-1 and Ox-3 lines. The protein accumulation of all three proteins in Ox-1 and Ox-3 was strongly increased comparing with WT (Fig. 6), indicating a good correlation of proteomics data.

JIOsPR10 enhances tolerance to rice blast fungus.

JIOsPR10 is a member of PR family genes, which provide biotic stress tolerance in rice. Therefore, the response of *M. oryzae* infection on transgenic lines was investigated. Spores of *M. oryzae* were sprayed on the leaves of WT and transgenic lines and the infection phenotype was analyzed after 72 h of spore inoculation. The number of infection lesions of *M. oryzae* was reduced in Ox-1 and Ox-3 lines in comparison with the WT (Fig. 7A). Average infected area on the leaves of Ox-1 and Ox-3 lines was reduced to 22.96% and 13.6%, respectively, from 39.72%, calculated in the WT (Fig. 7B).




Fig. 6. Overexpression of jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) induces highly accumulation of reactive oxygen species related protein. Same amount of protein was loaded on sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Antibody against DUF26-like protein, beta-1,3-glucanase, and Cu/Zn-SOD were used to detect the protein level expression in wild type (WT) and *JIOsPR10* overexpression plant. Equal loading of protein samples was checked by PonceauS staining.




Fig. 7. Jasmonic inducible pathogenesis-related class 10 (*JIOsPR10*) overexpression plants showing increased tolerance to rice blast fungus. (A) Phenotype of wild type (WT) and *JIOsPR10* overexpression plants infected with rice blast fungus at 3 days post inoculation (dpi). (B) Average percentage of infected area on WT, Ox-1, and Ox-3 leaves. Student's *t*-test was applied on the data analysis. * $P < 0.01$; ** $P < 0.05$.

Discussion

Crop productivity is strongly affected by the biotic and abiotic stresses in surrounding environments. Previous studies have shown that overexpression of stress responsive genes in plant may improve its adaption to various stresses and increase the yield (Li et al., 2014; Shi et al., 2014). Previously, *JIOsPR10* was reported to be involved in the biotic stress tolerance, such as *M. oryzae* infections (Kim et al., 2004). In this study, we showed that *JIOsPR10* was also induced by abiotic stresses, including drought and salt, suggesting *JIOsPR10* has multiple functions in rice and might be important for agricultural usage (Fig. 1).

Previous studies have shown that PR10 family protein has ribonuclease (Bantignies et al., 2000; Zhou et al., 2002) and cysteine protease inhibition activities (Andrade et al., 2010). Rice *JIOsPR10* was also reported to contain RNase activity, suggesting that biochemical activity might be essential for the role of *JIOsPR10* (Kim et al., 2008b). Therefore, characterization of downstream regulated proteins contributes the understanding of molecular mechanism of *JIOsPR10* induced stress tolerance. Proteomics, consists of protein separation and identification steps, is a widely used tool for large-scale identification of proteins. To investigate the mechanism of how *JIOsPR10* promotes rice growth, and enhances stress tolerance, we therefore applied a 2-DE based comparative proteomic analysis of WT and *JIOsPR10* overexpression plants. To increase the detection of low abundance proteins, a PEG mediated protein fractionation method, which depletes high-abundant proteins such as Rubisco, was used for protein extraction (Kim et al., 2001). The protein expression profile was observed by 2-DE gel analysis coupled with CBB staining. Differentially modulated protein spots from gel map of WT and transgenic plants were selected and identified by MALDI-TOF mass spectrometer, which revealed that proteins involved in metabolism, defense,

and oxidative stress were predominant. In particular, several proteins related with glycolytic process (enolase, spot 5), tricarboxylic acid cycle (cytoplasmic malate dehydrogenase, spot 10), and energy (nucleoside diphosphate kinase, spot 6) were down regulated in transgenic plants, indicating a down regulation of metabolism inside the plant. The metabolic and energy related pathways generate ROS as by product in cells in response to environmental stresses (Andreyev et al., 2005). Previous studies have suggested that the early down regulation of metabolism is one of the consistent trends of plants in response to abiotic stress (Cramer et al., 2011). Therefore, overexpression of *JIOsPR10* in rice may affect the balance between development and environmental stress adaption.

ROSs are key secondary signal molecules associated with both biotic and abiotic stresses in plants (Bailey-Serres and Mittler, 2006). Maintenance of ROS balance inside cell is essential for stress tolerance. Peroxidase has ability to generate oxidative burst and trigger immune responses in plants (Chittoor et al., 1997; Daudi et al., 2012; Sasaki et al., 2004). A peroxidase (spot 4) was highly expressed in Ox-3 lines as compared to the WT, suggesting overexpression of *JIOsPR10* may lead to an induction of ROS. Meanwhile, Cu/Zn-SOD (spot 8) was highly accumulated in *JIOsPR10* overexpression lines (Fig. 5). Cu/Zn-SOD has ability of catalyzing O_2^- into H_2O_2 . Higher accumulation of Cu/Zn-SOD in transgenic plants might be related with the maintenance of intracellular oxidative stress balances, therefore lead to a different phenotype in plant growth and stress resistance.

We also found that the expression of three proteins related with rice defense mechanism including TLP, DUF26-like protein, beta-1,3-glucanase was increased in the transgenic plants. TLP belongs to the PR-5 family based on the amino acid sequence and structural similarities (Edens et al., 1982). Overexpression of *TLP* enhances tolerance to different fungus (Acharya et al., 2013; Mackintosh et al., 2007; Munis et al., 2010), salt, drought,

and oxidative stresses (Munis et al., 2010). The DUF26-like protein is also involved in multiples stress tolerance, such as rice blast fungus (Kim et al., 2004), salt stress (Zhang et al., 2009), and ozone stress (Wrzaczek et al., 2010). In plants, beta-1,3-glucanases are related with the pathogen infection, ozone, H₂O₂, wounding stresses, and plant development (Chandler and Robertson, 1994; Cheong et al., 2000; Hwang et al., 2007). The induction of those proteins may be triggered by the peroxidase induced oxidative stresses. Taken together, we suggest that enhancement of biotic and abiotic stresses of *JIOsPR10* overexpression plant is probably because of its increased oxidative stress tolerance as compared to the WT.

Moreover, ROS were also emerged as important regulator of cell growth and plant development, and ROS are highly accumulated in plant during the germination and early development stages (Gapper and Dolan, 2006). In this study, we observed that overexpression of *JIOsPR10* promotes the root and shoot development in early development stages (Fig. 3) which could be the result of enhanced ROS scavenging ability in transgenic plants. Similarly, overexpression of *PR10* in *Brassica napus* enhanced seed germination under salinity conditions (Srivastava et al., 2004). Based on proteomics results, the up-regulated proteins such as TLP (Futamura et al., 2006), DUF26-like protein (Jiang et al., 2007), beta-1,3-glucanase (Cheong et al., 2000; Vögeli-Lange et al., 1994), and SOD (Kliebenstein et al., 1998) were also reported to be involved in plant development, indicating differential expression of those ROS and defense related proteins may be involved in plant growth promoting process. Furthermore, we also observed that the rice productivity was not changed in overexpression plants (Fig. 3D). However, these plants were grown in absence of any environmental stresses. In natural conditions, the overexpression plant may have high yield than WT due to its increase in tolerance to environmental stresses.

Here, we also confirmed that overexpression of *JIOsPR10* in rice enhances *M. oryzae* tolerance. Meanwhile, transgenic plants also showed drought and salt stress tolerance phenotype. These results together indicate that *JIOsPR10* is involved in the multiple processes of rice including early development, enhancement of host to fungal infection, and oxidative, drought and salt stress tolerance, demonstrating that *JIOsPR10* is a good candidate gene for agricultural application to protect the crops against biotic and abiotic stresses. However, we also found that the transcriptional expression of *JIOsPR10* in Ox-1 is higher than Ox-3 (Fig. 2), but Ox-3 rather than Ox-1 exhibits higher resistance against *M. oryzae* infection (Fig. 7). We also found that PR proteins such as DUF26-like protein and β -1,3-glucanase are differentially accumulated in Ox-1

and Ox-3 plant. Therefore, different correlation between transcriptional expression of *JIOsPR10* and phenotype may cause by the differential accumulation of other PR in different transgenic lines due to its RNase enzyme activity or post-transcriptional modification.

Acknowledgments

This work was supported by a 2-Year Research Grant of Pusan National University.

References

- Acharya, K., Pal, A. K., Gulati, A., Kumar, S., Singh, A. K. and Ahuja, P. S. 2013. Overexpression of *Camellia sinensis* thaumatin-like protein, *CsTLP* in potato confers enhanced resistance to *Macrophomina phaseolina* and *Phytophthora infestans* infection. *Mol. Biotechnol.* 54:609-622.
- Andrade, L. B., Oliveira, A. S., Ribeiro, J. K., Kiyota, S., Vasconcelos, I. M., de Oliveira, J. T. and de Sales, M. P. 2010. Effects of a novel pathogenesis-related class 10 (PR-10) protein from *Crotalaria pallida* roots with papain inhibitory activity against root-knot nematode *Meloidogyne incognita*. *J. Agric. Food Chem.* 58:4145-4152.
- Andreyev, A. Y., Kushnareva, Y. E. and Starkov, A. A. 2005. Mitochondrial metabolism of reactive oxygen species. *Biochemistry (Mosc.)* 70:200-214.
- Bailey-Serres, J. and Mittler, R. 2006. The roles of reactive oxygen species in plant cells. *Plant Physiol.* 141:311.
- Bantignies, B., Séguin, J., Muzac, I., Dédaldéchamp, F., Gulick, P. and Ibrahim, R. 2000. Direct evidence for ribonucleolytic activity of a PR-10-like protein from white lupin roots. *Plant Mol. Biol.* 42:871-881.
- Chandler, P. M. and Robertson, M. 1994. Gene expression regulated by abscisic acid and its relation to stress tolerance. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 45:113-141.
- Chen, Z. Y., Brown, R. L., Damann, K. E. and Cleveland, T. E. 2007. Identification of maize kernel endosperm proteins associated with resistance to aflatoxin contamination by *Aspergillus flavus*. *Phytopathology* 97:1094-1103.
- Cheong, Y. H., Kim, C. Y., Chun, H. J., Moon, B. C., Park, H. C., Kim, J. K., Lee, S., Han, C., Lee, S. Y. and Cho, M. J. 2000. Molecular cloning of a soybean class III beta-1,3-glucanase gene that is regulated both developmentally and in response to pathogen infection. *Plant Sci.* 154:71-81.
- Chittoor, J. M., Leach, J. E. and White, F. F. 1997. Differential induction of a peroxidase gene family during infection of rice by *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant-Microbe Interact.* 10:861-871.
- Cramer, G. R., Urano, K., Delrot, S., Pezzotti, M. and Shinozaki, K. 2011. Effects of abiotic stress on plants: a systems biology perspective. *BMC Plant Biol.* 11:163.
- Datta, K., Velazhahan, R., Oliva, N., Ona, I., Mew, T., Khush, G. S., Muthukrishnan, S. and Datta, S. K. 1999. Over-expression of the cloned rice thaumatin-like protein (*PR-5*) gene in transgenic rice plants enhances environmental friendly re-

- sistance to *Rhizoctonia solani* causing sheath blight disease. *Theor. Appl. Genet.* 98:1138-1145.
- Daudi, A., Cheng, Z., O'Brien, J. A., Mammarella, N., Khan, S., Ausubel, F. M. and Bolwell, G. P. 2012. The apoplastic oxidative burst peroxidase in *Arabidopsis* is a major component of pattern-triggered immunity. *Plant Cell* 24:275-287.
- de las Mercedes Dana, M., Pintor-Toro, J. A. and Cubero, B. 2006. Transgenic tobacco plants overexpressing chitinases of fungal origin show enhanced resistance to biotic and abiotic stress agents. *Plant Physiol.* 142:722-730.
- Duan, J., Zhang, M., Zhang, H., Xiong, H., Liu, P., Ali, J., Li, J. and Li, Z. 2012. *OsMIOX*, a *myo*-inositol oxygenase gene, improves drought tolerance through scavenging of reactive oxygen species in rice (*Oryza sativa* L.). *Plant Sci.* 196:143-151.
- Edens, L., Heslinga, L., Klok, R., Ledebor, A. M., Maat, J., Toonen, M. Y., Visser, C. and Verrips, C. T. 1982. Cloning of cDNA encoding the sweet-tasting plant protein thaumatin and its expression in *Escherichia coli*. *Gene* 18:1-12.
- Fernandes, H., Michalska, K., Sikorski, M. and Jaskolski, M. 2013. Structural and functional aspects of PR-10 proteins. *FEBS J.* 280:1169-1199.
- Flores, T., Alape-Girón, A., Flores-Díaz, M. and Flores, H. E. 2002. Ocatin. A novel tuber storage protein from the andean tuber crop oca with antibacterial and antifungal activities. *Plant Physiol.* 128:1291-1302.
- Fraire-Velázquez, S., Rodríguez-Guerra, R. and Sánchez-Calderón, L. 2011. Abiotic and biotic stress response crosstalk in plants. In: *Abiotic stress response in plants: physiological, biochemical and genetic perspectives*, eds. by A. K. Shanker and B. Venkateswarlu, pp. 3-26. InTech, Rijeka, Croatia.
- Futamura, N., Tani, N., Tsumura, Y., Nakajima, N., Sakaguchi, M. and Shinohara, K. 2006. Characterization of genes for novel thaumatin-like proteins in *Cryptomeria japonica*. *Tree Physiol.* 26:51-62.
- Gapper, C. and Dolan, L. 2006. Control of plant development by reactive oxygen species. *Plant Physiol.* 141:341-345.
- Gupta, R., Lee, S. E., Agrawal, G. K., Rakwal, R., Park, S., Wang, Y. and Kim, S. T. 2015. Understanding the plant-pathogen interactions in the context of proteomics-generated apoplastic proteins inventory. *Front. Plant Sci.* 6:352.
- Hashimoto, M., Kisseleva, L., Sawa, S., Furukawa, T., Komatsu, S. and Koshiba, T. 2004. A novel rice PR10 protein, RSOsPR10, specifically induced in roots by biotic and abiotic stresses, possibly via the jasmonic acid signaling pathway. *Plant Cell Physiol.* 45:550-559.
- He, C., Lin, Z., McElroy, D. and Wu, R. 2009. Identification of a rice *actin2* gene regulatory region for high-level expression of transgenes in monocots. *Plant Biotechnol. J.* 7:227-239.
- Hwang, D. H., Kim, S. T., Kim, S. G. and Kang, K. Y. 2007. Comprehensive analysis of the expression of twenty-seven beta-1, 3-glucanase genes in rice (*Oryza sativa* L.). *Mol. Cells* 23:207-214.
- Jiang, J., Li, J., Xu, Y., Han, Y., Bai, Y., Zhou, G., Lou, Y., Xu, Z. and Chong, K. 2007. RNAi knockdown of *Oryza sativa* root *meander curling* gene led to altered root development and coiling which were mediated by jasmonic acid signalling in rice. *Plant Cell Environ.* 30:690-699.
- Jwa, N. S., Agrawal, G. K., Rakwal, R., Park, C. H. and Agrawal, V. P. 2001. Molecular cloning and characterization of a novel Jasmonate inducible pathogenesis-related class 10 protein gene, *JIOsPR10*, from rice (*Oryza sativa* L.) seedling leaves. *Biochem. Biophys. Res. Commun.* 286:973-983.
- Kim, J. Y., Wu, J., Kwon, S. J., Oh, H., Lee, S. E., Kim, S. G., Wang, Y., Agrawal, G. K., Rakwal, R., Kang, K. Y., Ahn, I. P., Kim, B. G. and Kim, S. T. 2014. Proteomics of rice and *Cochliobolus miyabeanus* fungal interaction: insight into proteins at intracellular and extracellular spaces. *Proteomics* 14:2307-2318.
- Kim, S. G., Wang, Y., Lee, K. H., Park, Z. Y., Park, J., Wu, J., Kwon, S. J., Lee, Y. H., Agrawal, G. K., Rakwal, R., Kim, S. T. and Kang, K. Y. 2013. In-depth insight into in vivo apoplastic secretome of rice-*Magnaporthe oryzae* interaction. *J. Proteomics* 78:58-71.
- Kim, S. T., Cho, K. S., Jang, Y. S. and Kang, K. Y. 2001. Two-dimensional electrophoretic analysis of rice proteins by polyethylene glycol fractionation for protein arrays. *Electrophoresis* 22:2103-2109.
- Kim, S. T., Kang, S. Y., Wang, Y., Kim, S. G., Hwang, D. H. and Kang, K. Y. 2008a. Analysis of embryonic proteome modulation by GA and ABA from germinating rice seeds. *Proteomics* 8:3577-3587.
- Kim, S. T., Kim, S. G., Hwang, D. H., Kang, S. Y., Kim, H. J., Lee, B. H., Lee, J. J. and Kang, K. Y. 2004. Proteomic analysis of pathogen-responsive proteins from rice leaves induced by rice blast fungus, *Magnaporthe grisea*. *Proteomics* 4:3569-3578.
- Kim, S. T., Yu, S., Kang, Y. H., Kim, S. G., Kim, J. Y., Kim, S. H. and Kang, K. Y. 2008b. The rice pathogen-related protein 10 (JIOsPR10) is induced by abiotic and biotic stresses and exhibits ribonuclease activity. *Plant Cell Rep.* 27:593-603.
- Kliebenstein, D. J., Monde, R. A. and Last, R. L. 1998. Superoxide dismutase in *Arabidopsis*: an eclectic enzyme family with disparate regulation and protein localization. *Plant Physiol.* 118:637-650.
- Krishnaswamy, S., Baral, P. K., James, M. N. and Kav, N. N. 2011. Site-directed mutagenesis of histidine 69 and glutamic acid 148 alters the ribonuclease activity of pea ABR17 (PR10.4). *Plant Physiol. Biochem.* 49:958-962.
- Li, J. B., Luan, Y. S. and Yin, Y. L. 2014. *SpMYB* overexpression in tobacco plants leads to altered abiotic and biotic stress responses. *Gene* 547:145-151.
- Mackintosh, C. A., Lewis, J., Radmer, L. E., Shin, S., Heinen, S. J., Smith, L. A., Wyckoff, M. N., Dill-Macky, R., Evans, C. K., Kravchenko, S., Baldrige, G. D., Zeyen, R. J. and Muehlbauer, G. J. 2007. Overexpression of defense response genes in transgenic wheat enhances resistance to Fusarium head blight. *Plant Cell Rep.* 26:479-488.
- McGee, J. D., Hamer, J. E. and Hodges, T. K. 2001. Characterization of a *PR-10* pathogenesis-related gene family induced in rice during infection with *Magnaporthe grisea*. *Mol. Plant-Microbe Interact.* 14:877-886.
- Munis, M. F., Tu, L., Deng, F., Tan, J., Xu, L., Xu, S., Long, L. and Zhang, X. 2010. A thaumatin-like protein gene involved

- in cotton fiber secondary cell wall development enhances resistance against *Verticillium dahliae* and other stresses in transgenic tobacco. *Biochem. Biophys. Res. Commun.* 393:38-44.
- Park, C. J., Kim, K. J., Shin, R., Park, J. M., Shin, Y. C. and Paek, K. H. 2004. Pathogenesis-related protein 10 isolated from hot pepper functions as a ribonuclease in an antiviral pathway. *Plant J.* 37:186-198.
- Rizhsky, L., Davletova, S., Liang, H. and Mittler, R. 2004. The zinc finger protein Zat12 is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in *Arabidopsis*. *J. Biol. Chem.* 279:11736-11743.
- Sarowar, S., Kim, Y. J., Kim, E. N., Kim, K. D., Hwang, B. K., Islam, R. and Shin, J. S. 2005. Overexpression of a pepper basic pathogenesis-related protein 1 gene in tobacco plants enhances resistance to heavy metal and pathogen stresses. *Plant Cell Rep.* 24:216-224.
- Sasaki, K., Iwai, T., Hiraga, S., Kuroda, K., Seo, S., Mitsuhashi, I., Miyasaka, A., Iwano, M., Ito, H., Matsui, H. and Ohashi, Y. 2004. Ten rice peroxidases redundantly respond to multiple stresses including infection with rice blast fungus. *Plant Cell Physiol.* 45:1442-1452.
- Shi, W., Hao, L., Li, J., Liu, D., Guo, X. and Li, H. 2014. The *Gossypium hirsutum* WRKY gene *GhWRKY39-1* promotes pathogen infection defense responses and mediates salt stress tolerance in transgenic *Nicotiana benthamiana*. *Plant Cell Rep.* 33:483-498.
- Srivastava, S., Fristensky, B. and Kav, N. N. 2004. Constitutive expression of a PR10 protein enhances the germination of *Brassica napus* under saline conditions. *Plant Cell Physiol.* 45:1320-1324.
- Stintzi, A., Heitz, T., Prasad, V., Wiedemann-Merdinoglu, S., Kauffmann, S., Geoffroy, P., Legrand, M. and Fritig, B. 1993. Plant 'pathogenesis-related' proteins and their role in defense against pathogens. *Biochimie* 75:687-706.
- van Loon, L. C., Pierpoint, W. S., Boller, T. and Conejero, V. 1994. Recommendations for naming plant pathogenesis-related proteins. *Plant Mol. Biol. Report.* 12:245-264.
- van Loon, L. C. and van Strien, E. A. 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. Mol. Plant Pathol.* 55:85-97.
- Vögeli-Lange, R., Fründt, C., Hart, C. M., Nagy, F. and Meins, F. 1994. Developmental, hormonal, and pathogenesis-related regulation of the tobacco class I β -1,3-glucanase B promoter. *Plant Mol. Biol.* 25:299-311.
- Wang, Y., Kim, S. G., Wu, J., Kim, S. T. and Kang, K. Y. 2014. Differential proteome and secretome analysis during rice-pathogen interaction. *Methods Mol. Biol.* 1072:563-572.
- Wrzaczek, M., Brosché, M., Salojärvi, J., Kangasjärvi, S., Idänheimo, N., Mersmann, S., Robatzek, S., Karpiński, S., Karpińska, B. and Kangasjärvi, J. 2010. Transcriptional regulation of the CRK/DUF26 group of receptor-like protein kinases by ozone and plant hormones in *Arabidopsis*. *BMC Plant Biol.* 10:95.
- Xie, Y. R., Chen, Z. Y., Brown, R. L. and Bhatnagar, D. 2010. Expression and functional characterization of two pathogenesis-related protein 10 genes from *Zea mays*. *J. Plant Physiol.* 167:121-130.
- Zhang, L., Tian, L. H., Zhao, J. F., Song, Y., Zhang, C. J. and Guo, Y. 2009. Identification of an apoplastic protein involved in the initial phase of salt stress response in rice root by two-dimensional electrophoresis. *Plant Physiol.* 149:916-928.
- Zhou, X. J., Lu, S., Xu, Y. H., Wang, J. W. and Chen, X. Y. 2002. A cotton cDNA (*GaPR-10*) encoding a pathogenesis-related 10 protein with in vitro ribonuclease activity. *Plant Sci.* 162:629-636.