



HHS Public Access

Author manuscript

Alcohol Clin Exp Res. Author manuscript; available in PMC 2017 October 01.

Published in final edited form as:

Alcohol Clin Exp Res. 2016 October ; 40(10): 2056–2072. doi:10.1111/acer.13204.

Alcohol Use and Human Immunodeficiency Virus (HIV) Infection: Current Knowledge, Implications, and Future Directions

Emily C. Williams, PhD, MPH^{1,2}, Judith A. Hahn, PhD, MA³, Richard Saitz, MD, MPH^{4,5}, Kendall Bryant, PhD⁶, Marlene C. Lira, BA⁵, and Jeffrey H. Samet, MD, MA, MPH^{4,5}

¹Veterans Health Administration (VA) Health Services Research and Development, Center of Innovation for Veteran-Centered and Value-Driven Care, Seattle, WA

²University of Washington, Department of Health Services, Seattle, WA

³Departments of Medicine and Epidemiology and Biostatistics, University of California, San Francisco; San Francisco, CA

⁴Department of Community Health Sciences, Boston University School of Public Health, Boston, MA

⁵Clinical Addiction Research and Education (CARE) Unit, Section of General Internal Medicine, Department of Medicine, Boston Medical Center, Boston University School of Medicine, Boston, MA

⁶Consortiums for HIV/AIDS and Alcohol Research Translation (CHAART) National Institute on Alcohol Abuse and Alcoholism National Institutes of Health, Bethesda, MD

Abstract

Alcohol use is common among people living with human immunodeficiency virus (HIV). In this narrative review, we describe literature regarding alcohol's impact on transmission, care, co-infections and comorbidities that are common among people living with HIV (PLWH), as well as literature regarding interventions to address alcohol use and its influences among PLWH. This narrative review identifies alcohol use as a risk factor for HIV transmission, as well as a factor impacting the clinical manifestations and management of HIV. Alcohol use appears to have additive and potentially synergistic effects on common HIV-related comorbidities. We find that interventions to modify drinking and improve HIV-related risks and outcomes have had limited success to date, and we recommend research in several areas. Consistent with Office of AIDS Research/National Institutes of Health priorities, we suggest research to better understand how and at what levels alcohol influences comorbid conditions among PLWH, to elucidate the mechanisms by which alcohol use is impacting comorbidities, and to understand whether decreases in alcohol use improve HIV-relevant outcomes. This should include studies regarding whether state-of-the-art medications used to treat common co-infections are safe for PLWH who drink alcohol. We recommend that future research among PLWH include validated self-report measures of alcohol

Correspondence should be addressed to: Jeffrey H. Samet, MD, MA, MPH, Boston University Schools of Medicine and Public Health, Boston Medical Center, 801 Massachusetts Ave, 2nd Floor, Boston, MA 02118, Phone: 617-414-7288, Fax: 617-414-4676, jsamet@bu.edu.

Conflicts of Interest: The authors do not have conflicts of interest to declare.

use and/or biological measurements, ideally both. Additionally, subgroup variation in associations should be identified to ensure that the risks of particularly vulnerable populations are understood. This body of research should serve as a foundation for a next generation of intervention studies to address alcohol use from transmission to treatment of HIV. Intervention studies should inform implementation efforts to improve provision of alcohol-related interventions and treatments for PLWH in healthcare settings. By making further progress on understanding how alcohol use affects PLWH in the era of HIV as a chronic condition, this research should inform how we can mitigate transmission, achieve viral suppression, avoid exacerbating common comorbidities of HIV and alcohol use and make progress toward the 90-90-90 goals for engagement in the HIV treatment cascade.

Keywords

alcohol; alcohol use; substance use; HIV; HIV-related comorbidities

INTRODUCTION

Both human immunodeficiency virus (HIV) and alcohol use are common worldwide (UNAIDS, 2015). In some cases, such as Russia and Uganda, high per capita alcohol use overlaps with a high prevalence of HIV (Rehm et al., 2006; Rehm et al., 2009a). In the U.S., approximately 1.1 million people are living with HIV (PLWH) (Hall et al., 2009; Lansky et al., 2010; Centers for Disease Control and Prevention, 2011; Centers for Disease Control and Prevention, 2012), and 88% of adults report any past-year alcohol use (1 drink), 25% report past-month heavy drinking (5 drinks on 1 day), and 14% meet criteria for a past-year alcohol use disorder (Substance Abuse and Mental Health Services Administration, 2014; Grant et al., 2015).

A substantial proportion of PLWH consume alcohol, many at unhealthy levels (heavy drinking and/or alcohol use disorder) (Saitz, 2005). The prevalence of unhealthy alcohol use in PLWH ranges from 8% (Galvan et al., 2002) to 42% (Samet et al., 2004b; Lefevre et al., 1995). Based on data from the U.S. general population, sub-populations at high risk of having both HIV and unhealthy alcohol use include men who have sex with men (MSM), racial/ethnic minorities, persons who inject drugs, sex workers, and persons of low socioeconomic status (Hall et al., 2009; Lansky et al., 2010; Pellowski et al., 2013; Delker et al., 2016).

In this narrative review, we describe literature regarding alcohol's impact on HIV acquisition/transmission, care, co-infections and comorbidities common among PLWH, and morbidity and mortality, as well as literature regarding interventions to address alcohol use and its influences among PLWH.

Alcohol Use and Risk Behaviors for HIV Acquisition/Transmission

Alcohol use has a strong and consistent association with HIV incidence (Shuper et al., 2010; Scott-Sheldon et al., 2016). Associations between alcohol use and sex-risk behaviors have been identified consistently (Shuper et al., 2009; Samet et al., 2007b; Maisto and Simons,

2016; Maisto et al., 2012). A meta-analysis of 30 experimental studies found that participants randomly assigned to consume alcohol reported stronger intentions to engage in unprotected sex, weaker sexual communication and negotiation skills, and higher levels of sexual arousal than participants assigned to either placebo or no-alcohol control groups (Scott-Sheldon et al., 2016), suggesting a causal link between alcohol use and sexual risk taking (and thus HIV transmission).

Several studies have tested interventions focused either on reducing HIV risk among people who drink or on reducing alcohol use to reduce HIV transmission (Carrasco et al., 2016; Samet and Walley, 2010; Eaton et al., 2013; Kalichman et al., 2008; Samet et al., 2015; Samet et al., 2008; Monti et al., 2016; Kennedy et al., 2016) with limited success, especially with regards to sustained effects across multiple domains (e.g., reductions of both alcohol and sex-risk behaviors). A review of interventions tested in sub-Saharan Africa suggested a need for interventions that address alcohol use and HIV risk both at individual and policy levels (Carrasco et al., 2016).

Alcohol Use and the HIV Treatment Cascade

The HIV treatment cascade identifies five stages of HIV medical care—1) diagnosis, 2) linkage to medical care, 3) engagement with/retention in medical care, 4) treatment with antiretroviral therapy (ART), and 5) achievement of viral suppression—all considered essential targets to meet in order to effectively treat and prevent the spread of HIV (Joint United Nations Programme on HIV/AIDS, 2014; Skarbinski et al., 2015; Mugavero et al., 2013; Bor et al., 2015; Cohen et al., 2014; Park et al., 2007). World Health Organization (WHO) goals for 2020 are the achievement of 90-90-90 with regard to percent identification, engagement with ART initiation, and viral suppression (Joint United Nations Programme on HIV/AIDS, 2014)

Alcohol use has been shown to influence HIV care and outcomes at every stage of the cascade (Azar et al., 2010b; Vagenas et al., 2015); it is associated with lower receipt of HIV testing (Fatch et al., 2013; Vagenas et al., 2014; Bengtson et al., 2014), greater delay in engaging with medical care after testing positive (Samet et al., 1998), delay in, lower quality of, and worse retention in HIV treatment (Samet et al., 2001; Korthuis et al., 2012; Cunningham et al., 2006; Giordano et al., 2005; Samet et al., 2003; Monroe et al., 2016), and non-adherence to ART (Hendershot et al., 2009). Some studies have also identified associations between alcohol use and disease progression (see below) (Hahn and Samet, 2010). The high prevalence of alcohol use and the fact it is often unaddressed likely combine to result in suboptimal engagement. Reducing alcohol use may be fertile ground for interventions aimed at meeting WHO goals.

Alcohol and Treatment with ART—Adherence

The association between alcohol use and non-adherence to ART has received substantial attention regarding alcohol's role in the HIV treatment cascade (Hendershot et al., 2009). Some studies reveal any (versus no) alcohol use is associated with non-adherence (Hendershot et al., 2009; Samet et al., 2004a). Other studies have found threshold effects of heavy drinking (Samet et al., 2004a) or dose response relationships between the amount

consumed and the number of pills missed (Braithwaite et al., 2005; Parsons et al., 2008). The association between alcohol use and non-adherence is thought to reflect both failure to take medications as a result of complex dosing regimens and intoxication due to heavy drinking episodes, as well as intentional missed doses due to beliefs regarding potential toxic interactions between ART and alcohol (Kalichman et al., 2013a; Kalichman et al., 2013b; Parsons et al., 2007b; Pellowski et al., 2016). The extent to which alcohol and ART have toxic interactions and at what levels remains unclear, though potential biological mechanisms by which toxicity may exist have been described (Kumar et al., 2015; Hu et al., 2015; McCance-Katz et al., 2013) and one recent study identified potential interactive effects on liver disease (Bilal et al., 2016).

At least two studies have tested interventions aimed at simultaneously reducing alcohol use and improving ART adherence (Samet and Walley, 2010). One tested a nurse-led multi-component intervention that included 4 visits over 3 months, which had no effect on alcohol use or ART adherence (Samet et al., 2005). The other tested eight 1-hour individual sessions of motivational interviewing and cognitive behavioral skills over 3 months; positive intervention effects on self-reported medication adherence and markers of disease progression were observed at 3 months, but not sustained (Parsons et al., 2007a). Given the substantial influence of this modifiable risk factor on this key treatment goal, further research is warranted regarding how to influence ART adherence via reducing drinking, or in the context of continued drinking.

Alcohol and Achievement of Viral Suppression—Disease Progression

While multiple studies have identified associations between alcohol use and HIV disease progression, the extent to which alcohol use directly affects disease progression is not fully resolved (Hahn and Samet, 2010; Azar et al., 2010a). Identified associations may be mediated by behavioral mechanisms—the most common being non-adherence to ART—or they may be direct, resulting from biological mechanisms (Hahn and Samet, 2010).

Several plausible biological mechanisms may account for how alcohol may increase pre-ART HIV disease progression. Both HIV and alcohol are established causes of microbial translocation and systemic inflammation (Thurman, 1998; Brenchley et al., 2006; Douek, 2007), and both induce immune activation. Therefore, it is plausible that alcohol may accelerate the progression of HIV disease via these pathways. Alcohol use may contribute to T cell proliferative defects (Szabo et al., 2004). And both HIV and alcohol cause a similar disruption of the normal gut microbiome, which may contribute to systemic immune activation by either impairing gut barrier integrity or absorption of bacteria-derived metabolites that confer immune defects (Dillon et al., 2014). Markers of microbial translocation are associated with increased opportunistic co-infections (Cassol et al., 2010), HIV disease progression (Marchetti et al., 2008; Balagopal et al., 2008), and mortality (Sandler et al., 2010). The link between alcohol use and markers of microbial translocation in PLWH has been examined to a limited extent, and findings have been mixed (Monnig et al., 2016; Hunt et al., 2014; Cioe et al., 2015; Balagopal et al., 2008; Carrico et al., 2015). Experimental studies of the effect of alcohol on progression of the simian immunodeficiency virus (SIV) in macaques largely support the biologic plausibility of alcohol's influence on

disease progression (Amedee et al., 2014; Marcondes et al., 2008; Kumar et al., 2005; Bagby et al., 2006; Poonia et al., 2006; Molina et al., 2006). However, the most recent study, which randomized macaques to chronic heavy drinking (versus sucrose) and subsequently to ART (versus no ART), identified no association between chronic heavy drinking and viral load over 4 months, regardless of ART status (Molina et al., 2014a).

The results of human observational studies have been mixed, perhaps because of factors that can affect disease progression in people outside of tightly controlled experimental settings (Hahn and Samet, 2010). Of the several prospective studies conducted in the pre-highly active ART era (Kaslow et al., 1989; Penkower et al., 1995; Eskild and Petersen, 1994; Tang et al., 1997; Veugelers et al., 1994; Webber et al., 1999; Chandiwana et al., 1999), none found an association between alcohol consumption and the onset of the Acquired Immune Deficiency Syndrome (AIDS). In the highly-active ART era, after controlling for ART use, four prospective studies found no association between heavy drinking and HIV disease progression, as measured by CD4 cell count or HIV viral load (Ghebremichael et al., 2009; Cook et al., 2008; Chander et al., 2006b; Kowalski et al., 2012), and one identified an increased risk of virological failure associated with heavy alcohol use (Deiss et al., 2016). Two earlier prospective studies that separately examined HIV disease progression among those not on ART found that heavy alcohol consumption was associated with lower CD4 cell count (Samet et al., 2007a) and shorter time to CD4 cell count <200 cells/mm³ (Baum et al., 2010), but not differences in HIV viral load (Samet et al., 2007a; Baum et al., 2010). These studies both had small samples of persons not on ART (n=128 and n=87 respectively). A 7-year study among 5,067 Swiss PLWH, which stratified analyses by ART naïve and individuals initiating ART, found no association between alcohol use and virological failure or CD4 cell counts over time in either group (Conen et al., 2013). Findings to date suggest the possibility that alcohol use may not directly influence disease progression. However, further research is needed to explore longer term outcomes associated with heavy drinking in macaques, as well as influences of alcohol use on disease progression among larger samples of PLWH and among untreated PLWH.

Alcohol Use and HIV-Associated Comorbidities

Medical illnesses associated with alcohol use often seen in older adults *without* HIV infection (Cherpitel et al., 2015) are now occurring as common comorbid conditions in younger PLWH despite prolonged viral suppression (Crothers et al., 2009; Hasse et al., 2011; Simmons et al., 2013; Guaraldi et al., 2011; Sharma et al., 2015; Shiau et al., 2013; Hansen et al., 2012), which may reflect premature aging (High et al., 2012; Justice and Falutz, 2014). In addition to the direct, toxic pathophysiologic effects of both HIV and alcohol, mechanisms that are likely to lead to high rates of comorbid conditions among PLWH include immune senescence (the effects of aging on the function of the immune system), inflammation, and hypercoagulability (High et al., 2012; Deeks, 2009; Deeks, 2011; Katz et al., 2015). Because alcohol use impacts some of these same mechanisms, there has been investigation of the role of alcohol use in HIV-related comorbidities (Neuman et al., 2012; Bryant et al., 2010; Freiberg et al., 2010; Sarkar et al., 2015; World Health Organization, 2015), and the study of comorbidities is now a high priority for NIH (National

Institutes of Health, 2015). We review research to date on HIV-associated comorbidities in which alcohol may play a role.

Common Co-Infections

Hepatitis C and Tuberculosis

Approximately 25 – 30% of PLWH are co-infected with hepatitis C virus (HCV) (Alter, 2006; Platt et al., 2016). Chronic HCV infection is a major cause of morbidity and mortality among PLWH (Sulkowski, 2013). PLWH with HCV co-infection have accelerated progression of HCV-related liver disease (i.e., cirrhosis and liver cancer) (Sulkowski, 2007a; Sulkowski, 2007b; Sulkowski, 2008). Although little is known about how low-level drinking influences HCV outcomes, abstinence from alcohol use is recommended for PLWH with HCV co-infection (Tsui et al., 2013) because liver disease outcomes are also adversely impacted by alcohol use (Peters and Terrault, 2002; Morgan et al., 2003; Sulkowski, 2007a; Sulkowski, 2007b; Sulkowski, 2008; Cooper and Cameron, 2005; Safdar and Schiff, 2004; Schiff and Ozden, 2003; Fuster et al., 2014; Fuster et al., 2013; Tsui et al., 2013; Tsui et al., 2006).

Similarly, both alcohol use and HIV are associated with increased risk of tuberculosis (TB) infection (Lonnroth et al., 2008; Rehm et al., 2009b), which is the leading cause of mortality in PLWH worldwide (World Health Organization, 2014; UNAIDS, 2015). Alcohol use (any versus none) is associated with an increased risk (up to 3-fold) of having active TB and poorer TB outcomes (Volkmann et al., 2016), likely as a result of alcohol's impact on the immune system, poor TB treatment adherence, and social marginalization that increases risk for infection (Rehm et al., 2009b), all of which may be amplified among PLWH given potential overlapping mechanisms.

Alcohol use is relevant to each of these common co-infections not only due to its potential to speed their progression but also because it likely influences their care. Alcohol use was a contraindication to previously used treatments for HCV, and the WHO warns against the use of isoniazid preventive therapy—commonly used to prevent mortality and active TB among PLWH—in persons with “regular and heavy alcohol use” (World Health Organization, 2011). While the WHO recommendation was made based on prior reports of isoniazid toxicity among heavy drinkers, no studies have systematically assessed the safety of TB preventive therapy in heavy drinkers with or without HIV infection. Moreover, it is unknown whether emerging treatments for HCV—directly acting antivirals that are highly effective in HIV co-infected patients (Sulkowski, 2013)—have contraindications for persons who drink. This is a rapidly evolving story for both co-infections; studies are needed to assess whether alcohol use is associated with poor outcomes among PLWH who are treated for HCV or TB infection. In the meantime, historical contraindications and current WHO recommendations are likely to influence clinical decision making regarding provision of recommended treatments. Given that alcohol use is an established risk factor for decreased ART adherence (Hendershot et al., 2009) and data suggest risk for active TB treatment discontinuation (Kendall et al., 2013), it will also be important to assess alcohol's influence on adherence to and completion of treatments for both co-infections, as well as to identify interventions that may support reduced drinking and treatment adherence.

Cardiovascular Diseases

PLWH who drink heavily and/or meet criteria for alcohol use disorders are at increased risk for coronary heart disease (CHD) and other cardiovascular diseases (CVD) (Freiberg et al., 2013; Freiberg et al., 2010). In one study heavy drinking was a risk factor for CHD among PLWH but not those without HIV (Freiberg et al., 2010). Mechanisms underlying CHD risk among PLWH are unclear but have been linked to inflammation (Niaura et al., 2000; Armah et al., 2012). ART and HIV viral suppression do not eliminate the increased risk of CHD (Freiberg et al., 2013), nor does ART return increased inflammation to pre-HIV infection levels (Freiberg et al., 2015). A systematic review of 13 studies suggests an increased risk of CVD associated with any and heavy alcohol use among PLWH ranging from 37% (95% CI: 2% – 84%) to 78% (95% CI: 9% – 293%) (Kelso et al., 2015). While it is clear that alcohol increases risk for CVD among PLWH, alcohol's role in CVD risk merits further inquiry. For instance, the levels of alcohol use at which CVD risk is increased for PLWH are unclear: most studies have not compared non-drinking to low-level or "moderate" drinking, but one large observational study that did identified a protective effect of low-level drinking (Carrieri et al., 2012). In addition, while cardiomyopathy is a condition for which both alcohol and HIV can be underlying etiologies, the additive or synergistic impact of these combined exposures for this pathology has not been delineated.

Cancers

Rates of non-AIDS-defining cancer deaths are increasing among PLWH (Smith et al., 2014) and contributing an ever-greater health burden (Deeken et al., 2012; Shiels et al., 2011; Cutrell and Bedimo, 2013). Relative to uninfected persons, PLWH are at higher risk for many cancers, likely due to a combination of the direct toxic effects of the HIV virus, as well as chronic inflammation and immune suppression/activation, and high rates of co-infections and risk factors, such as smoking (Cutrell and Bedimo, 2013; Sigel et al., 2012; Bedimo et al., 2009; Borges et al., 2014; McGinnis et al., 2006; Park et al., 2016). Alcohol use may be another risk factor. Mechanisms by which rates of cancer may be elevated among PLWH (e.g., immune system effects, inflammation, and co-occurrence with other risk factors) overlap with mechanisms by which alcohol use adversely influences cancer outcomes. However, the specific role of alcohol use in cancer risk among PLWH is unknown (D'Souza et al., 2014). Alcohol use is a risk factor for liver and head and neck cancers among PLWH (McGinnis et al., 2006; Freedman et al., 2007; Viswanathan and Wilson, 2004), and findings from one study suggest that, together, alcohol use and HCV accounted for increased risk of liver cancer, but not non-Hodgkins lymphoma, among PLWH (McGinnis et al., 2006). Findings to date suggest alcohol's influence on cancers among PLWH is likely complex; alcohol may be a stronger contributor to some cancers than others and may have additive or synergistic effects on cancer-related outcomes.

Neurological Disorders

The severity of HIV-associated neurocognitive disorders (HAND) (Antinori et al., 2007) has declined in the post-ART era, but PLWH continue to face an array of neurological problems (Sacktor et al., 2016; Heaton et al., 2011; Heaton et al., 2010; Elbirt et al., 2015; Harezlak et

al., 2011). Alcohol use disorders are a known risk factor for neurocognitive dysfunction and are linked to impaired memory and decision making (Anand et al., 2010; Loeber et al., 2009). Alcohol use and HIV may have combined effects on neurocognitive decline (Anand et al., 2010; Persidsky et al., 2011; Thaler et al., 2015). However, the findings of studies assessing this question have been mixed. Some suggest that alcohol and HIV have additive adverse effects (Rothlind et al., 2005; Green et al., 2004), while a cross-sectional study of HIV-infected alcohol users found lower than average neurocognitive scores, but no significant relationship between number of heavy drinking days and neurocognitive performance (Attonito et al., 2014; Attonito, 2013). The interplay of alcohol and neurocognitive decline among PLWH merits further research.

Other related issues are also in need of further investigation. For instance, pain is common and associated with unhealthy alcohol use among PLWH (Merlin, 2015; Merlin et al., 2012a; Tsui et al., 2014). Because pain itself may interrupt engagement in HIV care (Berg et al., 2009; Merlin et al., 2012b) and may also induce alcohol use as a coping mechanism (Tsui et al., 2014), alcohol's role in pain management among PLWH, is an area ripe for further study. Another issue in need of further investigation is alcohol's role in neuropathy among PLWH. While both HIV and HIV treatments have known associations with neuropathy (Amruth et al., 2014), the role in these associations of a known etiology of peripheral neuropathy, alcohol use, has not been described.

Metabolic Complications

PLWH experience metabolic complications, which increase the risk of diabetes and cardiovascular disease (Crum-Cianflone et al., 2010; Crum-Cianflone et al., 2008; Samaras, 2012; Samaras et al., 2007; Calvo and Martinez, 2014). PLWH frequently gain weight following ART initiation (Lakey et al., 2013; Justice and Falutz, 2014). The prevalence of metabolic complications and diabetes is higher among PLWH than among those without HIV and is generally linked to lipid imbalances resulting from ART and chronic low-grade inflammation from HIV (Paula et al., 2013; Samaras et al., 2007; Jacobson et al., 2006; Mozumdar and Liguori, 2011; Rasmussen et al., 2012; De Wit et al., 2008). Alcohol use is associated with diabetes (Howard et al., 2004) and metabolic derangements consistent with lipodystrophy (Cheng et al., 2009). Alcohol use has also been linked to oxidative stress on muscles, bones and adipose tissue, and is thought to exacerbate lipid imbalances from HIV (Molina et al., 2014b). However, among PLWH and those without HIV, associations with alcohol use and both diabetes and metabolic syndrome appear to be U-shaped such that "moderate" alcohol use may be protective, whereas heavy drinking increases risk (Howard et al., 2004; Butt et al., 2009; Samaras et al., 2007; Tiozzo et al., 2015; Freiberg et al., 2004; Sidorenkov et al., 2010; Wakabayashi, 2014; Athyros et al., 2007). Research is needed regarding the potentially complex ways in which alcohol use influences metabolic complications among PLWH.

Falls and Injury

Up to 30% of PLWH experience falls annually (Erlandson et al., 2012). Falls can lead to fractures, which are 40–60% more common in PLWH relative to those without HIV (Sharma

et al., 2015; Shiau et al., 2013; Hansen et al., 2012), likely as a result of both high-energy injuries and fragility. Alcohol use is a potent risk factor for falls and fractures (Cherpitel et al., 2015; Cawthon et al., 2006; Kanis et al., 2004; Berg et al., 2008; Mukamal et al., 2007; Zhang et al., 2014) and it can also exacerbate inflammation, which could influence complications related to recovery from falls and fractures. Further research on the association between alcohol use and falls and injury among PLWH is needed, as is investigation of the role of alcohol reduction in fall prevention among PLWH.

Other Substance Use

More than 42% of PLWH in the US smoke (with some studies reporting 46–84%), compared with less than 20% of the general population (Lifson and Lando, 2012; Lifson et al., 2010; Mdodo et al., 2015; Marshall et al., 2009; Cooperman, 2016). Among PLWH, smoking is associated with increased risk for bacterial pneumonia, chronic obstructive pulmonary disease (COPD), coronary heart disease (CHD) and its associated complications, and decreased bone mineral density (Lifson et al., 2010; Cooperman, 2016). Though evidence is mixed (Kabali et al., 2011; Marshall et al., 2009), smoking may also be associated with HIV disease progression (Hile et al., 2016). Smoking is more strongly associated with poor health outcomes among PLWH than among those without HIV (Crothers et al., 2009; Crothers et al., 2005) and is likely an important contributor to the substantial comorbidity experienced by PLWH (Marshall et al., 2009). Smoking and alcohol use are correlated with one another in PLWH (Braithwaite et al., 2016) and, due to the potential for additive inflammatory effects of alcohol use and smoking (Niaura et al., 2000; Armah et al., 2012), PLWH who both smoke and drink alcohol may be at greater risk for adverse health outcomes, such as CHD and pneumonia. Despite the potential syndemic effects and the common co-occurrence of smoking and alcohol use among PLWH (Braithwaite et al., 2016), little research has investigated the synergistic effects of alcohol and tobacco use among PLWH, and interventions to improve both among PLWH.

Similarly, other drug use is common among PLWH (Green et al., 2010; Bing et al., 2001; Korthuis et al., 2012; Mathers et al., 2008) and often co-occurs with alcohol use (Green et al., 2010; Walley et al., 2008; Van Tieu and Koblin, 2009). This combination is likely to increase risk for multiple adverse outcomes. Given that drug use is associated with poorer engagement with the HIV treatment cascade, including medication adherence (Peretti-Watel et al., 2006; Mugavero et al., 2006), co-occurring alcohol and drug use may additively or synergistically adversely influence engagement with HIV care. Further, co-occurring alcohol and drug use may exacerbate severity of comorbid conditions, such as neurological function (Meyer et al., 2013; Weber et al., 2013; Hauser and Knapp, 2014), and outcomes of common co-infections among PLWH, which, in some cases, are disproportionately distributed among people who use drugs. For instance, an estimated 82% of people who inject drugs with HIV have HCV (Platt et al., 2016). Finally, alcohol use may enhance risk for drug overdose (Coffin et al., 2007; Shah et al., 2008; Coffin et al., 2003; Riley et al., 2016). Because similar levels of alcohol consumption may have greater effects for PLWH compared to those without HIV (McGinnis et al., 2016; Justice et al., 2016; McCance-Katz et al., 2012), alcohol's influence on overdose may be enhanced among PLWH. Research is needed to

more comprehensively understand the clustering of risks among PLWH and the combined effects of these risk profiles on adverse outcomes.

Mental Health Disorders

In addition to other substance use, mental health disorders, including depression, anxiety, trauma, and stress are common among PLWH (Sullivan et al., 2011; Sullivan et al., 2008; Chander et al., 2006a; Walkup et al., 2008; Braithwaite et al., 2016). The majority of research to date on alcohol use and mental health conditions among PLWH has focused on depression. Both alcohol use and depression are associated with reduced engagement with the HIV treatment cascade and adverse HIV-related consequences (Cruess et al., 2005; Cruess et al., 2003; Evans et al., 2002; Leserman, 2003; Leserman, 2008; Leserman et al., 2008; Ghebremichael et al., 2009; Kim et al., 2007; Goodness et al., 2014). Unhealthy (relative to low-risk) alcohol use is associated with higher risk of depressive symptoms (Sullivan et al., 2011), and longitudinal research among PLWH suggests that this association is uni-directional, such that alcohol use precedes depressive symptoms (Fergusson et al., 2009). Moreover, changes in drinking appeared to result in concomitant changes in depressive symptoms such that PLWH who switch to higher or lower level drinking experience a similar alteration in their depressive symptoms (Sullivan et al., 2011). While the association between alcohol use and depression appears to be similar for people with and without HIV (Sullivan et al., 2011), the common co-occurrence of these conditions and both of their influences on poor HIV-related outcomes is likely to have at least an additive effect on PLWH. Psychological trauma, stress, and anxiety, which often co-occur with heavy alcohol use (Pence et al., 2008), are associated with adverse HIV-related outcomes (Chander et al., 2006a; Leserman, 2003; Leserman, 2008; Leserman et al., 2008; Machtinger et al., 2012; Liebschutz et al., 2000). Research is needed to understand the potential synergistic effects of alcohol use with these common mental health conditions among PLWH, including investigation of issues related to temporality, mechanisms, prevention, and treatment in the context of co-occurring alcohol use and mental health conditions (Chander et al., 2006a). Finally, research on associations between alcohol use and co-occurring serious mental illness among PLWH is needed.

Morbidity and Mortality

While mortality rates among PLWH have declined (Simmons et al., 2013; Hasse et al., 2011), PLWH are still at higher risk for mortality, as well as physiological frailty, relative to those without HIV (Justice and Falutz, 2014). Alcohol use is associated with increased risk of mortality in general samples (Kinder et al., 2009; Rehm et al., 2001; Rehm et al., 2003), as well as specifically among PLWH (Neblett et al., 2011; Wandeler et al., 2016; Justice et al., 2016). While most large observational studies, including one among PLWH (Wandeler et al., 2016), have identified a J-shaped association between alcohol use and all-cause mortality (with a protective effect of low-level or “moderate” drinking), a recent meta-analysis (not HIV-specific) of 87 studies including over 3 million people, which adjusted for study design characteristics known to bias results, found no protective effect of low-level drinking (Stockwell et al., 2016). A recent study demonstrated an association between alcohol use and both physiological frailty and mortality among PLWH but did not include a non-drinker

group, and thus was not able to assess the effects of low-level or “moderate” drinking on morbidity and mortality (Justice et al., 2016). However, this study’s findings suggested that PLWH experienced risk at lower levels of alcohol use relative to uninfected individuals (>30 drinks per month for PLWH versus >70 drinks per month for those without HIV for mortality) (Justice et al., 2016). These findings are aligned with previous research, which has identified greater blood alcohol concentration (McCance-Katz et al., 2012) and increased likelihood of feeling “a buzz” (McGinnis et al., 2016) at similar levels of drinking for PLWH relative to those without HIV. Together these findings suggest the possibility that alcohol’s influence on health is greater among PLWH than it is among uninfected populations and further highlight the need to gain clarity and precision regarding how and at what levels alcohol use influences health among PLWH, as well as underlying mechanisms.

HIV and Alcohol in a Social Context, Intersectionality, and Potential Subgroup Variation

Both HIV and alcohol use are stigmatized conditions about which discrimination is common. Therefore, the social and environmental contexts in which PLWH who drink live is likely to influence outcomes (Stokols, 1992). Moreover, both alcohol use and HIV disproportionately impact vulnerable populations, including people living in poverty and poor resource settings, those who experience multiple forms of social disadvantage, racial/ethnic minorities, and MSM (Grant et al., 2015; Hall et al., 2009; Lansky et al., 2010; Mulia et al., 2009; Pellowski et al., 2013). Many of these characteristics overlap within individuals (e.g., racial/ethnic minorities disproportionately live in poor resource settings). Intersectionality theory and syndemic perspectives (Kelly, 2009; Talman et al., 2013; Pellowski et al., 2013) suggest that the chronic psychosocial stresses observed over the life course for these populations are likely to synergistically influence health over time and, thus, may result in health disparities. Research should investigate subgroup variation resulting from lived experiences in associations between alcohol use and HIV-related outcomes, to ensure risks specific to vulnerable populations are understood.

Addressing/Reducing Alcohol Use among PLWH

Despite its substantial burden on HIV and related illnesses, alcohol use is often overlooked in HIV care (Metsch et al., 2008; Conigliaro et al., 2003; Chander et al., 2016; Korthuis et al., 2008 ; Korthuis et al., 2011; Strauss et al., 2009a; Strauss et al., 2012). Research suggests substantial barriers to HIV care providers’ addressing alcohol use with their patients (Strauss et al., 2009a; Strauss et al., 2009b). Both PLWH and HIV care providers perceive alcohol use as a low priority for HIV clinical care relative to other domains of care (Fredericksen et al., 2015).

Efforts to describe and address deficiencies in alcohol-related care among PLWH are occurring in several domains. First, ongoing health services research is being conducted to assess whether, in healthcare settings in which alcohol-related care is common (e.g., the Veterans Health Administration), differences in receipt of alcohol-related care exist across HIV status (Williams and Bradley, 2015). Studies are also underway to evaluate the comparative effectiveness of alcohol treatments among PLWH (Kraemer, 2013). Second,

several studies of alcohol-related interventions have been tested among PLWH with unhealthy alcohol use (Papas et al., 2011; Samet et al., 2015; Hasin et al., 2013; Samet and Walley, 2010; Chander et al., 2015; World Health Organization, 2011), and others have been piloted (Edelman et al., 2012) or are ongoing (Edelman et al., 2016; Satre, 2012; McCaul, 2016; Armstrong et al., 1998; Parry et al., 2014; Papas et al., 2010).

Research to date is mixed regarding the efficacy of alcohol-related interventions among PLWH (Samet and Walley, 2010). Three interventions studied to date have had positive findings (Papas et al., 2011; Hasin et al., 2013; Chander et al., 2015). First, Papas, et al tested the preliminary efficacy of a culturally adapted 6-session cognitive-behavioral therapy intervention delivered by paraprofessionals relative to a usual care, assessment-only control condition among PLWH receiving care (n=102) at a Kenyan outpatient HIV clinic (Papas et al., 2011). The intervention was associated with a reduction in percent drinking days and number of drinking days after 30 days and greater abstinence after 90 days (Papas et al., 2011). In the U.S., Hasin et al recruited 258 heavy drinkers with HIV from a large urban HIV primary care clinic. They compared an active control condition (educational materials and advice to reduce drinking at baseline and 30- and 60-day follow-ups) to two intervention arms based on motivational interviewing (MI) (Hasin et al., 2013). The MI arm included a single 20–25 minute session of counselor-delivered MI with two 15-minute boosters at 30- and 60-days; the other arm (“MI+”) additionally included daily calls with “HealthCall”—an interactive voice-response (IVR) offering personalized feedback (Hasin et al., 2013). Overall group differences in the number of drinking days were observed at 60 days, with both MI and MI+ associated with fewer drinking days relative to the control group. No differences were observed between MI and MI+ in the full sample, although MI+ was associated with fewer drinking days relative to MI among participants meeting criteria for alcohol dependence at baseline (but not among those who did not) (Hasin et al., 2013). Most recently, Chander et al tested a 2-session (plus 2 booster calls) brief intervention relative to a control among 74 HIV+ women from an urban HIV clinic and found that the intervention was associated with reduced frequency of alcohol use at 90 days (Chander et al., 2015). While long-term outcomes, and outcomes beyond self-reported consumption (which is highly susceptible to social desirability bias) have not been measured, findings from these intervention studies suggest that alcohol use may be modifiable among PLWH who drink heavily. Moreover, the IVR component of Hasin’s MI+ has now been enhanced to be delivered via smartphones, with strong findings regarding acceptability and feasibility (Hasin et al., 2014). To the extent it is feasible, future research should measure long-term and objective clinical outcomes of these types of interventions, as well as to investigate subgroup differences in responses, and their potential for translation into real-world settings. Although likely to be dependent on specific contexts and settings, implementing multi-session interventions across settings may be met with substantial barriers (Papas et al., 2011; Papas et al., 2010). However, repeated brief interventions or mobile applications coupled with a single-session of MI may hold promise for translation into HIV clinics and resource poor settings.

Summary and Future Directions

Alcohol use is common among PLWH, and is associated with HIV acquisition and transmission, lack of viral suppression, common comorbid conditions, and ultimately morbidity and mortality via both biological (e.g., oxidative stress) and behavioral (e.g., ART adherence) mechanisms (see: Figure 1). Associations between alcohol use and HIV-related care and outcomes are likely to be influenced by independent and/or overlapping lived experiences of vulnerable sub-populations that are disproportionately represented among PLWH (e.g., sexual minorities, homeless populations). Interventions to modify drinking and improve HIV-related risks and outcomes have been tested in various populations with limited success to date.

Our review suggests further research in several areas. Consistent with NIH priorities (National Institutes of Health, 2015), substantial research is needed to better understand alcohol's influence on co-infections and comorbid conditions that are common among PLWH, including elucidating the mechanisms by which alcohol use is impacting comorbidities. While research suggests the possibility that no level of alcohol use may be considered "safe" for PLWH (Justice et al.), given the high prevalence of alcohol use among PLWH, it will be necessary to better understand which patterns of use substantially increase risks and how. These investigations will require measures of all levels of drinking (including non-drinking). This work should also quantify how changes in alcohol use can alter the course of HIV-related outcomes. Studies that assess whether state-of-the-art medications are safe for co-infected PLWH who drink alcohol are needed in order to ensure adequate treatment of common co-infections. While new medications have emerged that appear to be highly effective at controlling both HCV and TB, contraindications of previous medications, as well as resulting stigma are likely to be barriers to treatment for PLWH who drink and have common co-infections.

Future studies of alcohol's influence on common comorbidities among PLWH will need to explicitly address measurement issues plaguing the extant literature. While many studies of adverse outcomes among PLWH have not addressed alcohol use at all, many that have, have used alcohol measurements of varying quality. Future studies among PLWH, especially those of interventions which can be prone to mis-report, should include validated self-report measures of alcohol use. While diary-based methods of assessing consumption, such as Timeline Follow Back methods (Sobell et al., 1996), are considered state-of-the-art self-report measures, validated alcohol screens such as the Alcohol Use Disorders Identification Test (AUDIT) (Babor et al., 1989) and its abbreviated versions (i.e., the "AUDIT-C" made up of the first three questions regarding consumption (Bush et al., 1998; Bradley et al., 2007; Dawson et al., 2005), and the "AUDIT-3" comprised of the third question assessing heavy episodic drinking (Smith et al., 2009)) are also commonly-used self-report-based methods. When possible, due to limitations of self-report measures (e.g., social desirability bias, recall bias), biological measurements such as phosphatidylethanol (PEth) are recommended (though these also have limitations such as lack of sensitivity for low but still risky amounts) (Hahn et al., 2012; Wurst et al., 2005; Bajunirwe et al., 2014; Asiimwe et al., 2015), as well as objective measures of alcohol-related health consequences. Selection of optimal measures will depend on populations and contexts studied, and the purpose of the measurement. At all

phases of this research, subgroup variation in associations should be identified, such that risks specific to vulnerable populations that are over-represented among PLWH (e.g., racial/ethnic minorities, MSM, people with co-occurring substance use, mental health conditions, and homelessness, and the intersections of these identities) are understood.

This research should serve as foundational work that can be translated into a next generation of intervention studies. Future interventions should address alcohol use at individual and policy levels and at all phases from transmission to treatment of HIV. Because alcohol use is prevalent and persistent among PLWH even after exposure to interventions, future interventions should focus on improving engagement in and adherence to HIV care, reducing acute health care utilization, and mitigating comorbid complications in the context of continued drinking. For instance, given the effectiveness of ART for treating HIV, and that alcohol use is a potential modifiable risk factor for non-adherence, this can be considered a challenging but reasonable area for continued research. Given that reductions in alcohol use observed in earlier studies have not translated into improvements in ART, interventions aimed at clustered risks and focused on clearly identified mechanisms linking alcohol use and ART are likely needed. Future interventions should be designed to provide information regarding the potential magnitude of benefit in HIV-related care and outcomes that may be obtained by addressing alcohol use. Once effective interventions are identified, identifying ways to implement them in real world health care settings will be important, as well as understanding barriers to doing so.

By making further progress on understanding the ramifications of alcohol use among PLWH in the era of HIV as a chronic condition, research should inform how we can mitigate transmission, achieve viral suppression and avoid exacerbating common comorbidities of HIV and alcohol use while making progress toward the 90-90-90 goals set out by the WHO (Joint United Nations Programme on HIV/AIDS, 2014).

REFERENCES CITED

- Alter MJ. Epidemiology of viral hepatitis and HIV co-infection. *Journal of Hepatology*. 2006; 44:S6–S9. [PubMed: 16352363]
- Amedee AM, Nichols WA, Robichaux S, et al. Chronic alcohol abuse and HIV disease progression: studies with the non-human primate model. *Curr HIV Res*. 2014; 12:243–253. [PubMed: 25053367]
- Amruth G, Praveen-Kumar S, Nataraju B, et al. HIV Associated Sensory Neuropathy. *J Clin Diagn Res*. 2014; 8:MC04–MC07.
- Anand P, Springer SA, Copenhaver MM, et al. Neurocognitive impairment and HIV risk factors: a reciprocal relationship. *AIDS Behav*. 2010; 14:1213–1226. [PubMed: 20232242]
- Antinori A, Arendt G, Becker JT, et al. Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*. 2007; 69:1789–1799. [PubMed: 17914061]
- Armah KA, McGinnis K, Baker J, et al. HIV status, burden of comorbid disease, and biomarkers of inflammation, altered coagulation, and monocyte activation. *Clinical Infectious Diseases*. 2012; 55:126–136. [PubMed: 22534147]
- Armstrong MA, Midanik LT, Klatsky AL. Alcohol consumption and utilization of health services in a health maintenance organization. *Medical Care*. 1998; 36:1599–1605. [PubMed: 9821947]
- Asiimwe SB, Fatch R, Emenyonu NI, et al. Comparison of Traditional and Novel Self-Report Measures to an Alcohol Biomarker for Quantifying Alcohol Consumption Among HIV-Infected Adults in Sub-Saharan Africa. *Alcoholism, Clinical and Experimental Research*. 2015; 39:1518–1527.

- Athyros VG, Liberopoulos EN, Mikhailidis DP, et al. Association of drinking pattern and alcohol beverage type with the prevalence of metabolic syndrome, diabetes, coronary heart disease, stroke, and peripheral arterial disease in a Mediterranean cohort. *Angiology*. 2007; 58:689–697. [PubMed: 18216378]
- Attonito J. The influence of neurocognitive impairment, alcohol and other drug (AOD) use, and psychosocial factors on antiretroviral treatment adherence, service utilization and viral load among HIV-seropositive adults; Dissertation. Florida International University, FIU Digital Commons. 2013
- Attonito JM, Devieux JG, Lerner BD, et al. Exploring Substance Use and HIV Treatment Factors Associated with Neurocognitive Impairment among People Living with HIV/AIDS. *Front Public Health*. 2014; 2:105. [PubMed: 25157345]
- Azar MM, Springer SA, Meyer JP, et al. A systematic review of the impact of alcohol use disorders on HIV treatment outcomes, adherence to antiretroviral therapy and health care utilization. *Drug and Alcohol Dependence*. 2010a; 112:178–193. [PubMed: 20705402]
- Azar MM, Springer SA, Meyer JP, et al. A systematic review of the impact of alcohol use disorders on HIV treatment outcomes, adherence to antiretroviral therapy and health care utilization. *Drug and Alcohol Dependence*. 2010b; 112:178–193. [PubMed: 20705402]
- Babor, TF.; de la Fuente, JR.; Saunders, J., et al. AUDIT: Alcohol Use Disorders Identification Test: guidelines for use in primary health care. Geneva: World Health Organization; 1989.
- Bagby GJ, Zhang P, Purcell JE, et al. Chronic binge ethanol consumption accelerates progression of simian immunodeficiency virus disease. *Alcoholism, Clinical and Experimental Research*. 2006; 30:1781–1790.
- Bajunirwe F, Haberer JE, Boum Y 2nd, et al. Comparison of self-reported alcohol consumption to phosphatidylethanol measurement among HIV-infected patients initiating antiretroviral treatment in southwestern Uganda. *PLoS One*. 2014; 9:e113152. [PubMed: 25436894]
- Balogopal A, Philp FH, Astemborski J, et al. Human immunodeficiency virus-related microbial translocation and progression of hepatitis C. *Gastroenterology*. 2008; 135:226–233. [PubMed: 18457674]
- Baum MK, Rafie C, Lai S, et al. Alcohol use accelerates HIV disease progression. *AIDS Research and Human Retroviruses*. 2010; 26:511–518. [PubMed: 20455765]
- Bedimo RJ, McGinnis KA, Dunlap M, et al. Incidence of non-AIDS-defining malignancies in HIV-infected versus noninfected patients in the HAART era: impact of immunosuppression. *Journal of Acquired Immune Deficiency Syndromes*. 2009; 52:203–208. [PubMed: 19617846]
- Bengtson AM, L'Engle K, Mwarogo P, et al. Levels of alcohol use and history of HIV testing among female sex workers in Mombasa, Kenya. *AIDS Care*. 2014; 26:1619–1624. [PubMed: 25040114]
- Berg KM, Cooperman NA, Newville H, et al. Self-efficacy and depression as mediators of the relationship between pain and antiretroviral adherence. *AIDS Care*. 2009; 21:244–248. [PubMed: 19229695]
- Berg KM, Kunins HV, Jackson JL, et al. Association Between Alcohol Consumption and Both Osteoporotic Fracture and Bone Density. *The American journal of medicine*. 2008; 121:406–418. [PubMed: 18456037]
- Bilal U, Lau B, Lazo M, et al. Interaction Between Alcohol Consumption Patterns, Antiretroviral Therapy Type, and Liver Fibrosis in Persons Living with HIV. *Aids Patient Care and STDS*. 2016; 30:200–207. [PubMed: 27158847]
- Bing EG, Burnam MA, Longshore D, et al. Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Archives of General Psychiatry*. 2001; 58:721–728. [PubMed: 11483137]
- Bor J, Geldsetzer P, Venkataramani A, et al. Quasi-experiments to establish causal effects of HIV care and treatment and to improve the cascade of care. *Curr Opin HIV AIDS*. 2015; 10:495–501. [PubMed: 26371463]
- Borges AH, Dubrow R, Silverberg MJ. Factors contributing to risk for cancer among HIV-infected individuals, and evidence that earlier combination antiretroviral therapy will alter this risk. *Curr Opin HIV AIDS*. 2014; 9:34–40. [PubMed: 24225382]

- Bradley KA, DeBenedetti AF, Volk RJ, et al. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcoholism, Clinical and Experimental Research*. 2007; 31:1208–1217.
- Braithwaite RS, Fang Y, Tate J, et al. Do Alcohol Misuse, Smoking, and Depression Vary Concordantly or Sequentially? A Longitudinal Study of HIV-Infected and Matched Uninfected Veterans in Care. *AIDS Behav*. 2016; 20:566–572. [PubMed: 26187007]
- Braithwaite RS, McGinnis KA, Conigliaro J, et al. A temporal and dose-response association between alcohol consumption and medication adherence among veterans in care. *Alcoholism, Clinical and Experimental Research*. 2005; 29:1190–1197.
- Brenchley JM, Price DA, Schacker TW, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nature Medicine*. 2006; 12:1365–1371.
- Bryant KJ, Nelson S, Braithwaite RS, et al. Intergrating HIV/AIDS and alcohol research. *Alcohol Res Health*. 2010; 33:167–178. [PubMed: 23584058]
- Bush K, Kivlahan DR, McDonell MB, et al. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Archives of Internal Medicine*. 1998; 158:1789–1795. [PubMed: 9738608]
- Butt AA, McGinnis K, Rodriguez-Barradas MC, et al. HIV infection and the risk of diabetes mellitus. *AIDS*. 2009; 23:1227–1234. [PubMed: 19444074]
- Calvo M, Martinez E. Update on metabolic issues in HIV patients. *Curr Opin HIV AIDS*. 2014; 9:332–339. [PubMed: 24824886]
- Carrasco MA, Esser MB, Sparks A, et al. HIV-Alcohol Risk Reduction Interventions in Sub-Saharan Africa: A Systematic Review of the Literature and Recommendations for a Way Forward. *AIDS Behav*. 2016; 20:484–503. [PubMed: 26511865]
- Carrico AW, Hunt PW, Emenyonu NI, et al. Unhealthy Alcohol Use is Associated with Monocyte Activation Prior to Starting Antiretroviral Therapy. *Alcoholism, Clinical and Experimental Research*. 2015; 39:2422–2426.
- Carrieri MP, Protopopescu C, Le Moing V, et al. Impact of immunodepression and moderate alcohol consumption on coronary and other arterial disease events in an 11-year cohort of HIV-infected patients on antiretroviral therapy. *BMJ Open*. 2012;2.
- Cassol E, Malfeld S, Mahasha P, et al. Persistent microbial translocation and immune activation in HIV-1-infected South Africans receiving combination antiretroviral therapy. *Journal of Infectious Diseases*. 2010; 202:723–733. [PubMed: 20629534]
- Cawthon PM, Harrison SL, Barrett-Connor E, et al. Alcohol intake and its relationship with bone mineral density, falls, and fracture risk in older men. *Journal of the American Geriatrics Society*. 2006; 54:1649–1657. [PubMed: 17087690]
- Centers for Disease Control and Prevention. HIV Surveillance Report: Diagnoses of HIV Infection and AIDS in the United States and Dependent Areas. 2011;23.
- Centers for Disease Control and Prevention. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data - United States and 6 U.S. dependent areas - 2010 HIV Surveillance Supplemental Report. 2012
- Chander G, Himelhoch S, Moore RD. Substance abuse and psychiatric disorders in HIV-positive patients: epidemiology and impact on antiretroviral therapy. *Drugs*. 2006a; 66:769–789. [PubMed: 16706551]
- Chander G, Hutton HE, Lau B, et al. Brief Intervention Decreases Drinking Frequency in HIV-Infected, Heavy Drinking Women: Results of a Randomized Controlled Trial. *Journal of Acquired Immune Deficiency Syndromes*. 2015; 70:137–145. [PubMed: 25967270]
- Chander G, Lau B, Moore RD. Hazardous alcohol use: a risk factor for non-adherence and lack of suppression in HIV infection. *Journal of Acquired Immune Deficiency Syndromes*. 2006b; 43:411–417. [PubMed: 17099312]
- Chander G, Monroe AK, Crane HM, et al. HIV primary care providers-Screening, knowledge, attitudes and behaviors related to alcohol interventions. *Drug and Alcohol Dependence*. 2016
- Chandiwana SK, Sebit MB, Latif AS, et al. Alcohol consumption in HIV-I infected persons: a study of immunological markers, Harare, Zimbabwe. *Central African Journal of Medicine*. 1999; 45:303–308. [PubMed: 10892457]

- Cheng DM, Libman H, Bridden C, et al. Alcohol consumption and lipodystrophy in HIV-infected adults with alcohol problems. *Alcohol*. 2009; 43:65–71. [PubMed: 19185212]
- Cherpitel CJ, Ye Y, Bond J, et al. Alcohol attributable fraction for injury morbidity from the dose-response relationship of acute alcohol consumption: emergency department data from 18 countries. *Addiction*. 2015; 110:1724–1732. [PubMed: 26119350]
- Cioe PA, Baker J, Kojic EM, et al. Elevated Soluble CD14 and Lower D-Dimer Are Associated With Cigarette Smoking and Heavy Episodic Alcohol Use in Persons Living With HIV. *Journal of Acquired Immune Deficiency Syndromes*. 2015; 70:400–405. [PubMed: 26181818]
- Coffin PO, Galea S, Ahern J, et al. Opiates, cocaine and alcohol combinations in accidental drug overdose deaths in New York City, 1990-98. *Addiction*. 2003; 98:739–747. [PubMed: 12780362]
- Coffin PO, Tracy M, Bucciarelli A, et al. Identifying injection drug users at risk of nonfatal overdose. *Academic Emergency Medicine*. 2007; 14:616–623. [PubMed: 17554010]
- Cohen SM, Hu X, Sweeney P, et al. HIV viral suppression among persons with varying levels of engagement in HIV medical care, 19 US jurisdictions. *Journal of Acquired Immune Deficiency Syndromes*. 2014; 67:519–527. [PubMed: 25230292]
- Conen A, Wang Q, Glass TR, et al. Association of alcohol consumption and HIV surrogate markers in participants of the swiss HIV cohort study. *Journal of Acquired Immune Deficiency Syndromes*. 2013; 64:472–478. [PubMed: 23892243]
- Conigliaro J, Gordon AJ, McGinnis KA, et al. How harmful is hazardous alcohol use and abuse in HIV infection: do health care providers know who is at risk? *Journal of Acquired Immune Deficiency Syndromes*. 2003; 33:521–525. [PubMed: 12869842]
- Cook JA, Burke-Miller JK, Cohen MH, et al. Crack cocaine, disease progression, and mortality in a multicenter cohort of HIV-1 positive women. *AIDS*. 2008; 22:1355–1363. [PubMed: 18580615]
- Cooper CL, Cameron DW. Effect of alcohol use and highly active antiretroviral therapy on plasma levels of hepatitis C virus (HCV) in patients coinfecting with HIV and HCV. *Clinical Infectious Diseases*. 2005; (41 Suppl 1):S105–S109.
- Cooperman NA. Current research on cigarette smoking among people with HIV. *Curr Addict Rep*. 2016; 3:19–26.
- Crothers K, Goulet JL, Rodriguez-Barradas MC, et al. Impact of cigarette smoking on mortality in HIV-positive and HIV-negative veterans. *AIDS Education and Prevention*. 2009; 21:40–53. [PubMed: 19537953]
- Crothers K, Griffith TA, McGinnis KA, et al. The impact of cigarette smoking on mortality, quality of life, and comorbid illness among HIV-positive veterans. *Journal of General Internal Medicine*. 2005; 20:1142–1145. [PubMed: 16423106]
- Cruess DG, Douglas SD, Petitto JM, et al. Association of resolution of major depression with increased natural killer cell activity among HIV-seropositive women. *American Journal of Psychiatry*. 2005; 162:2125–2130. [PubMed: 16263853]
- Cruess DG, Petitto JM, Leserman J, et al. Depression and HIV infection: impact on immune function and disease progression. *CNS Spectr*. 2003; 8:52–58. [PubMed: 12627049]
- Crum-Cianflone N, Roediger MP, Eberly L, et al. Increasing rates of obesity among HIV-infected persons during the HIV epidemic. *PLoS One*. 2010; 5:e10106. [PubMed: 20419086]
- Crum-Cianflone N, Tejedor R, Medina S, et al. Obesity among patients with HIV: the latest epidemic. *Aids Patient Care and STDS*. 2008; 22:925–930. [PubMed: 19072098]
- Cunningham WE, Sohler NL, Tobias C, et al. Health services utilization for people with HIV infection: comparison of a population targeted for outreach with the U.S. population in care. *Medical Care*. 2006; 44:1038–1047. [PubMed: 17063136]
- Cutrell J, Bedimo R. Non-AIDS-defining cancers among HIV-infected patients. *Curr HIV/AIDS Rep*. 2013; 10:207–216. [PubMed: 23743643]
- D'Souza G, Carey TE, William WN Jr, et al. Epidemiology of head and neck squamous cell cancer among HIV-infected patients. *Journal of Acquired Immune Deficiency Syndromes*. 2014; 65:603–610. [PubMed: 24326607]
- Dawson DA, Grant BF, Stinson FS, et al. Effectiveness of the derived Alcohol Use Disorders Identification Test (AUDIT-C) in screening for alcohol use disorders and risk drinking in the US general population. *Alcoholism, Clinical and Experimental Research*. 2005; 29:844–854.

- De Wit S, Sabin CA, Weber R, et al. Incidence and risk factors for new-onset diabetes in HIV-infected patients: the Data Collection on Adverse Events of Anti-HIV Drugs (D:A:D) study. *Diabetes Care*. 2008; 31:1224–1229. [PubMed: 18268071]
- Deeken JF, Tjen ALA, Rudek MA, et al. The rising challenge of non-AIDS-defining cancers in HIV-infected patients. *Clinical Infectious Diseases*. 2012; 55:1228–1235. [PubMed: 22776851]
- Deeks SG. Immune dysfunction, inflammation, and accelerated aging in patients on antiretroviral therapy. *Topics in HIV medicine : a publication of the International AIDS Society, USA*. 2009; 17:118–123.
- Deeks SG. HIV infection, inflammation, immunosenescence, and aging. *Annual Review of Medicine*. 2011; 62:141–155.
- Deiss RG, Mesner O, Agan BK, et al. Characterizing the Association Between Alcohol and HIV Virologic Failure in a Military Cohort on Antiretroviral Therapy. *Alcoholism, Clinical and Experimental Research*. 2016; 40:529–535.
- Delker E, Brown Q, Hasin DS. Alcohol Consumption in Demographic Subpopulations: An Epidemiologic Overview. *Alcohol Research*. 2016; 38:7–15. [PubMed: 27159807]
- Dillon SM, Lee EJ, Kotter CV, et al. An altered intestinal mucosal microbiome in HIV-1 infection is associated with mucosal and systemic immune activation and endotoxemia. *Mucosal Immunol*. 2014; 7:983–994. [PubMed: 24399150]
- Douek D. HIV disease progression: immune activation, microbes, and a leaky gut. *Topics in HIV medicine : a publication of the International AIDS Society, USA*. 2007; 15:114–117.
- Eaton LA, Kalichman SC, Kenny DA, et al. A reanalysis of a behavioral intervention to prevent incident HIV infections: including indirect effects in modeling outcomes of Project EXPLORE. *AIDS Care*. 2013; 25:805–811. [PubMed: 23245226]
- Edelman EJ, Dinh A, Radulescu R, et al. Combining rapid HIV testing and a brief alcohol intervention in young unhealthy drinkers in the emergency department: a pilot study. *American Journal of Drug and Alcohol Abuse*. 2012; 38:539–543. [PubMed: 22794939]
- Edelman EJ, Hansen NB, Cutter CJ, et al. Implementation of integrated stepped care for unhealthy alcohol use in HIV clinics. *Addiction Science & Clinical Practice*. 2016; 11:1. [PubMed: 26763048]
- Elbirt D, Mahlab-Guri K, Bezalel-Rosenberg S, et al. HIV-associated neurocognitive disorders (HAND). *Isr Med Assoc J*. 2015; 17:54–59. [PubMed: 25739180]
- Erlanson KM, Allshouse AA, Jankowski CM, et al. Risk factors for falls in HIV-infected persons. *Journal of Acquired Immune Deficiency Syndromes*. 2012; 61:484–489. [PubMed: 23143526]
- Eskild A, Petersen G. Cigarette smoking and drinking of alcohol are not associated with rapid progression to acquired immunodeficiency syndrome among homosexual men in Norway. *Scandinavian Journal of Social Medicine*. 1994; 22:209–212. [PubMed: 7846480]
- Evans DL, Ten Have TR, Douglas SD, et al. Association of depression with viral load, CD8 T lymphocytes, and natural killer cells in women with HIV infection. *American Journal of Psychiatry*. 2002; 159:1752–1759. [PubMed: 12359683]
- Fatch R, Bellows B, Bagenda F, et al. Alcohol consumption as a barrier to prior HIV testing in a population-based study in rural Uganda. *AIDS Behav*. 2013; 17:1713–1723. [PubMed: 22878790]
- Fergusson DM, Boden JM, Horwood LJ. Tests of causal links between alcohol abuse or dependence and major depression. *Archives of General Psychiatry*. 2009; 66:260–266. [PubMed: 19255375]
- Fredericksen RJ, Edwards TC, Merlin JS, et al. Patient and provider priorities for self-reported domains of HIV clinical care. *AIDS Care*. 2015; 27:1255–1264. [PubMed: 26304263]
- Freedman ND, Schatzkin A, Leitzmann MF, et al. Alcohol and head and neck cancer risk in a prospective study. *British Journal of Cancer*. 2007; 96:1469–1474. [PubMed: 17387340]
- Freiberg, M.; Bebu, I.; Tracy, R., et al. D-dimer doesn't return to pre-HIV levels after therapy and is linked with HANA events; Conference on Retroviruses and Opportunistic Infections; Seattle, WA. 2015. p. 240-241.
- Freiberg MS, Cabral HJ, Heeren TC, et al. Alcohol consumption and the prevalence of the Metabolic Syndrome in the US.: a cross-sectional analysis of data from the Third National Health and Nutrition Examination Survey. *Diabetes Care*. 2004; 27:2954–2959. [PubMed: 15562213]

- Freiberg MS, Chang CC, Kuller LH, et al. HIV infection and the risk of acute myocardial infarction. *JAMA internal medicine*. 2013; 173:614–622. [PubMed: 23459863]
- Freiberg MS, McGinnis KA, Kraemer K, et al. The association between alcohol consumption and prevalent cardiovascular diseases among HIV-infected and HIV-uninfected men. *Journal of Acquired Immune Deficiency Syndromes*. 2010; 53:247–253. [PubMed: 20009766]
- Fuster D, Cheng DM, Quinn EK, et al. Chronic hepatitis C virus infection is associated with all-cause and liver-related mortality in a cohort of HIV-infected patients with alcohol problems. *Addiction*. 2014; 109:62–70. [PubMed: 24112091]
- Fuster D, Tsui JI, Cheng DM, et al. Impact of lifetime alcohol use on liver fibrosis in a population of HIV-infected patients with and without hepatitis C coinfection. *Alcoholism: Clinical and Experimental Research*. 2013
- Galvan FH, Bing EG, Fleishman JA, et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV cost and services utilization study. *Journal of Studies on Alcohol*. 2002; 63:179–186. [PubMed: 12033694]
- Ghebremichael M, Paintsil E, Ickovics JR, et al. Longitudinal association of alcohol use with HIV disease progression and psychological health of women with HIV. *AIDS Care*. 2009; 21:834–841. [PubMed: 20024739]
- Giordano TP, Visnegarwala F, White AC Jr, et al. Patients referred to an urban HIV clinic frequently fail to establish care: factors predicting failure. *AIDS Care*. 2005; 17:773–783. [PubMed: 16036264]
- Goodness TM, Palfai TP, Cheng DM, et al. Depressive symptoms and antiretroviral therapy (ART) initiation among HIV-infected Russian drinkers. *AIDS Behav*. 2014; 18:1085–1093. [PubMed: 24337725]
- Grant BF, Goldstein RB, Saha TD, et al. Epidemiology of DSM-5 Alcohol Use Disorder: Results From the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry*. 2015; 72:757–766. [PubMed: 26039070]
- Green JE, Saveanu RV, Bornstein RA. The effect of previous alcohol abuse on cognitive function in HIV infection. *American Journal of Psychiatry*. 2004; 161:249–254. [PubMed: 14754773]
- Green TC, Kershaw T, Lin H, et al. Patterns of drug use and abuse among aging adults with and without HIV: a latent class analysis of a US Veteran cohort. *Drug and Alcohol Dependence*. 2010; 110:208–220. [PubMed: 20395074]
- Guaraldi G, Orlando G, Zona S, et al. Premature Age-Related Comorbidities Among HIV-Infected Persons Compared With the General Population. *Clinical Infectious Diseases*. 2011; 53:1120–1126. [PubMed: 21998278]
- Hahn JA, Dobkin LM, Mayanja B, et al. Phosphatidylethanol (PEth) as a biomarker of alcohol consumption in HIV-positive patients in sub-Saharan Africa. *Alcoholism, Clinical and Experimental Research*. 2012; 36:854–862.
- Hahn JA, Samet JH. Alcohol and HIV disease progression: weighing the evidence. *Curr HIV/AIDS Rep*. 2010; 7:226–233. [PubMed: 20814765]
- Hall HI, Geduld J, Boulos D, et al. Epidemiology of HIV in the United States and Canada: current status and ongoing challenges. *Journal of Acquired Immune Deficiency Syndromes*. 2009; (51 Suppl 1):S13–S20. [PubMed: 19384096]
- Hansen AB, Gerstoft J, Kronborg G, et al. Incidence of low and high-energy fractures in persons with and without HIV infection: a Danish population-based cohort study. *AIDS*. 2012; 26:285–293. [PubMed: 22095195]
- Harezlak J, Buchthal S, Taylor M, et al. Persistence of HIV-associated cognitive impairment, inflammation, and neuronal injury in era of highly active antiretroviral treatment. *AIDS*. 2011; 25:625–633. [PubMed: 21297425]
- Hasin DS, Aharonovich E, Greenstein E. HealthCall for the smartphone: technology enhancement of brief intervention in HIV alcohol dependent patients. *Addiction Science & Clinical Practice*. 2014; 9:5. [PubMed: 24533631]
- Hasin DS, Aharonovich E, O’Leary A, et al. Reducing heavy drinking in HIV primary care: a randomized trial of brief intervention, with and without technological enhancement. *Addiction*. 2013; 108:1230–1240. [PubMed: 23432593]

- Hasse B, Ledergerber B, Furrer H, et al. Morbidity and Aging in HIV-Infected Persons: The Swiss HIV Cohort Study. *Clinical Infectious Diseases*. 2011; 53:1130–1139. [PubMed: 21998280]
- Hauser KF, Knapp PE. Interactions of HIV and drugs of abuse: the importance of glia, neural progenitors, and host genetic factors. *International Review of Neurobiology*. 2014; 118:231–313. [PubMed: 25175867]
- Heaton RK, Clifford DB, Franklin DR Jr, et al. HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology*. 2010; 75:2087–2096. [PubMed: 21135382]
- Heaton RK, Franklin DR, Ellis RJ, et al. HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *Journal of Neurovirology*. 2011; 17:3–16. [PubMed: 21174240]
- Hendershot CS, Stoner SA, Pantalone DW, et al. Alcohol use and antiretroviral adherence: review and meta-analysis. *Journal of Acquired Immune Deficiency Syndromes*. 2009; 52:180–202. [PubMed: 19668086]
- High KP, Brennan-Ing M, Clifford DB, et al. HIV and aging: state of knowledge and areas of critical need for research. A report to the NIH Office of AIDS Research by the HIV and Aging Working Group. *Journal of Acquired Immune Deficiency Syndromes*. 2012; (60 Suppl 1):S1–S18. [PubMed: 22688010]
- Hile SJ, Feldman MB, Alexy ER, et al. Recent Tobacco Smoking is Associated with Poor HIV Medical Outcomes Among HIV-Infected Individuals in New York. *AIDS Behav*. 2016
- Howard AA, Arnsten JH, Gourevitch MN. Effect of alcohol consumption on diabetes mellitus: a systematic review. *Annals of Internal Medicine*. 2004; 140:211–219. [PubMed: 14757619]
- Hu J, Han H, Lau MY, et al. Effects of combined alcohol and anti-HIV drugs on cellular stress responses in primary hepatocytes and hepatic stellate and kupffer cells. *Alcoholism, Clinical and Experimental Research*. 2015; 39:11–20.
- Hunt PW, Sinclair E, Rodriguez B, et al. Gut epithelial barrier dysfunction and innate immune activation predict mortality in treated HIV infection. *Journal of Infectious Diseases*. 2014; 210:1228–1238. [PubMed: 24755434]
- Jacobson DL, Tang AM, Spiegelman D, et al. Incidence of metabolic syndrome in a cohort of HIV-infected adults and prevalence relative to the US population (National Health and Nutrition Examination Survey). *Journal of acquired immune deficiency syndromes (1999)*. 2006; 43:458–466. [PubMed: 16980905]
- Joint United Nations Programme on HIV/AIDS. [accessed February 15, 2016] 90-90-90—an ambitious treatment target to help end the AIDS epidemic. 2014. <http://www.unaids.org/en/resources/documents/2014/90-90-90>
- Justice A, Falutz J. Aging and HIV: an evolving understanding. *Curr Opin HIV AIDS*. 2014; 9:291–293. [PubMed: 24871091]
- Justice AC, McGinnis KA, Tate JP, et al. Risk of mortality and physiologic injury evident with lower alcohol exposure among HIV infected compared with uninfected men. *Drug and Alcohol Dependence*. 2016; 161:95–103. [PubMed: 26861883]
- Kabali C, Cheng DM, Brooks DR, et al. Recent cigarette smoking and HIV disease progression: no evidence of an association. *AIDS Care*. 2011; 23:947–956. [PubMed: 21400309]
- Kalichman SC, Cherry C, Kalichman MO, et al. Randomized clinical trial of HIV treatment adherence counseling interventions for people living with HIV and limited health literacy. *Journal of Acquired Immune Deficiency Syndromes*. 2013a; 63:42–50. [PubMed: 23337369]
- Kalichman SC, Grebler T, Amaral CM, et al. Intentional non-adherence to medications among HIV positive alcohol drinkers: prospective study of interactive toxicity beliefs. *Journal of General Internal Medicine*. 2013b; 28:399–405. [PubMed: 23065532]
- Kalichman SC, Simbayi LC, Vermaak R, et al. Randomized trial of a community-based alcohol-related HIV risk-reduction intervention for men and women in Cape Town South Africa. *Annals of Behavioral Medicine*. 2008; 36:270–279. [PubMed: 18836789]
- Kanis JA, Johansson H, Johnell O, et al. Alcohol intake as a risk factor for fracture. *Osteoporosis International*. 2004; 16:737–742. [PubMed: 15455194]

- Kaslow RA, Blackwelder WC, Ostrow DG, et al. No evidence for a role of alcohol or other psychoactive drugs in accelerating immunodeficiency in HIV-1-positive individuals. A report from the Multicenter AIDS Cohort Study. *The Journal of the American Medical Association*. 1989; 261:3424–3429. [PubMed: 2524608]
- Katz PS, Siggins RW, Porretta C, et al. Chronic alcohol increases CD8+ T-cell immunosenescence in simian immunodeficiency virus-infected rhesus macaques. *Alcohol*. 2015; 49:759–765. [PubMed: 26603633]
- Kelly UA. Integrating intersectionality and biomedicine in health disparities research. *ANS. Advances in Nursing Science*. 2009; 32:E42–E56.
- Kelso NE, Sheps DS, Cook RL. The association between alcohol use and cardiovascular disease among people living with HIV: a systematic review. *American Journal of Drug and Alcohol Abuse*. 2015; 41:479–488. [PubMed: 26286352]
- Kendall EA, Theron D, Franke MF, et al. Alcohol, hospital discharge, and socioeconomic risk factors for default from multidrug resistant tuberculosis treatment in rural South Africa: a retrospective cohort study. *PLoS ONE [Electronic Resource]*. 2013; 8:e83480.
- Kennedy DP, Hunter SB, Osilla KC, et al. A computer-assisted motivational social network intervention to reduce alcohol, drug and HIV risk behaviors among Housing First residents. *Addiction Science & Clinical Practice*. 2016; 11:4. [PubMed: 26979982]
- Kim TW, Palepu A, Cheng DM, et al. Factors associated with discontinuation of antiretroviral therapy in HIV-infected patients with alcohol problems. *AIDS Care*. 2007; 19:1039–1047. [PubMed: 17852002]
- Kinder LS, Bryson CL, Sun H, et al. Alcohol screening scores and all-cause mortality in male Veterans Affairs patients. *Journal of Studies on Alcohol and Drugs*. 2009; 70:253–260. [PubMed: 19261237]
- Korthuis PT, Fiellin DA, McGinnis KA, et al. Unhealthy alcohol and illicit drug use are associated with decreased quality of HIV care. *Journal of Acquired Immune Deficiency Syndromes*. 2012
- Korthuis PT, Josephs JS, Fleishman JA, et al. Substance abuse treatment in human immunodeficiency virus: the role of patient-provider discussions. *Journal of Substance Abuse Treatment*. 2008; 35:294–303. [PubMed: 18329222]
- Korthuis PT, Saha S, Chander G, et al. Substance use and the quality of patient-provider communication in HIV clinics. *AIDS Behav*. 2011; 15:832–841. [PubMed: 20703792]
- Kowalski S, Colantuoni E, Lau B, et al. Alcohol consumption and CD4 T-cell count response among persons initiating antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes*. 2012; 61:455–461. [PubMed: 22955054]
- Kraemer KL. Comparative effectiveness of alcohol and drug treatment in HIV-infected Veterans, R01AA022886-02 University of Pittsburgh: National Institutes on Alcohol Abuse and Alcoholism. 2013
- Kumar R, Perez-Casanova AE, Tirado G, et al. Increased viral replication in simian immunodeficiency virus/simian-HIV-infected macaques with self-administering model of chronic alcohol consumption. *Journal of Acquired Immune Deficiency Syndromes*. 2005; 39:386–390. [PubMed: 16010157]
- Kumar S, Rao PS, Earla R, et al. Drug-drug interactions between anti-retroviral therapies and drugs of abuse in HIV systems. *Expert Opin Drug Metab Toxicol*. 2015; 11:343–355. [PubMed: 25539046]
- Lakey W, Yang LY, Yancy W, et al. Short communication: from wasting to obesity: initial antiretroviral therapy and weight gain in HIV-infected persons. *AIDS Research and Human Retroviruses*. 2013; 29:435–440. [PubMed: 23072344]
- Lansky A, Brooks JT, DiNenno E, et al. Epidemiology of HIV in the United States. *Journal of Acquired Immune Deficiency Syndromes*. 2010; (55 Suppl 2):S64–S68. [PubMed: 21406989]
- Lefevre F, O’Leary B, Moran M, et al. Alcohol consumption among HIV-infected patients. *Journal of General Internal Medicine*. 1995; 10:458–460. [PubMed: 7472704]
- Leserman J. HIV disease progression: depression, stress, and possible mechanisms. *Biological Psychiatry*. 2003; 54:295–306. [PubMed: 12893105]

- Leserman J. Role of depression, stress, and trauma in HIV disease progression. *Psychosomatic Medicine*. 2008; 70:539–545. [PubMed: 18519880]
- Leserman J, Barroso J, Pence BW, et al. Trauma, stressful life events and depression predict HIV-related fatigue. *AIDS Care*. 2008; 20:1258–1265. [PubMed: 18608079]
- Liebschutz JM, Feinman G, Sullivan L, et al. Physical and sexual abuse in women infected with the human immunodeficiency virus: increased illness and health care utilization. *Archives of Internal Medicine*. 2000; 160:1659–1664. [PubMed: 10847259]
- Lifson AR, Lando HA. Smoking and HIV: prevalence, health risks, and cessation strategies. *Curr HIV/AIDS Rep*. 2012; 9:223–230. [PubMed: 22618079]
- Lifson AR, Neuhaus J, Arribas JR, et al. Smoking-related health risks among persons with HIV in the Strategies for Management of Antiretroviral Therapy clinical trial. *American Journal of Public Health*. 2010; 100:1896–1903. [PubMed: 20724677]
- Loeber S, Duka T, Welzel H, et al. Impairment of cognitive abilities and decision making after chronic use of alcohol: the impact of multiple detoxifications. *Alcohol and Alcoholism*. 2009; 44:372–381. [PubMed: 19487491]
- Lonnroth K, Williams BG, Stadlin S, et al. Alcohol use as a risk factor for tuberculosis - a systematic review. *BMC Public Health*. 2008; 8:289. [PubMed: 18702821]
- Machtiger EL, Haberer JE, Wilson TC, et al. Recent trauma is associated with antiretroviral failure and HIV transmission risk behavior among HIV-positive women and female-identified transgenders. *AIDS Behav*. 2012; 16:2160–2170. [PubMed: 22426597]
- Maisto SA, Palfai T, Vanable PA, et al. The effects of alcohol and sexual arousal on determinants of sexual risk in men who have sex with men. *Archives of Sexual Behavior*. 2012; 41:971–986. [PubMed: 22009480]
- Maisto SA, Simons JS. Research on the Effects of Alcohol and Sexual Arousal on Sexual Risk in Men who have Sex with Men: Implications for HIV Prevention Interventions. *AIDS Behav*. 2016; (20 Suppl 1):158–172.
- Marchetti G, Bellistri GM, Borghi E, et al. Microbial translocation is associated with sustained failure in CD4+ T-cell reconstitution in HIV-infected patients on long-term highly active antiretroviral therapy. *AIDS*. 2008; 22:2035–2038. [PubMed: 18784466]
- Marcondes MC, Watry D, Zandonatti M, et al. Chronic alcohol consumption generates a vulnerable immune environment during early SIV infection in rhesus macaques. *Alcoholism, Clinical and Experimental Research*. 2008; 32:1583–1592.
- Marshall MM, McCormack MC, Kirk GD. Effect of cigarette smoking on HIV acquisition, progression, and mortality. *AIDS Education and Prevention*. 2009; 21:28–39. [PubMed: 19537952]
- Mathers BM, Degenhardt L, Phillips B, et al. Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review. *Lancet*. 2008; 372:1733–1745. [PubMed: 18817968]
- McCance-Katz EF, Gruber VA, Beatty G, et al. Interactions between alcohol and the antiretroviral medications ritonavir or efavirenz. *J Addict Med*. 2013; 7:264–270. [PubMed: 23666322]
- McCance-Katz EF, Lum PJ, Beatty G, et al. Untreated HIV infection is associated with higher blood alcohol levels. *Journal of Acquired Immune Deficiency Syndromes*. 2012; 60:282–288. [PubMed: 22495786]
- McCaul ME. ALCOHOL RESEARCH CONSORTIUM IN HIV, 5U24AA020801-05 Johns Hopkins University: National Institutes of Alcohol Abuse and Alcoholism. 2016
- McGinnis KA, Fiellin DA, Tate JP, et al. Number of Drinks to “Feel a Buzz” by HIV Status and Viral Load in Men. *AIDS Behav*. 2016; 20:504–511. [PubMed: 26936030]
- McGinnis KA, Fultz SL, Skanderson M, et al. Hepatocellular carcinoma and non-Hodgkin’s lymphoma: the roles of HIV, hepatitis C infection, and alcohol abuse. *Journal of Clinical Oncology*. 2006; 24:5005–5009. [PubMed: 17075119]
- Mdodo R, Frazier EL, Dube SR, et al. Cigarette smoking prevalence among adults with HIV compared with the general adult population in the United States: cross-sectional surveys. *Annals of Internal Medicine*. 2015; 162:335–344. [PubMed: 25732274]

- Merlin JS. Chronic Pain in Patients With HIV Infection: What Clinicians Need To Know. *Top Antivir Med.* 2015; 23:120–124. [PubMed: 26518396]
- Merlin JS, Cen L, Praestgaard A, et al. Pain and physical and psychological symptoms in ambulatory HIV patients in the current treatment era. *Journal of Pain and Symptom Management.* 2012a; 43:638–645. [PubMed: 22115794]
- Merlin JS, Westfall AO, Raper JL, et al. Pain, mood, and substance abuse in HIV: implications for clinic visit utilization, antiretroviral therapy adherence, and virologic failure. *Journal of Acquired Immune Deficiency Syndromes.* 2012b; 61:164–170. [PubMed: 22766967]
- Metsch LR, Pereyra M, Colfax G, et al. HIV-positive patients' discussion of alcohol use with their HIV primary care providers. *Drug and Alcohol Dependence.* 2008; 95:37–44. [PubMed: 18243580]
- Meyer VJ, Rubin LH, Martin E, et al. HIV and recent illicit drug use interact to affect verbal memory in women. *Journal of Acquired Immune Deficiency Syndromes.* 2013; 63:67–76. [PubMed: 23392462]
- Molina PE, Amedee AM, Veazey R, et al. Chronic Binge Alcohol Consumption Does Not Diminish Effectiveness of Continuous Antiretroviral Suppression of Viral Load in Simian Immunodeficiency Virus-Infected Macaques. *Alcoholism: Clinical and Experimental Research.* 2014a; 38:2335–2344.
- Molina PE, Bagby GJ, Nelson S. Biomedical consequences of alcohol use disorders in the HIV-infected host. *Curr HIV Res.* 2014b; 12:265–275. [PubMed: 25053365]
- Molina PE, McNurlan M, Rathmacher J, et al. Chronic alcohol accentuates nutritional, metabolic, and immune alterations during asymptomatic simian immunodeficiency virus infection. *Alcoholism, Clinical and Experimental Research.* 2006; 30:2065–2078.
- Monnig MA, Kahler CW, Cioe PA, et al. Alcohol use predicts elevation in inflammatory marker soluble CD14 in men living with HIV. *AIDS Care.* 2016:1–7.
- Monroe AK, Lau B, Mugavero MJ, et al. Heavy Alcohol Use is Associated with Worse Retention in HIV Care. *Journal of Acquired Immune Deficiency Syndromes.* 2016
- Monti PM, Mastroleo NR, Barnett NP, et al. Brief Motivational Intervention to Reduce Alcohol and HIV/Sexual Risk Behavior in Emergency Department Patients: A Randomized Controlled Trial. *Journal of Consulting and Clinical Psychology.* 2016
- Morgan TR, Brenner D, Everhart J, et al. Hepatitis C and alcohol: fundamental and translational research directions. *Alcoholism, Clinical and Experimental Research.* 2003; 27:726–731.
- Mozumdar A, Liguori G. Persistent increase of prevalence of metabolic syndrome among U.S. adults: NHANES III to NHANES 1999–2006. *Diabetes Care.* 2011; 34:216–219. [PubMed: 20889854]
- Mugavero M, Ostermann J, Whetten K, et al. Barriers to antiretroviral adherence: the importance of depression, abuse, and other traumatic events. *Aids Patient Care and STDS.* 2006; 20:418–428. [PubMed: 16789855]
- Mugavero MJ, Amico KR, Horn T, et al. The state of engagement in HIV care in the United States: from cascade to continuum to control. *Clinical Infectious Diseases.* 2013; 57:1164–1171. [PubMed: 23797289]
- Mukamal KJ, Robbins JA, Cauley JA, et al. Alcohol consumption, bone density, and hip fracture among older adults: the cardiovascular health study. *Osteoporosis International.* 2007; 18:593–602. [PubMed: 17318666]
- Mulia N, Ye Y, Greenfield TK, et al. Disparities in alcohol-related problems among white, black, and Hispanic Americans. *Alcoholism, Clinical and Experimental Research.* 2009; 33:654–662.
- National Institutes of Health. NIH HIV/AIDS Research Priorities and Guidelines for Determining AIDS Funding. 2015. <https://grants.nih.gov/grants/guide/notice-files/NOT-OD-15-137.html>
- Neblett RC, Hutton HE, Lau B, et al. Alcohol consumption among HIV-infected women: impact on time to antiretroviral therapy and survival. *J Womens Health (Larchmt).* 2011; 20:279–286. [PubMed: 21281111]
- Neuman MG, Schneider M, Nanau RM, et al. Alcohol Consumption, Progression of Disease and Other Comorbidities, and Responses to Antiretroviral Medication in People Living with HIV. *AIDS Res Treat.* 2012; 2012:751827. [PubMed: 22496971]

- Niaura R, Shadel WG, Morrow K, et al. Human immunodeficiency virus infection, AIDS, and smoking cessation: the time is now. *Clinical Infectious Diseases*. 2000; 31:808–812. [PubMed: 11017836]
- Papas RK, Sidle JE, Gakinya BN, et al. Treatment outcomes of a stage 1 cognitive-behavioral trial to reduce alcohol use among human immunodeficiency virus-infected out-patients in western Kenya. *Addiction*. 2011; 106:2156–2166. [PubMed: 21631622]
- Papas RK, Sidle JE, Martino S, et al. Systematic cultural adaptation of cognitive-behavioral therapy to reduce alcohol use among HIV-infected outpatients in western Kenya. *AIDS Behav*. 2010; 14:669–678. [PubMed: 19967441]
- Park LS, Hernandez-Ramirez RU, Silverberg MJ, et al. Prevalence of non-HIV cancer risk factors in persons living with HIV/AIDS: a meta-analysis. *AIDS*. 2016; 30:273–291. [PubMed: 26691548]
- Park WB, Choe PG, Kim SH, et al. One-year adherence to clinic visits after highly active antiretroviral therapy: a predictor of clinical progress in HIV patients. *Journal of Internal Medicine*. 2007; 261:268–275. [PubMed: 17305649]
- Parry CD, Morojele NK, Myers BJ, et al. Efficacy of an alcohol-focused intervention for improving adherence to antiretroviral therapy (ART) and HIV treatment outcomes - a randomised controlled trial protocol. *BMC Infect Dis*. 2014; 14:500. [PubMed: 25212696]
- Parsons JT, Golub SA, Rosof E, et al. Motivational interviewing and cognitive-behavioral intervention to improve HIV medication adherence among hazardous drinkers: a randomized controlled trial. *Journal of Acquired Immune Deficiency Syndromes*. 2007a; 46:443–450. [PubMed: 18077833]
- Parsons JT, Rosof E, Mustanski B. Patient-related factors predicting HIV medication adherence among men and women with alcohol problems. *Journal of health psychology*. 2007b; 12:357–370. [PubMed: 17284499]
- Parsons JT, Rosof E, Mustanski B. The temporal relationship between alcohol consumption and HIV-medication adherence: a multilevel model of direct and moderating effects. *Health Psychology*. 2008; 27:628–637. [PubMed: 18823189]
- Paula AA, Falcão MCN, Pacheco AG. Metabolic syndrome in HIV-infected individuals: underlying mechanisms and epidemiological aspects. *AIDS Research and Therapy*. 2013; 10:1. [PubMed: 23286882]
- Pellowski JA, Kalichman SC, Kalichman MO, et al. Alcohol-antiretroviral therapy interactive toxicity beliefs and daily medication adherence and alcohol use among people living with HIV. *AIDS Care*. 2016:1–8.
- Pellowski JA, Kalichman SC, Matthews KA, et al. A pandemic of the poor: social disadvantage and the U.S. HIV epidemic. *American Psychologist*. 2013; 68:197–209. [PubMed: 23688088]
- Pence BW, Thielman NM, Whetten K, et al. Coping Strategies and Patterns of Alcohol and Drug Use among HIV-Infected Patients in the United States Southeast. *Aids Patient Care and STDS*. 2008; 22:869–877. [PubMed: 19025481]
- Penkower L, Dew MA, Kingsley L, et al. Alcohol consumption as a cofactor in the progression of HIV infection and AIDS. *Alcohol*. 1995; 12:547–552. [PubMed: 8590617]
- Peretti-Watel P, Spire B, Lert F, et al. Drug use patterns and adherence to treatment among HIV-positive patients: evidence from a large sample of French outpatients (ANRS-EN12-VESPA 2003). *Drug and Alcohol Dependence*. 2006; (82 Suppl 1):S71–S79. [PubMed: 16769450]
- Persidsky Y, Ho W, Ramirez SH, et al. HIV-1 infection and alcohol abuse: neurocognitive impairment, mechanisms of neurodegeneration and therapeutic interventions. *Brain, Behavior, and Immunity*. 2011; (25 Suppl 1):S61–S70.
- Peters MG, Terrault NA. Alcohol use and hepatitis C. *Hepatology*. 2002; 36:S220–S225. [PubMed: 12407597]
- Platt L, Easterbrook P, Gower E, et al. Prevalence and burden of HCV co-infection in people living with HIV: a global systematic review and meta-analysis. *Lancet Infect Dis*. 2016
- Poonia B, Nelson S, Bagby GJ, et al. Chronic alcohol consumption results in higher simian immunodeficiency virus replication in mucosally inoculated rhesus macaques. *AIDS Research and Human Retroviruses*. 2006; 22:589–594. [PubMed: 16796534]

- Rasmussen LD, Mathiesen ER, Kronborg G, et al. Risk of diabetes mellitus in persons with and without HIV: a Danish nationwide population-based cohort study. *PLoS One*. 2012; 7:e44575. [PubMed: 22984529]
- Rehm J, Gmel G, Sempos CT, et al. Alcohol-related morbidity and mortality. *Alcohol Res Health*. 2003; 27:39–51. [PubMed: 15301399]
- Rehm J, Gutjahr E, Gmel G. Alcohol and all-cause mortality: a pooled analysis. *Contemporary Drug Problems*. 2001; 28:337–361.
- Rehm J, Mathers C, Popova S, et al. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet*. 2009a; 373:2223–2233. [PubMed: 19560604]
- Rehm J, Samokhvalov AV, Neuman MG, et al. The association between alcohol use, alcohol use disorders and tuberculosis (TB). A systematic review. *BMC Public Health*. 2009b; 9:450. [PubMed: 19961618]
- Rehm J, Taylor B, Patra J. Volume of alcohol consumption, patterns of drinking and burden of disease in the European region 2002. *Addiction*. 2006; 101:1086–1095. [PubMed: 16869838]
- Riley ED, Evans JL, Hahn JA, et al. A Longitudinal Study of Multiple Drug Use and Overdose Among Young People Who Inject Drugs. *American Journal of Public Health*. 2016; 106:915–917. [PubMed: 26985620]
- Rothlind JC, Greenfield TM, Bruce AV, et al. Heavy alcohol consumption in individuals with HIV infection: effects on neuropsychological performance. *Journal of the International Neuropsychological Society*. 2005; 11:70–83. [PubMed: 15686610]
- Sacktor N, Skolasky RL, Seaberg E, et al. Prevalence of HIV-associated neurocognitive disorders in the Multicenter AIDS Cohort Study. *Neurology*. 2016; 86:334–340. [PubMed: 26718568]
- Safdar K, Schiff ER. Alcohol and hepatitis C. *Seminars in Liver Disease*. 2004; 24:305–315. [PubMed: 15349807]
- Saitz R. Clinical practice. Unhealthy alcohol use. *New England Journal of Medicine*. 2005; 352:596–607. [PubMed: 15703424]
- Samaras K. The burden of diabetes and hyperlipidemia in treated HIV infection and approaches for cardiometabolic care. *Curr HIV/AIDS Rep*. 2012; 9:206–217. [PubMed: 22752405]
- Samaras K, Wand H, Law M, et al. Prevalence of metabolic syndrome in HIV-infected patients receiving highly active antiretroviral therapy using International Diabetes Foundation and Adult Treatment Panel III criteria: associations with insulin resistance, disturbed body fat compartmentalization, elevated C-reactive protein, and [corrected] hypoadiponectinemia. *Diabetes Care*. 2007; 30:113–119. [PubMed: 17192343]
- Samet JH, Cheng DM, Libman H, et al. Alcohol consumption and HIV disease progression. *Journal of Acquired Immune Deficiency Syndromes*. 2007a; 46:194–199. [PubMed: 17667330]
- Samet JH, Freedberg KA, Savetsky JB, et al. Discontinuation from HIV medical care: squandering treatment opportunities. *Journal of Health Care for the Poor and Underserved*. 2003; 14:244–255. [PubMed: 12739303]
- Samet JH, Freedberg KA, Savetsky JB, et al. Understanding delay to medical care for HIV infection: the long-term non-presenter. *AIDS*. 2001; 15:77–85. [PubMed: 11192871]
- Samet JH, Freedberg KA, Stein MD, et al. Trillion virion delay: time from testing positive for HIV to presentation for primary care. *Archives of Internal Medicine*. 1998; 158:734–740. [PubMed: 9554679]
- Samet JH, Horton NJ, Meli S, et al. A randomized controlled trial to enhance antiretroviral therapy adherence in patients with a history of alcohol problems. *Antivir Ther*. 2005; 10:83–93. [PubMed: 15751766]
- Samet JH, Horton NJ, Meli S, et al. Alcohol consumption and antiretroviral adherence among HIV-infected persons with alcohol problems. *Alcoholism, Clinical and Experimental Research*. 2004a; 28:572–577.
- Samet JH, Krupitsky EM, Cheng DM, et al. Mitigating risky sexual behaviors among Russian narcology hospital patients: the PREVENT (Partnership to Reduce the Epidemic Via Engagement in Narcology Treatment) randomized controlled trial. *Addiction*. 2008; 103:1474–1483. [PubMed: 18636998]

- Samet JH, Phillips SJ, Horton NJ, et al. Detecting alcohol problems in HIV-infected patients: use of the CAGE questionnaire. *AIDS Research and Human Retroviruses*. 2004b; 20:151–155. [PubMed: 15018702]
- Samet JH, Raj A, Cheng DM, et al. HERMITAGE--a randomized controlled trial to reduce sexually transmitted infections and HIV risk behaviors among HIV-infected Russian drinkers. *Addiction*. 2015; 110:80–90. [PubMed: 25170994]
- Samet JH, Walley AY. Interventions targeting HIV-infected risky drinkers: drops in the bottle. *Alcohol Res Health*. 2010; 33:267–279. [PubMed: 23584068]
- Samet JH, Walley AY, Bridden C. Illicit drugs, alcohol, and addiction in human immunodeficiency virus. *Panminerva Medica*. 2007b; 49:67–77. [PubMed: 17625483]
- Sandler, N.; Wand, H.; Nixon, D., et al. Plasma Levels of Soluble CD14 Predict Mortality in HIV Infection; 17th Conference on Retroviruses and Opportunistic Infections; San Francisco. 2010.
- Sarkar DK, Jung MK, Wang HJ. Alcohol and the Immune System. *Alcohol Research: Current Reviews*. 2015:37.
- Satre DD. Primary care-based interventions to reduce alcohol use among HIV patients; U01AA021997-04 University of California, San Francisco: National Institutes of Alcohol Abuse and Alcoholism. 2012
- Schiff ER, Ozden N. Hepatitis C and alcohol. *Alcohol Res Health*. 2003; 27:232–239. [PubMed: 15535451]
- Scott-Sheldon LA, Carey KB, Cunningham K, et al. Alcohol Use Predicts Sexual Decision-Making: A Systematic Review and Meta-Analysis of the Experimental Literature. *AIDS Behav*. 2016; (20 Suppl 1):19–39.
- Shah NG, Lathrop SL, Reichard RR, et al. Unintentional drug overdose death trends in New Mexico, USA, 1990–2005: combinations of heroin, cocaine, prescription opioids and alcohol. *Addiction*. 2008; 103:126–136. [PubMed: 18028518]
- Sharma A, Shi Q, Hoover DR, et al. Increased Fracture Incidence in Middle-Aged HIV-Infected and HIV-Uninfected Women: Updated Results From the Women's Interagency HIV Study. *Journal of Acquired Immune Deficiency Syndromes*. 2015; 70:54–61. [PubMed: 26322667]
- Shiau S, Broun EC, Arpadi SM, et al. Incident fractures in HIV-infected individuals: a systematic review and meta-analysis. *AIDS (London, England)*. 2013; 27:1949–1957.
- Shiels MS, Pfeiffer RM, Gail MH, et al. Cancer burden in the HIV-infected population in the United States. *Journal of the National Cancer Institute*. 2011; 103:753–762. [PubMed: 21483021]
- Shuper PA, Joharchi N, Irving H, et al. Alcohol as a correlate of unprotected sexual behavior among people living with HIV/AIDS: review and meta-analysis. *AIDS Behav*. 2009; 13:1021–1036. [PubMed: 19618261]
- Shuper PA, Neuman M, Kanteres F, et al. Causal considerations on alcohol and HIV/AIDS--a systematic review. *Alcohol and Alcoholism*. 2010; 45:159–166. [PubMed: 20061510]
- Sidorenkov O, Nilssen O, Grjibovski AM. Metabolic syndrome in Russian adults: associated factors and mortality from cardiovascular diseases and all causes. *BMC Public Health*. 2010; 10:582–2458–2410–2582. [PubMed: 20920226]
- Sigel K, Wisnivesky J, Gordon K, et al. HIV as an independent risk factor for incident lung cancer. *AIDS*. 2012; 26:1017–1025. [PubMed: 22382152]
- Simmons RD, Ciancio BC, Kall MM, et al. Ten-year mortality trends among persons diagnosed with HIV infection in England and Wales in the era of antiretroviral therapy: AIDS remains a silent killer. *HIV Med*. 2013; 14:596–604. [PubMed: 23672663]
- Skarbinski J, Rosenberg E, Paz-Bailey G, et al. Human immunodeficiency virus transmission at each step of the care continuum in the United States. *JAMA internal medicine*. 2015; 175:588–596. [PubMed: 25706928]
- Smith CJ, Ryom L, Weber R, et al. Trends in underlying causes of death in people with HIV from 1999 to 2011 (D:A:D): a multicohort collaboration. *Lancet*. 2014; 384:241–248. [PubMed: 25042234]
- Smith P, Schmidt SM, Allensworth-Davies D, et al. Primary Care Validation of a Single-Question Alcohol Screening Test. *Journal of General Internal Medicine*. 2009; 24:783–788. [PubMed: 19247718]

- Sobell LC, Brown J, Leo GI, et al. The reliability of the Alcohol Timeline Followback when administered by telephone and by computer. *Drug and Alcohol Dependence*. 1996; 42:49–54. [PubMed: 8889403]
- Stockwell T, Zhao J, Panwar S, et al. Do “Moderate” Drinkers Have Reduced Mortality Risk? A Systematic Review and Meta-Analysis of Alcohol Consumption and All-Cause Mortality. *Journal of Studies on Alcohol and Drugs*. 2016; 77:185–198. [PubMed: 26997174]
- Stokols D. Establishing and maintaining healthy environments. Toward a social ecology of health promotion. *American Psychologist*. 1992; 47:6–22. [PubMed: 1539925]
- Strauss SM, Munoz-Plaza C, Tiburcio NJ, et al. HIV Care Providers’ Role Legitimacy as Supporters of Their Patients’ Alcohol Reduction. *Open Infect Dis J*. 2009a; 3:13–20. [PubMed: 20556238]
- Strauss SM, Munoz-Plaza CE, Tiburcio NJ, et al. Barriers and facilitators in implementing “prevention for positives” alcohol-reduction support: the perspectives of directors and providers in hospital-based HIV care centers. *Journal of the Association of Nurses in AIDS Care*. 2012; 23:30–40. [PubMed: 21570321]
- Strauss SM, Tiburcio NJ, Munoz-Plaza C, et al. HIV care providers’ implementation of routine alcohol reduction support for their patients. *Aids Patient Care and STDS*. 2009b; 23:211–218. [PubMed: 19866539]
- Substance Abuse and Mental Health Services Administration. Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings. 2014. <http://www.samhsa.gov/data/sites/default/files/NSDUHresultsPDFWHTML2013/Web/NSDUHresults2013.pdf>
- Sulkowski MS. Hepatitis C virus infection in HIV-infected patients. *Curr Infect Dis Rep*. 2007a; 3:469–476. [PubMed: 24395487]
- Sulkowski MS. The HIV-coinfected patient: managing viral hepatitis. *Journal of Acquired Immune Deficiency Syndromes*. 2007b; (45 Suppl 2):S36–S37. [PubMed: 17704690]
- Sulkowski MS. Viral hepatitis and HIV coinfection. *Journal of Hepatology*. 2008; 48:353–367. [PubMed: 18155314]
- Sulkowski MS. Current management of hepatitis C virus infection in patients with HIV co-infection. *Journal of Infectious Diseases*. 2013; (207 Suppl 1):S26–S32. [PubMed: 23390302]
- Sullivan LE, Goulet JL, Justice AC, et al. Alcohol consumption and depressive symptoms over time: a longitudinal study of patients with and without HIV infection. *Drug and Alcohol Dependence*. 2011; 117:158–163. [PubMed: 21345624]
- Sullivan LE, Saitz R, Cheng DM, et al. The impact of alcohol use on depressive symptoms in human immunodeficiency virus-infected patients. *Addiction*. 2008; 103:1461–1467. [PubMed: 18637000]
- Szabo G, Dolganiuc A, Mandrekar P, et al. Inhibition of antigen-presenting cell functions by alcohol: implications for hepatitis C virus infection. *Alcohol*. 2004; 33:241–249. [PubMed: 15596093]
- Talman A, Bolton S, Watson JL. Interactions between HIV/AIDS and the environment: toward a syndemic framework. *American Journal of Public Health*. 2013; 103:253–261. [PubMed: 23237167]
- Tang AM, Graham NM, Chandra RK, et al. Low serum vitamin B-12 concentrations are associated with faster human immunodeficiency virus type 1 (HIV-1) disease progression. *Journal of Nutrition*. 1997; 127:345–351. [PubMed: 9039838]
- Thaler NS, Sayegh P, Kim MS, et al. Interactive effects of neurocognitive impairment and substance use on antiretroviral non-adherence in HIV disease. *Arch Clin Neuropsychol*. 2015; 30:114–121. [PubMed: 25589442]
- Thurman RG. II. Alcoholic liver injury involves activation of Kupffer cells by endotoxin. *American Journal of Physiology*. 1998; 275:G605–G611. [PubMed: 9756487]
- Tiozzo E, Konefal J, Adwan S, et al. A cross-sectional assessment of metabolic syndrome in HIV-infected people of low socio-economic status receiving antiretroviral therapy. *Diabetology & metabolic syndrome*. 2015; 7:1. [PubMed: 25810781]
- Tsui JI, Cheng DM, Coleman SM, et al. Pain is associated with risky drinking over time among HIV-infected persons in St. Petersburg, Russia. *Drug and Alcohol Dependence*. 2014; 144:87–92. [PubMed: 25220898]

- Tsui JI, Cheng DM, Libman H, et al. Risky alcohol use and serum aminotransferase levels in HIV-infected adults with and without hepatitis C. *Journal of Studies on Alcohol and Drugs*. 2013; 74:266–270. [PubMed: 23384374]
- Tsui JI, Pletcher MJ, Vittinghoff E, et al. Hepatitis C and hospital outcomes in patients admitted with alcohol-related problems. *Journal of Hepatology*. 2006; 44:262–266. [PubMed: 16226823]
- UNAIDS. *How AIDS Changed Everything: 2014 Global Statistics*. 2015
- Vagenas P, Azar MM, Copenhaver MM, et al. The Impact of Alcohol Use and Related Disorders on the HIV Continuum of Care: a Systematic Review : Alcohol and the HIV Continuum of Care. *Curr HIV/AIDS Rep*. 2015; 12:421–436. [PubMed: 26412084]
- Vagenas P, Ludford KT, Gonzales P, et al. Being unaware of being HIV-infected is associated with alcohol use disorders and high-risk sexual behaviors among men who have sex with men in Peru. *AIDS Behav*. 2014; 18:120–127. [PubMed: 23670711]
- Van Tieu H, Koblin BA. HIV, alcohol, and noninjection drug use. *Curr Opin HIV AIDS*. 2009; 4:314–318. [PubMed: 19532070]
- Veugelaers PJ, Page KA, Tindall B, et al. Determinants of HIV disease progression among homosexual men registered in the Tricontinental Seroconverter Study. *American Journal of Epidemiology*. 1994; 140:747–758. [PubMed: 7942776]
- Viswanathan H, Wilson JA. Alcohol--the neglected risk factor in head and neck cancer. *Clin Otolaryngol Allied Sci*. 2004; 29:295–300. [PubMed: 15270811]
- Volkman T, Moonan PK, Miramontes R, et al. Excess Alcohol Use and Death among Tuberculosis Patients in the United States, 1997–2012. *J Tuberc Res*. 2016; 4:18–22. [PubMed: 27054144]
- Wakabayashi I. Frequency of Heavy Alcohol Drinking and Risk of Metabolic Syndrome in Middle-Aged Men. *Alcoholism: Clinical and Experimental Research*. 2014; 38:1689–1696.
- Walkup J, Blank MB, Gonzalez JS, et al. The impact of mental health and substance abuse factors on HIV prevention and treatment. *Journal of Acquired Immune Deficiency Syndromes*. 2008; (47 Suppl 1):S15–S19.
- Walley AY, Cheng DM, Libman H, et al. Recent drug use, homelessness and increased short-term mortality in HIV-infected persons with alcohol problems. *AIDS*. 2008; 22:415–420. [PubMed: 18195568]
- Wandeler G, Kraus D, Fehr J, et al. The J-Curve in HIV: Low and Moderate Alcohol Intake Predicts Mortality but Not the Occurrence of Major Cardiovascular Events. *Journal of Acquired Immune Deficiency Syndromes*. 2016; 71:302–309. [PubMed: 26444500]
- Webber MP, Schoenbaum EE, Gourevitch MN, et al. A prospective study of HIV disease progression in female and male drug users. *AIDS*. 1999; 13:257–262. [PubMed: 10202832]
- Weber E, Morgan EE, Iudicello JE, et al. Substance use is a risk factor for neurocognitive deficits and neuropsychiatric distress in acute and early HIV infection. *Journal of Neurovirology*. 2013; 19:65–74. [PubMed: 23250704]
- Williams EC, Bradley KA. *Alcohol-related Care and Outcomes for Outpatients with HIV in a National VA Cohort, R21 AA022866* Group Health Research Institute: National Institute on Alcohol Abuse and Alcoholism. 2015
- World Health Organization. *Guidelines for intensified tuberculosis case-finding and isoniazid preventive therapy for people living with HIV in resource-constrained settings*. 2011.
- World Health Organization. *Global Tuberculosis Report*. Geneva: 2014.
- World Health Organization. *Alcohol*. 2015. <http://www.who.int/mediacentre/factsheets/fs349/en/>
- Wurst FM, Alling C, Aradottir S, et al. Emerging biomarkers: new directions and clinical applications. *Alcoholism, Clinical and Experimental Research*. 2005; 29:465–473.
- Zhang X, Yu Z, Yu M, et al. Alcohol consumption and hip fracture risk. *Osteoporosis International*. 2014; 26:531–542. [PubMed: 25266483]




Figure 1.
The Influence of Alcohol Use on HIV Care and Outcomes