



Cannabidiol Regulation of Learned Fear: Implications for Treating Anxiety-Related Disorders

Regimantas Jurkus¹, Harriet L. L. Day², Francisco S. Guimarães³, Jonathan L. C. Lee⁴, Leandro J. Bertoglio⁵ and Carl W. Stevenson^{2*}

¹ School of Mathematical Sciences, University of Nottingham, University Park, Nottingham, UK, ² School of Biosciences, University of Nottingham, Loughborough, UK, ³ Department of Pharmacology, University of São Paulo, São Paulo, Brazil, ⁴ School of Psychology, University of Birmingham, Birmingham, UK, ⁵ Department of Pharmacology, Federal University of Santa Catarina, Florianópolis, Brazil

OPEN ACCESS

Edited by:

Giuseppe Esposito,
Sapienza University of Rome, Italy

Reviewed by:

Benedict Green,
Poisonous Plant Research Laboratory,
Agricultural Research Service (USDA),
USA

Rajendra Karki,
St. Jude Children's Research Hospital,
USA

*Correspondence:

Carl W. Stevenson
carl.stevenson@nottingham.ac.uk

Specialty section:

This article was submitted to
Ethnopharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 01 October 2016

Accepted: 11 November 2016

Published: 24 November 2016

Citation:

Jurkus R, Day HLL, Guimarães FS,
Lee JLC, Bertoglio LJ and
Stevenson CW (2016) Cannabidiol
Regulation of Learned Fear:
Implications for Treating
Anxiety-Related Disorders.
Front. Pharmacol. 7:454.
doi: 10.3389/fphar.2016.00454

Anxiety and trauma-related disorders are psychiatric diseases with a lifetime prevalence of up to 25%. Phobias and post-traumatic stress disorder (PTSD) are characterized by abnormal and persistent memories of fear-related contexts and cues. The effects of psychological treatments such as exposure therapy are often only temporary and medications can be ineffective and have adverse side effects. Growing evidence from human and animal studies indicates that cannabidiol, the main non-psychotomimetic phytocannabinoid present in *Cannabis sativa*, alleviates anxiety in paradigms assessing innate fear. More recently, the effects of cannabidiol on learned fear have been investigated in preclinical studies with translational relevance for phobias and PTSD. Here we review the findings from these studies, with an emphasis on cannabidiol regulation of contextual fear. The evidence indicates that cannabidiol reduces learned fear in different ways: (1) cannabidiol decreases fear expression acutely, (2) cannabidiol disrupts memory reconsolidation, leading to sustained fear attenuation upon memory retrieval, and (3) cannabidiol enhances extinction, the psychological process by which exposure therapy inhibits learned fear. We also present novel data on cannabidiol regulation of learned fear related to explicit cues, which indicates that auditory fear expression is also reduced acutely by cannabidiol. We conclude by outlining future directions for research to elucidate the neural circuit, psychological, cellular, and molecular mechanisms underlying the regulation of fear memory processing by cannabidiol. This line of investigation may lead to the development of cannabidiol as a novel therapeutic approach for treating anxiety and trauma-related disorders such as phobias and PTSD in the future.

Keywords: cannabidiol, extinction, fear conditioning, reconsolidation

INTRODUCTION

Anxiety and trauma-related disorders will affect up to one in four people in their lifetime. Diseases like phobias and post-traumatic stress disorder (PTSD) show abnormal persistence of fear memories and can be debilitating, causing a huge societal and economic burden. Psychological treatments like exposure therapy are used to treat these disorders but they often show only limited or temporary effectiveness. Medications are also available but these are not fully effective in a

significant proportion of patients, can have adverse side effects, and may even interfere with the efficacy of psychological therapy. There is therefore an urgent need for better treatments for these disorders (Fineberg et al., 2013; Baldwin et al., 2014).

A promising area of research in this field focuses on repurposing existing and developing novel drugs to enhance the effectiveness of psychological therapies in alleviating fear-related symptoms (Myers and Davis, 2007; Steckler and Risbrough, 2012). One drug showing broad therapeutic potential in various psychiatric diseases is cannabidiol (CBD), the main non-psychoactive constituent of the *Cannabis sativa* plant (Izzo et al., 2009; see chemical structure of CBD in **Figure 1A**). Studies in humans demonstrate the promise of CBD for treating anxiety (Blessing et al., 2015) and preclinical studies in rodents are elucidating the pharmacological mechanisms underlying its acute anxiolytic effects. These mechanisms include potentiation of serotonin (5-HT) transmission via 5-HT_{1A} receptor (5-HT_{1A}R) activation and elevation of endocannabinoid levels via inhibition of their metabolism and re-uptake, which indirectly facilitates cannabinoid receptor type 1 (CB1R) activation (for a review see Campos et al., 2016).

As well as reducing anxiety in behavioral tests of unconditioned fear, emerging evidence indicates that CBD regulates fear learning and memory in paradigms that are translationally relevant to diseases such as phobias and PTSD, along with their psychological treatment. In this paper we review the recent studies on CBD regulation of fear memory processing, which have focused on contextual fear. We also present novel data on CBD regulation of auditory fear memory and its extinction, which forms the theoretical basis for exposure therapy. We then outline future directions for research on this topic to gain a broader perspective on the neural circuit, psychological, pharmacological, and cellular bases of the regulation of learned fear by CBD.


CBD REGULATION OF CONTEXTUAL FEAR MEMORY PROCESSING

Recent evidence indicates that CBD modulates fundamental neurobiological processes involved in Pavlovian fear conditioning, a form of associative learning by which certain stimuli or environments become predictive of threat and therefore enhance survival. During acquisition a neutral conditioned stimulus (CS) is associated with an aversive unconditioned stimulus (US), such as a mild footshock. The CS can be either discrete (i.e., cued), such as a light or tone, or the environment (i.e., context) where the US was presented. CS re-exposure after conditioning initially induces a fear response, which has frequently been inferred from behavioral (e.g.,

Abbreviations: AC, adenylyl cyclase; 5-HT, serotonin; 5-HT_{1A}R, 5-HT_{1A} receptor; Δ^9 -THC, Δ^9 -tetrahydrocannabinol; BDNF, brain derived neurotrophic factor; BNST, bed nucleus of the stria terminalis; CB1R, cannabinoid receptor type 1; CBD, cannabidiol; ERK, extracellular signal-regulated kinase; i.c.v., intracerebroventricular; IL, infralimbic; i.p., intraperitoneal; NFkB, nuclear factor kappa-light-chain-enhancer of activated B cells; PKC, protein kinase C; PL, prelimbic; PTSD, post-traumatic stress disorder; SHR, spontaneously hypertensive rat; TrkB, tyrosine receptor kinase B.

freezing) and/or autonomic (increased heart rate/blood pressure, decreased body temperature) changes (Fendt and Fanselow, 1999; Resstel et al., 2009). After its acquisition the CS-US association is consolidated into long-term fear memory. Later retrieval can render fear memory labile through destabilization of the memory trace, allowing for maintenance or updating of the memory through its reconsolidation (Lee, 2009). Extinction of fear memory occurs with longer durations or repeated sessions of retrieval. This form of inhibitory learning results in the encoding of a new CS-no US association which suppresses fear expression by competing with the original fear memory (Myers and Davis, 2007). **Figure 1B** depicts the different phases of fear memory and its possible reconsolidation or extinction after retrieval.

Accumulating evidence indicates that CBD regulates different contextual fear memory processes. An initial study by Resstel et al. (2006) showed that systemic CBD administration decreases



the freezing response and autonomic changes induced by exposure to an aversively conditioned context; this effect was similar to the positive control diazepam. Subsequent studies confirmed the CBD-induced reduction in conditioned freezing expression with acute administration before retrieval (Lemos et al., 2010) or acquisition (Levin et al., 2012). In contrast, ElBatsh et al. (2012) showed that repeated daily injections (14 days) of CBD increased freezing expression during contextual fear retrieval. Chronic treatment with CBD has, however, been shown to facilitate adult hippocampal neurogenesis (Wolf et al., 2010; Campos et al., 2013), which is involved in aversive learning and memory processing as its facilitation enhances contextual discrimination and related fear expression (Efstathopoulos et al., 2015). Mice with reduced neurogenesis, on the other hand, presented less contextual fear (Pan et al., 2012; Denny et al., 2014). Both associative (through facilitation of associative learning) and non-associative (by buffering non-associative, anxiogenic effects of the aversive experience) mechanisms seem to play a role in regulating fear learning by adult hippocampal neurogenesis (Seo et al., 2015). Since the animals were conditioned during chronic CBD treatment, facilitated contextual fear learning due to enhanced hippocampal neurogenesis could explain the results reported by ElBatsh et al. (2012). This idea is supported by studies which showed that chronic CBD treatment rescues deficits in other types of memory in animal models of cognitive impairment (Fagherazzi et al., 2012; Cheng et al., 2014).

In addition to reducing contextual fear acutely when given before retrieval, CBD has also been shown to have enduring effects on fear expression when given in conjunction with memory reactivation (leading to reconsolidation) or extinction. Systemic CBD administration immediately after briefly retrieving a contextual fear memory disrupted its reconsolidation, resulting in a lasting reduction in fear expression during later retrieval (Stern et al., 2012, 2015; Gazarini et al., 2014). This effect depended on the (indirect) activation of CB1Rs rather than 5-HT_{1A}Rs as prior antagonism of CB1Rs, but not 5-HT_{1A}Rs, blocked the disruptive effect of CBD on reconsolidation (Stern et al., 2012). In contrast to its effect on the reconsolidation of contextual fear memory, CBD potentiated contextual fear extinction but this also results in reduced fear expression at later retrieval. Intracerebroventricular infusion of CBD before three longer retrieval sessions facilitated the extinction of contextual fear, an effect mediated indirectly via CB1R activation as it was blocked by CB1R antagonist pretreatment (Bitencourt et al., 2008). Systemic CBD administration before a single long retrieval session also affects contextual fear extinction but this depends on the strength of prior fear conditioning. Whereas, CBD enhanced the extinction of contextual fear resulting from strong conditioning, it impaired the contextual fear extinction induced by weaker conditioning (Song et al., this issue). **Table 1** and **Figure 1C** summarize the reported effects of systemic CBD administration on contextual fear memory processing.

Recent studies have begun to determine the neural substrates mediating the effects of CBD on contextual fear memory processing. Lemos et al. (2010) initially investigated the brain sites in which CBD acts when attenuating the expression of

learned fear. They reported that systemic CBD pretreatment prevented the increase in c-Fos expression induced by re-exposure to the conditioned context in the prelimbic (PL) and infralimbic (IL) subregions of the medial prefrontal cortex (mPFC) and the bed nucleus of the stria terminalis (BNST), brain areas which play key roles in fear regulation (Dejean et al., 2015; Lebow and Chen, 2016). Subsequently, the effect of direct CBD infusion into these brain regions on contextual fear memory retrieval was investigated. There was reduced conditioned freezing expression after its infusion into the PL or BNST (Lemos et al., 2010; Gomes et al., 2012; Fogaça et al., 2014). When infused into the IL, however, CBD produced the opposite effect (Lemos et al., 2010; Marinho et al., 2015). Convergent evidence suggests that the PL and IL present distinct neuroanatomical connections and functional roles (Hurley et al., 1991; Condé et al., 1995; Vertes, 2004), including the regulation of learned fear expression (Vidal-Gonzalez et al., 2006; Fenton et al., 2014). Contrasting effects of CBD after PL or IL infusion have also been reported in unlearned tests of anxiety (e.g., elevated plus maze), in which it induced anxiogenic- or anxiolytic-like effects, respectively (Fogaça et al., 2014; Marinho et al., 2015). Restraint stress pre-exposure reversed these effects, indicating that the stress experience could modulate the response of these mPFC subregions to aversive stimuli (Fogaça et al., 2014; Marinho et al., 2015). CBD regulation of contextual fear expression in these regions depends on 5-HT_{1A}R activation as it is prevented by local 5-HT_{1A}R antagonist pretreatment (Gomes et al., 2012; Fogaça et al., 2014; Marinho et al., 2015). CBD disruption of fear memory reconsolidation depended on CB1R but not 5-HT_{1A}R activation in the PL (Stern et al., 2014). CBD infused into the IL cortex facilitated contextual fear extinction in a CB1R-dependent manner (Do Monte et al., 2013). **Table 2** summarizes the reported effects of central CBD infusion on contextual fear memory processing.

CBD REGULATION OF FEAR MEMORY PROCESSING INVOLVING DISCRETE CUES

In contrast to contextual fear memory processing, few studies have examined the effects of CBD on learned fear related to discrete cues. One study in humans showed that CBD enhanced the extinction of visual fear memory when given immediately after, but not before, extinction (Das et al., 2013). More recently it was found in rats that CBD infused into the nucleus accumbens shell impaired the encoding of olfactory fear memory, an effect blocked by 5-HT_{1A}R antagonism (Norris et al., 2016). To investigate this issue further we determined the effect of systemic CBD administration on the expression and extinction of auditory fear memory. Male Lister hooded rats were habituated to two distinct contexts (A and B), subjected to auditory fear conditioning (tone habituation and tone-shock pairings) in context A, treated with CBD (0, 5, 10, or 20 mg/kg, i.p.; $n = 10-11/\text{group}$) 30 min before undergoing extinction training (tone presentations) in context B, and underwent extinction recall testing (tone presentations) in context B (**Figure 2A**).

TABLE 1 | Summary of systemic CBD effects on contextual fear memory processing in male rats (Δ^9 -THC, Δ^9 -tetrahydrocannabinol; BDNF, brain derived neurotrophic factor; CB1R, cannabinoid type1 receptor; ERK1/2, extracellular signal-regulated kinase1/2; i.p., intraperitoneal; PL, prelimbic; SHR, spontaneously hypertensive rat; TrkB, tyrosine receptor kinase B).

Strain	Dosing details	Effect	Possible mechanism(s)	References
Wistar	10 mg/kg, i.p., pre-retrieval	Anxiolytic (\downarrow fear expression)	Not tested	Resstel et al., 2006
Lister hooded	10 mg/kg daily for 14 days, i.p., pre-acquisition and -retrieval	Anxiogenic (\uparrow fear expression) and/or potentiated fear conditioning	\downarrow hippocampal BDNF and TrkB expression, \downarrow frontal cortex phospho-ERK1/2 levels	ElBatsh et al., 2012
Wistar and SHR	1.0–15 mg/kg, i.p., pre-acquisition	Anxiolytic (\downarrow fear expression) and/or disrupted fear memory formation (in Wistar rats only)	Not tested	Levin et al., 2012
Wistar	3.0–30 mg/kg, i.p., post-retrieval	Disrupted memory reconsolidation (bell-shaped dose-response curve)	Indirect CB1R activation	Stern et al., 2012
Wistar	10 mg/kg, i.p., post-retrieval	Disrupted memory reconsolidation	Not tested	Gazarini et al., 2014
Wistar	10 mg/kg, i.p., post-retrieval	Disrupted memory reconsolidation	Indirect CB1R activation in PL	Stern et al., 2014
Wistar	1.0 mg/kg + Δ^9 -THC 0.1 mg/kg, i.p., post-retrieval	Disrupted memory reconsolidation	Not tested	Stern et al., 2015
Lister hooded	10 mg/kg, i.p., pre-extinction (after weak or strong fear conditioning)	Impaired extinction with weak fear conditioning and enhanced extinction with strong fear conditioning	Not tested	Song et al., this issue

We found no differences in freezing during fear conditioning between the groups to receive different doses of CBD before extinction training (**Figure 2B**). Despite the relatively low levels of freezing during tone-shock pairings, conditioning resulted in robust fear memory encoding as indicated by the high freezing levels in response to tones during early extinction in vehicle-treated rats (**Figure 2C**). During extinction the high dose of CBD significantly decreased tone-induced freezing early in the session, compared to vehicle and the low dose. However, there were no differences in freezing during tone presentations later in the session or during extinction recall testing the next day (**Figure 2D**). This suggests that the high CBD dose decreased auditory fear memory expression without affecting its extinction. Although extinction occurred in a separate context to fear conditioning, vehicle-treated rats did show freezing in the period before tone presentations, indicating contextual fear expression at the start of the session (**Figure 2E**). We found that all three doses of CBD significantly decreased this contextual fear in comparison to vehicle. However, there were no differences in contextual fear expression before extinction recall testing (**Figure 2F**), suggesting that CBD acted acutely to reduce contextual fear during extinction.

Taken together, our results broadly confirm previous findings demonstrating that acute CBD treatment reduces contextual fear memory expression and extend them by showing that CBD also has a similar effect on the expression of auditory fear memory. Given that previous studies have shown a bell-shaped dose-response curve in relation to CBD regulation of anxiety and fear memory processing (Guimarães et al., 1990; Lemos et al., 2010; Stern et al., 2012), it is worth noting that we found that (1) the effect of CBD on contextual fear expression showed no dose-dependency, and (2) the dose of CBD needed to reduce auditory fear expression was much higher than for contextual fear. Moreover, CBD had no effect on auditory fear extinction. Although, this result agrees with

the study of Das et al. (2013), which showed that CBD had no effect when given before cued fear extinction, it contrasts with the reported facilitatory effects of CBD on contextual fear extinction. The reasons for these discrepancies remain unclear but they could involve differences in the neural circuit and/or pharmacological mechanisms underlying CBD regulation of contextual vs. auditory fear memory expression (see below).


FUTURE DIRECTIONS

Neural Circuit and Psychological Mechanisms

As well as the mPFC, the neural circuitry underpinning fear memory processing comprises other inter-connected areas such as the hippocampus, amygdala, and periaqueductal gray (PAG) (Dejean et al., 2015). Infusing CBD into the PAG decreases anxiety in paradigms assessing innate fear (Campos and Guimarães, 2008). However, this area is also important for mediating freezing and other defensive behaviors in response to learned threats, suggesting that CBD regulation of fear memory expression may involve the PAG. In humans CBD reduces anxiety and autonomic arousal during the viewing of fearful facial expressions, which is accompanied by decreased activity in and functional connectivity between the amygdala and mPFC (Fusar-Poli et al., 2009, 2010). In mice CBD decreases amygdala activation as measured by *c-Fos* expression (Todd and Arnold, 2016). This raises the possibility that the amygdala plays a role in mediating CBD regulation of fear memory. CBD also reduces mPFC-hippocampus functional connectivity during cognitive processing (Bhattacharyya et al., 2015). This suggests that the hippocampus, which is crucial for contextual processing in particular, may also be involved in CBD regulation of certain aspects of learned fear. After extinction, fear expression is low when tested in the extinction context but fear renewal

TABLE 2 | Summary of central CBD effects on contextual fear memory processing in male rats (5-HT_{1A}R, serotonin1A receptor; BNST, bed nucleus of the stria terminalis; CB1R, cannabinoid type1 receptor; i.c.v., intracerebroventricular; IL, infralimbic; i.p., intraperitoneal; PL, prelimbic).

Strain	Dosing and site details	Effect	Possible mechanism	References
Wistar	6.4 nmol, i.c.v., pre-retrieval	Facilitated fear extinction	Indirect CB1R activation	Bitencourt et al., 2008
Wistar	30 nmol, PL, pre-retrieval	Anxiolytic (↓ fear expression)	Not tested	Lemos et al., 2010
Wistar	30 nmol, IL, pre-retrieval	Anxiogenic (↑ fear expression)	Not tested	Lemos et al., 2010
Wistar	30–60 nmol, BNST, pre-retrieval	Anxiolytic (↓ fear expression)	5-HT _{1A} R activation	Gomes et al., 2012
Long evans	1.3 nmol, IL, pre-extinction	Facilitated fear extinction	Indirect CB1R activation	Do Monte et al., 2013
Wistar	30 nmol, PL, pre-retrieval	Anxiolytic (↓ fear expression)	5-HT _{1A} R activation	Fogaça et al., 2014
Wistar	30 nmol, IL, pre-retrieval	Anxiogenic (↑ fear expression)	5-HT _{1A} R activation	Marinho et al., 2015



occurs outside of this context. This contextual regulation of fear extinction involves the hippocampus and its connections with the mPFC and amygdala (Maren et al., 2013). Although,

we found no effects of CBD on auditory fear extinction, CBD might reduce fear renewal and/or the spontaneous recovery of fear that occurs over time after extinction through its actions on

the hippocampus-mPFC-amygdala circuit. CBD modulation of this circuitry may also regulate learned fear in other paradigms with translational relevance. Phobias and PTSD are characterized by overgeneralization of fear to harmless discrete or contextual stimuli (Dunsmoor and Paz, 2015). A recent study showed that enhancing memory destabilization in combination with reconsolidation disruption by CBD reduced contextual fear generalization (Gazarini et al., 2014).

Cellular and Molecular Mechanisms

As mentioned above, CBD regulates contextual fear memory processing in a 5HT_{1A}R- and CB1R-dependent manner. It is unclear whether the action of CBD via 5-HT_{1A}Rs to reduce fear memory expression is mediated by intracellular mechanisms. While 5-HT_{1A}Rs certainly modulate intracellular kinases, the direction of modulation appears not to be consistent with fear inhibition. The 5-HT_{1A}R activates protein kinase C (PKC) and extracellular signal-regulated kinase (ERK) (Raymond et al., 2001), but it is inhibition of PKC (Vianna et al., 2000) and ERK (Szapiro et al., 2000; Chen et al., 2005) that impairs fear memory expression. It should be noted, however, that these observations were made with intra-hippocampal drug infusions, whereas the acute effect of CBD has been shown to be mediated by the PL and BNST. Nevertheless, the putative action of CBD at 5-HT_{1A}Rs might be mediated not via protein kinases, but by inhibition of adenylyl cyclase (AC). Such inhibition of AC and downstream dysregulation of second messenger systems might be expected ultimately to affect ongoing protein synthesis. Lopez et al. (2015) showed that inhibition of ongoing protein synthesis in the amygdala impaired fear memory expression, and so a similar perturbation of normal *de novo* protein synthesis by CBD via 5-HT_{1A}Rs might be expected to impact upon retrieval. Speculatively, a pharmacologically-induced increase in protein synthesis might also sufficiently dysregulate the cellular mechanisms of retrieval to reduce fear expression. Therefore, any impact of CBD on protein synthesis could account for its acute effect on contextual, and cued, fear memory expression.

The actions of CBD to enhance extinction and impair reconsolidation both depend upon CB1Rs. Therefore, the coupling of CB1R activation to intracellular cascades may provide the cellular mechanism of long-term fear reduction. At the cellular level, mechanisms of reconsolidation tend to be conserved with those of extinction, such that inhibition results in impairment of both with opposite behavioral outcomes. A primary example of this is the effect of protein synthesis inhibitors, such as anisomycin, which impair reconsolidation to reduce fear expression as well as disrupting extinction to enhance subsequent fear expression. Therefore, in general terms, CB1R activation by CBD might be expected to result in cellular activation that would enhance extinction, but also potentiate reconsolidation. That this is seemingly not the case suggests the recruitment of specific cellular mechanisms with dissociable impacts on extinction and reconsolidation. One primary candidate for such a role is the calcineurin-nuclear factor kappa-light-chain-enhancer of activated B cells (NFkB) signaling pathway. In the hippocampus, NFkB inhibition both impaired fear memory reconsolidation and enhanced extinction (De La Fuente et al., 2011). Moreover, inhibition of calcineurin, which

itself is a negative regulator of NFkB, impaired extinction and enhanced reconsolidation (De La Fuente et al., 2011, 2014). This pattern of results was partially replicated by calcineurin inhibition in the amygdala impairing cued fear memory extinction (Merlo et al., 2014). Therefore, if CB1R activation by CBD inhibits NFkB function, either directly (Curran et al., 2005) or via enhancement of calcineurin (Cannich et al., 2004), this would explain the behavioral reductions in fear expression, perhaps through downstream regulation of cytokine networks (Scholz et al., 2016).

SUMMARY

A growing body of literature provides compelling evidence that CBD has anxiolytic effects and recent studies have established a role for CBD in regulating learned fear by dampening its expression, disrupting its reconsolidation, and facilitating its extinction. The opposing effects of CBD on fear memory reconsolidation and extinction make it particularly attractive as a potential adjunct to psychological therapy as both may lead to lasting reductions in learned fear expression. Our novel data also suggests that CBD reduces the expression of fear memory related to both discrete and contextual cues. Although we found no effect of CBD on auditory fear extinction, decreasing fear expression during extinction without interfering in its encoding is still a useful property that has clinical implications. In this respect CBD might be an improvement over other available drugs used for treating the fear-related symptoms of phobias and PTSD, which can impair extinction (e.g., benzodiazepines) or have a less favorable side effect profile (e.g., antidepressants). As such, further research investigating the mechanisms underpinning CBD regulation of learned fear is warranted.

ETHICS STATEMENT

The study was conducted in accordance with ethical review by the Animal Welfare Ethical Review Board at the University of Nottingham and the Animals (Scientific Procedures) Act 1986, UK.

AUTHOR CONTRIBUTIONS

RJ: Conducted the experiment investigating the effects of CBD on auditory fear memory expression and extinction; RJ, HD, and CS: Analyzed the data from this experiment. FG, JL, LB, and CS: Drafted the paper. All authors contributed to interpreting the original data, revising the draft and approving the final version of the paper.

ACKNOWLEDGMENTS

This research was funded by a Biotechnology and Biological Sciences Research Council Doctoral Training Partnership with the University of Nottingham (BB/J014508/1) to RJ and HD, a FAPESP-University of Birmingham-University of Nottingham pump-priming award (2012/50896-8) to FG, JL, and CS, and a

Brazilian CNPq research fellowship (307895/2013-0) to LB. The funders had no other involvement in any aspect of this work. The data presented in **Figure 2** was first presented as an abstract at the 26th Annual International Cannabinoid Research Society Symposium on the Cannabinoids in June 2016.

REFERENCES

- Baldwin, D. S., Anderson, I. M., Nutt, D. J., Allgulander, C., Bandelow, B., den Boer, J. A., et al. (2014). Evidence-based pharmacological treatment of anxiety disorders, post-traumatic stress disorder and obsessive-compulsive disorder: a revision of the 2005 guidelines from the British Association for Psychopharmacology. *J. Psychopharmacol.* 28, 403–439. doi: 10.1177/0269881114525674
- Bhattacharyya, S., Falkenberg, I., Martin-Santos, R., Atakan, Z., Crippa, J. A., Giampietro, V., et al. (2015). Cannabinoid modulation of functional connectivity within regions processing attentional salience. *Neuropsychopharmacology* 40, 1343–1352. doi: 10.1038/npp.2014.258
- Bitencourt, R. M., Pamplona, F. A., and Takahashi, R. N. (2008). Facilitation of contextual fear memory extinction and anti-anxiogenic effects of AM404 and cannabidiol in conditioned rats. *Eur. Neuropsychopharmacol.* 18, 849–859. doi: 10.1016/j.euroneuro.2008.07.001
- Blessing, E. M., Steenkamp, M. M., Manzaneres, J., and Marmar, C. R. (2015). Cannabidiol as a potential treatment for anxiety disorders. *Neurotherapeutics* 12, 825–836. doi: 10.1007/s13311-015-0387-1
- Campos, A. C., Fogaça, M. V., Sonogo, A. B., and Guimarães, F. S. (2016). Cannabidiol, neuroprotection and neuropsychiatric disorders. *Pharmacol. Res.* 112, 119–127. doi: 10.1016/j.phrs.2016.01.033
- Campos, A. C., and Guimarães, F. S. (2008). Involvement of 5HT1A receptors in the anxiolytic-like effects of cannabidiol injected into the dorsolateral periaqueductal gray of rats. *Psychopharmacology (Berl.)* 199, 223–230. doi: 10.1007/s00213-008-1168-x
- Campos, A. C., Ortega, Z., Palazuelos, J., Fogaça, M. V., Aguiar, D. C., Díaz-Alonso, J., et al. (2013). The anxiolytic effect of cannabidiol on chronically stressed mice depends on hippocampal neurogenesis: involvement of the endocannabinoid system. *Int. J. Neuropsychopharmacol.* 16, 1407–1419. doi: 10.1017/S1461145712001502
- Cannich, A., Wotjak, C. T., Kamprath, K., Hermann, H., Lutz, B., and Marsicano, G. (2004). CB1 cannabinoid receptors modulate kinase and phosphatase activity during extinction of conditioned fear in mice. *Learn. Mem.* 11, 625–632. doi: 10.1101/lm.77904
- Chen, X., Garelick, M. G., Wang, H., Lil, V., Athos, J., and Storm, D. R. (2005). PI3 kinase signaling is required for retrieval and extinction of contextual memory. *Nat. Neurosci.* 8, 925–931. doi: 10.1038/nn1482
- Cheng, D., Low, J. K., Logge, W., Garner, B., and Karl, T. (2014). Chronic cannabidiol treatment improves social and object recognition in double transgenic APP^{swE}/PS1 Δ E9 mice. *Psychopharmacology (Berl.)* 231, 3009–3017. doi: 10.1007/s00213-014-3478-5
- Condé, F., Maire-Lepoivre, E., Audinat, E., and Crépel, F. (1995). Afferent connections of the medial frontal cortex of the rat. II. Cortical and subcortical afferents. *J. Comp. Neurol.* 352, 567–593. doi: 10.1002/cne.903520407
- Curran, N. M., Griffin, B. D., O'Toole, D., Brady, K. J., Fitzgerald, S. N., and Moynagh, P. N. (2005). The synthetic cannabinoid R(+)-WIN 55,212-2 inhibits the interleukin-1 signaling pathway in human astrocytes in a cannabinoid receptor-independent manner. *J. Biol. Chem.* 280, 35797–35806. doi: 10.1074/jbc.M507959200
- Das, R. K., Kamboj, S. K., Ramadas, M., Yogan, K., Gupta, V., Redman, E., et al. (2013). Cannabidiol enhances consolidation of explicit fear extinction in humans. *Psychopharmacology (Berl.)* 226, 781–792. doi: 10.1007/s00213-012-2955-y
- Dejean, C., Courtin, J., Rozeske, R. R., Bonnet, M. C., Dousset, V., Michelet, T., et al. (2015). Neuronal circuits for fear expression and recovery: recent advances and potential therapeutic strategies. *Biol. Psychiatry* 78, 298–306. doi: 10.1016/j.biopsych.2015.03.017
- de la Fuente, V., Federman, N., Fustiñana, M. S., Zalman, G., and Romano, A. (2014). Calcineurin phosphatase as a negative regulator of fear memory in hippocampus: control on nuclear factor-kappaB signaling in consolidation and reconsolidation. *Hippocampus* 24, 1549–1561. doi: 10.1002/hipo.22334
- De La Fuente, V., Freudenthal, R., and Romano, A. (2011). Reconsolidation or extinction: transcription factor switch in the determination of memory course after retrieval. *J. Neurosci.* 31, 5562–5573. doi: 10.1523/JNEUROSCI.6066-10.2011
- Denny, C. A., Kheirbek, M. A., Alba, E. L., Tanaka, K. F., Brachman, R. A., Laughman, K. B., et al. (2014). Hippocampal memory traces are differentially modulated by experience, time, and adult neurogenesis. *Neuron* 83, 189–201. doi: 10.1016/j.neuron.2014.05.018
- Do Monte, F. H., Souza, R. R., Bitencourt, R. M., Kroon, J. A., and Takahashi, R. N. (2013). Infusion of cannabidiol into infralimbic cortex facilitates fear extinction via CB1 receptors. *Behav. Brain Res.* 250, 23–27. doi: 10.1016/j.bbr.2013.04.045
- Dunsmoor, J. E., and Paz, R. (2015). Fear generalization and anxiety: behavioral and neural mechanisms. *Biol. Psychiatry* 78, 336–343. doi: 10.1016/j.biopsych.2015.04.010
- Efstathopoulos, P., Kourgiantaki, A., Karali, K., Sidiropoulou, K., Margioris, A. N., Gravanis, A., et al. (2015). Fingolimod induces neurogenesis in adult mouse hippocampus and improves contextual fear memory. *Transl. Psychiatry* 5, e685. doi: 10.1038/tp.2015.179
- ElBatsch, M. M., Assareh, N., Marsden, C. A., and Kendall, D. A. (2012). Anxiogenic-like effects of chronic cannabidiol administration in rats. *Psychopharmacology (Berl.)* 221, 239–247. doi: 10.1007/s00213-011-2566-z
- Fagherazzi, E. V., Garcia, V. A., Maurmann, N., Bervanger, T., Halmenschlager, L. H., Busato, S. B., et al. (2012). Memory-rescuing effects of cannabidiol in an animal model of cognitive impairment relevant to neurodegenerative disorders. *Psychopharmacology (Berl.)* 219, 1133–1140. doi: 10.1007/s00213-011-2449-3
- Fendt, M., and Fanselow, M. S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. *Neurosci. Biobehav. Rev.* 23, 743–760. doi: 10.1016/S0149-7634(99)00016-0
- Fenton, G. E., Pollard, A. K., Halliday, D. M., Mason, R., Bredy, T. W., and Stevenson, C. W. (2014). Persistent prelimbic cortex activity contributes to enhanced learned fear expression in females. *Learn. Mem.* 21, 55–60. doi: 10.1101/lm.033514.113
- Fineberg, N. A., Haddad, P. M., Carpenter, L., Gannon, B., Sharpe, R., Young, A. H., et al. (2013). The size, burden and cost of disorders of the brain in the UK. *J. Psychopharmacol.* 27, 761–770. doi: 10.1177/0269881113495118
- Fogaça, M. V., Reis, F. M., Campos, A. C., and Guimarães, F. S. (2014). Effects of intra-prelimbic prefrontal cortex injection of cannabidiol on anxiety-like behavior: involvement of 5HT1A receptors and previous stressful experience. *Eur. Neuropsychopharmacol.* 24, 410–419. doi: 10.1016/j.euroneuro.2013.10.012
- Fusar-Poli, P., Allen, P., Bhattacharyya, S., Crippa, J. A., Mechelli, A., Borgwardt, S., et al. (2010). Modulation of effective connectivity during emotional processing by Delta 9-tetrahydrocannabinol and cannabidiol. *Int. J. Neuropsychopharmacol.* 13, 421–432. doi: 10.1017/S1461145709990617
- Fusar-Poli, P., Crippa, J. A., Bhattacharyya, S., Borgwardt, S. J., Allen, P., Martin-Santos, R., et al. (2009). Distinct effects of [delta]9-tetrahydrocannabinol and cannabidiol on neural activation during emotional processing. *Arch. Gen. Psychiatry* 66, 95–105. doi: 10.1001/archgenpsychiatry.2008.519
- Gazarini, L., Stern, C. A. J., Piornedo, R. R., Takahashi, R. N., and Bertoglio, L. J. (2014). PTSD-like memory generated through enhanced noradrenergic activity is mitigated by a dual step pharmacological intervention targeting its reconsolidation. *Int. J. Neuropsychopharmacol.* 18:pyu026. doi: 10.1093/ijnp/pyu026
- Gomes, F. V., Reis, D. G., Alves, F. H., Corrêa, F. M., Guimarães, F. S., and Resstel, L. B. (2012). Cannabidiol injected into the bed nucleus of the stria terminalis reduces the expression of contextual fear conditioning via 5-HT1A receptors. *J. Psychopharmacol.* 26, 104–113. doi: 10.1177/0269881110389095

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fphar.2016.00454/full#supplementary-material>

- Guimarães, F. S., Chiaretti, T. M., Graeff, F. G., and Zuardi, A. W. (1990). Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology (Berl)*. 100, 558–559. doi: 10.1007/BF02244012
- Hurley, K. M., Herbert, H., Moga, M. M., and Saper, C. B. (1991). Efferent projections of the infralimbic cortex of the rat. *J. Comp. Neurol.* 308, 249–276. doi: 10.1002/cne.903080210
- Izzo, A. A., Borrelli, F., Capasso, R., Di Marzo, V., and Mechoulam, R. (2009). Non-psychotropic plant cannabinoids: new therapeutic opportunities from an ancient herb. *Trends Pharmacol. Sci.* 30, 515–527. doi: 10.1016/j.tips.2009.07.006
- Lebow, M. A., and Chen, A. (2016). Overshadowed by the amygdala: the bed nucleus of the stria terminalis emerges as key to psychiatric disorders. *Mol. Psychiatry* 21, 450–463. doi: 10.1038/mp.2016.1
- Lee, J. L. (2009). Reconsolidation: maintaining memory relevance. *Trends Neurosci.* 32, 413–420. doi: 10.1016/j.tins.2009.05.002
- Lemos, J. I., Resstel, L. B., and Guimarães, F. S. (2010). Involvement of the prelimbic prefrontal cortex on cannabidiol-induced attenuation of contextual conditioned fear in rats. *Behav. Brain Res.* 207, 105–111. doi: 10.1016/j.bbr.2009.09.045
- Levin, R., Almeida, V., Peres, F. F., Calzavara, M. B., da Silva, N. D., Suíama, M. A., et al. (2012). Antipsychotic profile of cannabidiol and rimonabant in an animal model of emotional context processing in schizophrenia. *Curr. Pharm. Des.* 18, 4960–4965. doi: 10.2174/138161212802884735
- Lopez, J., Gamache, K., Schneider, R., and Nader, K. (2015). Memory retrieval requires ongoing protein synthesis and NMDA receptor activity-mediated AMPA receptor trafficking. *J. Neurosci.* 35, 2465–2475. doi: 10.1523/JNEUROSCI.0735-14.2015
- Maren, S., Phan, K. L., and Liberzon, I. (2013). The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nat. Rev. Neurosci.* 14, 417–428. doi: 10.1038/nrn3492
- Marinho, A. L., Vila-Verde, C., Fogaça, M. V., and Guimarães, F. S. (2015). Effects of intra-infralimbic prefrontal cortex injections of cannabidiol in the modulation of emotional behaviors in rats: contribution of 5HT_{1A} receptors and stressful experiences. *Behav. Brain Res.* 286, 49–56. doi: 10.1016/j.bbr.2015.02.023
- Merlo, E., Milton, A. L., Goozee, Z. Y., Theobald, D. E., and Everitt, B. J. (2014). Reconsolidation and extinction are dissociable and mutually exclusive processes: behavioral and molecular evidence. *J. Neurosci.* 34, 2422–2431. doi: 10.1523/JNEUROSCI.4001-13.2014
- Myers, K. M., and Davis, M. (2007). Mechanisms of fear extinction. *Mol. Psychiatry* 12, 120–150. doi: 10.1038/sj.mp.4001939
- National Center for Biotechnology Information (2016). *National Center for Biotechnology Information. PubChem Compound Database CID=644019*. Available online at: <https://pubchem.ncbi.nlm.nih.gov/compound/644019> (Accessed November 6, 2016).
- Norris, C., Loureiro, M., Kramar, C., Zunder, J., Renard, J., Rushlow, W., et al. (2016). Cannabidiol modulates fear memory formation through interactions with serotonergic transmission in the mesolimbic system. *Neuropsychopharmacology* 41, 2839–2850. doi: 10.1038/npp.2016.93
- Pan, Y. W., Storm, D. R., and Xia, Z. (2012). The maintenance of established remote contextual fear memory requires ERK5 MAP kinase and ongoing adult neurogenesis in the hippocampus. *PLoS ONE* 7:e50455. doi: 10.1371/journal.pone.0050455
- Raymond, J. R., Mukhin, Y. V., Gelasco, A., Turner, J., Collinsworth, G., Gettys, T. W., et al. (2001). Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacol. Ther.* 92, 179–212. doi: 10.1016/S0163-7258(01)00169-3
- Resstel, L. B., Joca, S. R., Moreira, F. A., Corrêa, F. M., and Guimarães, F. S. (2006). Effects of cannabidiol and diazepam on behavioral and cardiovascular responses induced by contextual conditioned fear in rats. *Behav. Brain Res.* 172, 294–298. doi: 10.1016/j.bbr.2006.05.016
- Resstel, L. B., Moreira, F. A., and Guimarães, F. S. (2009). Endocannabinoid system and fear conditioning. *Vitam. Horm.* 81, 421–440. doi: 10.1016/S0083-6729(09)81016-9
- Scholz, B., Doidge, A. N., Barnes, P., Hall, J., Wilkinson, L. S., and Thomas, K. L. (2016). The regulation of cytokine networks in hippocampal CA1 differentiates extinction from those required for the maintenance of contextual fear memory after recall. *PLoS ONE* 11:e0153102. doi: 10.1371/journal.pone.0153102
- Seo, D. O., Carillo, M. A., Chih-Hsiung Lim, S., Tanaka, K. F., and Drew, M. R. (2015). Adult hippocampal neurogenesis modulates fear learning through associative and nonassociative mechanisms. *J. Neurosci.* 35, 11330–11345. doi: 10.1523/JNEUROSCI.0483-15.2015
- Steckler, T., and Risbrough, V. (2012). Pharmacological treatment of PTSD - established and new approaches. *Neuropharmacology* 62, 617–627. doi: 10.1016/j.neuropharm.2011.06.012
- Stern, C. A., Gazarini, L., Takahashi, R. N., Guimarães, F. S., and Bertoglio, L. J. (2012). On disruption of fear memory by reconsolidation blockade: evidence from cannabidiol treatment. *Neuropsychopharmacology* 37, 2132–2142. doi: 10.1038/npp.2012.63
- Stern, C. A., Gazarini, L., Vanvossen, A. C., Zuardi, A. W., Galve-Roperh, I., Guimarães, F. S., et al. (2015). Δ^9 -Tetrahydrocannabinol alone and combined with cannabidiol mitigate fear memory through reconsolidation disruption. *Eur. Neuropsychopharmacol.* 25, 958–965. doi: 10.1016/j.euroneuro.2015.02.001
- Stern, C. A., Gazarini, L., Vanvossen, A. C., Zuardi, A. W., Guimarães, F. S., Takahashi, R. N., et al. (2014). Involvement of the prelimbic cortex in the disruptive effect of cannabidiol on fear memory reconsolidation. *Eur. Neuropsychopharmacol.* 24(Suppl. 2), S322. doi: 10.1016/S0924-977X(14)70511-5
- Szapiro, G., Izquierdo, L. A., Alonso, M., Barros, D., Paratcha, G., Ardenghi, P., et al. (2000). Participation of hippocampal metabotropic glutamate receptors, protein kinase A and mitogen-activated protein kinases in memory retrieval. *Neuroscience* 99, 1–5. doi: 10.1016/S0306-4522(00)00236-0
- Todd, S. M., and Arnold, J. C. (2016). Neural correlates of interactions between cannabidiol and Δ^9 -tetrahydrocannabinol in mice: implications for medical cannabis. *Br. J. Pharmacol.* 173, 53–65. doi: 10.1111/bph.13333
- Vertes, R. P. (2004). Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse* 51, 32–58. doi: 10.1002/syn.10279
- Vianna, M. R., Barros, D. M., Silva, T., Choi, H., Madche, C., Rodrigues, C., et al. (2000). Pharmacological demonstration of the differential involvement of protein kinase C isoforms in short- and long-term memory formation and retrieval of one-trial avoidance in rats. *Psychopharmacology (Berl)*. 150, 77–84. doi: 10.1007/s002130000396
- Vidal-Gonzalez, I., Vidal-Gonzalez, B., Rauch, S. L., and Quirk, G. J. (2006). Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. *Learn. Mem.* 13, 728–733. doi: 10.1101/lm.306106
- Wolf, S. A., Bick-Sander, A., Fabel, K., Leal-Galicia, P., Tauber, S., Ramirez-Rodriguez, G., et al. (2010). Cannabinoid receptor CB1 mediates baseline and activity-induced survival of new neurons in adult hippocampal neurogenesis. *Cell Commun. Signal.* 8:12. doi: 10.1186/1478-811X-8-12

Conflict of Interest Statement: FG is co-inventor of the patent “Fluorinated CBD compounds, compositions and uses thereof. Pub. No.: WO/2014/108899. International Application No: PCT/IL2014/050023”; Def. US no. Reg. 62193296; 29/07/2015; INPI in 19/08/2015 (BR1120150164927).

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Jurkus, Day, Guimarães, Lee, Bertoglio and Stevenson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.