

Reducing neuroinflammation by delivery of IL-10 encoding lentivirus from multiple-channel bridges

Daniel J. Margul^{1,2} | Jonghyuck Park² | Ryan M. Boehler³ | Dominique R. Smith^{1,2} | Mitchell A. Johnson² | Dylan A. McCreedy^{2,3} | Ting He³ | Aishani Ataliwala⁴ | Todor V. Kukushliev³ | Jesse Liang⁴ | Alireza Sohrabi⁴ | Ashley G. Goodman³ | Christopher M. Walthers⁴ | Lonnie D. Shea^{2,5*} | Stephanie K. Seidlits^{3,4,6,7*}

¹Dept. of Biomedical Engineering, Northwestern University, Evanston, IL, 48109

²Dept. of Biomedical Engineering, University of Michigan, Ann Arbor, MI, 48109

³Dept. of Chemical and Biological Engineering, Northwestern University, Evanston, IL, 48109

⁴Dept. of Bioengineering, University of California Los Angeles, Los Angeles, CA, 90095

⁵Dept. of Chemical Engineering, University of Michigan, Ann Arbor, MI, 48109

⁶Brain Research Institute, University of California Los Angeles, Los Angeles, CA, 90095

⁷Jonsson Comprehensive Cancer Center, University of California Los Angeles, Los Angeles, CA, 90024

*Stephanie K. Seidlits and Lonnie D. Shea are co-corresponding authors.

Correspondence

Lonnie D. Shea, Dept. of Biomedical Engineering, University of Michigan, 2200 Bonisteel Blvd, 1119 Carl A. Gerstacker Building, Ann Arbor, MI, 48109.

Email: ldshea@umich.edu or

Stephanie K. Seidlits, Dept. of Bioengineering, University of California Los Angeles, 420 Westwood Plaza, Engineering V room 5121H, Los Angeles, CA 90095.

Email: seidlits@g.ucla.edu.

Funding information: Financial support for this research was provided by the NIH (RO1 EB005678). This work was supported by the Northwestern University – Flow Cytometry Core Facility supported by Cancer Center Support Grant (NCI CA060553).

Abstract

The spinal cord is unable to regenerate after injury largely due to growth-inhibition by an inflammatory response to the injury that fails to resolve, resulting in secondary damage and cell death. An approach that prevents inhibition by attenuating the inflammatory response and promoting its resolution through the transition of macrophages to anti-inflammatory phenotypes is essential for the creation of a growth permissive microenvironment. Viral gene delivery to induce the expression of anti-inflammatory factors provides the potential to provide localized delivery to alter the host inflammatory response. Initially, we investigated the effect of the biomaterial and viral components of the delivery system to influence the extent of cell infiltration and the phenotype of these cells. Bridge implantation reduces antigen-presenting cell infiltration at day 7, and lentivirus addition to the bridge induces a transient increase in neutrophils in the spinal cord at day 7 and macrophages at day 14. Delivery of a lentivirus encoding IL-10, an anti-inflammatory factor that inhibits immune cell activation and polarizes the macrophage population towards anti-inflammatory phenotypes, reduced neutrophil infiltration at both day 7 and day 28. Though IL-10 lentivirus did not affect macrophages number, it skewed the macrophage population toward an anti-inflammatory M2 phenotype and altered macrophage morphology. Additionally, IL-10 delivery resulted in improved motor function, suggesting reduced secondary damage and increased sparing. Taken together, these results indicate that localized expression of anti-inflammatory factors, such as IL-10, can modulate the inflammatory response following spinal cord injury, and may be a key component of a combinatorial approach that targets the multiple barriers to regeneration and functional recovery.

KEYWORDS

immune modulation, gene delivery, multiple-channel bridge, spinal cord injury

1 | INTRODUCTION

To date, the only treatment for treating spinal cord injury (SCI) with any degree of success in clinical trials is methylprednisolone, which mitigates the inflammatory response after injury.¹ Inflammation after SCI initiates a damaging secondary injury response that establishes multiple barriers (e.g., accumulation of myelin debris, and reactive gliosis) to regeneration and restoration of function. The inflammatory cascade begins with upregulated production of inflammatory cytokines, chemokines, and reactive oxygen species (ROS),^{2–5} which leads to recruitment of peripheral immune cells⁶ and activation of local microglia, oligodendrocytes, and astrocytes.^{7–10} Early recruitment of neutrophils (PMNs)—and later macrophages—comprises the innate immune response to SCI. PMNs and macrophages recruited to the injury can remove pathogens; however, they also propagate secondary injury, which results in death of neurons and oligodendrocytes.^{11,12} Macrophages, in particular, also phagocytose debris left behind by necrotic or apoptotic cells, including inhibitory myelin debris whose slow clearance contributes to the inability of axons to regenerate.^{13,14} In addition, inflammatory cytokines produced by PMNs and macrophages induce astrocytes to adopt a reactive state in which they proliferate and can increase the secretion of chondroitin sulfate proteoglycans (CSPGs). These reactive astrocytes then form an inhibitory glial scar that surrounds the injury site and prevents regenerating axons from crossing.^{9,14–19}

In normal wound healing, the inflammatory response resolves over time, leaving behind regenerated, functional tissue. In contrast, SCI results in a chronic immune response characterized by the persistence of inflammatory cells, insufficient clearance of cellular debris at the injury, and formation of a robust glial scar that creates a barrier that limits regeneration.^{12,20,21} In peripheral tissues, a transition in macrophage phenotype from inflammatory (M1 or classically activated) to anti-inflammatory (M2 or alternatively activated) is largely responsible for dampening and resolving the immune response after injury.^{22–24} However, this transition does not occur on a sufficient scale after SCI. Instead, M1 macrophages chronically persist in zones of axon degeneration, where they propagate inflammation and inhibit axon regeneration.^{25–30} Though a preponderance of evidence has shown that M1 macrophages predominate within the injury site after SCI, M2 macrophages are present—typically peaking at day 14 after injury in rodents.³¹ M2 macrophages generally have enhanced phagocytic capabilities^{32,33} and can promote axon growth across inhibitory boundaries *in vitro*.²⁵ They stimulate tissue repair through attenuated production of inflammatory cytokines, reduced ROS production, and expression of pro-resolving cytokines such as IL-10.³⁴ In contrast, M1 macrophages are characterized by increased production of pro-inflammatory cytokines, ROS and NO—each of which leads to neuron and oligodendrocyte toxicity.^{25,35} Several factors promote M2 phenotypes including IL-4, IL-13, glucocorticoids, and IL-10. Of these, IL-10 was selected for this study due to its ability to promote neuron survival while reducing leukocyte infiltration and activation.^{36–38}

Macrophage depletion has been investigated as a potential solution to mitigate the inflammatory response, but there have been mixed results. Treatment with liposome-encapsulated clodronate to deplete hematogenous macrophages improved partial hindlimb recovery and tissue repair in

one study³⁹ and decreased fibrotic scarring while increasing axon numbers in a second study.⁴⁰ In contrast, antibody-mediated depletion of CD11c⁺ monocytes/macrophages or conditional ablation by diphtheria toxin each reduced functional recovery.⁴¹ These seemingly contradictory results suggest that specific macrophage phenotypes may have different effects on tissue repair and that eliminating or enhancing specific subtypes may be more beneficial than targeting the entire macrophage population. Additionally, there may be unintended systemic or local consequences of depleting the entire population, such as increased susceptibility to infection and disease and reduced clearance of inhibitory debris.

While macrophage phenotype has often been depicted as a binary system where a cell is either M1 or M2 at any given time, it is now thought that macrophages exist along a continuous spectrum of activation states, with M1 and M2 as polar opposites.^{23,42} For example, some reports have sub-divided M2 macrophages into additional subtypes (e.g., M2a, M2b, and M2c), each of which has a role in suppressing inflammation. The M2a (alternative) macrophages are involved in initial wound healing and the Th2-type (T-cell mediated) inflammatory response, while M2b (type 2) macrophages are believed to be immunoregulatory. Finally, M2c (deactivated) macrophages are immunosuppressive and facilitate matrix deposition and issue remodeling.^{43,44} Despite observations of these macrophage subtypes *in vitro*, it remains unclear how this correlates with macrophage activation *in vivo*. Macrophages retain inherent capacity for plasticity along this activation spectrum, dynamically responding to changes in their local environment.^{36,45} This dynamic plasticity makes macrophages a compelling target for resolving inflammation in the spinal cord to enable regeneration.

In addition to addressing neuroinflammation, therapies for spinal cord regeneration must provide physical support and guidance for regenerating axons across the injury site. SCI therapies focused solely on resolution of neuroinflammation have largely failed due to both lack of a growth-promoting substrate and accumulation of inhibitory factors at the injury.^{46,47} While immunomodulatory strategies can reduce inflammation and promote axonal sparing, guidance of regenerating axons requires a permissive substrate. Initial attempts to provide a permissive bridge across spinal cord lesions implanted donor peripheral nerve grafts.⁴⁸ More recently, synthetic multiple-channel bridges have been developed to replicate the synergistic advantages of peripheral nerve grafts—physical support and guidance and a biologically active, pro-regenerative microenvironment.^{49–54} We have previously demonstrated that multiple-channel bridges made from biodegradable poly(lactide-co-glycolide) (PLG) support robust crossing of axons across the lesion site, resulting in formation of regenerated axon bundles after complete bridge degradation 6 months after SCI.^{54,55} Longitudinal, macroscale channels within these bridges act as conduits for axons regenerating across the injury site, while scaffold microporosity supports host cell infiltration and tissue integration.^{55,56}

Herein, we investigate the hypothesis that localized lentiviral expression of IL-10 from multiple-channel PLG bridges will modulate the numbers, phenotypes, and proportions of leukocyte populations infiltrating the bridge to promote a “resolving” anti-inflammatory environment thought to be more permissive to regeneration. In addition to providing a substrate for regeneration, PLG bridges have also served as

a platform for localized delivery of gene therapy vectors that can induce the expression of therapeutic proteins.^{54,57} Biomaterial-mediated lentivirus delivery after SCI efficiently transduces infiltrating cells to yield a localized, stable pattern of gene expression.^{54,58} Lentiviral particles associated with the heparin-modified PLG bridges, which functions to retain the vector locally and can increase its half-life for enhanced gene transfer. Infiltrating host cells are transduced by the lentivirus, with peak transgene expression occurring approximately 7 days after bridge transplantation with sustained expression for at least 8 weeks.^{54,58} Expression of IL-10 will be investigated due to its neuroprotective and anti-apoptotic properties, as well as its ability to skew macrophages towards anti-inflammatory phenotypes.^{36,59} The studies focus on the first four weeks after injury comprising the acute, subacute, and intermediate phases of recovery, which encompasses the time over which many of the barriers to regeneration become established.

2 | MATERIALS AND METHODS

2.1 | Virus production

Lentivirus was produced by co-transfecting HEK-293T cells with third generation lentiviral packaging vectors (pMDL-GagPol, pRSV-Rev, pVSV-G,⁶⁰ and the gene of interest (pLenti-CMV-Luciferase or pLenti-CMV-IL-10) using Lipofectamine 2000 (Life Technologies, Grand Island, NY, USA). After, lentiviral particles were purified using the Lenti-X Maxi Purification Kit (Clontech Laboratories, Mountain View, CA, USA) and then concentrated using Vivaspin centrifugal concentrators (Sartorius, Göttingen, Germany). Viral titers used throughout the study were 2E10 IU/ml as determined by the Lentivirus qPCR Titer Kit (Applied Biological Materials, Richmond, BC, Canada).

2.2 | Fabrication of multiple-channel bridges

Bridges were fabricated using a sacrificial template variation⁶¹ of the gas foaming/particulate leaching technique,⁵⁴ as previously described.⁶² Briefly, PLG (75:25 lactide:glycolide; inherent viscosity 0.76 dl/g; Lakeshore Biomaterials, Birmingham, AL, USA) was dissolved in dichloromethane (6% w/w) and emulsified in 1% poly(vinyl alcohol) using a homogenizer (PolyTron 3100; Kinematica AG, Littau, Switzerland) at 3000 rpm to create microspheres (z-average diameter \sim 1 μ m). D-sucrose was caramelized, cooled, and drawn from solution with a Pasteur pipette to make sugar fibers. These fibers were coated with a mixture of PLG microspheres and salt (63–106 μ m) and pressed into a salt-lined aluminum mold. The materials were then equilibrated with CO₂ gas (800 psi) for 16 hrs and then gas foamed in a custom-made pressure vessel. Bridges were subsequently cut into 2.25 mm sections and leached for 2 hrs to remove porogens. The bridges are dried overnight and stored in a desiccator.

2.3 | Heparinization of nerve bridges

Heparin coating of bridges has been shown to enhance lentiviral loading and transduction from bridges *in vivo*.⁶³ PLG bridges were incubated with chitosan (Sigma Aldrich, St. Louis, MO, USA; 25 μ g/ μ l in 2% glacial acetic acid) for 10 min, followed in a N-(3-dimethylamino-

propyl)-N'-ethylcarbodiimide/N-hydroxysuccinimide (Sigma Aldrich) mixture dissolved in 2-(N-morpholino)ethanesulfonic acid (EDC/NHS in MES; 1.5:1:1 mg/mg/ml, Sigma Aldrich) for 2 hrs. To conjugate chitosan, bridges were washed 3 times with water, dried and subsequently incubated with heparin (Sigma Aldrich; 25 μ g/ μ l in 1M MES) for 10 min followed by EDC/NHS in MES for an additional 2 hrs for covalent conjugation. Finally, bridges were washed in water three times and dried.

2.4 | Virus loading onto heparinized nerve bridges

Multiple additions of viruses were adsorbed onto bridges in an iterative manner in order to increase lentiviral loading. Prior to virus addition, bridges were disinfected in 70% ethanol and washed with water. After 12 min of drying time, bridges were saturated with 2 μ l of virus. Bridges were then dried for an additional 12 min followed by another 2 μ l of virus. Bridges were then dried for 14 additional min followed by a final 2 μ l of virus. After a final 5 min of drying, bridges were stored at -80° C until used for surgery.

2.5 | Mouse spinal cord hemisection

A hemisection model of SCI was performed as previously described⁶² on female C57/BL6 mice (4–6 weeks-old; Charles River Laboratories, Wilmington, MA, USA), according to the Animal Care and Use Committee guidelines at Northwestern University. A laminectomy was performed at T9–T10 to allow for a 2.25 mm lateral hemisection for bridge implantation. The injury site was covered using Gelfoam (Pfizer, New York, NY, USA) followed by suturing together of the muscle and stapling of skin. Postoperative care consisted of administration of enrofloxacin (2.5 mg/kg; daily for 2 weeks), buprenorphine (0.1 mg/kg; twice daily for 3 days), and Lactated Ringer's solution (5 ml/100 g; daily for 5 days). Bladders were expressed twice daily until function recovered.

2.6 | Tissue processing and immunofluorescence

Spinal cord tissue was collected at days 7, 14, and 28, which were chosen to represent acute, subacute and intermediate phases of regeneration respectively. It may be noted that days 7 and 14 are at the tail end of their respective phases; however, it is not possible to extract the spinal cord and bridge together prior to day 7 as the bridge falls out due to a weak interface tissue-bridge interface. Additionally, cellular infiltration is lower at earlier time points, making flow cytometry challenging. For immunofluorescence, spinal cord segments were then snap frozen in isopentane and embedded in Tissue Tek O.C.T. Compound (Sakura Finetek, Torrance, CA, USA) with 20% sucrose. Cords were cryosectioned transversely in 18- μ m-thick sections.

Antibodies against the following antigens were used for immunofluorescence: neurofilament 200 (NF200, Sigma Aldrich), myelin basic protein (MBP, Santa Cruz Biotech, Dallas, TX, USA) P-zero myelin protein (P0) (Aves Labs, Tigard, OR) F4/80 (AbD Serotec, Raleigh, NC, USA) arginase I (clone N20, Santa Cruz Biotech) and Ly-6G (clone RB6-8C5, Biolegend).

Using stained tissue sections, immuno-positive cells and axons within the bridge area were quantified manually by two blinded researchers independently. Co-staining for multiple markers was assessed by evaluating pixel overlap of different channels in NIH ImageJ (Bethesda, MD,

USA). Total numbers of F4/80⁺ cells and numbers of F4/80⁺/arginase⁺ cells were evaluated to determine numbers of total macrophages and M2 macrophages, respectively. Macrophages were identified by localizing Hoechst nuclear staining to F4/80⁺ immunofluorescence and labeling them in ImageJ. Subsequently, each identified F4/80⁺ cell was binned based on having arginase immunofluorescence visible over background. Cell counts were then normalized to counted area in each tissue section. Macrophage shape was characterized for each positive cell and cells were binned as fibrous/elongated, round and not ruffled with a bright F4/80⁺ perimeter, or as multinucleated foreign body giant cell (FBGC). To assess the numbers of regenerated and myelinated axons, NF200 was used to identify axons, NF200⁺/MBP⁺ to determine the number of myelinated axons, and NF200⁺/MBP⁺/PO⁺ to determine the amount of myelin derived from infiltrating Schwann cells.⁵⁷

2.7 | Flow cytometry of digested tissue samples

Flow cytometry was performed on spinal cord bridge implants loaded with no lentivirus, lentivirus encoding firefly luciferase (FLUC) or IL-10 to determine the identity of infiltrating cells 7, 14, and 28 days after injuries. Spinal cord hemisection with no bridge implants were also performed. To isolate sufficient cell numbers for flow cytometry, implant sites from 3 individual mice were pooled together per condition. Bridge tissue samples were mechanically dissociated by 16g needle and syringe and digested in a solution containing collagenase (1 mg/ml; Worthington Biochemical Corp., Lakewood, NJ, USA) and trypsin (0.5 mg/ml; Life Technologies) for 20 min at 37°C. After incubation, 1 ml of fetal bovine serum (Sigma Aldrich) was immediately added to inhibit digestion and solution was triturated with fire-polished glass pipettes of at least three successively decreasing diameters to achieve a single cell suspension. Digested tissues were pushed through 70 μm strainers and washed with Hank's Balanced Salt Solution without calcium and magnesium (VWR, Radnor, PA, USA). Cells were then separated from myelin using the OptiPrep gradient system (Sigma Aldrich) in MOPS buffer, as previously described by Beck et al.²⁰ Cells were subsequently blocked with a solution containing 1% normal mouse and rat serum (Sigma Aldrich) and anti-mouse CD16/32 (eBioscience) and stained for viability using fixable violet dead cell stain (Invitrogen) and stained with the specific antibodies listed above. Data were acquired on a BD LSR II cytometer, and analyzed using FloJo software. Fluorescence minus one staining was used as a negative control.

The following flow cytometry antibodies were used: v500-conjugated 30-F11 against CD45, PE(phycoerythrin)/Cy7-conjugated RM4-5 against CD4, and Alexa Fluor 647-conjugated 1B4 against GFAP (glial fibrillary acidic protein) (BD Biosciences, San Jose, CA, USA), PE-conjugated N418 against CD11c, APC/eFluor780-conjugated BM8 against F4/80 (eBioscience, San Diego, CA, USA), PerCP-conjugated RB6-8C5 against Gr-1 (high affinity Ly-6G, low affinity Ly-6C,^{64,65} Biologend, San Diego, CA, USA). Unconjugated antibody against CNPase (2',3'-cyclic-nucleotide 3'-phosphodiesterase) was purchased from Abcam and conjugated to fluorescein isocyanathionate (FITC) using an EasyLink Conjugation kit (Abcam, Cambridge, MA, USA).

2.8 | mRNA isolation and qRT-PCR analysis

To isolate mRNA, spinal cord tissues and bridge implants were explanted with 2 mm of spinal cord both rostral and caudal to the bridge. Explanted tissues were homogenized using 1 ml of Trizol reagent (Life Technologies) with a tissue grinder. RNA isolation was followed by chloroform extraction and isopropanol precipitation.⁶⁶ The extracted RNA was dissolved in 30 μl of RNase-free distilled water and RNA concentration was measured using a NanoDrop 2000C (ThermoFisher Scientific, Newark, DE, USA) and to assure sufficient purity (A260/A280 ratios between 1.9 and 2.1 for all samples). Total isolated RNA was stored at -80°C freezer until use. cDNA was synthesized using iScript™ cDNA Synthesis kit (Bio-Rad, Hercules, CA, USA) according to the manufacturer's instructions using 1 μg of RNA per sample.

Primers for qRT-PCR (quantitative real-time polymerase chain reaction) quantification of arginase I expression were chosen based on a previous study²⁵: forward 5'-GAACACGGCAGTGGCTTAAAC-3', and reverse 5'-TGCTTAGCTCTGTCTGCTTTGC-3'. 18s-rRNA was used as an internal control with following sequences: forward 5'-GCAATTATCCCATGAACG-3', and reverse 5'-GGCCTCACTAAAC-CATCCAA-3'.⁶⁷ The qRT-PCR products were measured using the accumulation level of iQ™ SYBR Green Supermix (Bio-Rad) fluorescence following a manufacturer's protocol on CFX Connect™ Real-Time PCR Detection System (Bio-Rad). The gene expression level of arginase I mRNA was normalized to that of 18s-rRNA and differences in gene expression were presented as fold ratios from sham (laminectomy only) spinal cord samples. Relative quantification was calculated as $X = 2^{-\Delta\Delta C_t}$, where $\Delta\Delta C_t = \Delta E - \Delta C_t$ and $\Delta E = C_{t,exp} - C_{t,18s-rRNA}$, $\Delta C_t = C_{t,sham} - C_{t,18s-rRNA}$.⁶⁸

2.9 | Behavioral analysis

The Basso mouse scale (BMS) open-field locomotor test (range 0–9) was used to assess functional recovery for a period 24 weeks after SCI as previously described.⁶⁹ A baseline was determined prior to SCI, and mice were tested 3, 7, 14, 21, and 28 days. Observations and BMS scoring were performed by two trained observers at 4-min intervals.

2.10 | Statistical analysis

For multiple comparisons, statistical significance between groups was determined by one-way or two-way ANOVA with Tukey's post hoc testing or Šidák post hoc testing. For single comparisons, the statistical significance between pairs was determined by unpaired t test. For, statistical analysis of macrophage histology, data sets were standardized using an $\ln(x+1)$ transformation to eliminate skewness and normalize to a Gaussian distribution as verified with D'Agostino-Pearson omnibus test.⁷⁰ All statistics test significance using a *p* value of .05 unless otherwise noted. Error bars represent standard error in all figures. Prism 7 (GraphPad Software, La Jolla, CA, USA) software was used for all data analysis.




FIGURE 1 Time course of immune cell infiltration into spinal cord bridges. Infiltrating cells from the central nervous system and peripheral immune system were identified in bridges implanted into a hemisection spinal cord injury ($n = 6-21$ mice) (A) CD45⁺ immune cell, GFAP⁺ astrocyte, and CNPase⁺ oligodendrocyte infiltration into bridges were quantified as a percentage of total cells using flow cytometry at days 7, 14, and 28. (B) Immune sub-populations, including CD11c⁺ dendritic cells, F4/80⁺ macrophages, Gr-1⁺ neutrophils, and CD4⁺ helper T cells, in the bridge were quantified as a percentage of total immune cells using flow cytometry at days 7, 14, and 28. Statistical analysis completed using a one-way ANOVA with Tukey's post hoc test. Significantly different groups denoted by letter with "a" and "b" denoting statistical significance between the two groups ($p < .05$)

3 | RESULTS

3.1 | Cell infiltration into bridges

Cell infiltration into the bridges was initially investigated to identify the cell populations and their abundance at multiple time points. No statistically significant changes in the numbers of CD45⁺ immune cells or GFAP⁺ reactive astrocytes were observed within the bridges between days 7 to 28 (Figure 1A). Additionally, although no significant decrease in CNPase⁺ oligodendrocytes was observed during this time, a trend towards decreasing abundance was seen (Figure 1A). When subpopulations of CD45⁺ immune cells were assessed, a significant increase in the percentage of F4/80⁺ macrophages was observed from day 7 to day 14 (Figure 1B). Numbers of Gr-1⁺ neutrophils (PMNs) and CD11c⁺ dendritic cells (DCs) did not significantly change over time. Finally, CD4⁺ helper T (T_H) cells had a significant increase in abundance from days 14 to 28, indicating that the adaptive immune response is activated at later time points.

3.2 | IL-10 gene delivery from bridges reduces PMN infiltration

The immune response to bridges with and without delivery of IL-10-encoding lentivirus was investigated to determine if the infiltration of inflammatory cells can be modulated through localized cytokine expression. In previous reports, we have demonstrated lentiviral IL-10 expression³⁶ and sustained transgene expression for at least 8 weeks in the injured spinal cord.⁶³ The flow cytometry studies were performed at days 7 and 28, to capture differences from the acute to intermediate phases of recovery observed in Figure 1. First, though differences in infiltrating immune cells (CD45⁺) as a percentage of total cells were not statistically different at day 7, the addition of a bridge to the SCI resulted in a 38% decrease and the addition of lentivirus to the bridge resulted in a 23–34% increase over the bridge alone (Figure 2). This result indicates that the bridge may participate in stabilizing the injury and reducing inflammation at early time points. From day 7 to 28, hemisection alone

(no bridge implantation) and bridges loaded with either lentivirus had a reduction of greater than 50% ($p < .05$) in CD45⁺ cells as a percentage of total cells. Bridges without lentivirus had only a 33% reduction over this same time. At day 28, immune cells as a percentage of total cells were similar for all conditions. Subpopulations of CD45⁺ immune cells were characterized at days 7 and 28 with all bridge conditions having a 2-fold decrease in the percentage of CD11c⁺ antigen-presenting cells (APCs, including dendritic cells and macrophages) relative to no bridge implantation at day 7 (Figure 3A). This reduction did not persist to day 28, at which point APC numbers were similar across all conditions.




FIGURE 2 Bridge implantation and lentivirus delivery do not increase immune cell infiltration. The infiltration of immune cells into the injury site after hemisection spinal cord injury were quantified as a percentage of total cells by flow cytometry at days 7 and 28 ($n = 9-15$ mice). A mock implantation consisting of the hemisection injury without bridge implantation (no bridge) was compared to bridge implantation, luciferase control lentivirus (vCtrl) loaded bridge implantation, and IL-10 lentivirus (vIL-10) loaded bridge implantation. Statistical analysis was completed using a two-way ANOVA with a Šidák correction for multiple comparisons. Significant differences between days 7 and 28 notated with ** ($p < .01$) or * ($p < .05$)




FIGURE 3 IL-10 delivery from bridge reduces neutrophil infiltration. The infiltration of immune cell sub-populations into the injury site was investigated for no bridge, bridge, luciferase control lentivirus (vCtrl) loaded bridges, and IL-10 lentiviral (vIL-10) loaded bridges ($n = 6-21$ mice). Infiltration of sub-populations of innate immune cells were quantified by flow cytometry at day 7 and day 28 including (A) CD11c⁺ dendritic cells, (B) F4/80⁺ macrophages, and (C) Gr-1⁺ neutrophils. Statistical analysis completed using a one-way ANOVA with Tukey's post hoc test. Significantly different groups denoted by letter with "a" and "b" denoting statistical significance between the two groups $p < .05$ for Gr-1 and $p < .005$ for all others

Interestingly, the hemisection-only condition had a dramatic 3.5-fold ($p < .01$) decrease in APC infiltration. The percentage of infiltrating F4/80⁺ macrophages as percent of CD45⁺ cells was reduced approximately 2-fold in the bridge conditions relative to the no bridge control at day 7; however, by day 28, the macrophage percentage increased in the bridge conditions such that they were no longer significantly less than the no bridge condition (Figure 3B). Notably, these changes represent the percentage of CD45⁺ cells that were also F4/80⁺; thus, the increase in macrophage percentage at day 28 does not imply an increase in the total number of macrophages. Finally, delivery of IL-10-encoding lentivirus reduced numbers of infiltrating PMNs by 3.5-fold when compared to the bridges loaded with control lentivirus at Day 7 ($p < .05$) (Figure 3C). Bridges loaded with IL-10-encoding lentivirus had similar numbers of infiltrating PMNs as in hemisections with no bridge implant. Localized IL-10 expression at the bridge did sustain a 3.5-fold reduction of PMN infil-

tration through day 28 (relative to the no bridge control) ($p < .05$) (Figure 3C). Additionally, IL-10 expression reduced the mean level of PMN infiltration by 2.4-fold compared to all other bridges, though this decrease was not statistically significant (Figure 3C).

3.3 | IL-10 gene delivery from bridges increases M2 macrophage infiltration

Immunofluorescence staining of tissue sections was applied to investigate morphological features and distribution of immune cells, which complements the flow cytometry characterization of the relative populations of immune cells infiltrating the bridge.

Immunofluorescently stained sections were initially analyzed to quantify densities of total macrophages (F4/80⁺) and M2 macrophages (F4/80⁺/arginase⁺) (Figure 4). A statistical difference ($p < .05$) was




FIGURE 4 IL-10 delivery increases the density of M2 macrophages. (A) Immunofluorescence (F4/80⁺/arginase⁺/Hoechst⁺: Red/Green/Blue, respectively) was used to characterize the phenotype of infiltrating macrophages within the bridge (scale bar 200 μ m). Line indicates the bridge area used for quantification and dashed line indicates higher magnification regions in (B) F4/80⁺ total macrophages (C) arginase⁺ cells, and (D) merge (scale bar 50 μ m). Brightness and contrast were adjusted for clarity. Arrows note arginase⁻ macrophages (white) and arginase⁺ macrophages (yellow). (E) Quantification of density (cells/mm²) of total F4/80⁺ macrophages as counted manually. (F) Quantification of density (cells/mm²) of F4/80⁺/arginase⁺ M2 macrophages as counted manually. For all quantification, mean \pm SEM was plotted and $n \geq 3$ where each replicate is an average of 1-4 tissues from an individual animal. Statistical analysis was completed on log normalized data using a two-way ANOVA with Tukey's post hoc test with statistical significance between groups indicated by a single asterisk ($p < .05$). Statistical significance of individual conditions between time points indicated by letter with "a" and "b" denoting statistical significance between the two groups ($p < .05$)




FIGURE 5 IL-10 alters macrophage morphology. Immunofluorescence (F4/80⁺/arginase⁺/Hoechst⁺: red/green/blue, respectively), was used to characterize the morphology of infiltrating macrophages within the bridge as either (A) fibrous, (B) round, or (C) FBGC (scale bar 50 μ m). Arrows note arginase⁻ macrophages (white) and arginase⁺ macrophages (yellow). (D) Manual quantification of density (cells/mm²) of fibrous macrophages. (E) Manual quantification of density (cells/mm²) of round macrophages. (F) Manual quantification of density (fused nuclei/mm²) of FBGC macrophages. For all quantification, mean \pm SEM was plotted and $n \geq 3$ where each replicate is an average of 1–4 tissues from an individual animal. Statistical analysis was completed on log-normalized data using a two-way ANOVA with Šidák correction for multiple comparisons with statistical significance indicated by a single asterisk ($p < .05$)

observed in macrophage density between days 14 and 28 for delivery of lentivirus (encoding FLUC or IL-10) from the bridge. In agreement with the flow cytometry results, macrophage number was similar between days 7 and 28 for all conditions (Figure 4E). We subsequently characterized the density of F4/80⁺ cells that also expressed arginase, a marker for the M2 macrophage phenotype. A significant increase in arginase⁺ macrophage density was observed at day 14 in the bridges loaded with IL-10-encoding lentivirus relative to control bridge implants (no lentivirus) (Figure 4F), with a similar trend observed relative to FLUC loaded bridges.

The morphology of the F4/80⁺ cells was subsequently characterized to further represent macrophage phenotype. The F4/80⁺ cells were categorized according three distinct morphologies. The first morphology consisted of fibrous, elongated cells that appeared to have multiple processes (Figure 5A). A second morphology was round and had a prominent F4/80⁺ border, and a majority of these round cells were arginase⁺ (Figure 5B). The third F4/80⁺ morphology consisted of multinucleated foreign body giant cells (FBGCs) (Figure 5C), and nearly every nucleus in the FBGCs was surrounded by arginase⁺ immuno-

staining. Each macrophage was binned according to arginase expression and morphology (fibrous, round, or FBGC). For the fibrous macrophages, the density was largely consistent across conditions and time points, with the singular exception of a 5.5-fold decrease in density between days 14 and day 28 for the control virus condition (Figure 5D). For the round macrophages, no significant differences were observed, though a strong trend ($p = .088$) toward an increase in arginase⁺ round macrophages was identified for IL-10 expression relative to FLUC across all time points (Figure 5E). Finally, for the FBGCs, a greater density of the arginase⁺ cells was observed at Day 14 for the IL-10 conditions relative to the FLUC virus control (Figure 5F).

3.4 | IL-10 delivery results in elevated arginase expression

mRNA levels for arginase were measured by qRT-PCR to complement the arginase⁺ cells density assessment by immunofluorescence. At day 7, qRT-PCR revealed a 5-fold or greater increase in arginase mRNA in the IL-10 group ($p < .001$) (Figure 6). Increased mRNA at day 7 may




FIGURE 6 IL-10 results in elevated arginase mRNA. qRT-PCR revealed elevated arginase levels at 7 and 14 days after SCI. Levels statistically returned to baseline by day 28, but a strong trend remained ($0.08 < p < 0.13$). $N = 4-6$ per group for PCR. Statistical analysis was completed using a one-way ANOVA with Tukey's post hoc test at each time point. Significantly different groups were denoted with letters with "a" and "b" denoting statistical significance between the two groups: day 7 ($p < .001$) and day 14 ($p < .0001$)

correlate to increased protein observed in tissue sections at Day 14 (Figure 4F). At day 14, arginase expression in IL-10 bridges was elevated almost 5-fold compared to bridges loaded with FLUC-encoding lentivirus ($p < .0001$). At day 28, arginase mRNA levels remained more than 2-fold higher for the IL-10 condition, but this difference was not statistically significant ($.08 < p < .13$)

3.5 | Axon regeneration and myelination

Axon numbers and myelination were quantified at day 28 to assess whether IL-10 overexpression would prove detrimental, which has

been reported in the peripheral nervous system.⁷¹ Myelinated (NF200⁺/MBP⁺) and unmyelinated (NF200⁺/MBP⁻) axons were seen throughout the bridges in animals receiving no lentivirus, FLUC-encoding control lentivirus, or IL-10-encoding lentivirus (Figure 7A–D) 28 days after SCI. NF200⁺ axons were typically observed as bundles as previously reported for multichannel PLG bridges.^{54,57,72} Empty bridges had approximately 800 neurites/mm², and both lentiviral conditions had approximately 1100 neurites/mm². While the data suggest a trend toward greater densities of regenerating axons with lentivirus delivery, these differences were not statistically significant. Similarly, the percentage of axons that were myelinated (22–34%) or the percentage of myelination that was derived from Schwann cells (15–35%) did not vary between conditions (Figure 7E).

3.6 | Motor function is improved with IL-10

Motor function in the ipsilateral hindlimb was characterized over the first 28 days after SCI using the BMS (Figure 8). Prior to surgery, all mice had full function, and 3 days post injury, no movement was observed in the ipsilateral hindlimb of any mice. From day 14 onward, mice receiving IL-10 lentivirus had substantially improved function in comparison mice that received a bridge without lentivirus. Mice receiving IL-10 lentivirus received a BMS score of ~4.4, where 4 indicates the capacity to take a step (defined as having weight support at lift off, forward limb advancement, and re-establishment of weight support at initial contact) and 5 indicates that stepping occurs at least 50% of the time. In contrast, without IL-10, mice scored an average of ~3, which only indicates plantar placement of the foot while moving.




FIGURE 7 Myelinated axons 4 weeks after injury. (A) Immunofluorescence of myelinated (NF200⁺/MBP⁺/PO red/green/blue, respectively) and total (NF200⁺: red) axons from bridge implants (scale bar 200 μ m). Dashed line indicates border between bridge implant and host tissue. NF200⁺/MBP⁺ Immunofluorescence at higher magnification from bridges delivering (B) No virus (empty), (C) F-LUC, or (D) IL-10 (scale bar 50 μ m). Brightness and contrast were adjusted for clarity. Arrows note unmyelinated axons (red), oligodendrocyte myelinated axons (yellow), and Schwann cell myelinated axons (blue). (E) Quantification of total axon numbers, total myelinated axons, and Schwann cell myelinated axons. (mean \pm SEM)




FIGURE 8 Functional recovery induced by IL-10 overexpression. The BMS was used to test for differences in motor function in the hindlimb ipsilateral to SCI ($n = 15$ per group). Statistical analysis was completed using a two-way ANOVA with repeated measures and a Sidák correction for multiple comparisons. Significant differences notated with *** ($p < .001$) or **** ($p < .0001$)

4 | DISCUSSION

The studies reported herein employ lentiviral delivery from bridges in order to induce sustained expression of the anti-inflammatory cytokine IL-10 to modulate the inflammatory response. Previous reports of lentiviral delivery from a bridge implanted into the injured spinal cord have demonstrated sustained expression for at least 8 weeks, with maximal expression localized within the bridge and decreased expression both rostral and caudal to the bridge.^{60,63} IL-10 has been reported in some studies to be neuroprotective, reduce secondary inflammation, and reduce lesion volume; however, others have suggested that IL-10 has insignificant effects on spinal cord regeneration. This discrepancy in responses may result from the different models of SCI, or from differences in delivery. Notably, IL-10 does not pass the blood-brain barrier (BBB), and must be delivered either locally or pass through a disrupted BBB following injury. IL-10 also is unstable with a short half-life of 1.1–2.6 hrs, necessitating repeated doses or sustained expression approaches.⁷⁴ Systemic overexpression of IL-10 has been investigated, yet this approach is associated with peripheral neuropathy and demyelination of the sciatic nerve.⁷¹ Systemic delivery of IL-10 protein has had conflicting results: either increasing tissue damage with impaired locomotion or reducing lesion size.^{47,75} IL-10 poliovirus elevated IL-10 for 4 days with modestly improved function, and intrathecal injection of HSV-encoding IL-10 improved axonal sparing and functional recovery.^{37,38} Despite several studies investigating IL-10 delivery for SCI, the effects of sustained localized delivery on dampening and resolving the immune response to promote regeneration have not previously been studied.⁵⁹ and are the focus of the studies herein.

We initially demonstrate that delivery of IL-10-encoding lentivirus can decrease PMN infiltration into biomaterial bridges implanted following SCI, and that though the loading of lentivirus to bridges may promote PMN infiltration, IL-10 over-expression reverses this effect. Although PMNs are transient participants in the immune response fol-

lowing injury to peripheral tissues, they can persist for weeks after injury to the spinal cord.^{12,20} Furthermore, though virus delivery from biomaterials can be used to enhance nerve regeneration,^{54,76} addition of virus to the biomaterial platform can increase the extent of cell infiltration.^{77–79} Secretion of chemokines from the increased immune cells may be responsible for the increase in PMN infiltration at day 7 and 28 reported here (Figure 3C), though increased cytokines would also be expected to simultaneously influence macrophage infiltration, which was not observed and is in agreement with a previous report.⁵⁸ Due to their primarily bactericidal role in wound healing, PMNs are not expected to provide neuroprotection and are primarily detrimental in SCI.¹² PMNs (and macrophages) release cytokines, free radicals, eicosanoids, and proteases, which are toxic to neurons and glia. In particular, superoxide, nitric oxide, and peroxynitrite are highly toxic and create irreversible damage to cellular components, inducing apoptosis.⁸⁰ However, tail vein administration of the RB6-8C5 Ly6G/Gr-1 antibody, which selectively depleted the hematogenous PMN population by more than 90%, resulted in less glial scarring, decreased spared tissue, and worse BMS functional results.⁸¹ Cytokines released by neutrophils may be necessary for sufficient formation of reactive astrocytes to seal the injury and re-establish the BBB. In contrast, reducing infiltration of PMNs ($\downarrow 70\%$) and macrophages ($\downarrow 36\%$) by administration of antibodies against CD11d/CD18 results in less ROS, decreased Caspase 3, and improved functional recovery.^{82–84} These results suggest that though PMNs are essential for the initiation of proper wound healing in the acute injury stage, strategies that mitigate PMN presence in later stages of injury may improve regeneration. Notably, the decrease reported herein regarding the chronic persistence of PMNs, a feature characteristic of mouse and human SCI, has not been previously reported.^{12,20,25} These reports have demonstrated PMNs lasting through 180 days in mice and 12 months in humans.^{20,85} Neutrophils are generally believed to aggravate SCI, though there may be some beneficial function that remains unidentified.²⁰ Taken together, the sustained production of IL-10 through lentivirus transduction alone, or preferably as part of a combinatorial therapy, may prove crucial to reducing the secondary damage and cell death associated with PMN infiltration¹¹ and enhancing regeneration.

Macrophage phenotype, and to some extent number, was influenced by lentiviral delivery of IL-10. Flow cytometry and immunofluorescence staining demonstrated no changes in F4/80⁺ macrophage numbers at day 7 and 28 between conditions. At day 14, quantification revealed a 1.7-fold increase in the number of F4/80⁺ macrophages with expression of IL-10 relative to the control conditions ($p < .05$). These results were expected, as macrophage number is known to peak around Day 7–10 and plateau for several days before decreasing throughout the intermediate and chronic phase.^{12,20} More importantly, IL-10 production altered macrophage polarity with a substantial (3-fold) increase in M2 (F4/80⁺/arginase⁺) macrophage numbers at Day 14 relative to empty bridges ($p < .05$). While the number of arginase⁺ macrophages was not elevated at Day 7 or Day 28, arginase mRNA was upregulated at Day 7 and 14, and trended upward at Day 28. Macrophages can have varying degrees of arginase expression as seen in

Figure 4, which may depend on the specific subtype and phase of regeneration (inflammatory, proliferative, or remodeling).⁴⁴ IL-10 is known for its ability to upregulate expression of IL-4R α , which may synergize with IL-4 dependent arginase expression.⁸⁶ Additionally, alternatively activated (M2) macrophages produce IL-10, potentially creating a feed forward process when IL-10 is overexpressed.

IL-10 expression influenced the relative distribution of three macrophage morphologies: elongated, fibrous cells, round cells with a prominent F4/80 border, and FBGCs. The significance of these morphologies is unclear, and the literature has little information about the relationship between macrophage morphology and phenotype. *In vitro* studies, in particular, have been conflicting. Human macrophages polarized to M1/M2 with LPS/IFN γ and IL-4 respectively that were cultured on tissue culture plastic or within collagen gels exhibited elongated morphologies for M1 macrophages and round, less adherent morphologies for M2 macrophages.⁸⁷ In contrast, when C57BL/6 macrophages are cultured on fibronectin-coated polydimethylsiloxane molds, M1 macrophages have a rounded morphology while M2 macrophages have a fibroblast-like morphology.⁸⁸ Moreover, McWhorter et al. demonstrated that substrates patterned with lines (width 20 μ m) could elongate macrophages leading to upregulated arginase expression.

In vivo characterization of the relationship between macrophage phenotype and morphology has similarly been conflicting. One hypothesis is that the elongated F4/80⁺ cells are activated hematogenous macrophages, while the round cells are activated resident microglia. Microglia are well known to proceed from a highly ramified morphology toward an amoeboid morphology as they become activated and phagocytic after injury.⁸⁹ Unfortunately, antibodies have not been identified to accurately distinguish macrophages from microglia using immunohistochemistry, though future studies using genetic models such as *cr2^{flp}::Cx3cr1^{flp}* mice⁹⁰ have the potential to distinguish cell morphology, phenotype, and lineage. Alternatively, these different morphologies may represent distinct M2 phenotypes. Shechter et al. selectively ablated monocyte-derived macrophages followed by adoptive transfer of monocytes.⁴¹ The exogenous monocytes differentiated into macrophages with an activated morphology, which was manifested by a large cell body with few to no processes, arginase expression, and IL-10 release. These factors suggested an anti-inflammatory phenotype, and the monocytes contributed to regeneration.⁴¹ A subset of infiltrating monocyte-derived, anti-inflammatory macrophages may be essential for recovery, though these cells represent a specific subset of M2 macrophages and will require further characterization.^{91,92}

The round macrophages with prominent F4/80⁺ borders may be undergoing fusogenesis. FBGCs have not been extensively studied in the spinal cord and their impact on spinal cord regeneration is unknown. Fusion has been proposed to begin with the creation of fusion-competent macrophages by IL-4/IL-13 and/or DAP12/TREM-2 interactions between adjacent macrophages,⁹³ followed by CCL2-mediated chemotaxis and E-cadherin-based macrophage adhesion. Despite this simple model, the general requirements for fusion are not fully understood in that it is not clear what role phag-

ocytic capability, macrophage subtype, and material interactions play in their formation.⁹³ Additionally, though IL-4 and IL-13 are typically thought of as cytokines associated with macrophage fusion,⁹³ IL-10 co-localizes with M2 macrophages at sites of macrophage fusion during FBGC development.⁹⁴ Whether IL-10 induces or facilitates fusion or if FBGCs are formed by IL-10-producing macrophages remains unclear.

Although IL-10 did not increase axon numbers in the bridge, IL-10 robustly improved motor function after SCI (Figure 8). IL-10 may act to moderate the deleterious immune response, resulting in improved axon sparing. Increased sparing with increased IL-10 expression is consistent with previous reports, which have demonstrated that IL-10 reduces loss of neurons and oligodendrocytes directly through trophic support³⁸ and indirectly by limiting the immune response.⁵⁹ Additionally, in our past work, differences in numbers of regenerating axons between conditions become more pronounced at later time points.⁵⁴ Thus, improved motor function observed in the current study suggests improved sparing and plasticity instead of regeneration. While these data suggest that IL-10 overexpression from PLG bridges effectively improves function after SCI, delivery of IL-10 in combination with neurotrophic factors known to promote axon regeneration, such as neurotrophin-3 (NT-3),^{54,57} may act synergistically to further improve both sparing and regeneration.

The present study indicates that immobilization of IL-10 lentivirus onto multiple channel bridges alters the immune response to create a microenvironment that is permissive to regeneration and shows promise as a translatable strategy. The bridges used are made of PLG, a biomaterial that has been utilized for decades in FDA-approved applications, including biodegradable sutures and drug delivery vehicles.⁹⁵ PLG is easily sterilized for clinical use using standard techniques, most commonly γ -irradiation.⁹⁶ Though safety has been cited as a concern for lentiviral vectors, there are ongoing clinical trials using lentivirus without adverse effects or evidence of insertional mutagenesis.⁹⁷⁻⁹⁹ Pyrogens from virus production can be limited by purification through gravity-flow columns to yield endotoxin-free, concentrated plasmid, as done in these studies. Furthermore, good manufacturing practices (GMP) have been developed for large-scale preparation of lentivirus for clinical use.^{98,100}

Finally, the hemisection model of SCI, while not perfectly representative of all SCI, is a translatable platform for studying the injury site and developing treatments. While contusive injuries represent the majority of SCI in the developed world, at least 28% of cases in the US military are penetrating,¹⁰¹ and in South Africa, more than 60% of SCI is categorized as penetrating.¹⁰² Moreover, the initial deficits resulting from penetrating SCI exhibit substantially less improvement over time than those from contusive injuries.¹⁰³ For more severe contusions, even if tissue architecture is initially preserved, secondary damage results in cavitation, loss of parenchyma, and glial scarring. Severe contusions and penetrating injuries may require a biomaterial bridge to provide a permissive pathway for regenerating neurons to cross the injury site and reconnect spinal pathways through true axonal regeneration, as opposed to local sprouting.¹⁰⁴

5 | CONCLUSIONS

We report that delivery of IL-10-encoding lentivirus from multiple-channel bridges to reduce the innate immune response following injury. IL-10 overexpression reduces PMN infiltration into the injured spinal cord compared to the delivery of control lentivirus. IL-10 overexpression induced arginase expression in macrophages and substantially altered macrophage morphology. Additionally, IL-10 lentivirus resulted in substantially improved motor function. This result indicates that IL-10 will be an essential component of subsequent combinatorial gene therapies in order to reduce cell death and inhibition.

ACKNOWLEDGMENTS

Advice and verification of statistical methods was provided by The University of Michigan Center for Statistical Consultation and Research (CSCAR). The authors would like to thank Center for Comparative Medicine (CCM) at Northwestern University and Unit for Laboratory Animal Medicine (ULAM) at the University of Michigan for assistance with animal care and health maintenance. Finally, the authors would like to acknowledge contributions by Hannah Tongthaworn and Paula Lahera (UCLA HSSEAS HSSRP).

LITERATURE CITED

- [1] Hurlbert RJ, Hadley MN, Walters BC, et al. Pharmacological therapy for acute spinal cord injury. *Neurosurgery*. 2015;76(suppl 1): S71-83.
- [2] Ahn KS, Aggarwal BB. Transcription factor NF-kappaB: a sensor for smoke and stress signals. *Ann NY Acad Sci*. 2005;1056:218-233.
- [3] Young SH, Ye J, Frazer DG, Shi X, Castranova V. Molecular mechanism of tumor necrosis factor-alpha production in 1->3-beta-glucan (zymosan)-activated macrophages. *J Biol Chem*. 2001; 6(23): 20781-20787.
- [4] Bethea JR, Castro M, Keane RW, Lee TT, Dietrich WD, Yeziarski RP. Traumatic spinal cord injury induces nuclear factor-kappaB activation. *J Neurosci*. 1998;18(9):3251-3260.
- [5] Chen CJ, Kono H, Golenbock D, Reed G, Akira S, Rock KL. Identification of a key pathway required for the sterile inflammatory response triggered by dying cells. *Nat Med*. 2007;13(7):851-856.
- [6] Anderson AJ. Mechanisms and pathways of inflammatory responses in CNS trauma: spinal cord injury. *J Spinal Cord Med*. 2002;25(2):70-79; discussion 80.
- [7] Rhodes KE, Raivich G, Fawcett JW. The injury response of oligodendrocyte precursor cells is induced by platelets, macrophages and inflammation-associated cytokines. *Neuroscience*. 2006;140(1): 87-100.
- [8] Lau LT, Yu AC. Astrocytes produce and release interleukin-1, interleukin-6, tumor necrosis factor alpha and interferon-gamma following traumatic and metabolic injury. *J Neurotrauma*. 2001;18(3): 351-359.
- [9] Fitch MT, Doller C, Combs CK, Landreth GE, Silver J. Cellular and molecular mechanisms of glial scarring and progressive cavitation: in vivo and in vitro analysis of inflammation-induced secondary injury after CNS trauma. *J Neurosci*. 1999;19(19):8182-8198.
- [10] Qin L, Li G, Qian X, et al. Interactive role of the toll-like receptor 4 and reactive oxygen species in LPS-induced microglia activation. *Glia*. 2005;52(1):78-84.
- [11] Pineau I, Lacroix S. Proinflammatory cytokine synthesis in the injured mouse spinal cord: multiphasic expression pattern and identification of the cell types involved. *J Comp Neurol*. 2007;500 (2):267-285.
- [12] Donnelly DJ, Popovich PG. Inflammation and its role in neuroprotection, axonal regeneration and functional recovery after spinal cord injury. *Exp Neurol*. 2008;209(2):378-388.
- [13] Filbin MT. Myelin-associated inhibitors of axonal regeneration in the adult mammalian CNS. *Nat Rev Neurosci*. 2003;4(9):703-713.
- [14] Yiu G, He Z. Glial inhibition of CNS axon regeneration. *Nat Rev Neurosci*. 2006;7(8):617-627.
- [15] Fitch MT, Silver J. CNS injury, glial scars, and inflammation: inhibitory extracellular matrices and regeneration failure. *Exp Neurol*. 2008;209(2):294-301.
- [16] Tom VJ, Steinmetz MP, Miller JH, Doller CM, Silver J. Studies on the development and behavior of the dystrophic growth cone, the hallmark of regeneration failure, in an in vitro model of the glial scar and after spinal cord injury. *J Neurosci*. 2004;24(29):6531-6539.
- [17] Schwab ME, Bartholdi D. Degeneration and regeneration of axons in the lesioned spinal cord. *Physiol Rev*. 1996;76(2):319-370.
- [18] Lang BT, Cregg JM, DePaul MA, et al. Modulation of the proteoglycan receptor PTPsigma promotes recovery after spinal cord injury. *Nature*. 2015;518(7539):404-408.
- [19] Busch SA, Silver J. The role of extracellular matrix in CNS regeneration. *Curr Opin Neurobiol*. 2007;17(1):120-127.
- [20] Beck KD, Nguyen HX, Galvan MD, Salazar DL, Woodruff TM, Anderson AJ. Quantitative analysis of cellular inflammation after traumatic spinal cord injury: evidence for a multiphasic inflammatory response in the acute to chronic environment. *Brain*. 2010; 133(Pt 2):433-447.
- [21] Waxman SG. Demyelination in spinal cord injury. *J Neurol Sci*. 1989;91(1-2):1-14.
- [22] Mosser DM. The many faces of macrophage activation. *J Leukoc Biol*. 2003;73(2):209-212.
- [23] Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nat Rev Immunol*. 2011;11(11):723-737.
- [24] Sindrilaru A, Peters T, Wieschalka S, et al. An unrestrained proinflammatory M1 macrophage population induced by iron impairs wound healing in humans and mice. *J Clin Invest*. 2011;121(3):985-997.
- [25] Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, Popovich PG. Identification of two distinct macrophage subsets with divergent effects causing either neurotoxicity or regeneration in the injured mouse spinal cord. *J Neurosci*. 2009;29(43):13435-13444.
- [26] Pruss H, Kopp MA, Brommer B, et al. Non-resolving aspects of acute inflammation after spinal cord injury (SCI): indices and resolution plateau. *Brain Pathol*. 2011;21(6):652-660.
- [27] Kitayama M, Ueno M, Itakura T, Yamashita T. Activated microglia inhibit axonal growth through RGMa. *PLoS One*. 2011;6(9):e25234.
- [28] Busch SA, Horn KP, Silver DJ, Silver J. Overcoming macrophage-mediated axonal dieback following CNS injury. *J Neurosci*. 12 2009;29(32):9967-9976.
- [29] Horn KP, Busch SA, Hawthorne AL, van Rooijen N, Silver J. Another barrier to regeneration in the CNS: activated macrophages induce extensive retraction of dystrophic axons through direct physical interactions. *J Neurosci*. 2008;28(38):9330-9341.
- [30] Popovich PG, Guan Z, McLaughly V, Fisher L, Hickey WF, Basso DM. The neuropathological and behavioral consequences of intraspinal microglial/macrophage activation. *J Neuropathol Exp Neurol*. 2002;61(7):623-633.

- [31] Shakhbazou A, Mishra M, Chu TH, et al. Fluorescent phosphorus dendrimer as a spectral nanosensor for macrophage polarization and fate tracking in spinal cord injury. *Macromol Biosci*. 2015;15(11):1523–1534.
- [32] Porcheray F, Viaud S, Rimaniol AC, et al. Macrophage activation switching: an asset for the resolution of inflammation. *Clin Exp Immunol*. 2005;142(3):481–489.
- [33] Ogden CA, Pound JD, Bath BK, et al. Enhanced apoptotic cell clearance capacity and B cell survival factor production by IL-10-activated macrophages: implications for Burkitt's lymphoma. *J Immunol*. 2005;174(5):3015–3023.
- [34] Galli SJ, Borregaard N, Wynn TA. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat Immunol*. 2011;12(11):1035–1044.
- [35] Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci*. 2007;8(1):57–69.
- [36] Boehler RM, Kuo R, Shin S, et al. Lentivirus delivery of IL-10 to promote and sustain macrophage polarization towards an anti-inflammatory phenotype. *Biotechnol Bioeng*. 2014;111(6):1210–1221.
- [37] Zhou Z, Peng X, Insolera R, Fink DJ, Mata M. IL-10 promotes neuronal survival following spinal cord injury. *Exp Neurol*. 2009;220(1):183–190.
- [38] Zhou Z, Peng X, Insolera R, Fink DJ, Mata M. Interleukin-10 provides direct trophic support to neurons. *J Neurochem*. 2009;110(5):1617–1627.
- [39] Popovich PG, Guan Z, Wei P, Huitinga I, van Rooijen N, Stokes BT. Depletion of hematogenous macrophages promotes partial hindlimb recovery and neuroanatomical repair after experimental spinal cord injury. *Exp Neurol*. 1999;158(2):351–365.
- [40] Zhu Y, Soderblom C, Krishnan V, Ashbaugh J, Bethea JR, Lee JK. Hematogenous macrophage depletion reduces the fibrotic scar and increases axonal growth after spinal cord injury. *Neurobiol Dis*. 2015;74:114–125.
- [41] Shechter R, London A, Varol C, et al. Infiltrating blood-derived macrophages are vital cells playing an anti-inflammatory role in recovery from spinal cord injury in mice. *PLoS Med*. 2009;6(7):e1000113.
- [42] Murray PJ, Wynn TA. Obstacles and opportunities for understanding macrophage polarization. *J Leukoc Biol*. 2011;89(4):557–563.
- [43] Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep*. 2014;6:13.
- [44] Gensel JC, Zhang B. Macrophage activation and its role in repair and pathology after spinal cord injury. *Brain Res*. 2015;1619:1–11.
- [45] Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol*. 2008;8(12):958–969.
- [46] Weaver LC, Gris D, Saville LR, et al. Methylprednisolone causes minimal improvement after spinal cord injury in rats, contrasting with benefits of an anti-integrin treatment. *J Neurotrauma*. 2005;22(12):1375–1387.
- [47] Takami T, Oudega M, Bethea JR, Wood PM, Kleitman N, Bunge MB. Methylprednisolone and interleukin-10 reduce gray matter damage in the contused Fischer rat thoracic spinal cord but do not improve functional outcome. *J Neurotrauma*. 2002;19(5):653–666.
- [48] Richardson PM, McGuinness UM, Aguayo AJ. Axons from CNS neurons regenerate into PNS grafts. *Nature*. 1980;284(5753):264–265.
- [49] Yang Y, De Laporte L, Rives CB, et al. Neurotrophin releasing single and multiple lumen nerve conduits. *J Control Release*. 2005;104(3):433–446.
- [50] He L, Zhang Y, Zeng C, et al. Manufacture of PLGA multiple-channel conduits with precise hierarchical pore architectures and in vitro/vivo evaluation for spinal cord injury. *Tissue Eng Part C Methods*. 2009;15(2):243–255.
- [51] de Ruyter GC, Onyeneho IA, Liang ET, et al. Methods for in vitro characterization of multichannel nerve tubes. *J Biomed Mater Res A*. 2008;84(3):643–651.
- [52] Krych AJ, Rooney GE, Chen B, et al. Relationship between scaffold channel diameter and number of regenerating axons in the transected rat spinal cord. *Acta Biomater*. 2009;5(7):2551–2559.
- [53] Chen BK, Knight AM, Madigan NN, et al. Comparison of polymer scaffolds in rat spinal cord: a step toward quantitative assessment of combinatorial approaches to spinal cord repair. *Biomaterials*. 2011;32(32):8077–8086.
- [54] Tuinstra HM, Aviles MO, Shin S, et al. Multifunctional, multichannel bridges that deliver neurotrophin encoding lentivirus for regeneration following spinal cord injury. *Biomaterials*. 2012;33(5):1618–1626.
- [55] Tuinstra HM, Margul DJ, Goodman AG, et al. Long-term characterization of axon regeneration and matrix changes using multiple channel bridges for spinal cord regeneration. *Tissue Eng Part A*. 2014;20(5-6):1027–1037.
- [56] Pawar K, Cummings BJ, Thomas A, et al. Biomaterial bridges enable regeneration and re-entry of corticospinal tract axons into the caudal spinal cord after SCI: association with recovery of forelimb function. *Biomaterials*. 2015;65:1–12.
- [57] Thomas AM, Seidlits SK, Goodman AG, et al. Sonic hedgehog and neurotrophin-3 increase oligodendrocyte numbers and myelination after spinal cord injury. *Integr Biol (Camb)*. 2014;6(7):694–705.
- [58] Abdellatif AA, Pelt JL, Benton RL, et al. Gene delivery to the spinal cord: comparison between lentiviral, adenoviral, and retroviral vector delivery systems. *J Neurosci Res*. 2006;84(3):553–567.
- [59] Thompson CD, Zurko JC, Hanna BF, Hellenbrand DJ, Hanna A. The therapeutic role of interleukin-10 after spinal cord injury. *J Neurotrauma*. 2013;30(15):1311–1324.
- [60] Dull T, Zufferey R, Kelly M, et al. A third-generation lentivirus vector with a conditional packaging system. *J Virol*. 1998;72(11):8463–8471.
- [61] Li J, Rickett TA, Shi R. Biomimetic nerve scaffolds with aligned intraluminal microchannels: a “sweet” approach to tissue engineering. *Langmuir: the ACS journal of surfaces and colloids*. 2009;25(3):1813–1817.
- [62] Thomas AM, Kubilius MB, Holland SJ, et al. Channel density and porosity of degradable bridging scaffolds on axon growth after spinal injury. *Biomaterials*. 2013;34(9):2213–2220.
- [63] Thomas AM, Shea LD. Polysaccharide-modified scaffolds for controlled lentivirus delivery in vitro and after spinal cord injury. *J Control Release*. 2013;170(3):421–429.
- [64] Fleming T, Fleming M, Malek T. Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. *J Immunol*. 1993;151(5):2399–2408.
- [65] Daley JM, Thomay AA, Connolly MD, Reichner JS, Albina JE. Use of Ly6G-specific monoclonal antibody to deplete neutrophils in mice. *J Leukoc Biol*. 2008;83(1):64–70.
- [66] Due MR, Park J, Zheng L, et al. Acrolein involvement in sensory and behavioral hypersensitivity following spinal cord injury in the rat. *J Neurochem*. 2014;128(5):776–786.
- [67] Uchida S, Hara K, Kobayashi A, et al. Early life stress enhances behavioral vulnerability to stress through the activation of REST4-mediated gene transcription in the medial prefrontal cortex of rodents. *J Neurosci*. 2010;30(45):15007–15018.

- [68] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)⁻(Delta Delta C) method. *Methods*. 2001;25(4):402–408.
- [69] Basso DM, Fisher LC, Anderson AJ, Jakeman LB, McTigue DM, Popovich PG. Basso Mouse Scale for locomotion detects differences in recovery after spinal cord injury in five common mouse strains. *J Neurotrauma*. 2006;23(5):635–659.
- [70] Tabachnick BG, Fidell LS. *Using Multivariate Statistics*. 6th ed. Boston: Pearson Education; 2013.
- [71] Dace DS, Khan AA, Stark JL, Kelly J, Cross AH, Apte RS. Interleukin-10 overexpression promotes Fas-ligand-dependent chronic macrophage-mediated demyelinating polyneuropathy. *PLoS One*. 2009;4(9):e7121.
- [72] McCreedy DA, Margul DJ, Seidlits SK, et al. Semi-automated counting of axon regeneration in poly(lactide co-glycolide) spinal cord bridges. *J Neurosci Methods*. 2016;263:15–22.
- [73] Shin S, Tuinstra HM, Salvay DM, Shea LD. Phosphatidylserine immobilization of lentivirus for localized gene transfer. *Biomaterials*. 2010;31(15):4353–4359.
- [74] Mueller C, Braag SA, Martino AT, Tang Q, Campbell-Thompson M, Flotte TR. The pros and cons of immunomodulatory IL-10 gene therapy with recombinant AAV in a Cfr^{-/-} dependent allergy mouse model. *Gene Ther*. 2009;16(2):172–183.
- [75] Pearse DD, Marcillo AE, Oudega M, Lynch MP, Wood PM, Bunge MB. Transplantation of Schwann cells and olfactory ensheathing glia after spinal cord injury: does pretreatment with methylprednisolone and interleukin-10 enhance recovery? *J Neurotrauma*. 2004;21(9):1223–1239.
- [76] Hendriks WT, Ruitenber MJ, Blits B, Boer GJ, Verhaagen J. Viral vector-mediated gene transfer of neurotrophins to promote regeneration of the injured spinal cord. *Prog Brain Res*. 2004;146:451–476.
- [77] Sakurai H, Sakurai F, Kawabata K, et al. Comparison of gene expression efficiency and innate immune response induced by Ad vector and lipoplex. *J Control Release*. 2007;117(3):430–437.
- [78] Sakurai H, Kawabata K, Sakurai F, Nakagawa S, Mizuguchi H. Innate immune response induced by gene delivery vectors. *Int J Pharm*. 2008;354(1-2):9–15.
- [79] Liu Q, Muruve DA. Molecular basis of the inflammatory response to adenovirus vectors. *Gene Ther*. 2003;10(11):935–940.
- [80] Xiong Y, Rabchevsky AG, Hall ED. Role of peroxynitrite in secondary oxidative damage after spinal cord injury. *J Neurochem*. 2007;100(3):639–649.
- [81] Stirling DP, Liu S, Kubers P, Yong VW. Depletion of Ly6G/Gr-1 leukocytes after spinal cord injury in mice alters wound healing and worsens neurological outcome. *J Neurosci*. 2009;29(3):753–764.
- [82] Bao F, Dekaban GA, Weaver LC. Anti-CD11d antibody treatment reduces free radical formation and cell death in the injured spinal cord of rats. *J Neurochem*. 2005;94(5):1361–1373.
- [83] Saville LR, Pospisil CH, Mawhinney LA, et al. A monoclonal antibody to CD11d reduces the inflammatory infiltrate into the injured spinal cord: a potential neuroprotective treatment. *J Neuroimmunol*. 2004;156(1-2):42–57.
- [84] Geremia NM, Bao F, Rosenzweig TE, et al. CD11d Antibody Treatment Improves Recovery in Spinal Cord-Injured Mice. *J Neurotrauma*. 2012;29(3):539–550.
- [85] Fleming JC, Norenberg MD, Ramsay DA, et al. The cellular inflammatory response in human spinal cords after injury. *Brain*. 2006;129(Pt 12):3249–3269.
- [86] Lang R, Patel D, Morris JJ, Rutschman RL, Murray PJ. Shaping gene expression in activated and resting primary macrophages by IL-10. *J Immunol*. 2002;169(5):2253–2263.
- [87] Zajac E, Schweighofer B, Kupriyanova TA, et al. Angiogenic capacity of M1- and M2-polarized macrophages is determined by the levels of TIMP-1 complexed with their secreted proMMP-9. *Blood*. 2013;122(25):4054–4067.
- [88] McWhorter FY, Wang T, Nguyen P, Chung T, Liu WF. Modulation of macrophage phenotype by cell shape. *Proc Natl Acad Sci USA*. Oct 22 2013;110(43):17253–17258.
- [89] Kettenmann H, Ransom BR. *Neuroglia*. 2nd ed. New York: Oxford University Press; 2005.
- [90] Yamasaki R, Lu H, Butovsky O, et al. Differential roles of microglia and monocytes in the inflamed central nervous system. *J Exp Med*. 2014;211(8):1533–1549.
- [91] Jung S, Schwartz M. Non-identical twins - microglia and monocyte-derived macrophages in acute injury and autoimmune inflammation. *Front Immunol*. 2012;3:89.
- [92] London A, Cohen M, Schwartz M. Microglia and monocyte-derived macrophages: functionally distinct populations that act in concert in CNS plasticity and repair. *Front Cell Neurosci*. 2013;7:34.
- [93] Helming L, Gordon S. Molecular mediators of macrophage fusion. *Trends Cell Biol*. 2009;19(10):514–522.
- [94] Higgins DM, Basaraba RJ, Hohnbaum AC, Lee EJ, Grainger DW, Gonzalez-Juarrero M. Localized immunosuppressive environment in the foreign body response to implanted biomaterials. *Am J Pathol*. 2009;175(1):161–170.
- [95] Whittlesey KJ, Shea LD. Delivery systems for small molecule drugs, proteins, and DNA: the neuroscience/biomaterial interface. *Exp Neurol*. 2004;190(1):1–16.
- [96] Abuhanoglu G, Ozer AY. Radiation sterilization of new drug delivery systems. *Interv Med Appl Sci*. 2014;6(2):51–60.
- [97] Palfi S, Gurruchaga JM, Ralph GS, et al. Long-term safety and tolerability of ProSavin, a lentiviral vector-based gene therapy for Parkinson's disease: a dose escalation, open-label, phase 1/2 trial. *Lancet*. 29 2014;383(9923):1138–1146.
- [98] Biffi A, Montini E, Lorioli L, et al. Lentiviral hematopoietic stem cell gene therapy benefits metachromatic leukodystrophy. *Science*. 2013;341(6148):1233158.
- [99] Hacein-Bey Abina S, Gaspar HB, Blondeau J, et al. Outcomes following gene therapy in patients with severe Wiskott-Aldrich syndrome. *JAMA*. 21 2015;313(15):1550–1563.
- [100] Merten OW, Charrier S, Laroudie N, et al. Large-scale manufacture and characterization of a lentiviral vector produced for clinical ex vivo gene therapy application. *Hum Gene Ther*. 2011;22(3):343–356.
- [101] Blair JA, Possley DR, Petfield JL, et al. Military penetrating spine injuries compared with blunt. *Spine J*. 2012;12(9):762–768.
- [102] Velmahos GC, Degiannis E, Hart K, Souter I, Saadia R. Changing profiles in spinal cord injuries and risk factors influencing recovery after penetrating injuries. *J Trauma*. 1995;38(3):334–337.
- [103] Rhee P, Kuncir EJ, Johnson L, et al. Cervical spine injury is highly dependent on the mechanism of injury following blunt and penetrating assault. *J Trauma*. 2006;61(5):1166–1170.
- [104] Straley KS, Foo CW, Heilshorn SC. Biomaterial design strategies for the treatment of spinal cord injuries. *J Neurotrauma*. 2010;27(1):1–19.