



Published in final edited form as:

*Diabetologia*. 2016 October ; 59(10): 2181–2192. doi:10.1007/s00125-016-4053-x.

## Oral AGE restriction ameliorates insulin resistance in obese individuals with the metabolic syndrome: a randomised controlled trial

Helen Vlassara<sup>1,2</sup>, Weijing Cai<sup>2</sup>, Elizabeth Tripp<sup>2</sup>, Renata Pyzik<sup>2</sup>, Kalle Yee<sup>2</sup>, Laurie Goldberg<sup>2</sup>, Laurie Tansman<sup>2</sup>, Xue Chen<sup>2</sup>, Venkatesh Mani<sup>3</sup>, Zahi A. Fayad<sup>3</sup>, Girish Nadkarni<sup>1</sup>, Gary E. Striker<sup>1,2</sup>, John C. He<sup>1</sup>, and Jaime Uribarri<sup>1</sup>

<sup>1</sup>Department of Medicine, Division of Nephrology, The Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1147, New York, NY 10021, USA

<sup>2</sup>Department of Geriatrics, Division of Experimental Diabetes, The Icahn School of Medicine at Mount Sinai, New York, NY, USA

<sup>3</sup>Translational and Molecular Imaging Institute, The Icahn School of Medicine at Mount Sinai, New York, NY, USA

### Abstract

**Aims/hypothesis**—We previously reported that obese individuals with the metabolic syndrome (at risk), compared with obese individuals without the metabolic syndrome (healthy obese), have elevated serum AGEs that strongly correlate with insulin resistance, oxidative stress and inflammation. We hypothesised that a diet low in AGEs (L-AGE) would improve components of the metabolic syndrome in obese individuals, confirming high AGEs as a new risk factor for the metabolic syndrome.

**Methods**—A randomised 1 year trial was conducted in obese individuals with the metabolic syndrome in two parallel groups: L-AGE diet vs a regular diet, habitually high in AGEs (Reg-AGE). Participants were allocated to each group by randomisation using random permuted blocks.

---

Corresponding author: Jaime Uribarri, Department of Medicine, Division of Nephrology, The Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1147, New York, NY 10021, USA, Jaime.uribarri@mssm.edu.

**Duality of interest:** The authors declare that there is no duality of interest associated with this manuscript.

**Contribution statement:** HV designed the study, interpreted the data, and wrote, reviewed and edited the manuscript. WC participated in the experimental design, the acquisition and interpretation of data, and reviewed the manuscript. ET, the study dietitian, participated in the design of the dietary protocol and the interpretation of nutritional data, and reviewed the manuscript. RP, the study coordinator, participated in the design of the clinical protocol and the acquisition and interpretation of clinical data, and reviewed the manuscript. KY, the study coordinator, participated in the design of the clinical protocol and the acquisition of clinical data, and reviewed manuscript. LG, the study dietitian, participated in the design of the dietary protocol and the acquisition of nutritional data, and reviewed and edited the manuscript. LT, a study dietitian, participated in the design of the dietary protocol and the acquisition and interpretation of nutritional data, and reviewed and edited the manuscript. XC participated in the experimental design and the acquisition of data, and reviewed and edited the manuscript. VM participated in the study design and performance and in the interpretation of data from MRI studies, and edited the final manuscript. ZAF participated in the original study design and performance and in the interpretation of data from MRI studies, and edited the final manuscript. GN participated in the study design and in all the statistical analyses, and reviewed and edited the manuscript. GES participated in the study design and interpretation of data, and reviewed and edited the manuscript. JCH participated in the study design and interpretation of data and reviewed the final manuscript. JU participated in the design of the study, the acquisition and interpretation of data and in the statistical analyses, and reviewed and edited the manuscript. JU and HV are the guarantors of this work. All authors have revised the final updated version of the manuscript.

At baseline and at the end of the trial, we obtained anthropometric variables, blood and urine samples, and performed OGTTs and MRI measurements of visceral and subcutaneous abdominal tissue and carotid artery. Only investigators involved in laboratory determinations were blinded to dietary assignment. Effects on insulin resistance (HOMA-IR) were the primary outcome.

**Results**—Sixty-one individuals were randomised to a Reg-AGE diet and 77 to an L-AGE diet; the data of 49 and 51, respectively, were analysed at the study end in 2014. The L-AGE diet markedly improved insulin resistance; modestly decreased body weight; lowered AGEs, oxidative stress and inflammation; and enhanced the protective factors sirtuin 1, AGE receptor 1 and glyoxalase I. The Reg-AGE diet raised AGEs and markers of insulin resistance, oxidative stress and inflammation. There were no effects on MRI-assessed measurements; no side effects from the intervention were identified. HOMA-IR came down from  $3.1 \pm 1.8$  to  $1.9 \pm 1.3$  ( $p < 0.001$ ) in the L-AGE group, while it increased from  $2.9 \pm 1.2$  to  $3.6 \pm 1.7$  ( $p < 0.002$ ) in the Reg-AGE group.

**Conclusions/interpretation**—L-AGE ameliorates insulin resistance in obese people with the metabolic syndrome, and may reduce the risk of type 2 diabetes, without necessitating a major reduction in adiposity. Elevated serum AGEs may be used to diagnose and treat ‘at-risk’ obesity.

**Trial registration**—ClinicalTrials.gov NCT01363141

**Funding**—The study was funded by the National Institute of Diabetes and Digestive and Kidney Diseases (DK091231)

### Keywords

AGER1; Cardiovascular disease; Diabetes; Glycotoxins; Inflammation; Innate defence; RAGE; SIRT1

## Introduction

Current efforts to arrest the epidemics of type 2 diabetes and cardiovascular disease, as well as dementia or Alzheimer's disease, have had limited success [1-3]. Obesity is a prevalent component of type 2 diabetes, often coupled with insulin resistance, hypertension and dyslipidaemia in a cluster known as the metabolic syndrome, associated with chronic oxidative stress and inflammation [4-6]. Overnutrition is thought to be a major cause of obesity [5]. However, many obese people do not develop the metabolic syndrome and have been labelled ‘healthy obese’ [7-9]. Although excessive consumption of fatty foods is considered an important cause of the metabolic syndrome [4, 5], this view has recently come under new scrutiny [10, 11]. Thus, there is a need to identify alternative causes and treatments for ‘at-risk’ obese people.

Thermally prepared foods are rich in pro-inflammatory AGEs, which are also flavourful, a fact that enhances palatability and consumption and therefore may promote weight gain [12, 13]. Excessive intake of dietary AGEs has directly and causally been linked in both humans and mice to high serum AGEs, oxidative stress and inflammation, reduced innate defences, and insulin resistance [12-19]. In a cross-sectional study of obese individuals with and without the metabolic syndrome, we recently found a direct correlation between AGEs and insulin resistance. Obese individuals with the metabolic syndrome had significantly higher

dietary AGE intake and serum AGEs, as well as insulin resistance, oxidative stress and inflammation, compared with healthy or obese individuals without the metabolic syndrome [20]. Two distinct serum AGE markers, carboxymethyllysine (CML) and methylglyoxal (MG) derivatives, were markedly elevated in this population and strongly correlated with elevated HOMA-IR, oxidative stress and inflammation, and decreased protective or host defence factors, such as sirtuin 1 (SIRT1) and AGE receptor 1 (AGER1) [20].

Based on these observations, we hypothesised that high serum AGE levels in at-risk obese adults contribute to the risk of type 2 diabetes. We proposed that a 1 year randomised interventional trial with dietary AGE restriction (L-AGE) would improve metabolic syndrome risk factors.

## Methods

### Clinical protocol

Selection criteria to participate in the study have already been published [20]. Briefly, volunteers aged 50 years or above with at least two of the following five criteria of the metabolic syndrome, based on the Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) (NCEP ATP III) [21], were recruited from the New York City urban community surrounding the Icahn School of Medicine at Mount Sinai: (1) waist circumference  $\geq 102$  cm in men and  $\geq 88$  cm in women; (2) BP  $\geq 130/85$  mmHg (or use of BP-lowering medication); (3) HDL-cholesterol  $<1.04$  mmol/l in men or  $<1.29$  mmol/l in women; (4) triacylglycerol  $\geq 1.69$  mmol/l (or use of medication for high triacylglycerol, such as fibrates or nicotinic acid); and (5) fasting blood glucose  $\geq 5.55$  mmol/l, but an HbA<sub>1c</sub>  $<6.5\%$ / $47.5$  mmol/mol, or use of metformin. Volunteers were screened with a 3 day AGE food record and those whose daily intake was  $\geq 12$  AGE equivalents per day were invited to participate in the study. These participants were then randomised either to an L-AGE diet or to their usual diet (Reg-AGE) and used as controls for the next 12 months (Fig. 1). Routine blood tests were performed in the hospital clinical laboratory. Only investigators involved in laboratory determinations were blinded to the dietary assignment. Participant recruitment started in 2011 and the trial finished in 2014.

All volunteers signed a consent form approved by the Icahn School of Medicine at Mount Sinai Institutional Review Board. The study was registered in [www.clinicaltrials.gov](http://www.clinicaltrials.gov) (NCT01363141).

L-AGE participants prepared their own food at home after being individually instructed on how to reduce dietary AGE intake by modifying the cooking time and temperature without changing the quantity, quality or composition of food. They were specifically instructed to avoid frying, baking or grilling, and they were encouraged to prepare their food by boiling, poaching, stewing or steaming (Table 1). It has been previously demonstrated that switching to these suggested methods of cooking limits new AGE formation in foods, especially animal food products [15]. Participants received a personal interview with the study dietitian every 3 months at the clinic to emphasise instructions and review records. In addition, a dietitian contacted them regularly via telephone (twice/week) to evaluate and promote

dietary compliance. Together these measures helped avoid undue changes in calorie consumption while adhering to L-AGE.

All participants underwent a physical examination and provided a medical history at baseline. A fasting blood sample, a 24 h urine sample and previously validated 3 day food records were obtained at baseline and at 12 months (end of study). During these visits, participants also underwent an OGTT (75 g oral glucose load, followed by serum samples at 0, 60 and 120 min) and abdominal and neck MRI to define subcutaneous and visceral adipose fat distribution and carotid wall artery variables. Routine blood tests were performed in the hospital clinical laboratory. HOMA-IR was calculated from fasting blood glucose and insulin as previously published [22].

**Dietary intake**—Assessment of daily dietary AGE content was based on 3 day food records and estimated from a database of ~560 foods which lists AGE values [15] expressed as AGE equivalents per day (1 AGE equivalent=1,000 kilounits). The 3 day food record is based on established guidelines developed to assist in estimating portions [15]. Nutrient intakes, important in monitoring and preventing undue changes in calorie consumption, were then estimated from food records using a nutrient software program (Food Processor, version 10.1; ESHA Research, Salem, OR, USA).

**Imaging studies (MRI)**—Subcutaneous and visceral abdominal fat deposits were assessed as previously described [23, 24] (electronic supplementary material [ESM] Methods).

**Materials**—See ESM Methods.

**AGE determination**—AGEs (CML and MG) in serum, urine and peripheral blood mononuclear cell (PMNC) lysates were determined by well-validated, competitive ELISAs based on monoclonal antibodies for protein-bound CML (4G9) [25, 26] and protein-bound MG derivatives, i.e. arginine-MG-H1, characterised by HPLC [12], shown to detect pathologically relevant AGEs in multiple studies [12-15, 27] (ESM Methods).

**RNA isolation and qRT-PCR**—Total RNA was isolated from PMNCs using TRIzol reagent according to the manufacturer's protocol (Sigma-Aldrich, St Louis, MO, USA). First-strand cDNA synthesis was performed using SuperScript III Reverse Transcriptase (Roche, Indianapolis, IN, USA). *AGER1* (also known as *DDOST*), receptor for AGEs (*RAGE*, also known as *AGER*) and *SIRT1* mRNA expression were analysed by quantitative SYBR Green real-time PCR. The transcript copy number of target genes was determined based on their  $C_t$  values [17, 20, 27] (ESM Methods).

## Ex vivo studies

**Cell culture**—PMNCs from individuals with the metabolic syndrome were separated from fasting, EDTA anticoagulated blood by Ficoll-Hypaque Plus gradient (GE Healthcare Bioscience, Pittsburgh, PA, USA) and incubated in serum-free culture media AIM-V (Invitrogen, Carlsbad, CA, USA) at 37°C with 5% CO<sub>2</sub> for 1 h, followed by the addition of MG-BSA (60 µg/ml) [12, 14, 16]. MG-BSA contained 22 MG-modified arginine residues per mole based on HPLC. MG-BSA was passed through an endotoxin-binding affinity

column (Pierce, Rockford, IL, USA) to remove endotoxins [12, 14] (ESM Methods). Cells were incubated with or without the SIRT1 inhibitor Sirtinol (10  $\mu\text{mol/l}$ ; Calbiochem, La Jolla, CA, USA) or a SIRT1 activator, SRT1720 (1  $\mu\text{mol/l}$ ; Selleckchem, Houston, TX, USA). After 72 h, the cells were harvested for western blotting, the culture medium was centrifuged at 1000  $g$  for 10 min and the supernatant fraction was collected for testing human TNF $\alpha$  levels with an ELISA kit (Invitrogen) [28]. In a separate study, PMNCs from normal volunteers without the metabolic syndrome, described previously [20], were used as controls for PMNCs from individuals with the metabolic syndrome. All baseline cellular data were collected from PMNCs obtained at study commencement.

**Conditioned medium**—PMNCs freshly isolated at baseline and at study end were plated in serum-free culture media at 37°C with 5% CO<sub>2</sub> for 24 h. The media were collected and centrifuged to remove cells and other particles and concentrated by Amicon Ultra centrifugal filter units (Sigma-Aldrich) for co-culture experiments [14].

**Adipocyte culture and treatment**—3T3-L1 cells (ATCC, Manassas, VA, USA), cultured and differentiated, as described [12], were incubated with DMEM or PMNC conditioned medium and diluted at 1:500 with DMEM for 18-24 h. After washing and replacing media with Hanks' Balanced Salt Solution for 1 h, cells were stimulated with insulin (100 nmol/l) for 30 min before they were harvested.

**Western blotting**—After incubating PMNCs with MG-BSA and with or without Sirtinol or SRT1720 for 72 h, cells were disrupted in lysis buffer. The cellular proteins were separated on 8% SDS-PAGE gels, transferred on to polyvinylidene fluoride membranes and immunoblotted with the indicated antibodies. To test for NF $\kappa$ B acetylation, 250  $\mu\text{g}$  protein lysate was immunoprecipitated by an anti-NF $\kappa$ B p65 antibody at 4°C overnight. Protein A/G agarose beads 60  $\mu\text{l}$  were added. The immunoprecipitates were immunoblotted with a specific anti-acetyl-NF $\kappa$ B p65 (lysine 310) antibody, 3T3-L1 adipocyte lysates (50  $\mu\text{g}$ ) were immunoblotted by anti-phospho-Akt (Ser473) antibody or immunoprecipitated (400  $\mu\text{g}$ ) by anti-insulin receptor- $\beta$  antibody, followed by immunoblotting with anti-phosphotyrosine antibody (4G10; Millipore, Billerica, MA, USA).

## Statistical analysis

Of the 383 individuals originally screened for eligibility, 138 were randomised (Fig. 1). Participants were considered eligible for analysis if they attended at least the 6 month clinic appointment. The pre-specified statistical analysis plan defined the study outcomes to be the mean differences between the final recorded values minus the baseline values for the Reg-AGE and L-AGE groups. The primary outcome variable was taken to be HOMA-IR. Sample size was established on the basis of prior studies showing an effect of L-AGE on healthy individuals who had a difference in HOMA-IR between those on L-AGE and those on normal diets of 4.57 [17]. To find a difference of 4 with 80% power would require a sample size of 98. Allowing for ~20% dropout, a sample size of 120 was proposed (60 in each group, to maximise power). Due to a higher per cent dropout in the L-AGE group, the total recruitment was increased to 138 (Fig. 1).

The study statistician performed the randomisation and all statistical analyses (Stata software, version 12; StataCorp LP, College Station, TX, USA). Descriptive analyses summarised continuous variables at baseline through their mean (SD) and median (first quartile, third quartile). Categorical variables were summarised using percentages. On the whole, the outcome variables (the differences between the mean within-intervention group differences over the length of the trial) were not greatly skewed. Thus, although the primary analyses were from *t* tests, sensitivity analyses were conducted using both general linear models, to adjust for sex, race, BMI and baseline age, and non-parametric Wilcoxon tests in case of skewed variables.

## Results

### Human data

**General description**—Baseline characteristics of cohorts randomised to L-AGE and Reg-AGE who finished the study (Fig. 1) are shown in Table 2. Both groups included participants who had two or more features of the metabolic syndrome and displayed no baseline differences in any variables, except for BMI, serum vascular cell adhesion molecule 1 (VCAM1) and glyoxalase I. We have previously compared this population with a cohort of age- and sex-matched volunteers without the metabolic syndrome [20].

### Interventional data

**AGE restriction markedly improves insulin resistance and modestly reduces body weight:** The effects of the intervention on the different variables are depicted in Table 3 and ESM Table 1. The L-AGE compared with the Reg-AGE diet led to a marked decrease in HOMA, fasting plasma insulin and insulin levels at 120 min of an OGTT (Fig. 2a,b). There were no end-of-study effects on fasting blood glucose, HbA<sub>1c</sub> levels or blood glucose at 60 and 120 min of the OGTT. However, HOMA decreased in 80% of participants on the L-AGE diet compared with 31% on the Reg-AGE diet.

Both diets were associated with a modest weight loss, which reached significance only in the L-AGE group (Table 3; ESM Table 1). Of note, markers of insulin resistance (HOMA and fasting plasma insulin), as well as serum AGEs and markers of oxidative stress and inflammation, including 8-isoprostanes, RAGE and TNF $\alpha$ , were reduced even in those participants on the L-AGE diet who did not lose body weight (n=12). By comparison, there were no such end-of-study changes in participants on the Reg-AGE diet who lost weight (n=25) (ESM Table 2).

We observed no end-of-study effects on abdominal visceral and subcutaneous fat deposits or on carotid wall variables by MRI, and there was no effect on BP, plasma triacylglycerols or HDL (Table 3; ESM Table 1).

The effect of the L-AGE diet on HOMA-IR remained highly significant, after adjusting for baseline BMI and intake of calories, protein, carbohydrate and fat, as well as age, sex and race.



**Dietary AGE restriction improves inflammatory and host defence markers:** There were marked decreases in circulating AGE markers (serum CML and MG) (Fig. 2c), as well as in intracellular AGEs (CML and MG) of PMNCs and in urinary AGEs (CML and MG) in the L-AGE group (Table 2). By contrast, in the Reg-AGE cohort, serum AGEs, intracellular AGEs and urinary AGEs continued to increase (Table 2; Fig. 2c).

Levels of 8-isoprostanes, as well as TNF $\alpha$ , VCAM1 and RAGE, were decreased in the L-AGE cohort at the end of the study (Table 2; Fig. 2c,d). On the other hand, the levels of anti-oxidative stress and inflammatory SIRT1, AGER1, adiponectin and glyoxalase I were increased in this group at the end of the study. By comparison, and similar to earlier studies [17, 19], in the Reg-AGE group, 8-isoprostanes, RAGE, HOMA-IR and leptin levels continued to increase (Table 3; Fig. 2a). There were no differences in lipid levels or renal function before and after the study in either group (Table 3).

**Sex, race, ethnicity and AGEs:** None of the above effects of the L-AGE diet were modified after adjusting for sex, age, race or ethnicity. There were no significant side effects reported by either group during the intervention, although more participants in the L-AGE group were lost to follow-up (Fig. 1).

### Ex vivo data

**L-AGE intervention ameliorates PMNC host defence deficiency and inflammation, and abrogates dysmetabolic effects on adipocytes—**PMNCs from metabolic syndrome volunteers were compared with PMNCs from healthy individuals, as previously described [20]. Cells from those with the metabolic syndrome had lower SIRT1 and AGER1 and increased TNF $\alpha$  protein levels, consistent with a pro-inflammatory metabolic syndrome PMNC phenotype [20], compared with PMNCs from participants without the metabolic syndrome (Fig. 3a,b).

Ex vivo stimulation of baseline PMNCs with MG-BSA suppressed SIRT1 and AGER1 protein levels (Fig. 3c) and increased inflammatory markers (via JNK phosphorylation), marked by increased TNF $\alpha$  secretion (Fig. 3d,e). These responses were modulated by a SIRT1 agonist and a SIRT1 inhibitor (Fig. 3a–c), directly implicating SIRT1 in these events [29].

At study end, PMNCs from L-AGE, but not Reg-AGE, participants, displayed lower TNF $\alpha$  secretion (Fig. 4a; ESM Table 1). Moreover, when insulin-stimulated 3T3-L1 adipocytes were exposed to the L-AGE conditioned medium from PMNCs obtained at study end, they revealed an improved Akt and insulin receptor phosphorylation pattern compared with adipocytes exposed to Reg-AGE conditioned medium (Fig. 4b,c).

### Discussion

This 1 year intervention shows that dietary L-AGE can improve insulin resistance, a major risk factor for type 2 diabetes, in obese people with the metabolic syndrome. It also provides clinical evidence that elevated levels of AGEs, related to AGE-rich diets, are a causal factor of insulin resistance. Since maintaining weight reduction is a difficult objective, finding that

L-AGE mitigates insulin resistance in at-risk obese people introduces a novel additional approach (e.g. to calorie restriction or drugs) in the treatment of insulin resistance, and, thus, prevention or delay of onset of type 2 diabetes. Since this was possible without a modification in energy intake, or substantial weight reduction, L-AGE may offer a feasible treatment goal of risk reduction in obese people.

We recently reported that obese individuals with the metabolic syndrome consume significantly greater amounts of dietary AGEs compared with healthy obese individuals even when consuming similar amounts of calories [20]. Herein, after 1 year of L-AGE, insulin resistance markers including HOMA and plasma insulin (fasting and 2 h OGTT) as well as leptin markedly decreased to near-normal levels [20]. There was also a modest weight loss in the L-AGE group, the impact of which on insulin resistance is uncertain. However, insulin resistance changes were independent of BMI, as well as total calorie and nutrient consumption. Furthermore, insulin resistance (HOMA, fasting plasma insulin and leptin) improved even in those on the L-AGE diet who did not have any change in body weight. Moreover, those on the Reg-AGE diet who lost body weight were found at study end to have increased levels of markers of insulin resistance or of oxidative stress and inflammation (ESM Table 2). Together these findings are consistent with the view that insulin resistance is partly independent of adiposity.

In the absence of overt diabetes or renal failure, dietary AGE consumption is shown to promote high serum AGE levels, which can influence insulin sensitivity [17-19]; thus, dietary AGEs are a risk factor for insulin resistance and the metabolic syndrome, distinct from overnutrition. Diets rich in flavourful AGEs [13] may promote food consumption and can lead to obesity [20].

The marked reductions in serum and PMNC (intracellular) AGEs, due to the L-AGE diet, together with reduced TNF $\alpha$  as well as RAGE levels, in this non-diabetic cohort with the metabolic syndrome (normal HbA<sub>1c</sub> and fasting blood glucose) confirm the view that dietary AGEs can foment dysmetabolism via oxidative stress and inflammation [20, 29]. Indeed in the L-AGE group oxidative stress and inflammation decreased in tandem with insulin resistance, thus delaying progression to type 2 diabetes. By comparison, worsening insulin resistance and markers of oxidative stress and inflammation in the Reg-AGE group, also noted in shorter trials [17], were possibly due to an expanding net AGE pool from both exogenous and endogenous sources. Thus, this 1 year trial temporally documents increased risk of progression to type 2 diabetes linked to an AGE-rich diet.

The L-AGE diet benefited several defensive factors which were found to be suppressed at baseline [20], including SIRT1, a master regulator of inflammation, insulin action, fat mobilisation [7, 30, 31] and cognitive function [32]; AGER1, an AGE receptor that antagonises AGEs, oxidative stress and inflammation [33]; and the anti-inflammatory adiponectin [34]. Restored SIRT1 and AGER1 have been implicated in the benefits of L-AGE on the metabolic syndrome as well as on dementia or Alzheimer's disease in mice [14, 24, 35]. L-AGE increased mRNA levels of *GLO1*, encoding glyoxalase I, an MG-degrading enzyme (thus a protective factor) [36], which is an altogether new and novel finding. Taken together, levels of seven genes important in the balance of oxidative stress and inflammation



were largely restored to normal by the L-AGE diet at the end of the study. Greater bioavailability of native antioxidants under the L-AGE diet might also have contributed to those effects.

The effects of L-AGE on PMNCs were likely reflected in peripheral tissues, although we did not directly assess adipose tissue [14, 27]. At study end, PMNCs from L-AGE individuals secreted less TNF $\alpha$ , such that PMNCs no longer impaired adipocyte responses to insulin. Indeed, PMNCs, like macrophages, participate in adipocyte dysfunction [28, 36], and chronic dietary AGE excess by inciting PMNC activation can impair adipocyte insulin sensitivity [12]. The pro-inflammatory phenotype of PMNCs was reversed at study end by the L-AGE diet, strongly concurring at the cellular level with the systemic findings. The net effect of L-AGE on the balance of oxidative stress and inflammation involved a robust improvement in insulin resistance.

The absence of changes at study end on central fat or carotid wall area by MRI in either group may be partly because there were no detectable abnormalities at baseline. Similarly, there were no effects on BP, plasma HDL or triacylglycerols, possibly due to the concurrent use of BP-and lipid-lowering drugs. We did not find that age, race, ethnicity and sex had an effect on the efficacy of the intervention.

AGEs may be estimated by immunochemical (ELISA) [25, 26, 37], chromatographic and mass-spectroscopy methods [18, 38]. Herein, AGEs were assessed by ELISA, a standard method used extensively to demonstrate their role in chronic disease as well as the benefits of AGE modulation in restoring health [12-15, 17-20, 25-27, 29, 33].

Weaknesses of this trial include the relatively small cohort size and the use of HOMA-IR instead of direct assessment methods of insulin resistance and sensitivity [39]. This 1 year L-AGE trial did not directly demonstrate an effect on the rate of progression to overt type 2 diabetes; however, insulin resistance, and thus risk of type 2 diabetes, decreased in the majority (80%) of L-AGE participants, compared with only 31% in the Reg-AGE group. More withdrawals in the L-AGE group might have resulted from dietary behavioural changes requiring significant effort and time commitment only in this subgroup.

In conclusion, L-AGE is effective against insulin resistance in obese individuals with the metabolic syndrome. Longer trials employing L-AGE alone or combined with other interventions should determine efficacy on risk of cardiovascular disease and other features of the metabolic syndrome.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

**Funding:** This work was supported by the National Institutes of Health (grant DK091231 to HV) and by the National Institute of Research Resources (grant MO1-RR-00071 to the General Clinical Research Center at Mount Sinai School of Medicine) for clinical and statistical support.

## References


1. Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabet Med.* 1997; 14(Suppl 5):S1–S85.
2. Fox CS, Pencina MJ, Meigs JB, Vasan RS, Levitzky YS, D'Agostino RB Sr. Trends in the incidence of type 2 diabetes mellitus from the 1970s to the 1990s: the Framingham Heart Study. *Circulation.* 2006; 113:2914–2918. [PubMed: 16785337]
3. Jason J, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC. Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes.* 2004; 53:474–481. [PubMed: 14747300]
4. Ford ES, Giles WH, Mokdad AH. Increasing prevalence of the metabolic syndrome among U.S. Adults. *Diabetes Care.* 2004; 27:2444–2449. [PubMed: 15451914]
5. Lutsey PL, Steffen LM, Stevens J. Dietary intake and the development of the metabolic syndrome: the Atherosclerosis Risk in Communities study. *Circulation.* 2008; 117:754–761. [PubMed: 18212291]
6. Hotamisligil GS. Inflammation and metabolic disorders. *Nature.* 2006; 444:860–867. [PubMed: 17167474]
7. Roberson LL, Aneni EC, Maziak W, et al. Beyond BMI: The 'metabolically healthy obese' phenotype, its association with clinical/subclinical cardiovascular disease and all-cause mortality—a systematic review. *BMC Public Health.* 2014; 14:14. [PubMed: 24400816]
8. Stefan N, Haring HU, Hu FB, Schulze MB. Metabolically healthy obesity: epidemiology, mechanisms, and clinical implications. *Lancet Diabetes Endocrinol.* 2013; 1:152–162. [PubMed: 24622321]
9. Hankinson AL, Daviglus ML, Van Horn L, et al. Diet composition and activity level of at risk and metabolically healthy obese American adults. *Obesity (Silver Spring).* 2013; 21:637–643. [PubMed: 23592673]
10. Mozzafarian D, Ludwig D. The 2015 US dietary guidelines: lifting the ban on total dietary fat. *JAMA.* 2015; 313:2421–2422. [PubMed: 26103023]
11. Chowdhury R, Warnakula S, Kunutsor S, et al. Association of dietary, circulating, and supplement fatty acids with coronary risk: a systematic review and meta-analysis. *Ann Intern Med.* 2014; 160:398–406. [PubMed: 24723079]
12. Cai W, Gao QD, Zhu L, Peppas M, He C, Vlassara H. Oxidative stress-inducing carbonyl compounds from common foods: novel mediators of cellular dysfunction. *Mol Med.* 2002; 8:337–346. [PubMed: 12393931]
13. Vlassara H, Striker GE. AGE restriction in diabetes mellitus: a paradigm shift. *Nat Rev Endocrinol.* 2011; 7:526–539. [PubMed: 21610689]
14. Cai W, Ramdas M, Zhu L, Chen X, Striker GE, Vlassara H. Oral advanced glycation endproducts (AGEs) promote insulin resistance and diabetes by depleting the antioxidant defenses AGE receptor-1 and sirtuin 1. *Proc Natl Acad Sci U S A.* 2012; 109:15888–15893. [PubMed: 22908267]
15. Uribarri J, Woodruff S, Goodman S, et al. Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc.* 2010; 110:911–916. [PubMed: 20497781]
16. Birlouez-Aragon I, Saavedra G, Tessier FJ, et al. A diet based on high-heat-treated foods promotes risk factors for diabetes mellitus and cardiovascular diseases. *Am J Clin Nutr.* 2010; 91:1220–1226. [PubMed: 20335546]
17. Uribarri J, Cai W, Ramdas M, et al. Restriction of advanced glycation end products improves insulin resistance in human type 2 diabetes: potential role of AGER1 and SIRT1. *Diabetes Care.* 2011; 34:1610–1616. [PubMed: 21709297]
18. Mark AB, Poulsen MW, Andersen S, et al. Consumption of a diet low in advanced glycation end products for 4 weeks improves insulin sensitivity in overweight women. *Diabetes Care.* 2014; 37:88–95. [PubMed: 23959566]
19. Vlassara H, Cai W, Goodman S, et al. Protection against loss of innate defenses in adulthood by low advanced glycation end products (AGE) intake: role of the antiinflammatory AGE receptor-1. *J Clin Endocrinol Metab.* 2009; 94:4483–4491. [PubMed: 19820033]

20. Uribarri J, Cai W, Woodward M, et al. Elevated serum advanced glycation endproducts in obese indicate risk for the metabolic syndrome: a link between healthy and unhealthy obese? *J Clin Endocrinol Metab.* 2015; 100:1957–1966. [PubMed: 25695886]
21. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation.* 2002; 106:3143–3421. [PubMed: 12485966]
22. Matthews DR, Hosker JP, Rudenski AS, et al. Homeostasis model assessment: insulin resistance and beta cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985; 28:412–419. [PubMed: 3899825]
23. Bonekamp S, Ghosh P, Crawford S, et al. Quantitative comparison and evaluation of software packages for assessment of abdominal adipose tissue distribution by magnetic resonance imaging. *Int J Obes (Lond).* 2008; 32:100–111. [PubMed: 17700582]
24. Fayad ZA, Mani V, Woodward M, et al. Safety and efficacy of dalcetrapib on atherosclerotic disease using novel non-invasive multimodality imaging (dal-PLAQUE): a randomised clinical trial. *Lancet.* 2011; 378:1547–59. [PubMed: 21908036]
25. Makita Z, Vlassara H, Cerami A, Bucala R. Immunochemical detection of advanced glycosylation end products in vivo. *J Biol Chem.* 1992; 267:5133–5138. [PubMed: 1371995]
26. Mitsuhashi T, Vlassara H, Founds HW, Li YM. Standardizing the immunological measurement of advanced glycation endproducts using normal human serum. *J Immunol Methods.* 1997; 207:79–88. [PubMed: 9328589]
27. Cai W, Uribarri J, Zhu L, et al. Oral glycotoxins are a modifiable cause of dementia and the metabolic syndrome in mice and humans. *Proc Natl Acad Sci U S A.* 2014; 111:4940–4945. [PubMed: 24567379]
28. Yoshizaki T, Schenk S, Imamura T, et al. SIRT 1 inhibits inflammatory pathways in macrophages and modulates insulin sensitivity. *Am J Physiol Endocrinol Metab.* 2010; 298:E419–E428. [PubMed: 19996381]
29. Uribarri J, Cai W, Pyzik R, et al. Suppression of native defense mechanisms, SIRT1 and PPAR $\gamma$ , by dietary glycoxidants precedes disease in adult humans; relevance to lifestyle-engendered chronic diseases. *Amino Acids.* 2014; 46:301–309. [PubMed: 23636469]
30. Liang F, Kume S, Koya D. SIRT1 and insulin resistance. *Nat Rev Endocrinol.* 2009; 5:367–373. [PubMed: 19455179]
31. Yoshizaki T, Milne JC, Imamura T, et al. SIRT1 exerts anti-inflammatory effects and improves insulin sensitivity in adipocytes. *Mol Cell Biol.* 2009; 29:1363–1374. [PubMed: 19103747]
32. Guarente L, Franklin H, Franklin H. Epstein lecture. Sirtuins, aging, and medicine. *N Engl J Med.* 2011; 364:2235–2244. [PubMed: 21651395]
33. Torreggiani M, Liu H, Wu J, et al. Advanced glycation end product receptor-1 transgenic mice are resistant to inflammation, oxidative stress, and post-injury intimal hyperplasia. *Am J Pathol.* 2009; 175:1722–1732. [PubMed: 19779136]
34. Takemura Y, Ouchi N, Shibata R, et al. Adiponectin modulates inflammatory reactions via calreticulin receptor-dependent clearance of early apoptotic bodies. *J Clin Invest.* 2007; 117:375–386. [PubMed: 17256056]
35. de Kreutzenberg SV, Ceolotto G, Papparella I, et al. Downregulation of the longevity-associated protein sirtuin 1 in insulin resistance and metabolic syndrome: potential biochemical mechanisms. *Diabetes.* 2010; 59:1006–1015. [PubMed: 20068143]
36. Brouwers O, Niessen PM, Ferreira I, et al. Overexpression of glyoxalase-I reduces hyperglycemia-induced levels of advanced glycation end products and oxidative stress in diabetic rats. *J Biol Chem.* 2011; 286:1374–1380. [PubMed: 21056979]
37. Bucala R, Makita Z, Koschinsky T, et al. Lipid advanced glycosylation: pathway for lipid oxidation in vivo. *Proc Natl Acad Sci USA.* 1993; 90:6634–6638.
38. Scheijen JL, Clevers E, Engelen L, et al. Analysis of advanced glycation endproducts in selected food items by ultra-performance liquid chromatography tandem mass spectrometry: presentation of a dietary AGE database. *Food Chem.* 2016; 190:1145–50. [PubMed: 26213088]

39. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care*. 1999; 22:1462–1470. [PubMed: 10480510]


## Abbreviations

<b>AGER1</b>	AGE receptor 1
<b>CML</b>	Carboxymethyllysine
<b>L-AGE</b>	Restricted AGE intake
<b>MG</b>	Methylglyoxal
<b>PMNC</b>	Peripheral blood mononuclear cell
<b>qRT-PCR</b>	Quantitative real-time reverse transcription PCR
<b>RAGE</b>	Receptor for AGEs
<b>Reg-AGE</b>	Regular AGE intake
<b>SIRT1</b>	Sirtuin 1
<b>VCAM1</b>	Vascular cell adhesion molecule 1



**Fig. 1. CONSORT study flow diagram**


a-d

**Fig. 2.**

Effects of L-AGE diet (black bars) vs Reg-AGE diet (white bars) on metabolic variables and systemic markers of AGEs, oxidative stress and inflammation (also see Table 2). **(a)** Markers of insulin resistance: per cent changes (means  $\pm$  SEM) in levels of HOMA, leptin and fasting insulin between baseline and end of study (Reg-AGE,  $n=43$ ; L-AGE,  $n=51$ ;  $*p < 0.05$ ). **(b)** Plasma insulin after OGTT: per cent changes (means  $\pm$  SEM) in plasma insulin levels 0 and 120 min after OGTT between baseline and end of study (Reg-AGE,  $n=43$ ; L-AGE,  $n=51$ ;  $*p < 0.05$ ). **(c)** AGEs and pro-oxidative stress and inflammation markers: per cent changes (means  $\pm$  SEM) in markers of serum (s) AGEs (CML, MG), plasma 8-isoprostanes (8-iso) and PMNC inflammatory factors (TNF $\alpha$  protein and RAGE mRNA) between baseline and end of study (Reg-AGE,  $n=33$ ; L-AGE,  $n=21$ ;  $*p < 0.05$ ). **(d)** Anti-oxidative stress and inflammation markers: per cent changes (means  $\pm$  SEM) in levels of antioxidants and markers of host defence, SIRT1, AGER1 and GLO1 mRNA, between baseline and end of study (Reg-AGE,  $n=33$ ; L-AGE,  $n=21$ ;  $*p < 0.05$ )




a-e

**Fig. 3.**

Baseline ex vivo cellular data: AGEs suppress host defences and promote inflammatory activation of PMNCs in individuals with the metabolic syndrome (MS) compared with normal individuals (NL). **(a)** Expression levels of anti-oxidative stress and inflammation factors, SIRT1 (white bars) and AGER1 (black bars), and **(b)** of the pro-oxidative stress and inflammation cytokine TNF $\alpha$  protein in PMNCs freshly obtained at study commencement (baseline), from obese study MS participants (MS-PMNCs) (black bar) compared with that in PMNCs simultaneously obtained from NL (white bar). Data in **(a)** are from western blots and densitometry, shown as AU or ratio of target protein to  $\beta$ -actin (means  $\pm$  SEM) and in **(b)** as (means  $\pm$  SEM) pg/mg cell protein by ELISA ( $n=10$ /group, each in triplicate; \* $p$  0.05 vs NL). **(c)** Baseline MS-PMNCs were exposed ex vivo to MG-BSA 60  $\mu$ g/ml for 72 h, in the presence or absence of a SIRT1 activator, SRT1720 (1  $\mu$ mol/l), or a SIRT1 inhibitor, sirtinol (10  $\mu$ mol/l). Western blots, performed on cell extracts, and density analysis (means  $\pm$  SEM) ( $n=3-5$ ) are shown as the ratio of SIRT1 (white bars) or AGER1 (black bars) to  $\beta$ -actin; \* $p$  0.05 vs non-stimulated MS-PMNCs (control, CL);  $\dagger p$  0.05 vs MG alone. **(d)** MS-PMNC extracts, prepared as in **(c)**, were also probed for levels of p-JNK (black bars) and acetylated NF $\kappa$ B p65 at lys310 (AcL-p65) (white bars) by western blots, using the respective antibodies and density analyses. Total JNK and p65 served as internal controls (CL). Density data (means  $\pm$  SEM,  $n=3-5$ ) indicate the ratios of phosphorylated or acetylated protein to total protein; \* $p$  0.05 vs non-stimulated CL;  $\dagger p$  0.05 vs MG alone. **(e)** TNF $\alpha$  secreted into the culture medium at 72 h from non-stimulated MS-PMNCs (CL, white bar) or MG-stimulated MS-PMNCs (black bars) at baseline, in the presence or absence of SRT1720 (1  $\mu$ mol/l) or sirtinol (10  $\mu$ mol/l), as above. Assays were performed in triplicate for each sample. Data (means  $\pm$  SEM) are shown as per cent of control ( $n=7$ , \* $p$  0.05 vs CL;  $\dagger p$  0.05 vs MG alone). All PMNC samples were collected at study baseline

a-c

**Fig. 4.**

End-of-study ex vivo cellular data: L-AGE diet mitigates inflammatory activation of PMNCs from individuals with the metabolic syndrome (MS-PMNCs) and improves their impact on insulin responses of adipocytes in vitro. **(a)** TNF $\alpha$  from MS-PMNCs secreted into the conditioned media, after 1 year on L-AGE (black bar,  $n=5$ ) or Reg-AGE diet (white bar,  $n=5$ ). Data are shown as means  $\pm$  SEM pg/ml medium, each in triplicate; \* $p$  0.05 L-AGE vs Reg-AGE diet. **(b)** JNK phosphorylation in differentiated 3T3-L1 adipocytes after overnight incubation with conditioned medium either from baseline (BL) or end-of-study PMNCs from L-AGE (black bar,  $n=5$ ) and Reg-AGE (white bar,  $n=5$ ) treatment groups. Cell lysates were subjected to western blot and densitometry. As additional control, 3T3-L1 adipocytes were incubated with normal culture medium, shown as 0. Density data are means  $\pm$  SEM ( $n=3-5$  independent experiments); \* $p$  0.05, L-AGE or Reg-AGE vs BL; †  $p$  0.05, L-AGE vs Reg-AGE diet. **(c)** Serine phosphorylation (at Ser473) of Akt and tyrosine phosphorylation of insulin receptor (IR) in 3T3-L1 adipocytes after exposure to conditioned medium (CM) overnight as in **(b)**. Cells were incubated in the presence (black bars) or absence of insulin (Ins) (white bars) (100 nmol/l, 30 min) and subjected to western blot and densitometry (means  $\pm$  SEM,  $n=3$  independent experiments); \* $p$  0.05 L-AGE vs BL. PMNC samples ( $n=7$ ) were from the same individuals evaluated at baseline (Fig. 2,  $n=7$ ). 3T3-L1 adipocytes incubated with normal CM are shown as 0

**Table 1**  
**Sample of daily dietary AGE intake from a participant on the L-AGE diet**

Meal	Baseline high AGE diet			Intervention L-AGE diet		
	Item	Portion	AGEs <sup>a</sup>	Item	Portion	AGEs <sup>a</sup>
Breakfast	Fresh fruit cup	0.12 l (1/2 cup)	15	Fresh fruit cup	0.12 l (½ cup)	15
	Fried eggs	1	1,200	Boiled egg	1	75
	Toasted bagel	112 g	200	Fresh bagel	112 g	120
	Cream cheese	5 ml	500	Cream cheese	5 ml	500
	Skimmed milk	240 ml	2	Skimmed milk	240 ml	2
	Coffee	240 ml	19	Coffee	240 ml	19
	Orange juice	120 ml	3	Orange juice	120 ml	3
Lunch	Grilled chicken breast	84 g	5,200	Poached chicken breast	84 g	1,000
	Green salad	0.24 l (1 cup)	0	Green salad	0.24 l (1 cup)	0
	Caesar dressing	30 ml	220	Caesar dressing	30 ml	200
	Bread, white	1 slice	10	Bread, white	1 slice	10
	Margarine	5 ml	900	Margarine	5 ml	900
	Iced tea	360 ml	5	Iced tea	360 ml	5
	Apple	1 medium	15	Apple	1 medium	15
Dinner	Cantaloupe wedge	¼ small	20	Cantaloupe wedge	¼ small	20
	Grilled steak	84 g	6,600	Beef stew	84 g	2,200
	Mashed potato	1	20	Mashed potato	1	20
	Carrots	0.12 l (½ cup)	10	Carrots	0.12 l (½ cup)	10
	Coffee with milk	240 ml	5	Coffee with milk	240 ml	5
	Muffin, bran	1	102	Muffin, bran	1	102
	Total AGEs		15,046			5,221
Total energy, kJ/day		7.94			7.77	

<sup>a</sup> Arbitrary AGE units are expressed in AGE kilounits based on data from reference [15]

**Table 2**

**Characteristics of study groups at baseline**

Characteristic	Reg-AGE diet			L-AGE diet			p value <sup>d</sup>
	n	Mean	SD	n	Mean	SD	
Age, years	49	60.7	6.9	51	60.8	7.3	0.949
Men, %	49	30.6	-	51	39.2	-	0.668
Systolic BP, kPa	49	17.2	1.9	51	16.8	1.8	0.322
Diastolic BP, kPa	49	10.0	1.4	51	10.0	1.3	0.629
Weight, kg	49	90.7	19.2	51	84.8	13.0	0.079
BMI, kg/m <sup>2</sup>	49	33.3	6.3	51	31.2	4.2	0.050
Waist circumference, cm	49	110.4	12.3	51	106.3	8.2	0.052
Body fat (BIA), %	49	39.8	7.3	51	38.3	7.6	0.297
SAT, cm <sup>2</sup>	39	229.6	94.3	43	212.7	91.7	0.815
VAT, cm <sup>2</sup>	39	137.9	81.3	43	116.8	65.4	0.165
Fasting plasma glucose, mmol/l	49	4.8	0.6	51	4.9	0.7	0.143
Glucose AUC (OGTT)	49	261.7	66.1	51	283.6	68.7	0.107
Fasting plasma insulin, pmol/l	49	95.8	42.4	50	94.8	54.2	0.852
Insulin AUC (OGTT)	49	146.6	92.8	50	125.2	85.0	0.204
HbA <sub>1c</sub> , %	49	5.8	0.5	51	6.0	0.5	0.068
HbA <sub>1c</sub> , mmol/mol	49	39.9	-	51	42.1	-	0.068
HOMA-IR	49	3.0	1.3	50	3.1	1.8	0.714
Triacylglycerol, mmol/l	19	1.53	1.07	51	1.49	1.02	0.852
HDL-cholesterol, mmol/l	49	1.47	0.46	51	1.41	0.39	0.477
Serum CML, U/ml	49	17.1	6.8	51	17.3	7.1	0.928
Serum MG, nmol/ml	49	2.28	0.48	51	2.28	0.59	0.981
iCML, U/mg	40	9.15	1.54	41	8.85	1.61	0.194
iMG, nmol/mg	40	1.05	0.30	41	1.01	0.31	0.604
Plasma 8-isoprostanol, pg/ml	49	214.9	114.4	51	205.7	110.8	0.684
Serum VCAM1, ng/ml	14	831.1	171.2	11	709.9	183.7	0.018
Serum leptin, ng/ml	49	29.5	11.8	51	26.5	13.4	0.243
Plasma adiponectin, µg/ml	49	8.79	3.01	51	9.40	3.09	0.378

Characteristic	Reg-AGE diet			L-AGE diet			p value <sup>a</sup>
	n	Mean	SD	n	Mean	SD	
<i>AGER</i> mRNA	47	177.6	72.1	51	184.7	88.1	0.659
<i>RAGE</i> mRNA	47	509.7	203.4	51	480.3	221.8	0.521
<i>SIRT1</i> mRNA	47	247.6	110.7	51	259.4	108.2	0.595
<i>GLO1</i> mRNA	47	28.6	9.9	51	23.5	11.2	0.018
TNF $\alpha$ , pg/mg protein	48	15.0	4.5	48	14.5	4.7	0.639
Dietary energy, kJ/day	49	8,880	2,801	51	8,060	2,613	0.072
Dietary AGEs, AGE Eq/day	49	18.6	10.9	51	17.4	11.3	0.498
eGFR, ml min <sup>-1</sup> 1.73 m <sup>-2</sup>	49	79	17	51	76	75	0.399
Urine CML excretion, U/day	49	90,480	62,904	51	97,492	64,516	0.344
Urine MG excretion, nmol/day	49	1,056	893	51	1,190	1161	0.307
Average carotid wall area, mm <sup>2</sup>	36	23.7	6.3	42	23.9	5.0	0.826
Average carotid wall thickness, mm	36	0.991	0.180	42	0.993	0.126	0.816

<sup>a</sup> Comparison between means at baseline between groups was carried out using Student's *t* test

BIA, bioelectrical impedance; eGFR, estimated GFR by Modification of Diet in Renal Disease study equation; Eq, equivalent; iCML, intracellular CML per mg protein in PMNCs; iMG, intracellular MG per mg protein in PMNCs; SAT, subcutaneous abdominal tissue; VAT, visceral abdominal tissue

**Table 3**

**Changes from baseline to 12 months by intervention group**

Variable	Reg-AGE diet			L-AGE diet		
	Month 0	Month 12	p value <sup>a</sup>	Month 0	Month 12	p value <sup>b</sup>
Systolic BP, kPa	17.2±1.9	17.3±1.9	0.553	16.9±1.7	17.2±1.7	0.330
Diastolic BP, kPa	10.1±1.2	10.3±1.3	0.428	10±1.3	10±1.3	0.850
Weight, kg	90.7±19	90.2±19	0.359	84.5±13	82.8±13	0.001
BMI, kg/m <sup>2</sup>	33.3±6.3	33.2±6.2	0.468	31.2±4.2	30.0±6.0	0.091
Waist circumference, cm	110±12	107±13	0.001	106±8	104±10	0.012
Body fat (BIA), %	40±7	40±8	0.618	38±7	37±9	0.232
SAT, cm <sup>2</sup>	225±93	215±90	0.132	212±94	200±92	0.009
VAT, cm <sup>2</sup>	142±82	134±80	0.044	113±66	101±65	0.010
Fasting plasma glucose, mmol/l	4.9±0.6	4.8±0.6	0.635	5.1±0.7	4.9±0.7	0.105
Glucose AUC (OGTT)	263±66	265±67	0.713	286±69	279±70	0.270
Fasting plasma insulin, pmol/l	95.8±41.7	113.2±45.8	0.002	93.8±55.6	60.4±38.9	0.001
Insulin AUC (OGTT)	149±98	147±105	0.906	128±87	124±111	0.974
HbA <sub>1c</sub> , %	5.8±0.5	5.8±0.5	0.850	6.0±0.5	5.9±0.6	0.073
HbA <sub>1c</sub> , mmol/mol	39.9	39.9	0.850	42.1	41.0	0.073
HOMA-IR	2.9±1.2	3.6±1.7	0.002	3.1±1.8	1.9±1.3	0.001
Triacylglycerol, mmol/l	1.54±1.06	1.37±0.82	0.042	1.54±1.03	1.60±1.13	0.605
HDL-cholesterol, mmol/l	1.47±0.47	1.50±0.41	0.477	1.40±0.36	1.47±0.47	0.006
Serum CML, U/ml	17±7	23±8	0.001	17±7	13±7	0.001
Serum MG, nmol/ml	2.2±0.4	2.9±0.7	0.001	2.3±0.6	1.7±0.7	0.001
iCML, U/mg	9.0±1.6	10.7±2.9	0.007	8.7±1.7	6.9±2.2	0.001
iMG, nmol/mg	1.0±0.3	1.2±0.4	0.039	1.0±0.3	0.7±0.2	0.001
Plasma 8-isoprostanes, pg/ml	207±100	291±115	0.001	205±112	127±77	0.001
Serum VCAM1, ng/ml	1,066±104	1,172±77	0.559	813±157	589±162	0.041
Serum leptin, ng/ml	29±10	38±12	0.001	27±14	16±10	0.001
Plasma adiponectin, µg/ml	8.9±31	7.8±2.6	0.072	9.5±3.1	16.7±7.2	0.001
AGER mRNA	180±71	159±65	0.110	186±86	347±149	0.001



Variable	Reg-AGE diet			L-AGE diet		
	Month 0	Month 12	<i>p</i> value <sup>a</sup>	Month 0	Month 12	<i>p</i> value <sup>b</sup>
RAGE mRNA	500±209	606±185	0.002	471±217	322±163	0.001
SIRT7 mRNA	251±110	216±87	0.052	262±108	403±152	0.001
GLO1 mRNA	28±9	26±10	0.262	23±11	36±15	0.001
TNFr $\alpha$ , pg/mg protein	14.6±4.8	16.2±4.0	0.051	13.8±4.5	8.9±3.3	0.001
Dietary energy, kJ/day	8,901±2826	8,051±2349	0.033	7,813±2855	6,381±1997	0.001
Dietary AGEs, AGE Eq/day	19±9	20±11	0.459	17±12	7±6	0.001
eGFR ml min <sup>-1</sup> 1.73 m <sup>-2</sup>	79±17	79±20	0.979	76±18	77±16	0.358
Urine CML excretion, U/day	90.5±62.9	111.5±70.6	0.057	99.0±66.1	66.1±43.2	0.001
Urine MG excretion, nmol/day	1,056±893	1,723±1235	0.001	1,201±1191	832±753	0.029
Average carotid wall area, mm <sup>2</sup>	24±6	25±5	0.170	24±5	25±4	0.156
Average carotid wall thickness, mm <sup>2</sup>	0.98±0.16	1.03±0.13	0.173	1.00±0.13	1.04±0.12	0.116

All values expressed as mean ± SD

Both *p* values were obtained by paired *t* test analyses

<sup>a</sup>Difference between baseline and end of study for the L-AGE diet

<sup>b</sup>Difference between baseline and end of study for the Reg-AGE diet

iCML, intracellular CML per mg protein in PMNCs; iMG, intracellular MG per mg protein in PMNCs; Eq, equivalent