



CrossMark  
click for updates

## Comment

**Cite this article:** Postma E. 2016 Why we should *not* dismiss a relationship between attractiveness and performance: a comment on Smoliga & Zavorsky (2015). *Biol. Lett.* **12**: 20160068.

<http://dx.doi.org/10.1098/rsbl.2016.0068>

Received: 27 January 2016

Accepted: 23 May 2016

### Author for correspondence:

Erik Postma

e-mail: [erik.postma@ieu.uzh.ch](mailto:erik.postma@ieu.uzh.ch)

The accompanying reply can be viewed at  
<http://dx.doi.org/10.1098/rsbl.2016.0856>.

## Evolutionary biology

# Why we should *not* dismiss a relationship between attractiveness and performance: a comment on Smoliga & Zavorsky (2015)

Erik Postma

Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

EP, 0000-0003-0856-1294

Smoliga & Zavorsky (S&Z) [1] dismiss a series of studies reporting a relationship between facial attractiveness and sports performance because the proportion of variance explained is small and the effect may not be generalizable to the general population. They, therefore, conclude that such studies has “questionable biological importance” and “are not valid for studying evolution” [1].

While few will disagree with S&Z when they write that statistical significance does not equal biological significance, their suggestion that biological meaningfulness can be equated to the proportion of variation explained (measured by  $r^2$ ; see their first recommendation for future research) is open to debate. Although the low  $r^2$  reported in, for example [2], indeed means that physical appearance alone poorly predicts performance of a Tour de France rider, the prediction of whether a Darwin’s finch is going to survive to the following year on the basis of its beak size is similarly imprecise ( $r^2 = 0.06\text{--}0.09$ ), and this despite a significant relationship between beak size and survival [3]. Their definition of biological meaningfulness would thus lead S&Z to dismiss a textbook example of natural selection.

Fitness components such as survival, reproductive success and attractiveness are complex traits, and any single variable will—by definition—explain only a small amount of variation. Hence,  $r^2$  is a poor measure of the strength of selection, which is instead measured by the selection differential, i.e. the covariance between some component of relative fitness ( $w$ ) and the trait of interest ( $z$ ) (see e.g. [4]). If  $z$  is standardized to have a variance of one, a standardized selection differential can be obtained by regressing  $w$  on  $z$ . Importantly, whereas the slope is given by the covariance between  $w$  and  $z$ , divided by the variance in  $z$  (which is equal to one if  $z$  is standardized), the  $r^2$  is equal to the covariance between  $w$  and  $z$  squared, divided by the product of the variances in  $w$  and  $z$ . Hence, even if the slope is steep (and selection, therefore, strong),  $r^2$  will be low whenever variation in  $w$  is large and attributable to a multitude of factors other than  $z$ . Given the complex and multidimensional nature of both endurance performance and attractiveness, their shared component will, therefore, always be small, and expecting  $r^2$  to be any higher would be naive. The low  $r^2$  of a relationship between facial attractiveness and performance is, therefore, a poor reason to dismiss its evolutionary relevance.

Whereas [2] reports the slope of untransformed attractiveness on performance, the standardized estimate of the strength of sexual selection within the 2012 Tour de France peloton, estimated as the slope of the regression line of *relative* attractiveness on *variance-standardized* performance, is 0.056. This means that an increase in performance by 1 s.d. comes with a 6% increase in attractiveness. Albeit weaker than the median strength of linear sexual selection observed in non-human animals (0.18) [5], assuming attractiveness is correlated with reproductive success, theory predicts (a preference for) performance to evolve. Although there are various reasons why we have to be careful making such predictions [6], the low

proportion of variance that performance and attractiveness have in common is not among them.

S&Z, furthermore, make the obvious point that the Tour de France peloton is not a random sample of the general population, capturing only a fraction of all variation in performance that exists. How the absolute and relative importance of genes (talent) and environment (training) in shaping variation in performance (sensu [2]) differs between the Tour de France peloton and the general population is an outstanding question. However, assuming that it is the variation of non-genetic origin, attributable to, for example variation in training quality and volume, that is reduced in particular, performance variation within the peloton may arguably be more representative of the variation that selection has acted upon during our evolutionary history [7,8]. If this indeed is the case, testing for a relationship between attractiveness and performance in

the general population, including both couch potatoes and ambitious athletes, addresses an interesting, but fundamentally different question and dismissing the pattern observed in [2] by extrapolating it to the general population would be fallacious.

S&Z and I agree that an evolutionary perspective may provide novel insights into the nature of human physical fitness, and it is beyond doubt that a conclusive demonstration of endurance performance being subject to sexual selection, now or in our evolutionary past, will require more research. It is, therefore, unfortunate that several of their recommendations for future studies are misguided and, therefore, unlikely to bring us closer to an answer.

**Competing interests.** I have no competing interests.

## References

1. Smoliga JM, Zavorsky GS. 2015 Faces and fitness: attractive evolutionary relationship or ugly hypothesis? *Biol. Lett.* **11**, 20150839. (doi:10.1098/rsbl.2015.0839).
2. Postma E. 2014 A relationship between attractiveness and performance in professional cyclists. *Biol. Lett.* **10**, 20130966. (doi:10.1098/rsbl.2013.0966)
3. Price TD, Grant PR, Gibbs HL, Boag PT. 1984 Recurrent patterns of natural selection in a population of Darwin finches. *Nature* **309**, 787–789. (doi:10.1038/309787a0)
4. Brodie ED, Moore AJ, Janzen FJ. 1995 Visualizing and quantifying natural selection. *Trends Ecol. Evol.* **10**, 313–318. (doi:10.1016/S0169-5347(00)89117-X)
5. Kingsolver JG, Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hill CE, Hoang A, Gibert P, Beerli P. 2001 The strength of phenotypic selection in natural populations. *Am. Nat.* **157**, 245–261. (doi:10.1086/319193)
6. Morrissey MB, Kruuk LEB, Wilson AJ. 2011 The danger of applying the breeder's equation in observational studies of natural populations. *J. Evol. Biol.* **23**, 2277–2288. (doi:10.1111/j.1420-9101.2010.02084.x)
7. Bramble DM, Lieberman DE. 2004 Endurance running and the evolution of Homo. *Nature* **432**, 345–352. (doi:10.1038/nature03052)
8. Longman D, Wells JCK, Stock JT. 2015 Can persistence hunting signal male quality? A test considering digit ratio in endurance athletes. *PLoS ONE* **10**, e0121560. (doi:10.1371/journal.pone.0121560)