



Published in final edited form as:

*Cell Microbiol.* 2016 June ; 18(6): 844–858. doi:10.1111/cmi.12554.

## ***Porphyromonas gingivalis* Initiates a Mesenchymal-like Transition through ZEB1 in Gingival Epithelial Cells**

Maryta N. Sztukowska<sup>1</sup>, Akintunde Ojo<sup>1</sup>, Saira Ahmed<sup>1</sup>, Anne L. Carenbauer<sup>1</sup>, Qian Wang<sup>1</sup>, Brain Shumway<sup>2</sup>, Howard F. Jenkinson<sup>3</sup>, Huizhi Wang<sup>1</sup>, Douglas S. Darling<sup>1</sup>, and Richard J. Lamont<sup>1,\*</sup>

<sup>1</sup>Department of Oral Immunology and Infectious Diseases, University of Louisville School of Dentistry, Louisville, Kentucky, United States of America

<sup>2</sup>Department of Surgical and Hospital Dentistry, University of Louisville School of Dentistry, Louisville, Kentucky, United States of America

<sup>3</sup>School of Oral and Dental Sciences, University of Bristol, Bristol, United Kingdom

### **Summary**

The oral anaerobe *Porphyromonas gingivalis* is associated with the development of cancers including oral squamous cell carcinoma (OSCC). Here we show that infection of gingival epithelial cells with *P. gingivalis* induces expression and nuclear localization of the ZEB1 transcription factor which controls epithelial-mesenchymal transition (EMT). *P. gingivalis* also caused an increase in ZEB1 expression as a dual species community with *Fusobacterium nucleatum* or *Streptococcus gordonii*. Increased ZEB1 expression was associated with elevated ZEB1 promoter activity and did not require suppression of the miR-200 family of micro RNAs. *P. gingivalis* strains lacking the FimA fimbrial protein were attenuated in their ability to induce ZEB1 expression. ZEB1 levels correlated with an increase in expression of mesenchymal markers, including vimentin and MMP-9, and with enhanced migration of epithelial cells into matrigel. Knockdown of ZEB1 with siRNA prevented the *P. gingivalis*-induced increase in mesenchymal markers and epithelial cell migration. Oral infection of mice by *P. gingivalis* increased ZEB1 levels in gingival tissues, and intracellular *P. gingivalis* were detected by antibody staining in biopsy samples from OSCC. These findings indicate that FimA-driven ZEB1 expression could provide a mechanistic basis for a *P. gingivalis* contribution to OSCC.

### **Introduction**

Once considered implausible, the concept that bacteria can be associated with cancer development is now well established. Indeed, a causal relationship between *Helicobacter pylori* and gastric cancer has been demonstrated (Kim *et al.*, 2011), and a growing body of evidence supports the relationship between specific bacteria and various types of cancer

\*Corresponding author: 570 South Preston Street, University of Louisville, Louisville, KY, 40202, Phone: 502-852-2112, rich.lamont@louisville.edu.

#### **Conflict of Interest**

The authors have no conflict of interest to declare.

(Garrett, 2015, Sahingur and Yeudall, 2015). For example, *Fusobacterium nucleatum*, a common inhabitant of the oral cavity, is over-represented in colorectal carcinoma (Castellarin *et al.*, 2012, Kostic *et al.*, 2012) and can induce colorectal carcinogenesis by activating E-cadherin/ $\beta$ -catenin signaling (Rubinstein *et al.*, 2013). *F. nucleatum* can also inhibit natural killer (NK) cell cytotoxicity and killing of various tumors (Gur *et al.*, 2015). High levels of antibodies to *Porphyromonas gingivalis*, a keystone pathogen in periodontal diseases, correlate with a greater than 2-fold increased risk of pancreatic cancer (Michaud, 2013). *P. gingivalis* is also associated with oral squamous cell carcinoma (OSCC). The surfaces of OSCCs harbor higher levels of *Porphyromonas* compared to contiguous healthy mucosa (Nagy *et al.*, 1998), and *P. gingivalis* can be detected within gingival carcinomas by immunohistochemistry (Katz *et al.*, 2011). Moreover, recent studies have established that combined infection with *P. gingivalis* and *F. nucleatum* promotes tumor progression in an oral-specific chemical carcinogenesis mouse model (Gallimidi *et al.*, 2015).

*P. gingivalis* and oral epithelial cells engage in an intricate molecular dialog, one consequence of which is entry of bacterial cells into the cytoplasm of the host cell (Lamont and Hajishengallis, 2015, Lamont *et al.*, 1995). Primary cultures of epithelial cells containing *P. gingivalis* do not undergo apoptotic cell death and indeed *P. gingivalis* can suppress several proapoptotic pathways. In response to *P. gingivalis* infection Jak1/Akt/Stat3 signaling is activated with resultant increase in Bcl2 and inhibition of intrinsic mitochondrial apoptotic pathways (Yilmaz *et al.*, 2004, Mao *et al.*, 2007). By an independent mechanism *P. gingivalis* upregulates the level of miR-203 which suppresses expression of SOCS3, consequently impeding apoptosis (Moffatt and Lamont, 2011). In tandem with suppression of apoptosis, *P. gingivalis* promotes acceleration of primary epithelial cells through the S-phase of the cell cycle by impacting cyclin/CDK activities and reducing the amount of p53 (Kuboniwa *et al.*, 2008). The process is dependent on the major fimbriae of *P. gingivalis* as a mutant deficient in FimA, the structural fimbrial subunit protein, does not induce increased cell proliferation.

While inhibition of apoptosis and enhanced replication of cells can contribute directly to tumor development, it is unknown if *P. gingivalis* is capable of initiating the malignant transformation or oncogenic progression of epithelial cells. The epithelial-mesenchymal transition (EMT) is a process by which epithelial cells change shape and acquire a motile phenotype (Lamouille *et al.*, 2014). The EMT is required for normal development and wound healing; however, it is also associated with the generation of self-renewing tumor-initiating cells, and in a malignant tumor it gives rise to a population of migratory and invasive cancer cells (Lamouille *et al.*, 2014). This switch in cell differentiation and behavior is controlled by a group of transcription factors including the zinc-finger E-box-binding homeobox 1 and 2 proteins (ZEB1/2), SNAIL and TWIST (Vandewalle *et al.*, 2009, Scanlon *et al.*, 2013). The ZEB1 ( $\delta$ EF1, Zfhx1a, Zfh1) and ZEB2 (SIP1) transcription factors are critical EMT activators that bind to 5'-CACCTG sequences and repress transcription of epithelial specific genes such as E-cadherin (*cdh1*) (Vandewalle *et al.*, 2009). ZEB can also positively regulate genes associated with the mesenchymal phenotype such as those encoding vimentin and matrix-metalloproteinases (Vandewalle *et al.*, 2009, Lamouille *et al.*, 2014). ZEB1/2 are in the TGF $\beta$  signaling pathway, binding SMADs and having essential effects on embryonic development (Gheldof *et al.*, 2012). ZEB1 has been implicated in

activating EMT and metastasis in several type of cancers (Sanchez-Tillo *et al.*, 2012, Jia *et al.*, 2012). Moreover, *ZEB1/2* are linked to the miR-200 family in a reciprocal negative feedback loop whereby each regulates the expression of the other (Brabletz and Brabletz, 2010).

*H. pylori* has been shown to upregulate expression of *ZEB1* which can initiate an EMT and cancer stem-cell properties in infected gastric epithelial cells (Baud *et al.*, 2013, Bessede *et al.*, 2014). In this study we show that *P. gingivalis* can increase *ZEB1* levels in gingival epithelial cells in a fimbriae dependent manner. Upregulation of *ZEB1* was dependent on increased promoter activity. Elevated expression of *ZEB1* was associated with a partial mesenchymal phenotype in *P. gingivalis*-infected gingival epithelial cells, including increased migration. We also detected *P. gingivalis* antigens in oral carcinoma in situ and poorly differentiated cancer, and mice orally infected with *P. gingivalis* had an increase in *ZEB1* mRNA expression in gingival tissues. The results suggest a novel mechanism by which oral bacteria such as *P. gingivalis* can contribute to a mesenchymal phenotype, and potentially drive the progression of cancer.

## Results

### *P. gingivalis* upregulates *ZEB1* in gingival epithelial cells

We investigated the impact of *P. gingivalis* on *ZEB1* expression in TIGK cells using qRT-PCR and immunoblotting. As shown in Figure 1A, *P. gingivalis* increased *ZEB1* mRNA levels in a time and dose dependent manner, with maximal induction occurring after 24 h infection with a MOI of 100. An increase in the amount of *ZEB1* protein was also observed at 24 h following *P. gingivalis* infection at both MOI 50 and 100 (Fig 1B). As *P. gingivalis* infections of oral tissue are chronic conditions, we further examined *ZEB1* activity 72 h after *P. gingivalis* infection. MOIs of 1, 10 and 50 were used as at MOI 100 the proteases of *P. gingivalis* can cause detachment of cells from the substratum. While an MOI 1 did not affect *ZEB1* expression, mRNA levels were increased by *P. gingivalis* at MOI 10 and 50 (Figure 1C). The ability of *P. gingivalis* at MOI 10 to increase *ZEB1* expression after 72 h, but not earlier, indicates that infection of epithelial cells with low numbers of the organism has the potential to elevate *ZEB1* over extended times, possibly as a result of intracellular *P. gingivalis* replication and cell to cell spread (Lamont *et al.*, 1995, Yilmaz *et al.*, 2006). To corroborate the nuclear location of the *ZEB1* transcription factor following *P. gingivalis* infections, TIGKs were examined by CLSM with quantitative image analysis (Figures 1D and E). After *P. gingivalis* infection there was increased expression of *ZEB1* protein in the nucleus where it is functionally active.

### *ZEB1* responses to *P. gingivalis* are strain and fimbriae dependent

*P. gingivalis* is a host adapted organism with a nonclonal population structure, and isolates from different individuals often vary considerably (Tribble *et al.*, 2013). Hence, we next examined the ability of different strains of *P. gingivalis* to enhance *ZEB1* mRNA levels. As shown in Figure 1F, an additional ATCC strain (49417) and two low passage clinical isolates (11029 and 10512) induced *ZEB1* expression to a similar degree as the type strain 33277. In contrast, the commonly used laboratory strain W83 did not significantly increase *ZEB1*

expression. One of the major differences among *P. gingivalis* strains is in the expression of fimbriae (Nadkarni *et al.*, 2014). The two ATCC strains, along with the two low passage isolates, all expressed FimA, the structural subunit protein of the major fimbriae (Supporting Information Figure S1). Strain W83 does not express FimA (Nishikawa and Duncan, 2010), which prompted us to speculate that FimA may be an effector protein for ZEB1 induction. This concept was corroborated by the failure of an isogenic *fimA* mutant of 33277 to increase ZEB1 mRNA levels (Figure 1G). We also found that induction of *ZEB1* expression required direct contact between *P. gingivalis* and epithelial cells (Figure 1G), consistent with a role for the FimA adhesin. Fimbriated *P. gingivalis* activate JNK signaling (Watanabe *et al.*, 2001) which has been reported to increase transcription of ZEB1 (Zhang *et al.*, 2012). However, siRNA knockdown of JNK in TIGK cells did not impede the ability of *P. gingivalis* to upregulate ZEB1 (Supporting Information Figure S2). In addition, pharmacological inhibition of Akt with LY294002 also failed to reduce *P. gingivalis*-mediated ZEB1 induction (Supporting Information Figure S3). Hence the signaling pathways activated by *P. gingivalis* fimbriae that converge on Zeb1 remain to be determined, and this topic is under active investigation in our laboratory.

The FimA fimbriae are required for maximal invasion of *P. gingivalis* into gingival epithelial cells (Lamont and Jenkinson, 1998). However, invasion per se was not required for ZEB1 induction as a mutant of *P. gingivalis* that is invasion-deficient, due to disruption of the gene encoding the serine phosphatase SerB (Takeuchi *et al.*, 2013), retained the ability to upregulate ZEB1 (Supporting Information Figure S4). Nonetheless, the spatial definition of *P. gingivalis* initiation of ZEB1 activation requires further study. Moreover, when *P. gingivalis* was separated from the epithelial cells by a semi-permeable membrane, *ZEB1* levels were lower than the control uninfected condition, indicating that there may be components secreted by *P. gingivalis* that can antagonize *ZEB1* regulation in the absence of FimA mediated contact. Thus, multiple effectors of *P. gingivalis* may be capable of impacting *ZEB1* expression, with the effect of whole cells representing the collective output of several distinct interactions and signaling pathways.

### ***P. gingivalis* communities regulate ZEB1 expression**

On mucosal surfaces bacteria rarely exist as monospecies accumulations but rather as complex multispecies communities. *P. gingivalis* engages in synergistic community formation with *S. gordonii* and *F. nucleatum*, common inhabitants of the oral microbiota, and *in vivo* these organisms can be found in close association (Benitez-Paez *et al.*, 2014, Valm *et al.*, 2011, Wright *et al.*, 2014, Hendrickson *et al.*, 2014). Individually, neither *S. gordonii* nor *F. nucleatum* were capable of regulating ZEB1 expression, indicating that of these three widespread oral species, *P. gingivalis* has the most potential to effect an EMT through ZEB1 (Figure 2). Importantly, *P. gingivalis* remained effective at elevating ZEB1 mRNA in the context of a community with either *S. gordonii* or *F. nucleatum*, consistent with recent reports demonstrating that a community of *P. gingivalis* and *F. nucleatum* can promote tumor progression in animal models (Gallimidi *et al.*, 2015). Thus, the tumorigenic properties of *P. gingivalis* can prevail in the presence of co-colonizing organisms, an important principle for *in vivo* relevance.

### ***P. gingivalis* upregulates ZEB1 promoter activity and downregulates miR200**

An increase in the amount of steady state mRNA levels can result from an increase in transcription or a decrease in degradation. To begin to distinguish between these possibilities, we first examined transcriptional activity from the *ZEB1* promoter using a series of *ZEB1* upstream regulatory regions promoting transcription of the *Luc* gene. These human *ZEB1* promoter constructs contain sequences important for regulation of ZEB1 expression in several cell types (Manavella *et al.*, 2007, Liu *et al.*, 2007). *P. gingivalis* stimulated the activity of all of these promoter constructs (Figure 3A), indicating that the increase of ZEB1 mRNA induced by *P. gingivalis* can occur through an elevated transcription rate. These data also localize the response element(s) within the first 400 bp of the promoter. An additional mechanism by which ZEB1 is controlled posttranscriptionally is through the action of the miR-200 family of microRNAs (Brabletz and Brabletz, 2010). miR-200 family members target conserved recognition sites on the 3' UTR of ZEB1 mRNA (Brabletz and Brabletz, 2010), and thus a decrease in miR-200 leads to higher levels of ZEB1 mRNA. However, we did not observe a reduction in the amount of miR-200b, miR-200c or miR-205 in cells at 6 h after *P. gingivalis* infection when levels of ZEB1 mRNA begin to rise (Figure 3B–D). Indeed there was a slight increase in miR-200 family expression, indicating that the increase in ZEB1 levels at 6 h is not the consequence of decreased miR-200 expression. The action of ZEB1, in turn, represses the transcription of the miR-200 family, and consistent with this at 24 and 48 h after *P. gingivalis* infection, miR-200b, miR-200c and miR-205 levels were reduced. A control microRNA, miR-21, which is not involved in *ZEB1* feedback regulation, did not show a significant decrease in expression (Supporting Information Figure S5). Hence, the pattern of miRNA expression is consistent with the results from the promoter-reporter constructs in pointing toward increased mRNA synthesis as the initial cause of the elevated levels of steady state mRNA for ZEB1.

### **Changes in epithelial and mesenchymal marker expression upon *P. gingivalis* infection**

The expression pattern of ZEB1 targets in TIGKs infected with *P. gingivalis* was assessed by qRT-PCR (Table 1). Mesenchymal markers N-cadherin, vimentin and matrix metalloproteinase (MMP)-9 were upregulated at 24 h after infection by *P. gingivalis* at MOI 50 and 100, while fibronectin levels were increased by *P. gingivalis* at MOI 100. An increase in vimentin protein expression was confirmed by immunoblotting (Supporting Information Figure S6), and elevated MMP-9 amounts following infection with fimbriated *P. gingivalis* was corroborated by zymography (Figure 4). While both pro and active forms of MMP-9 were increased by *P. gingivalis* wild type, there was no difference in the ratio of active MMP-9 to total (cleaved and pro-MMP9) between parental and fimbrial deficient mutant strains. Under these infection conditions, therefore, fimbriated *P. gingivalis* elevate the amount of MMP-9 produced by TIGK cells but do not modulate MMP-9 activation. In contrast expression of MMP-2, which may be more predominantly regulated by Twist (Yang *et al.*, 2013), was not impacted by *P. gingivalis* infection. Of the epithelial markers tested, collagen 1 (COL1A1) and cytokeratin 19 (KRT19) were suppressed by *P. gingivalis*. Collectively, these results support the concept that infection by *P. gingivalis* can contribute to the process of transition toward a mesenchymal phenotype. Although the mesenchymal marker integrin  $\alpha 5$  (ITGA5) and the epithelial marker cytokeratin 7 (KRT7) were unaffected

by *P. gingivalis*, variability in expression of important cell proteins is not unexpected as control of expression by ZEB1 is cell and context dependent (Lamouille *et al.*, 2014). One of the major targets of ZEB1 is E-cadherin, and ZEB1 mediated repression of E-cadherin, with associated disruption of E-cadherin dependent junctions, is an important marker of EMT. We did not observe differential regulation of E-cadherin following *P. gingivalis* infection (not shown). However, the gingival epithelium is highly porous with only sparse interconnections (Bosshardt and Lang, 2005) and expression of E-cadherin is very low (Heymann *et al.*, 2001). Thus a reduction of E-cadherin may not be as important for the EMT of gingival epithelial cells as in other cell types.

To corroborate the role ZEB1 in the differential regulation of mesenchymal markers, siRNA mediated knockdown was performed. Reduction of ZEB1 mRNA and protein following siRNA transfection was confirmed by qRT-PCR and immunoblotting, respectively (Figure 5A,B). TIGKs with diminished ZEB1 expression were then infected with *P. gingivalis* MOI 50 or 100 over 24 h. As shown in Figure 5C and D, ZEB1 deficiency prevented *P. gingivalis* induced modulation of expression of vimentin and MMP-9.

### ***P. gingivalis* promotes migration of epithelial cells**

Cells that acquire an EMT phenotype display an invasive behavior *in vitro*, and thus we tested the ability of *P. gingivalis* to increase the migration of TIGK cells into matrigel. Figure 6 shows that *P. gingivalis* infection resulted in a greater than 2-fold increase in TIGK cell invasion into the gel compared to control cells. Knockdown of ZEB1 prevented *P. gingivalis*-induced TIGK cell migration, verifying the importance of ZEB1 in this aspect of the *P. gingivalis*-dependent partial mesenchymal phenotype.

### ***P. gingivalis* elevates ZEB1 levels *in vivo***

To determine whether *P. gingivalis*' ability to increase ZEB1 levels also occurs *in vivo*, mice were orally infected with *P. gingivalis*, and gingival tissue recovered 1, 3 and 8 days following the final inoculation. Levels of *P. gingivalis* on the gingival tissues were determined by qPCR (Supporting Information Figure S7) and remained constant over the 8-day period. As shown in Figure 7, colonization with *P. gingivalis* induced an increase in gingival tissue expression of ZEB1 mRNA over 8 days compared to sham infected animals. Thus, *P. gingivalis* has the potential to stimulate ZEB1 and contribute to an EMT in an animal model.

### **Presence of *P. gingivalis* in human OSCC**

*P. gingivalis* could exacerbate carcinogenesis at several stages through its ability to increase ZEB1 expression, but only if the bacteria are physically associated with the developing cancer. We investigated whether *P. gingivalis* bacteria are present within oral squamous cell carcinoma biopsy samples. Immunofluorescence microscopy with a specific polyclonal *P. gingivalis* antiserum labeled discrete speckles in the cells of a poorly differentiated OSCC sample and a carcinoma in situ, whereas antibodies to *S. gordonii* showed little or no labeling (Figure 8A–B). Similar results were seen with two carcinoma in situ cases and two poorly differentiated carcinomas. Confocal microscopy more clearly detected the particles which were fluorescently labeled by the *P. gingivalis* antibodies. Serial optical sections were

taken at 0.4 microns, and individual particles were found to persist in 5 to 7 adjacent optical slices (Supporting Information Figure S8). This estimates the fluorescent particles to be 2.0 to 2.8 microns in size, consistent with intact *P. gingivalis*. As described in primary gingival epithelial cells, *P. gingivalis* was observed in both the cytoplasm and in the nuclei (Belton *et al.*, 1999).

## Discussion

Typically, the gingival epithelium provides a major physical barrier to oral pathogens. Disruption of the gingival barrier by inducing an EMT may enhance the ability of *P. gingivalis* to invade the tissue and enhance access to nutrients derived from inflammatory tissue breakdown (Hajishengallis, 2014). Hence, up-regulation of ZEB1 by *P. gingivalis* can be seen as providing an evolutionary advantage to the organism. Beyond this, the ability of *P. gingivalis* to stimulate ZEB1 expression could have several distinct clinically relevant effects. ZEB1 influences multiple stages of carcinogenesis, including the initial transformation, progression, EMT leading to metastasis, and resistance to therapy (Sanchez-Tillo *et al.*, 2012, Zhang *et al.*, 2014). Therefore, the presence of *P. gingivalis*, interacting with other environmental effectors, may enhance the initiation of oral cancer within the pre-cancerous field, or increase carcinogenic progression.

The ability to manipulate ZEB1 location and function constitutes an important attribute of bacteria with a potential role in carcinogenesis. *P. gingivalis* is a keystone member of dysbiotic oral communities, which in combination with its ability to spread systemically, and enhance cell survival and proliferation, supports epidemiological evidence of an association with cancers such as OSCC (Whitmore and Lamont, 2014, Lamont and Hajishengallis, 2015). Moreover, in established invasive OSCC lines, *P. gingivalis* activates the ERK1/2-Ets1, p38/HSP27, and PAR2/NF- $\kappa$ B pathways to promote cellular invasion (Inaba *et al.*, 2014). The results of the present study indicate the *P. gingivalis* may induce nontransformed gingival epithelial cells to undergo a partial EMT through the upregulation of ZEB1. We show that *P. gingivalis* increases the transcriptional activity of the ZEB1 gene and increases ZEB1 protein levels in the nucleus. Infection of epithelial cells with *P. gingivalis* upregulated expression of genes associated with the mesenchymal phenotype and knockdown of ZEB1 attenuated this effect. *P. gingivalis* also induced a migratory phenotype in epithelial cells which was ZEB1-dependent. Oral infection of mice with *P. gingivalis* stimulated ZEB1 expression in the gingival tissues and biopsy tissue from human OSCC carcinoma in situ and poorly differentiated cancer showed the presence of *P. gingivalis*.

On the hard and soft tissues of the oral cavity *P. gingivalis* is an inhabitant of multispecies communities. Organisms such as *S. gordonii* and *F. nucleatum* provide mutual physiological support, and *P. gingivalis* in the context of a community is phenotypically distinct from single species accumulations (Wright *et al.*, 2013). In addition, infection of epithelial cells with the early colonizing streptococci can reprogram specific signaling pathways such that they do not respond to the later colonizing *P. gingivalis* (Handfield *et al.*, 2005). We found here that while neither *S. gordonii* nor *F. nucleatum* modulated ZEB1 mRNA levels, combinations of *P. gingivalis* with either species retained the capacity to upregulate ZEB1. As porphyromonads, fusobacteria and streptococci are all found in higher numbers on the

surfaces of OSCC compared to contiguous healthy mucosa (Nagy *et al.*, 1998), it is likely therefore that these microbial communities contribute to the EMT.

*P. gingivalis* strains exhibit extensive genetic variation as a result of genomic rearrangements (Naito *et al.*, 2008), and horizontal gene transfer is considered an adaptive strategy for long term survival in the oral environment (Nadkarni *et al.*, 2014, Tribble *et al.*, 2007). Most strains of *P. gingivalis* express fimbriae comprised of FimA major fimbrial subunit proteins, although in some strains, such as the commonly used lab strain W83, FimA production is very low due to a mutation in the FimS histidine kinase component of the FimS/FimR TCS that controls transcription of the *fimA* operon (Nishikawa and Duncan, 2010). Results with a variety of strains and isogenic mutants of *P. gingivalis* indicate that FimA is the effector of *P. gingivalis* responsible for upregulation of ZEB1. FimA is a major antigen on the *P. gingivalis* surface and can also incite the production of proinflammatory cytokines (Lamont and Jenkinson, 1998, Bostanci and Belibasakis, 2012). FimA is capable of manipulating a number of signal transduction pathways and transcription factors in different cell types (Zhou and Amar, 2007, Hajishengallis *et al.*, 2012), and the processes that lead to increased ZEB1 promoter activity require further study. The potential importance of FimA expressing *P. gingivalis* lineages in the events that can lead to tumor development is corroborated by the role of this protein in the acceleration of the epithelial cell cycle (Kuboniwa *et al.*, 2008). FimA fimbriae, which do not share significant homology to other fimbrial proteins (Enersen *et al.*, 2013), may thus constitute an attractive target for novel biomarkers or therapeutics.

ZEB1 can be regulated at the transcriptional level and posttranscriptionally regulated by the miR-200 family through a double negative feedback loop (Brabletz and Brabletz, 2010). In epithelial cells miR-200s inhibit ZEB expression and maintain the epithelial phenotype. By contrast, in mesenchymal cells elevated ZEB activity suppresses expression of the miR-200s. Our results show that the increased levels of ZEB1 were associated with a reduction in the amounts of the miR200 family, potentially facilitating a stable transition to the partial mesenchymal phenotype. The initial upregulation of *ZEB1* in epithelial cells, however, was not associated with a decrease in the amounts of the miR200 family, but with an increase in ZEB1 promoter activity. NF- $\kappa$ B activation also promotes ZEB1 transcription (Vandewalle *et al.*, 2009); however, in epithelial cells *P. gingivalis* suppresses the activation of NF- $\kappa$ B by dephosphorylation of the p65 subunit at the S536 residue (Takeuchi *et al.*, 2013). It is unlikely, therefore, that NF- $\kappa$ B is involved in *P. gingivalis*-induced ZEB1 upregulation.

*P. gingivalis* induced expression of the mesenchymal markers and decreased expression of the epithelial markers. siRNA knockdown of ZEB1 abrogated the ability of *P. gingivalis* to regulate epithelial and mesenchymal gene expression, establishing ZEB1 as a major transcriptional effector of the *P. gingivalis*-induced partial EMT. Epithelial markers down regulated by *P. gingivalis* included cytokeratin 19 which is characteristically expressed in cells of the junctional epithelium within the gingival tissues, and has also been reported to be suppressed in OSCC (Khanom *et al.*, 2012). The mesenchymal relevant genes induced by *P. gingivalis* included N-cadherin, vimentin, fibronectin and MMP-9. N-cadherin is a calcium dependent cell-cell adhesion glycoprotein, which is upregulated in EMT and some studies



have found associated with OSCC (Zhao *et al.*, 2012). Vimentin is a cytoskeletal intermediate filament protein involved in maintaining cell shape and stabilizing cytoskeletal interactions. Expression of vimentin is associated with OSCC tumorigenesis (Lee *et al.*, 2015), and vimentin has been proposed as a predictor of the malignant potential of high risk oral lesions (Sawant *et al.*, 2014). Fibronectin is a component of the cell matrix involved in cell migration processes including metastasis, and expression of alternatively spliced segments of fibronectin is related to OSCC tumorigenesis (Kamarajan *et al.*, 2010). MMP-9 is secreted as inactive proproteins which are activated by proteolytic cleavage. As a gelatinase, MMP-9 can degrade collagen IV in the basement membrane and extracellular matrix facilitating tumor growth, invasion, metastasis, and angiogenesis (Westermarck and Kahari, 1999). MMP-9 plays a crucial role in the development of several human malignancies, including OSCC (Kruger *et al.*, 2005, Bedal *et al.*, 2014). Moreover, epithelial cells infected with *P. gingivalis* showed ZEB1-dependent increased migration into matrigel, a phenotype consistent with increased MMP-9 activity and with an overall partial mesenchymal phenotype.

To begin to translate our results from reductionist *in vitro* models to the *in vivo* situation, we orally infected mice with *P. gingivalis* and examined ZEB1 expression in gingival tissues. Although *P. gingivalis* is not a normal member of the mouse oral microbiota, it does colonize transiently and causes alveolar bone loss (Hajishengallis *et al.*, 2015). Our results show for the first time that *P. gingivalis* colonization of the gingival tissues *in vivo* leads to upregulation of ZEB1. Further *in vivo* evidence for a role of *P. gingivalis* in oral tumor development was provided by IF analysis of OSCC biopsy samples. Antigenic based detection of *P. gingivalis* within biopsy samples from OSCC poorly differentiated cancer and carcinoma in situ corroborates a similar study in which *P. gingivalis* antigens were detected in ten gingival squamous cell carcinomas of differing degrees of differentiation (Katz *et al.*, 2011). The use of optical sectioning in the current study established that the size of particles detected by *P. gingivalis* antibodies was in the 2–3  $\mu\text{m}$  range, consistent with whole organisms, rather than shed antigens or outer membrane vesicles. Intimate association of *P. gingivalis* with OSCC lesions shows that the organism has the opportunity as well as the capability, to contribute to EMT *in vivo*.

Oral cancers are among the most prevalent (Jemal *et al.*, 2008), and despite considerable advances in diagnosis and therapeutic options, the 5-year survival rate has remained stable at approximately 50% among all tumor stages during the past decades (Wikner *et al.*, 2014). The early phase of OSCC is often asymptomatic, therefore the identification of both novel biomarkers and contributing etiological agents is important for improving survival rates. The results of the current study suggest that infection with FimA-positive *P. gingivalis* can induce a ZEB1 dependent partial EMT. The detection of *P. gingivalis*, or of FimA, in early erythroplakia or leukoplakia lesions, therefore, may have utility for the early detection of lesion likely to progress to malignant status.

## Materials and Methods

### Bacterial strains, eukaryotic cells, and growth and infection conditions

*Porphyromonas gingivalis* strain ATCC 33277 and its isogenic mutant *fimA*, strains ATCC 49417, W83, and low passage clinical isolates 11029, 10512 (laboratory strains), were cultured in trypticase soy broth (TSB) supplemented with yeast extract (1 mg/ml), hemin (5 µg/ml) and menadione (1 µg/ml). Tetracycline (1 µg/ml) was incorporated into the medium for the growth of *fimA*. *Fusobacterium nucleatum* ATCC 25586 was cultured in brain heart infusion (BHI) broth supplemented with hemin (5 µg/ml) and menadione (1 µg/ml). *Streptococcus gordonii* DL1 was grown in BHI supplemented with yeast extract (5 µg/ml). All bacterial strains were cultured anaerobically at 37°C. Human telomerase immortalized keratinocytes (TIGKs) derived from gingival epithelium were maintained at 37°C and 5% CO<sub>2</sub> in Dermalife-K serum free culture medium (Lifeline Cell Technology, Carlsbad, CA) as described (Moffatt-Jauregui *et al.*, 2013). TIGKs between passages 10 and 20 at 70% confluence were stimulated with bacteria as described for individual experiments. For transwell (Corning, Corning NY) assays, TIGKs were cultured in the lower compartment and *P. gingivalis* added to the upper chamber.

### Immunoblotting

TIGK cells were solubilized in cold cell lysis reagent (Cell Signaling, Danvers, MA) containing Protease Inhibitor and PhosSTOP Phosphatase Inhibitor (Roche, Indianapolis, IN). Proteins (40 ng) were separated by 10% SDS-polyacrylamide gel electrophoresis, blotted onto a PVDF membrane and blocked in 5% BSA in TBS with 0.1% Tween20. Blots were probed at 4°C overnight with primary antibodies followed by 1 h with HRP-conjugated secondary antibody at room temperature. Antigen-antibody binding were detected using ECL Substrate (Thermoscientific, Hudson NH). Primary antibodies targeted ZEB1 (Novus, Littleton, CO) or vimentin (Cell Signaling). Duplicate blots were probed with GAPDH antibodies (Cell Signaling) as a loading control

### RNA extraction and quantitative reverse transcription-PCR (qRT-PCR)

Total RNA from TIGK cells and from homogenized gingival tissue was isolated and purified with PerfectPure RNA kit (5Prime, Gaithersburg, MD). miRNA was isolated and purified from TIGKs with PureLink miRNA isolation kit (Invitrogen, Carlsbad, CA). RNA concentrations were determined by spectrophotometry (NanoDrop Technology, Wilmington, DE). cDNA from total RNA and miRNA was synthesized (2 µg RNA/reaction volume) using a High Capacity cDNA Reverse Transcription kit and a TaqMan MicroRNA Reverse Transcription kit (Applied Biosystems, Grand Island, NY), respectively. Real time RT-PCRs used TaqMan Fast universal PCR mastermix and gene expression assays for Zeb1, vimentin, MMP-9, ITGA5, fibronectin, KRT7, COL-1A1 and GAPDH. Negative RT reactions were included in each assay. TaqMan microRNA assays were used to quantify the mature miRNA-200b, mi-RNA-200c, miRNA-205, miRNA-21 and RNU48. Real time qPCR was performed on an Applied Biosystems StepOne Plus cyler with StepOne software V2.2.2 and autocalculated threshold cycle selected. The cycle threshold (C<sub>t</sub>) values were determined, mRNA and miRNA expression levels were normalized to GAPDH and RNU48, respectively, and expressed relative to controls following the 2<sup>-CT</sup> method.

### Luciferase reporter assay

TIGK cells were transfected with ZEB1 promoter constructs: Z1p 1000-Luc (–912 bp to +43 bp of the ZEB1 gene), Z1p.400-Luc (–367 bp to +43 bp) and Z1p.196-Luc (–212 bp to +43 bp); at  $2 \mu\text{g}/10^5$  cells using FuGENE6 Transfection Reagent (Promega, Madison, WI). Following 48 h in transfection media, cells were returned to TIGK medium for further 24 h, prior to the stimulation with *P. gingivalis*. Cells were lysed and the reporter activity was determined with the Dual-Glo Luciferase Assay System (Promega). Firefly luciferase activity was normalized on the basis of Renilla luciferase activity in the same samples.

### Zymography

The activities of MMP2 and MMP9 in culture supernatant collected from control uninfected and *P. gingivalis*-infected TIGK cells, were determined using gelatin zymography as described (Inaba *et al.*, 2014). Samples were mixed with SDS sample buffer without reducing reagents, then separated on 10% SDS-polyacrylamide gels containing 0.1% gelatin. The gels were incubated at 37°C with 2.5% Triton X-100 for 1 h, and then in 20 mM Tris-HCl (pH 7.5) containing 200 mM NaCl and 5 mM  $\text{Ca Cl}_2$  for 48 h. After staining with 5% Coomassie Brilliant Blue R-250, gelatinolytic activities were visualized as clear bands against a blue background and quantified using ImageJ.

### RNA interference

TIGKs were transfected with predesigned 30 nM siRNA (siGENOME SMARTpool siRNA) targeting ZEB1 or control siRNA (GE Healthcare Dharmacon, Lafayette, CO) for 24 h using LipoJet (SigmaGen, Gaithersburg, MD) transfection reagent. At 48 h after transfection, the medium was replaced and cells were incubated for a further 24 h prior to infection.

### Immunofluorescence and confocal laser scanning microscopy

TIGK cells were grown on glass coverslips, washed twice in phosphate-buffered saline (PBS) and fixed with 4% paraformaldehyde for 10 min. Permeabilization was with 0.2% TritonX-100 for 10 min at room temperature prior to blocking in 10% goat serum for 20 min. ZEB1 was detected by reacting with primary antibodies at 1:100 overnight at 4°C, followed by Alexa Fluor 488-conjugated anti-rabbit secondary antibodies (Life Technologies) at 1:200 for 3 h in the dark. Following a 20 min blocking in 0.1% goat serum, actin was labeled with Texas Red-phalloidin (Life Technologies) for 2 h at room temperature in the dark. Coverslips were mounted with on glass slides using ProLong Gold with DAPI (4'-6-diamidino-2-phenylindole) mounting medium (Invitrogen) prior to imaging with a Leica SP8 confocal inverted fluorescence microscope. Images were analyzed using Volocity 6.3 Software (PerkinElmer, Waltham, MA).

### Matrigel invasion assay

Cell motility was measured by assessment of the migration rate of TIGKs using a BD BioCoat Matrigel Invasion Chamber (BD Biosciences, San Jose, CA). Cells ( $2 \times 10^5$ ) were plated on transwell filters coated with matrigel. The lower compartment of the invasion chambers contained cell culture medium. After 18 h incubation at 37°C, cells remaining on the upper surface of the filter were removed, and the cells that migrated through the filter

were fixed with 1% methanol and stained with toluidine blue. Cells were enumerated from three random 20× fields for each filter using a Nikon Eclipse TS100 microscope.

### Animal infection

BALB/c mice were orally infected with  $10^7$  cfu *P. gingivalis* 33277 five times at 2-day intervals as described previously (Daep *et al.*, 2011) and approved by the University of Louisville Institutional Animal Care and Use Committee. The levels of *P. gingivalis* colonization were determined by qPCR with the *P. gingivalis* 16SrRNA gene as described (Daep *et al.*, 2011). On days 1, 3 and 8 after the last infection, mice were euthanized and the upper and lower jaw with gingival tissue were recovered. After RNA extraction, the ratio of ZEB1 mRNA to GAPDH mRNA for each mouse was determined by qRT-PCR using the 2-CT method.

### Human oral tissue immunofluorescent and immunohistochemical staining

Paraffin embedded human tongue biopsy samples were sectioned at 4  $\mu$ m, dewaxed and rehydrated. The slides were blocked with 10% goat serum for 1 h, and reacted with *P. gingivalis* 33277 or *S. gordonii* antibodies 1:1000 for 2 h at room temperature. Secondary Alexa Fluor 488 conjugated anti-rabbit antibody (1:500) was for 1 h, following which slides were treated with DAPI (1:2000). Slides were mounted with VectaShield (Vector Labs, Burlingame, CA) and imaged with a Zeiss Axiocam MRc5 fluorescence microscope. Procedures were approved by the University of Louisville Institutional Review Board.

### Statistical analyses

Statistical analyses were conducted using the GraphPad Prism software. Data were evaluated by one-way ANOVA with Tukey's multiple comparison test. Experimental data presented are representative of at least three biological replicates.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

### Acknowledgments

Grant Sponsors: We thank the NIH/NIDCR for support through DE011111, DE017921, DE016690 and DE023193.

### References

- Baud J, Varon C, Chabas S, Chambonnier L, Darfeuille F, Staedel C. *Helicobacter pylori* initiates a mesenchymal transition through ZEB1 in gastric epithelial cells. PLoS One. 2013; 8:e60315. [PubMed: 23565224]
- Bedal KB, Grassel S, Oefner PJ, Reinders J, Reichert TE, Bauer R. Collagen XVI induces expression of MMP9 via modulation of AP-1 transcription factors and facilitates invasion of oral squamous cell carcinoma. PLoS One. 2014; 9:e86777. [PubMed: 24466237]
- Belton CM, Izutsu KT, Goodwin PC, Park Y, Lamont RJ. Fluorescence image analysis of the association between *Porphyromonas gingivalis* and gingival epithelial cells. Cell Microbiol. 1999; 1:215–223. [PubMed: 11207554]
- Benitez-Paez A, Belda-Ferre P, Simon-Soro A, Mira A. Microbiota diversity and gene expression dynamics in human oral biofilms. BMC Genomics. 2014; 15:311. [PubMed: 24767457]


- Bessede E, Staedel C, Acuna Amador LA, Nguyen PH, Chambonnier L, Hatakeyama M, et al. *Helicobacter pylori* generates cells with cancer stem cell properties via epithelial-mesenchymal transition-like changes. *Oncogene*. 2014; 33:4123–4131. [PubMed: 24096479]
- Bosshardt DD, Lang NP. The junctional epithelium: from health to disease. *J Dent Res*. 2005; 84:9–20. [PubMed: 15615869]
- Bostanci N, Belibasakis GN. *Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen. *FEMS Microbiol Lett*. 2012; 333:1–9. [PubMed: 22530835]
- Brabletz S, Brabletz T. The ZEB/miR-200 feedback loop--a motor of cellular plasticity in development and cancer? *EMBO Rep*. 2010; 11:670–677. [PubMed: 20706219]
- Castellarin M, Warren RL, Freeman JD, Dreolini L, Krzywinski M, Strauss J, et al. *Fusobacterium nucleatum* infection is prevalent in human colorectal carcinoma. *Genome Res*. 2012; 22:299–306. [PubMed: 22009989]
- Daep CA, Novak EA, Lamont RJ, Demuth DR. Structural dissection and in vivo effectiveness of a peptide inhibitor of *Porphyromonas gingivalis* adherence to *Streptococcus gordonii*. *Infect Immun*. 2011; 79:67–74. [PubMed: 21041492]
- Enersen M, Nakano K, Amano A. *Porphyromonas gingivalis* fimbriae. *J Oral Microbiol*. 2013; 5 doi: 10.3402.
- Gallimidi AB, Fischman S, Revach B, Bulvik R, Maliutina A, Rubinstein AM, et al. Periodontal pathogens *Porphyromonas gingivalis* and *Fusobacterium nucleatum* promote tumor progression in an oral-specific chemical carcinogenesis model. *Oncotarget*. 2015; 6:22613–22623. [PubMed: 26158901]
- Garrett WS. Cancer and the microbiota. *Science*. 2015; 348:80–86. [PubMed: 25838377]
- Gheldorf A, Hulpiau P, van Roy F, De Craene B, Berx G. Evolutionary functional analysis and molecular regulation of the ZEB transcription factors. *Cell Mol Life Sci*. 2012; 69:2527–2541. [PubMed: 22349261]
- Gur C, Ibrahim Y, Isaacson B, Yamin R, Abed J, Gamliel M, et al. Binding of the Fap2 protein of *Fusobacterium nucleatum* to human inhibitory receptor TIGIT protects tumors from immune cell attack. *Immunity*. 2015; 42:344–355. [PubMed: 25680274]
- Hajishengallis G. The inflammophilic character of the periodontitis-associated microbiota. *Mol Oral Microbiol*. 2014; 29:248–257. [PubMed: 24976068]
- Hajishengallis G, Krauss JL, Liang S, McIntosh ML, Lambris JD. Pathogenic microbes and community service through manipulation of innate immunity. *Adv Exp Med Biol*. 2012; 946:69–85. [PubMed: 21948363]
- Hajishengallis G, Lamont RJ, Graves DT. The enduring importance of animal models in understanding periodontal disease. *Virulence*. 2015; 6:229–235. [PubMed: 25574929]
- Handfield M, Mans JJ, Zheng G, Lopez MC, Mao S, Progulsk-Fox A, et al. Distinct transcriptional profiles characterize oral epithelium-microbiota interactions. *Cell Microbiol*. 2005; 7:811–823. [PubMed: 15888084]
- Hendrickson EL, Wang T, Beck DA, Dickinson BC, Wright CJ, Lamont RJ, Hackett M. Proteomics of *Fusobacterium nucleatum* within a model developing oral microbial community. *Microbiologyopen*. 2014; 3:729–751. [PubMed: 25155235]
- Heymann R, Wroblewski J, Terling C, Midtvedt T, Obrink B. The characteristic cellular organization and CEACAM1 expression in the junctional epithelium of rats and mice are genetically programmed and not influenced by the bacterial microflora. *J Periodontol*. 2001; 72:454–460. [PubMed: 11338297]
- Inaba H, Sugita H, Kuboniwa M, Iwai S, Hamada M, Noda T, et al. *Porphyromonas gingivalis* promotes invasion of oral squamous cell carcinoma through induction of proMMP9 and its activation. *Cell Microbiol*. 2014; 16:131–145. [PubMed: 23991831]
- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, Thun MJ. Cancer statistics, 2008. *CA Cancer J Clin*. 2008; 58:71–96. [PubMed: 18287387]
- Jia B, Liu H, Kong Q, Li B. Overexpression of ZEB1 associated with metastasis and invasion in patients with gastric carcinoma. *Mol Cell Biochem*. 2012; 366:223–229. [PubMed: 22466758]

- Kamarajan P, Garcia-Pardo A, D'Silva NJ, Kapila YL. The CS1 segment of fibronectin is involved in human OSCC pathogenesis by mediating OSCC cell spreading, migration, and invasion. *BMC Cancer*. 2010; 10:330. [PubMed: 20579373]
- Katz J, Onate MD, Pauley KM, Bhattacharyya I, Cha S. Presence of *Porphyromonas gingivalis* in gingival squamous cell carcinoma. *Int J Oral Sci*. 2011; 3:209–215. [PubMed: 22010579]
- Khanom R, Sakamoto K, Pal SK, Shimada Y, Morita K, Omura K, et al. Expression of basal cell keratin 15 and keratin 19 in oral squamous neoplasms represents diverse pathophysiologicals. *Histol Histopathol*. 2012; 27:949–959. [PubMed: 22648550]
- Kim SS, Ruiz VE, Carroll JD, Moss SF. *Helicobacter pylori* in the pathogenesis of gastric cancer and gastric lymphoma. *Cancer Lett*. 2011; 305:228–238. [PubMed: 20692762]
- Kostic AD, Gevers D, Pedamallu CS, Michaud M, Duke F, Earl AM, et al. Genomic analysis identifies association of *Fusobacterium* with colorectal carcinoma. *Genome Res*. 2012; 22:292–298. [PubMed: 22009990]
- Kruger A, Arlt MJ, Gerg M, Kopitz C, Bernardo MM, Chang M, et al. Antimetastatic activity of a novel mechanism-based gelatinase inhibitor. *Cancer Res*. 2005; 65:3523–3526. [PubMed: 15867341]
- Kuboniwa M, Hasegawa Y, Mao S, Shizukuishi S, Amano A, Lamont RJ, Yilmaz O. *P. gingivalis* accelerates gingival epithelial cell progression through the cell cycle. *Microbes Infect*. 2008; 10:122–128. [PubMed: 18280195]
- Lamont RJ, Chan A, Belton CM, Izutsu KT, Vasel D, Weinberg A. *Porphyromonas gingivalis* invasion of gingival epithelial cells. *Infect Immun*. 1995; 63:3878–3885. [PubMed: 7558295]
- Lamont RJ, Hajishengallis G. Polymicrobial synergy and dysbiosis in inflammatory disease. *Trends Mol Med*. 2015; 21:172–183. [PubMed: 25498392]
- Lamont RJ, Jenkinson HF. Life below the gum line: pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol Mol Biol Rev*. 1998; 62:1244–1263. [PubMed: 9841671]
- Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol*. 2014; 15:178–196. [PubMed: 24556840]
- Lee CH, Chang JS, Syu SH, Wong TS, Chan JY, Tang YC, et al. IL-1beta promotes malignant transformation and tumor aggressiveness in oral cancer. *J Cell Physiol*. 2015; 230:875–884. [PubMed: 25204733]
- Liu Y, Costantino ME, Montoya-Durango D, Higashi Y, Darling DS, Dean DC. The zinc finger transcription factor ZFH1A is linked to cell proliferation by Rb-E2F1. *Biochem J*. 2007; 408:79–85. [PubMed: 17655524]
- Manavella PA, Roqueiro G, Darling DS, Cabanillas AM. The ZFH1A gene is differentially autoregulated by its isoforms. *Biochem Biophys Res Commun*. 2007; 360:621–626. [PubMed: 17610840]
- Mao S, Park Y, Hasegawa Y, Tribble GD, James CE, Handfield M, et al. Intrinsic apoptotic pathways of gingival epithelial cells modulated by *Porphyromonas gingivalis*. *Cell Microbiol*. 2007; 9:1997–2007. [PubMed: 17419719]
- Michaud DS. Role of bacterial infections in pancreatic cancer. *Carcinogenesis*. 2013; 34:2193–2197. [PubMed: 23843038]
- Moffatt-Jauregui CE, Robinson B, de Moya AV, Brockman RD, Roman AV, Cash MN, et al. Establishment and characterization of a telomerase immortalized human gingival epithelial cell line. *J Periodontol Res*. 2013; 48:713–721. [PubMed: 23441958]
- Moffatt CE, Lamont RJ. *Porphyromonas gingivalis* induction of microRNA-203 expression controls suppressor of cytokine signaling 3 in gingival epithelial cells. *Infect Immun*. 2011; 79:2632–2637. [PubMed: 21536793]
- Nadkarni MA, Chhour KL, Chapple CC, Nguyen KA, Hunter N. The profile of *Porphyromonas gingivalis* *kgp* biotype and *fimA* genotype mosaic in subgingival plaque samples. *FEMS Microbiol Lett*. 2014; 361:190–194. [PubMed: 25353706]
- Nagy KN, Sonkodi I, Szoke I, Nagy E, Newman HN. The microflora associated with human oral carcinomas. *Oral Oncol*. 1998; 34:304–308. [PubMed: 9813727]
- Naito M, Hirakawa H, Yamashita A, Ohara N, Shoji M, Yukitake H, et al. Determination of the genome sequence of *Porphyromonas gingivalis* strain ATCC 33277 and genomic comparison with

- strain W83 revealed extensive genome rearrangements in *P. gingivalis*. *DNA Res.* 2008; 15:215–225. [PubMed: 18524787]
- Nishikawa K, Duncan MJ. Histidine kinase-mediated production and autoassembly of *Porphyromonas gingivalis* fimbriae. *J Bacteriol.* 2010; 192:1975–1987. [PubMed: 20118268]
- Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW. *Fusobacterium nucleatum* promotes colorectal carcinogenesis by modulating E-cadherin/beta-catenin signaling via its FadA adhesin. *Cell Host Microbe.* 2013; 14:195–206. [PubMed: 23954158]
- Sahingur SE, Yeudall WA. Chemokine function in periodontal disease and oral cavity cancer. *Front Immunol.* 2015; 6:214. [PubMed: 25999952]
- Sanchez-Tillo E, Liu Y, de Barrios O, Siles L, Fanlo L, Cuatrecasas M, et al. EMT-activating transcription factors in cancer: beyond EMT and tumor invasiveness. *Cell Mol Life Sci.* 2012; 69:3429–3456. [PubMed: 22945800]
- Sawant SS, Vaidya M, Chaukar DA, Alam H, Dmello C, Gangadaran P, et al. Clinical significance of aberrant vimentin expression in oral premalignant lesions and carcinomas. *Oral Dis.* 2014; 20:453–465. [PubMed: 23865921]
- Scanlon CS, Van Tubergen EA, Inglehart RC, D’Silva NJ. Biomarkers of epithelial-mesenchymal transition in squamous cell carcinoma. *J Dent Res.* 2013; 92:114–121. [PubMed: 23128109]
- Takeuchi H, Hirano T, Whitmore SE, Morisaki I, Amano A, Lamont RJ. The serine phosphatase SerB of *Porphyromonas gingivalis* suppresses IL-8 production by dephosphorylation of NF-kappaB RelA/p65. *PLoS Pathog.* 2013; 9:e1003326. [PubMed: 23637609]
- Tribble GD, Kerr JE, Wang BY. Genetic diversity in the oral pathogen *Porphyromonas gingivalis*: molecular mechanisms and biological consequences. *Future Microbiol.* 2013; 8:607–620. [PubMed: 23642116]
- Tribble GD, Lamont RJ, Progulske-Fox A, Lamont RJ. Conjugal transfer of chromosomal DNA contributes to genetic variation in the oral pathogen *Porphyromonas gingivalis*. *J Bacteriol.* 2007; 189:6382–6388. [PubMed: 17573478]
- Valm AM, Mark Welch JL, Rieken CW, Hasegawa Y, Sogin ML, Oldenbourg R, et al. Systems-level analysis of microbial community organization through combinatorial labeling and spectral imaging. *Proc Natl Acad Sci U S A.* 2011; 108:4152–4157. [PubMed: 21325608]
- Vandewalle C, Van Roy F, Bex G. The role of the ZEB family of transcription factors in development and disease. *Cell Mol Life Sci.* 2009; 66:773–787. [PubMed: 19011757]
- Watanabe K, Yilmaz O, Nakhjiri SF, Belton CM, Lamont RJ. Association of mitogen-activated protein kinase pathways with gingival epithelial cell responses to *Porphyromonas gingivalis* infection. *Infect Immun.* 2001; 69:6731–6737. [PubMed: 11598045]
- Westermarck J, Kahari VM. Regulation of matrix metalloproteinase expression in tumor invasion. *FASEB J.* 1999; 13:781–792. [PubMed: 10224222]
- Whitmore SE, Lamont RJ. Oral bacteria and cancer. *PLoS Pathog.* 2014; 10:e1003933. [PubMed: 24676390]
- Wikner J, Grobe A, Pantel K, Riethdorf S. Squamous cell carcinoma of the oral cavity and circulating tumour cells. *World J Clin Oncol.* 2014; 5:114–124. [PubMed: 24829858]
- Wright CJ, Burns LH, Jack AA, Back CR, Dutton LC, Nobbs AH, et al. Microbial interactions in building of communities. *Mol Oral Microbiol.* 2013; 28:83–101. [PubMed: 23253299]
- Wright CJ, Xue P, Hirano T, Liu C, Whitmore SE, Hackett M, Lamont RJ. Characterization of a bacterial tyrosine kinase in *Porphyromonas gingivalis* involved in polymicrobial synergy. *Microbiologyopen.* 2014; 3:383–394. [PubMed: 24811194]
- Yang G, Yuan J, Li K. EMT transcription factors: implication in osteosarcoma. *Med Oncol.* 2013; 30:697. [PubMed: 23975634]
- Yilmaz O, Jungas T, Verbeke P, Ojcius DM. Activation of the phosphatidylinositol 3-kinase/Akt pathway contributes to survival of primary epithelial cells infected with the periodontal pathogen *Porphyromonas gingivalis*. *Infect Immun.* 2004; 72:3743–3751. [PubMed: 15213114]
- Yilmaz O, Verbeke P, Lamont RJ, Ojcius DM. Intercellular spreading of *Porphyromonas gingivalis* infection in primary gingival epithelial cells. *Infect Immun.* 2006; 74:703–710. [PubMed: 16369027]

- Zhang P, Cai Y, Soofi A, Dressler GR. Activation of Wnt11 by transforming growth factor-beta drives mesenchymal gene expression through non-canonical Wnt protein signaling in renal epithelial cells. *J Biol Chem.* 2012; 287:21290–21302. [PubMed: 22556418]
- Zhang P, Wei Y, Wang L, Debeb BG, Yuan Y, Zhang J, et al. ATM-mediated stabilization of ZEB1 promotes DNA damage response and radioresistance through CHK1. *Nat Cell Biol.* 2014; 16:864–875. [PubMed: 25086746]
- Zhao D, Tang XF, Yang K, Liu JY, Ma XR. Over-expression of integrin-linked kinase correlates with aberrant expression of Snail, E-cadherin and N-cadherin in oral squamous cell carcinoma: implications in tumor progression and metastasis. *Clin Exp Metastasis.* 2012; 29:957–969. [PubMed: 22638656]
- Zhou Q, Amar S. Identification of signaling pathways in macrophage exposed to *Porphyromonas gingivalis* or to its purified cell wall components. *J Immunol.* 2007; 179:7777–7790. [PubMed: 18025224]





**Figure 1. *P. gingivalis* up-regulates ZEB1 expression in TIGK cells in a FimA-dependent manner**

A. TIGKs were infected with *P. gingivalis* 33277 at the MOI and time indicated. ZEB1 mRNA levels were measured by qRT-PCR. Data were normalized to GAPDH mRNA and are expressed relative to noninfected (NI) controls. Results are means  $\pm$  SD; n = 3; \* P < 0.05; \*\*\* P < 0.001.

B. Immunoblot of lysates of TIGK cells infected with *P. gingivalis* 33277 for 24 h at the MOI indicated. Control cells were uninfected (NI). Duplicate blots were probed with antibodies to ZEB1 or GAPDH (loading control).


C. ZEB1 mRNA levels in TIGKs after 72 h infection with *P. gingivalis* 33277 at MOI indicated. qRT-PCR data were normalized to GAPDH mRNA and are expressed relative to noninfected (NI) controls. Results are means  $\pm$  SD, n = 3; \*\* P < 0.01; \*\*\* P < 0.001.

D. Fluorescent confocal microscopy of TIGK cells infected with *P. gingivalis* 33277 (Pg) at MOI 50 or MOI 100 for 24 h. Control cells were noninfected (NI). Cells were fixed and probed with ZEB1 antibodies (green). Actin (red) was stained with Texas Red-phalloidin, and nuclei (blue) stained with DAPI. Cells were imaged at magnification  $\times$ 63, and shown are representative merged images of projections of z-stacks obtained with Volocity software. Bar = 10  $\mu$ m.

E. Nuclear localization of ZEB1 calculated by Pearson's correlation coefficient from images in C (n=100 cells) using Volocity software. Results are mean  $\pm$  SD; \*\* P < 0.01; \*\*\* P < 0.001.

F. qRT-PCR of ZEB1 mRNA expression in TIGK cells infected with *P. gingivalis* (Pg) strains at MOI 100 for 24 h. qRT-PCR data were normalized to GAPDH mRNA and are expressed relative to noninfected (NI) controls. Results are means  $\pm$  SD, n = 3; \* P < 0.05; \*\*\* P < 0.001.

G. qRT-PCR of ZEB1 mRNA expression in TIGK cells infected with *P. gingivalis* 33277 (Pg WT), *fimA* mutant, or W83, or Pg WT in the presence of membrane insert. qRT-PCR data were normalized to GAPDH mRNA and are expressed relative to noninfected (NI) controls. Results are means  $\pm$  SD, n = 3; \*\*\* P < 0.001 compared to NI; ### P < 0.001 compared to Pg WT.




**Figure 2. *P. gingivalis* communities regulate ZEB1 expression**

A. qRT-PCR of ZEB1 mRNA expression in TIGK cells infected with *P. gingivalis* 33277 (Pg), *S. gordonii* (Sg) or a combination of Pg and Sg at MOI 100 for 24 h.

B. qRT-PCR of ZEB1 mRNA expression in TIGK cells infected with *P. gingivalis* 33277 (Pg), *F. nucleatum* (Fn) or a combination of Pg and Fn at MOI 100 for 24 h.


Data were normalized to GAPDH mRNA and are expressed relative to noninfected (NI) controls. Results are means  $\pm$  SD; n = 3; \*\* P < 0.01; \*\*\* P < 0.001.



**Figure 3. *P. gingivalis* regulates ZEB1 promoter activity and increases in ZEB1 levels are not dependent on the miRNA 200 family**


A. TIGK cells were transiently transfected with ZEB1 promoter-luciferase plasmids: Z1p 1000-Luc (−912 bp to +43 bp), Z1p.400-Luc (−367 bp to +43 bp) or Z1p.196-Luc (−212 bp to +43 bp); or with a constitutively-expressing Renilla luciferase reporter. Transfected cells were infected with *P. gingivalis* 33277 (Pg) at MOI 100 for 24 h. Control cells were noninfected (Ctr). Luciferase activity was normalized to the level of Renilla luciferase. Results are mean ± SD, n = 3; \* P < 0.01; \*\*\* P < 0.001.

B–D. Expression of mature miRNA-200b (B), miRNA-200c (C), or miRNA-205 (D) in TIGK cells infected with *P. gingivalis* 33277 MOI 100 for the times indicated. miRNA levels were measured by qRT-PCR, normalized to RNU48 miRNA, and expressed relative to noninfected (NI) controls. Results are means ± SD, n = 3; \* P < 0.05; \*\* P < 0.01; \*\*\* P < 0.001.



**Figure 4. *P. gingivalis* induces MMP9 expression in a FimA-dependent manner**

TIGKs were infected with *P. gingivalis* 33277 (WT) or *fimA* mutant at MOI 10 for 24 h, or left uninfected. A. Culture supernatants were analyzed for MMP9 and MMP2 by gelatin zymography. B. Quantitative analysis of 4 independent zymograms using ImageJ. \* P < 0.05; \*\*\* P < 0.001.




**Figure 5. ZEB1 knockdown suppresses TIGK responses to *P. gingivalis***

A. TIGK cells were transiently transfected with siRNA to ZEB1 (si Zeb1) or scrambled siRNA (si ctr). Control (ctr) cells were nontransfected. ZEB1 mRNA levels in transfected cells were measured by qRT-PCR. Data were normalized to GAPDH mRNA and are expressed relative to ctr. Results are means  $\pm$  SD,  $n = 3$ ; \*\*\*  $P < 0.001$ .


B. Immunoblot of lysates of TIGK cells transfected (as in A) and probed with antibodies to ZEB1 or GAPDH (loading control).

C–D. Transfected TIGK cells (as in A) were infected with *P. gingivalis* 33277 for 24 h at the MOI indicated. The Effect of ZEB1 knockdown on expression of vimentin (B) and MMP9 (C) was measured by qRT-PCR. Data were normalized to GAPDH mRNA and are expressed relative to the noninfected (NI) control. Results are means  $\pm$  SD,  $n = 3$ ; \*  $P < 0.05$ ; \*\*  $P < 0.01$ ; \*\*\*  $P < 0.001$  compared to NI. ###  $P < 0.001$  compared to si ctr.



**Figure 6. *P. gingivalis* increases TIGK migration in a ZEB1-dependent manner**


Quantitative analysis of TIGK migration through matrigel-coated transwells. TIGK cells were transiently transfected with siRNA to ZEB1 (si Zeb1) or scrambled siRNA (si ctr). Control cells were nontransfected. TIGKs were infected with *P. gingivalis* 33277 for 24 h at the MOI indicated. Data are presented as the average number of cells invading through inserts coated with matrigel. Results are means  $\pm$  SD, n = 3; \*\*\* P < 0.001 compared to NI. ### P < 0.001 compared to si ctr.



**Figure 7. *P. gingivalis* induces ZEB1 expression in vivo**

A. qRT-PCR of ZEB1 mRNA expression relative to GAPGH control in gingival tissues from mice infected with *P. gingivalis* or sham infected (NI). Tissue samples were collected at days 1, 3 and 8 after infection. Each point represents the determination from a single animal. \*  $P < 0.05$ ; \*\*  $P < 0.01$ .





**Figure 8. *P. gingivalis* antigens are present in OSCC**

Tissue biopsy of (A) poorly differentiated carcinoma, and (B) a tongue carcinoma in situ. Biopsy sections were stained with H&E, *P. gingivalis* 33277 antibodies (Anti-Pg 1:1000) or *S. gordonii* antibodies (Anti-Sg 1:1000) in the presence or absence of DAPI. Controls had no primary antibody. Red blood cells in the connective tissue are autofluorescent. Samples were imaged with a Zeiss AxioCam MRC5 fluorescence microscope at the magnification indicated.

**Table 1**

Changes in expression of mesenchymal and epithelial markers in TIGK cells infected with *P. gingivalis* 33277.

		Fold change induced by <i>P. gingivalis</i> 33277	
		MOI 50	MOI 100
Mesenchymal markers	Vimentin	1.72 ± 0.14 **	3.4 ± 0.41 **
	ITGA5	1.06 ± 0.12	1.29 ± 0.23
	MMP-9	8.943 ± 0.38 ***	12.77 ± 0.75 ***
	Fibronectin	1.16 ± 0.05	3.13 ± 0.16 ***
	N-cadherin	2.55 ± 0.34 **	2.4 ± 0.17 **
Epithelial markers	KRT7	0.97 ± 0.09	1.29 ± 0.09
	KRT19	1.2 ± 0.07	0.63 ± 0.003 ***
	COL-1A1	0.63 ± 0.06 **	0.67 ± 0.035 ***

Data represent qRT-PCR results of the individual genes normalized to that of GAPDH mRNA and expressed relative to noninfected cells. Results are means ± SD, n = 6;

\*\*  
P < 0.01;

\*\*\*  
P < 0.001