

RESEARCH

Open Access



Increased Caspase-6 activity in the human anterior olfactory nuclei of the olfactory bulb is associated with cognitive impairment

Benedicte Foveau¹, Steffen Albrecht², David A. Bennett³, José A. Correa⁴ and Andrea C. LeBlanc^{1,5*}

Abstract

Abnormally elevated hippocampal Caspase-6 (Casp6) activity is intimately associated with age-related cognitive impairment in humans and in mice. In humans, these high levels of Casp6 activity are initially localized in the entorhinal cortex, the area of the brain first affected by the formation of neurofibrillary tangles, according to Braak staging. The reason for the high vulnerability of entorhinal cortex neurons to neurofibrillary tangle pathology and Casp6 activity is unknown. Casp6 activity is involved in axonal degeneration, therefore, one possibility to explain increased vulnerability of the entorhinal cortex neurons would be that the afferent neurons of the olfactory bulb, some of which project their axons to the entorhinal cortex, are equally degenerating. To examine this possibility, we examined the presence of Casp6 activity, neurofibrillary tangle formation and amyloid deposition by immunohistochemistry with neopeptide antisera against the p20 subunit of active Casp6 and Tau cleaved by Casp6 (Tau Δ Casp6), phosphorylated Tau paired helical filament (PHF-1) antibodies and anti- β -amyloid antiserum, respectively, in brains from individuals with no or mild cognitive impairment and Alzheimer disease (AD) dementia. Co-localization of Casp6 activity, PHF-1 and β -amyloid was detected mostly in the anterior olfactory nucleus (AON) of the olfactory bulb. The levels of active Casp6 in the AON, which were the highest in the AD brains, correlated with PHF-1 levels, but not with β -amyloid levels. AON Tau Δ Casp6 levels correlated with entorhinal cortex Casp6 activity and PHF-1 levels. Multiple regression analyses demonstrated that AON Casp6 activity was associated with lower global cognitive function, mini mental state exam, episodic memory and semantic memory scores. These results suggest that AON Casp6 activity could lead to Casp6-mediated degeneration in the entorhinal cortex, but cannot exclude the possibilities that entorhinal cortex degeneration signals degeneration in the AON or that the pathologies occur in both regions independently. Nevertheless, AON Casp6 activity reflects that of the entorhinal cortex.

(Continued on next page)

* Correspondence: andrea.leblanc@mcgill.ca

¹Bloomfield Center for Research in Aging, Lady Davis Institute for Medical Research, Jewish General Hospital, 3755 ch. Côte Ste-Catherine, Montreal, QC, Canada

⁵Department of Neurology and Neurosurgery, McGill University, Montreal, QC, Canada

Full list of author information is available at the end of the article



(Continued from previous page)

Keywords: Alzheimer Disease, Caspase-6, Olfactory bulb, Entorhinal cortex, Hippocampus, CA1, Memory scores, Cognitive scores, Neurofibrillary tangles, Amyloid beta peptide, Mild cognitive impairment, Non-cognitively impaired

Introduction

Identification of specific early molecular mechanisms underlying neurodegeneration and leading to age-related cognitive impairment and AD dementia may provide novel therapeutic targets against which efficient therapies can be developed. Our laboratory has focused on Casp6, a cytosolic cysteinyl protease, which is almost undetectable in normal aged human brain tissues [25], but extremely abundant and active in both sporadic [27] and familial [2] AD. From early to late stages of AD, Casp6 activity is detected in neurofibrillary tangles (NFT), neuritic plaques and neuropil threads, the main pathological lesions of AD [3, 27]. The activation of Casp6 in human primary neuron cultures increases amyloid beta peptide production [32, 36, 38]. Furthermore, Casp6 activity is implicated in age-related cognitive impairment. In aged humans, higher levels of Casp6 activity in the entorhinal cortex (ERC) and Cornu Ammonis 1 (CA1) regions of the hippocampus predict lower performance in episodic memory, the first type of memory to be decreased in AD [3, 47]. Post-natal expression of a self-activated form of human Casp6 in the hippocampal CA1 region of mouse brains induces age-related spatial and episodic memory impairment and is associated with early inflammation and neuronal loss [39]. Unlike the other two effector caspases, Casp3 and Casp7, Casp6 does not induce cell death when activated [26, 33]. However, Casp6 causes axonal degeneration. Casp6 proteolytically cleaves cytoskeletal proteins crucial to neuronal integrity and function, such as alpha-tubulin, microtubule-associated protein Tau and post-synaptic density proteins regulating the actin cytoskeleton of the dendritic spines in synapses [29, 33, 56]. Casp6 is activated and associated with axonal degeneration in nerve growth factor-deprived mouse sensory neuron cultures, coordinately with cellular apoptosis generated by effector Casp3 activation [45, 52, 53, 64]. In human primary neurons transfected to overexpress wild type or Swedish and London mutant amyloid precursor protein (APP), three conditions associated with familial AD [11, 48], Casp6 is activated in the absence of Casp3 activity and causes axonal degeneration [37, 55]. This is consistent with the absence of significant amounts of either active Casp3 or Casp7 in AD brains that exhibit high amounts of active Casp6 [37, 50, 57].

The early vulnerability of the ERC and CA1 to age-related Casp6 activation is commensurate with early NFT formation in pre-clinical Braak stages I and II of AD [12]. In humans, neurons from the ERC project their

axons towards the CA1 region and receive input from the olfactory bulb neurons. Therefore, degenerating neurons of the olfactory bulb could cause degeneration of input neurons of the ERC, which require synaptic signals for survival. Specifically, mitral and tufted neurons of the olfactory bulb receive input from neurons of the olfactory epithelium where the olfactory nerves of the epithelium cross the cribriform plate of the ethmoid bone. Mitral and tufted neurons direct their axons to the olfactory tract and can form clusters of axons called the AON in the olfactory bulb and the olfactory tract, or project their axons to the piriform cortex, the olfactory tubercle and the lateral ERC [51].

The possibility that olfactory bulb degeneration is related to hippocampal degeneration is supported by the fact that odor identification impairment in aging is associated with lower cognition [67]. Furthermore, impaired odor identification predicts the progression of non-cognitively impaired (NCI) to mild cognitively impaired (MCI) and of MCI to AD, and is associated with decreased episodic and semantic memory as well as decreased perceptual speed [68]. Impaired odor identification is also associated with NFT pathology in the ERC [66]. Furthermore, the levels of Tau and amyloid pathology in the olfactory bulb echoes the presence and severity of AD brain pathologies found in other regions, suggesting that brain biopsies of the olfactory bulb could be used for early diagnosis or to monitor the effects of drug treatments [6]. Atrophy of the olfactory bulb is detected in MCI and in AD and correlate with lower mini mental state exam (MMSE) scores [59–61]. The AON is the area most affected by neuronal loss in AD [22]. Cholinergic neurons of the olfactory system decrease by one third [35, 54] and mitral cells are reduced significantly in AD olfactory bulb [28, 58]. The AON is also the area where NFT and senile plaques occur [16, 19, 30, 35, 41, 44, 46], and the number of NFT is compatible with a diagnosis of AD with an accuracy of 93.3% [34]. The NFT pathology of the AON correlates with Braak stages, Lewy body pathology and the Apo E4 genotype [5, 23, 63].

Given the projection of olfactory tracts to the ERC, here, we examined Casp6 activation in the olfactory bulb of NCI, MCI and AD individuals. Casp6 activity was mainly observed in the AON region of the olfactory bulb. Levels of Casp6 activity correlated with PHF-1 NFT pathology in the AON. Casp6 activity and PHF-1 NFT pathology in the AON correlated with the levels of Casp6 activity in the

ERC. In all AD dementia cases and a few MCI cases, Casp6 activity in the AON region was higher than that observed in the ERC. A similar trend was observed for PHF-1. By contrast, levels of β -amyloid immunopositive pathology were generally similar in the AON and the ERC. Levels of AON Casp6 activity were associated with lower cognition in four independent cognitive measures. These results suggest that AON Casp6 activity could precede and induce Casp6-dependent ERC degeneration but do not exclude the possibility that ERC degeneration triggers AON degeneration. However, the results show an association between AON and ERC degeneration.

Materials and methods

Human brain tissues

Olfactory bulb and hippocampal tissue sections were obtained from the Religious Orders Study (ROS). The study was approved by the Institutional Review Board of Rush University Medical Center. All subjects signed informed consent and an Anatomic Gift Act. The McGill University Institutional Review Board approved the use of these human tissues in research. Details of the clinical and pathologic evaluation have been previously reported [9]. Yearly clinical evaluations included a medical history, neurological and cognitive assessment and a clinical diagnosis. Each case underwent MMSE to describe the cohort. Tests of visuospatial ability, perceptual speed and episodic, semantic and working memory evaluations yearly and these were converted to Z scores (using the mean and S.D. at baseline). A global cognitive score was developed by averaging the 19 cognitive function tests used to assess the five cognitive domains. Diagnostic classifications were provided by a clinician following the National Institute of Neurological and Communicative Disorders and Stroke and the AD and Related Disorders Association (NINCDS-ADRDA) criteria, without consideration of the pathological evaluation [42]. The diagnosis of AD dementia was assigned to cases with a history of cognitive decline and evidence of impairment of episodic memory and one other cognitive domain. MCI referred to cases of cognitive impairment without dementia. In the absence of dementia or MCI, cases were assigned a NCI diagnosis. At autopsy, brains underwent standard neuropathological evaluations, including Braak Stage determination. Neuropathological examination of the autopsied brains was done by a board-certified neuropathologist according to a modified Consortium to establish a registry for Alzheimer's Disease (CERAD) criteria, without access to clinical information [8]. Brain tissue sections were stained with a Bielschowski silver stain to reveal neuritic and diffuse plaques and NFT. Braak stages were determined based on NFT severity and distribution [31]. Other pathologies were examined. Only 2 AD from all subjects examined had typical hippocampal sclerosis. Lewy body disease (alpha-synuclein), TDP-43,

Table 1 Demographics and clinical characteristics of the study groups

Characteristic	NCI (n = 18)	MCI (n = 9)	AD (n = 7)
Age at death (yrs)	86.3 (6.4)	84.9 (6.8)	92.4 (4.7)
Male : Female	6 : 12	5 : 4	2 : 5
Education (yrs)	17.8 (3.2)	18.0 (1.8)	17.9 (2.1)
Post mortem interval, median (IQR)	5.6 (5.3–6.3)	5.1 (4.5–7.3)	5.8 (4.5–13.3)
Apolipoprotein E ϵ 4, n (%)	3 (16.7%)	2 (11.1%)	0
Mini mental state exam (MMSE)	28.3 (1.8)	26.7 (1.9)	14.4 (8.3)
Cognitive test scores			
Global Cognitive Scores	0.03 (0.4)	-0.4 (0.5)	-1.6 (0.8)
Episodic memory	0.4 (0.6)	-0.3 (0.9)	-2.2 (1.3)
Working memory	-0.02 (0.5)	-0.4 (0.8)	-0.8 (0.4)
Semantic memory	-0.1 (0.6)	-0.3 (0.6)	-1.3 (0.9)
Visuospatial activity	-0.1 (0.6)	-0.4 (0.9)	-1.2 (0.3)
Perceptual Speed	-0.3 (0.5)	-0.9 (0.7)	-1.6 (0.4)
Braak Stages, n (%)			
0	1 (5.6%)	0	0
I	1 (5.6%)	1 (11.1%)	0
II	0	2 (22.2%)	1 (14.3%)
III	10 (55.6%)	3 (33.3%)	1 (14.3%)
IV	6 (33.3%)	3 (33.3%)	1 (14.3%)
V	0	0	4 (57.1%)

Data represent the mean and (SD). For Braak stage the number of cases and the % in each group (NCI, MCI or AD)

atherosclerosis, cerebral amyloid angiopathy, cerebral arteriosclerosis, gross chronic infarction and gross cerebral infarcts were present and mostly equivalent in the brain of NCI, MCI and AD individual. Chronic microinfarcts and micro infarctions were present only in NCI and AD cases. Note that the clinical diagnosis was not always in agreement with the neuropathological diagnosis [8] and that our designation of AD, MCI and NCI in this paper is based on the cognitive assessment.

A total of 34 cases comprising 18 NCI, 9 MCI and 7 AD dementia were studied. Table 1 shows the demographic and the clinical characteristics of the study group including cognitive scores taken within a year of death and the Braak stages of NFT pathology determined at autopsy. The number of years of education and post-mortem interval (PMI) were similar among the three groups. The AD cases were slightly older and had lower cognitive scores than the NCI and MCI groups.

Immunohistochemistry

Paraformaldehyde-fixed, paraffin-embedded 4 μ m thick olfactory bulb and hippocampal formation tissue sections were deparaffinized, hydrated and treated with

Tris-EDTA antigen retrieval buffer (10 mmol/L Tris Base, 1 mmol/L ethylenediaminetetraacetic acid (EDTA), 0.05% Tween 20, pH 9) for 20 min at 97 °C in the Pascal Dako Cytomation apparatus. The Dako Autostainer Plus automated slide processor and the EnVision Flex system (Dako, Burlington, ON, Canada) were used to perform all the immunostaining. Brain cases with no pathology were added as negative control, and previously identified immunopositive AD hippocampus or cortical tissue sections added as positive controls for each immunostaining. The following antisera or antibodies were used: neopeptide antisera p20Casp6 10,630 antiserum (1/2000) recognizing the processed active p20 subunit of Casp6 and 10,635 TauΔCasp6 (1/25,000) produced in our laboratory [27], F25276 antiserum (1/1000) raised against Aβ₁₋₄₀ peptide [36] and PHF-1 antibody (1/5000) against phosphorylated Tau kindly provided by Dr Peter Davies (Department of Neuroscience, Albert Einstein College of Medicine, New York, NY). Slides were counterstained with hematoxylin, dehydrated and coverslipped with Permount mounting medium (Fisher Scientific, Ottawa, ON, Canada).

The neuronal density of AON and ERC tissue was evaluated in three AD cases where both AON and ERC were available, by counting the neurons within a defined area in the counterstained hematoxylin/eosin Casp6 immunostained tissue sections. Neurons were recognized according to their morphological characteristics as defined [24]. At least 500 neuronal nuclei were counted per 3-4 different regions of interest for each tissue section.

Cresyl violet staining (Nissl staining)

Olfactory bulb sections were deparaffinized with xylene, hydrated in descending series of ethanol concentrations and treated with a 0.1% cresyl violet in water (Sigma-Aldrich, St Louis, MO, USA) solution for 15 min followed by a few seconds of differentiation in 95% ethanol. After staining, the sections were dehydrated in absolute ethanol, treated with xylene and coverslipped using Permount mounting medium (Fisher Scientific, Ottawa, ON, Canada).

Quantitation of the immunopositive regions of the olfactory bulb

Stained tissue sections were scanned digitally with the MIRAX Scanner (Zeiss, Oberkochen, Germany). Consecutive cresyl violet stained olfactory bulb tissue sections were used to delineate the AON, and the identification of the AON was determined by a neuropathologist (SA) as described [51]. The AON was defined as a group of large pyramidal neurons slightly stained with cresyl violet. The *Atlas of the Human Brain* was used as a reference to identify specific brain regions [40]. Five regions of interest at least were photographed at magnification × 20 with the MIRAX Viewer Program

(Zeiss, Oberkochen, Germany). Immunostaining density was obtained with the ImageJ software (NIH, Bethesda, MD, USA) and results were expressed as μm² immunopositive staining/mm² of tissue.


Statistical analyses

Descriptive statistics are reported for the variables of interest. For categorical variables, we report counts and percentages. For continuous variables whose distribution of values showed evidence of being Normal, we report mean and standard deviation (SD); otherwise we report median and inter-quartile range (IQR). Spearman's correlations were run to determine monotonic relationship between the different pathologies present in the AON or the ERC and the relationship between the presence in AON and ERC for each marker. The distributions of values for active Casp6, TauΔCasp6, PHF-1 and Aβ between NCI, MCI and AD were compared with the Kruskal-Wallis (KW) test. When warranted by a conclusion of significant differences from the KW test, post-hoc pairwise comparisons were performed with Dunn's multiple comparisons test [20, 21]. These comparisons were done separately for the AON and the ERC. Multiple linear regressions were performed to investigate the influence of pathological characteristics present in AON or ERC (these analyses were done separately for the AON and the ERC) on cognitive scores measured. The models were adjusted for sex, age, PMI and years of education. All statistical tests of hypothesis were two-sided and carried out at the level of significance of 0.05. The multiple linear regressions were done using the R software (R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>). All other statistical analyses were performed using SAS, version 9.3 (SAS Institute, Cary, NC, USA).

Results

The AON is the region of the olfactory bulb most immunopositive to active Casp6, TauΔCasp6, PHF-1 Tau and Aβ


The AON area of the olfactory bulb whose neurites extend to the ERC (Fig. 1a) was chosen for quantification of Casp6 activity since the AON represents the area that is most strongly immunopositive to the neopeptide antisera against the p20 active subunit of Casp6 and TauΔCasp6, phosphorylated Tau PHF-1 antibodies and anti-Aβ antiserum (Fig. 1b & c). Strong anti-active Casp6 immunostaining was detected in neurons resembling pre-NFT and mature NFT, and in neuropil threads, as observed previously in AD cortex [27]. TauΔCasp6 immunoreactivity was detected in NFT-containing neurons and neuropil threads of adjacent tissue sections, and was more abundant than the immunoreactivity of



the anti-active Casp6 as expected since this antiserum is considerably stronger and Tau is more abundant in neurons than the Casp6 enzyme. Casp6 activity was not associated with apoptotic neuronal cell death based on the normal appearance of the nuclei of immunopositive neurons. PHF-1 immunostained NFT, although more sparsely than the Tau Δ Casp6 antiserum. Conversely, PHF-1 immunostained neuropil threads more abundantly than anti-active Casp6 and Tau Δ Casp6 neopeptide antisera. Anti-A β immunostained β -amyloid deposits resembling

classical β -amyloid plaques and some smaller diffuse deposits. Together, these results show that both the processed form of active Casp6 and Tau Δ Casp6, that define Casp6 activity, co-exist with classical AD pathology in the AON of the olfactory bulb.

Levels of active Casp6 correlate with Tau Δ Casp6 and PHF-1, but not with A β , in the AON of the olfactory bulb
Spearman's correlation showed a strong positive relationship between the anti-active Casp6 and the Tau Δ Casp6 in



the AON of the olfactory bulb ($r_s = 0.61$, $n = 21$, $p = 0.003$) (Fig. 2a). Both the anti-active Casp6 and the TauΔCasp6 immunostaining showed a strong positive relationship with PHF-1 immunostaining ($r_s = 0.63$, $n = 20$, $p = 0.003$ and $r_s = 0.69$, $n = 20$, $p = 0.001$, respectively) indicating, as observed previously in hippocampus [47], an association between Tau NFT pathology and Casp6 activity. By contrast, neither the anti-active Casp6 nor TauΔCasp6 immunostaining showed a relationship with Aβ immunostaining ($r_s = 0.44$, $n = 19$, $p = 0.06$ and $r_s = 0.18$, $n = 22$, $p = 0.47$, respectively). The level of PHF-1 showed a weak positive relationship with Aβ immunostaining in the AON ($r_s = 0.47$, $n = 22$, $p = 0.03$).

Spearman's correlation also showed a strong positive relationship between anti-active Casp6 and TauΔCasp6 immunopositivity in the ERC ($r_s = 0.69$, $n = 21$, $p = 0.001$) (Fig. 2b). As observed in the AON, both the anti-active Casp6 and the TauΔCasp6 immunostaining showed a strong positive relationship with the PHF-1 immunostaining in the ERC ($r_s = 0.69$, $n = 18$, $p = 0.001$ and $r_s = 0.85$, $n = 19$, $p < 0.0001$, respectively). Active Casp6 did not show a relationship with Aβ immunostaining in the ERC ($r_s = 0.19$, $n = 19$, $p = 0.42$), but TauΔCasp6 did ($r_s = 0.52$, $n = 21$, $p = 0.02$). Levels of Aβ immunostaining

showed a strong positive relationship with PHF-1 immunostaining only in the ERC ($r_s = 0.68$, $n = 19$, $p = 0.002$). Together, these results indicate that Casp6 activity is strongly associated with NFT pathology, but not with Aβ pathology, in the AON of the olfactory bulb and in the ERC.

AON active Casp6 and ERC TauΔCasp6 increase with AD progression

Table 2 shows descriptive statistics for pathological markers by brain region and clinical diagnosis. A KW test showed statistical significant differences in the levels of active Casp6 in the AON (χ^2 (2 degrees of freedom) = 6.05, $p = 0.04$). A post-hoc Dunn's test, showed a difference between AD dementia and NCI, as well as between AD dementia and MCI. By contrast, although TauΔCasp6 in the AON showed higher amounts in AD dementia as compared to NCI and MCI, the KW test showed that there were no statistically significant differences. This may be explained by the fact that the variability in TauΔCasp6 values, as measured by the IQR, is greater within AD dementia, NCI and MCI. Similarly for PHF-1 and Aβ, there were no statistically significant differences between the clinical diagnostic groups.

Table 2 AON and ERC immunostaining levels in NCI, MCI and AD

Region of interest	Clinical Diagnosis		Active Casp6	TauΔCasp6	PHF-1	Aβ
AON	NCI	Median	837.7	472.9	3655.9	403.3
		IQR	289.0–1481.8	263.0–4182.6	1248.2–23248.5	109.9–658.2
		n	11	11	11	10
	MCI	Median	226.1	980.7	7494.4	452.9
		IQR	118.1–2705.2	458.0–9493.4	651.1–67082.5	136.3–2993.1
		n	6	6	7	7
	AD	Median	23982.4	20400.0	75146.7	27355.0
		IQR	11445.0–26221.2	3087.2–54521.9	41473.2–151857.4	1222.9–55633.7
		n	4	4	5	5
ERC	NCI	Median	69.7	29.3	4138.2	940.7
		IQR	53.8–518.3	10.8–1138.3	615.9–16672.1	106.5–3406.6
		n	8	9	9	9
	MCI	Median	662.4	1035.8	15562.1	96.1
		IQR	14.4–1639.4	246.8–3901.1	9394.8–42729.6	54.2–4787.4
		n	7	7	7	7
	AD	Median	1022.1	7859.5	56051.4	14342.6
		IQR	497.2–1453.1	3181.4–18251.4	53590.7–70360.4	11159.6–24689.1
		n	4	5	3	5

Data represent the median (μm^2 staining/ μm^2 tissue) and the Inter-Quartile Range (25% percentile–75% percentile). n indicates the number of cases studied with each immunostaining in this region of interest. Comparisons of distributions for pathological markers by clinical diagnosis were done separately in the AON and in the ERC using the Kruskal-Wallis test, followed by a Dunn's post-hoc test where warranted (results and *p* values are reported in the text)

In the ERC, there were no statistically significant differences in the levels of active Casp6 between NCI, MCI and AD brains. However, differences in levels of ERC TauΔCasp6 were observed (KW test, χ^2 (2 df) = 8.37, *p* = 0.01). A post-hoc Dunn's test, showed a difference between AD and NCI. Similarly, statistically significant differences in the levels of ERC PHF-1 (KW test, χ^2 (2 df) = 7.13, *p* = 0.02) and Aβ levels (KW test, χ^2 (2) = 8.32, *p* = 0.01) were obtained and post-hoc Dunn's test confirmed a difference between AD and NCI, and between AD and MCI. Together, these results indicate higher levels of active Casp6, but not TauΔCasp6, PHF-1 or Aβ, in the AD AON than in the NCI or MCI AON. By contrast, levels of active Casp6 are not different in NCI, MCI or AD ERC, TauΔCasp6 is increased in AD ERC relative to NCI ERC levels, but PHF-1 and Aβ levels are increased in AD ERC compared to both NCI and MCI ERC.

Levels of Casp6 activity in the AON correlate with the levels of Casp6 activity and PHF-1 positive NFT in the ERC

It is expected that if degeneration in the ERC occurs because of the degeneration of input neurons in the AON, then Casp6 activity in the AON should show a relationship with that of the ERC. Spearman's correlation showed that levels of anti-active Casp6 in the AON did not have a significant relationship with those in the ERC

(r_s = 0.15, *n* = 14, *p* = 0.60) (Fig. 3a). However, levels of TauΔCasp6 immunoreactivity in the AON were associated positively with those in the ERC (r_s = 0.70, *n* = 14, *p* = 0.01) (Fig. 3b). Since cytosolic caspase enzyme levels are lower than the cytoskeletal Tau protein and because Casp6 undergoes rapid turnover by the proteasome [62], we consider TauΔCasp6 to be a stronger indicator of Casp6 activity than active Casp6. Furthermore, PHF-1 immunostaining in AON were associated positively with that of the ERC (r_s = 0.75, *n* = 14, *p* = 0.002) (Fig. 3c). By contrast, Aβ levels in the AON did not show a relationship with those in the ERC (r_s = 0.41, *n* = 15, *p* = 0.12) (Fig. 3d). These results indicate that degeneration in the AON parallels that of the ERC but does not give any information as to whether degeneration in the AON preceded that in the ERC. To assess this possibility, we compared immunopositive levels where staining were obtained in both the AON and ERC of the same case (Fig. 4). All 3 AD dementia cases and one MCI case showed higher active Casp6 in the AON than in the ERC (Fig. 4a). Two out of 3 AD dementia cases and 2 MCI cases showed higher TauΔCasp6 in the AON than in the ERC (Fig. 4b). Similarly, PHF-1 levels in AON were higher in AD and 2 MCI cases (Fig. 4c). By contrast, Aβ was higher in AON in only one AD case, other cases showed similar levels.




Fig. 3 AON TauΔCasp6 and PHF-1 staining shows a positive monotonic relationship with ERC staining. Scatterplots and Spearman's correlations (r_s) between the active Casp6 (a), TauΔCasp6 (b), PHF-1 (c) and Aβ (d) immunostaining (μ^2 staining/ mm^2 of tissue) in the AON versus the ERC




Fig. 4 Casp6 and Tau pathologies are more abundant in AON than in the ERC for AD cases. Active Casp6 (a), TauΔCasp6 (b), PHF-1 (c) and Aβ (d) immunostaining values (μ^2 staining/ mm^2 of tissue) in the AON and ERC for 5 NCI, 3 MCI and 3 AD cases where both the AON and ERC were stained from the same individual case

Together, these results suggest that either the AON is more susceptible to Casp6 activation and PHF-1 immunopositive AD pathology than the ERC or that the AON pathology precedes the ERC pathology and this result in a greater accumulation of Casp6 activity and PHF-1 pathology.

Higher levels of active Casp6 in the AON correlate with lower cognitive performance

To investigate the association between the pathologies of the AON or ERC and the cognitive performance of individuals, we performed separate regression analyses for five cognitive domains assessed within a year of death. Analyses were done separately for the AON and the ERC. In each regression, we adjusted for sex, age,

PMI and years of education. Spearman’s correlation results discussed above showed that active Casp6, TauΔ-Casp6 and PHF-1 immunostaining were highly associated with one another. Multicollinearity of these three factors was confirmed by their high values of the variance inflation factor found in a preliminary for-diagnosis multiple regression model. To adjust for this multicollinearity, we used a ridge regression approach, which allows for the inclusion of these factors in the same model. The value of the ridge parameter λ was chosen as the one for which the coefficients were relatively stable from the ridge traces. The values selected were λ = 20 and λ = 100 for AON and ERC, respectively. Furthermore, due to the small sample sizes, *p* values for the regression coefficients were computed using a

Table 3 Ridge regression coefficients of cognitive performance scores from AON and ERC pathologies

Predicting cognition	AON (n = 19)			ERC (n = 18)		
	Marker	Coefficient	<i>p</i> value	Marker	Coefficient	<i>p</i> value
GCS	Active Casp6	-0.18	0.01	Active Casp6	-0.02	0.40
	TauΔCasp6	-0.08	0.29	TauΔCasp6	-0.05	0.06
	PHF-1	-0.06	0.33	PHF-1	-0.03	0.27
	Aβ	-0.18	0.04	Aβ	-0.03	0.27
MMSE	Active Casp6	-1.97	0.001	Active Casp6	-0.29	0.23
	TauΔCasp6	-1.36	0.05	TauΔCasp6	-0.72	0.01
	PHF-1	-0.92	0.06	PHF-1	-0.41	0.04
	Aβ	-0.90	0.21	Aβ	-0.30	0.14
Episodic Memory	Active Casp6	-0.31	0.01	Active Casp6	-0.02	0.67
	TauΔCasp6	-0.13	0.32	TauΔCasp6	-0.07	0.07
	PHF-1	-0.14	0.17	PHF-1	-0.05	0.24
	Aβ	-0.28	0.05	Aβ	-0.05	0.27
Working memory	Active Casp6	-0.06	0.26	Active Casp6	-0.02	0.21
	TauΔCasp6	-0.03	0.59	TauΔCasp6	-0.01	0.64
	PHF-1	-0.004	0.93	PHF-1	-0.004	0.81
	Aβ	-0.09	0.16	Aβ	0.001	0.95
Semantic memory	Active Casp6	-0.15	0.03	Active Casp6	-0.01	0.69
	TauΔCasp6	-0.11	0.14	TauΔCasp6	-0.05	0.05
	PHF-1	-0.07	0.25	PHF-1	-0.02	0.55
	Aβ	-0.16	0.06	Aβ	-0.03	0.28
VSA	Active Casp6	-0.09	0.12	Active Casp6	-0.01	0.62
	TauΔCasp6	-0.01	0.91	TauΔCasp6	-0.01	0.84
	PHF-1	0.02	0.72	PHF-1	-0.01	0.57
	Aβ	-0.06	0.41	Aβ	-0.03	0.21
PS	Active Casp6	-0.08	0.23	Active Casp6	-0.05	0.05
	TauΔCasp6	-0.07	0.32	TauΔCasp6	-0.05	0.04
	PHF-1	-0.02	0.71	PHF-1	-0.02	0.29
	Aβ	-0.04	0.64	Aβ	-0.02	0.31

Data represent the ridge regression analysis of the different markers in the AON and the ERC and the cognitive scores. Model adjusted for sex, age, PMI and years of education, in bold *p* values less or equal to 0.05

permutation method with 10,000 permutations [17]. All ridge regressions and permutation tests were performed using the package MASS in R [65].

Higher levels of active Casp6 in the AON were associated with lower global cognitive, MMSE, episodic and semantic memory scores (Table 3). Tau Δ Casp6 and A β levels in the AON were marginally negatively associated with the MMSE score and GCS, respectively. By contrast, PHF-1 levels in the AON were not associated with any of the cognitive measures. In the ERC, active Casp6 and Tau Δ Casp6 levels were marginally negatively associated with perceptual speed, and episodic memory or perceptual speed, respectively. Tau Δ Casp6 levels were however negatively associated with MMSE scores, a feature observed also for PHF-1, although the association was weak. A β levels did not associate with any of the cognitive measures. These results indicate that AON active Casp6 levels are associated with cognitive decline, and interestingly, the association is with episodic and semantic memory, two types of memory function initially altered in AD.

Discussion

In this study, the levels of Casp6 activity were assessed in the olfactory bulb of NCI, MCI and AD dementia individuals to determine if previously observed Casp6 activity in the ERC was associated with degeneration in the olfactory bulb since olfactory bulb neurons project their axons to the ERC. The results of the study revealed that (1) Casp6 activity is highest in and mostly restricted to the AON of the olfactory bulb and co-exists with Tau, but not A β , AD pathology, (2) Casp6 activity increases in the AON with the progression of disease and appears to be higher than that of Casp6 activity in the ERC, and (3) active Casp6 levels in the AON are negatively associated with cognitive performance.

The co-existence of Casp6 activity with Tau pathology in the AON of the olfactory bulb is consistent with our previous observations of an intimate relationship between Casp6 activity and PHF-1 Tau pathology in the ERC and CA1 regions of the hippocampus [47]. The fact that both Tau and Casp6 are localized in the neuronal cytosol and are involved in axonal degeneration suggests that they could both be part of a neurodegeneration pathway in AD neurons [18, 45, 55]. Casp6 cleavage of Tau at amino acid 402 [27] may impact Tau's polymerization, as observed for cleaved Tau at amino acids 391 and 421 [1, 10]. Cleavage of Tau may also alter Tau's function in the stabilization of microtubules, in a manner similar to the phosphorylation of Tau at amino acid 396 [13]. Whether NFT formation and Casp6 activity occur as parallel pathways of degeneration or as part of the same pathway remains to be determined, and will be an important question to resolve eventually as the answer would have a

significant implication in the development of treatments against neurodegeneration in AD.

In contrast to its association with NFT pathology, Casp6 activity in the AON is not associated with A β extracellular deposits. The lack of association between Casp6 activity and A β is consistent with our previous observations in the hippocampus [3, 27, 47]. Our results differ from several reports indicating β -amyloid toxicity associated with caspase activation [7, 14, 15]. Nevertheless, our analyses cannot exclude the possibility of toxic A β oligomers triggering neuronal Casp6 activation. It is also important to consider that A β production can be a consequence of Casp6 activation in primary cultures of human CNS neurons [32, 36, 38], a fact that would be consistent with Casp6 activity preceding A β overproduction and deposition as extracellular plaques.

This study shows that Casp6 activity in the AON could be added as an additional biomarker of neurodegeneration associated with AD. First, the AON active Casp6 levels were significantly higher in AD compared to AON NCI or MCI, indicating increased levels of enzyme with progression of disease. A similar trend was observed with Tau Δ Casp6 but did not reach statistical significance because of the large variability amongst samples. Second, the levels of Tau Δ Casp6 in the AON were positively and strongly associated with the levels of Tau Δ Casp6 in the ERC. Similar results were observed for PHF-1 pathology. Third, the higher levels of Casp6 activity, determined by both the presence of active Casp6 and Tau Δ Casp6, and PHF-1 Tau, in the AON than in the ERC of individual MCI and AD cases suggest that degeneration in the AON could precede degeneration in the ERC. However, our approach cannot exclude that ERC degeneration leads to AON degeneration or that the two pathologies occur concomitantly. More intensive studies will be required to determine which of these three possibilities explains the relationship between AON and ERC degeneration. The number of available cases evaluated limited our study. Furthermore, a longitudinal study would be required to determine if pathology in one area of the brain precedes another. This longitudinal study would require the ability to observe the degeneration in live patients. This is likely to soon become possible with the evolution of PET ligands against tangles and A β , but is not yet possible for Casp6 activity. Furthermore, as proposed by Attems [4–6], olfactory biopsies could be done to assess the diagnostic potential of the olfactory bulb, and to evaluate the efficiency of treatments against AD.

This study shows an association between higher levels of AON active Casp6 and lower memory performance. Indeed, higher levels of Casp6 associate with lower global cognitive, MMSE and episodic and semantic memory scores. These results are consistent with the association of

impaired odor identification and lower global cognition and a more rapid rate of episodic and semantic memory decline in MCI [68]. The fact that episodic and semantic memory impairment occurs before AD dementia [43] suggests the involvement of Casp6 activity at a very early stage of the disease. In other studies, an association between olfactory bulb and tract atrophy and MMSE scores was observed [60, 61]. Given the role of Casp6 in axonal degeneration [45, 55], it would be of interest to determine eventually if Casp6 activity may be responsible for the atrophy of the olfactory bulb and tract. Surprisingly, AON TauΔCasp6 showed only a marginal association with a decline in MMSE score and no association with the other cognitive measures. Similarly, no association was observed between AON PHF-1 levels and cognitive measures. These results suggest that the presence of Tau pathology in the AON is not associated with cognitive performance. Similarly, the AON levels of Aβ associated very weakly with lower GCS and not with any of the other measures of cognition. However, with larger numbers, significance may be achieved. For example, an association between olfactory Tau pathology and clinical dementia in AD and MCI subjects has been reported [4, 5].

In the ERC, TauΔCasp6 and PHF-1 levels associated negatively with the MMSE score. Active Casp6 and TauΔCasp6 levels both associated negatively but very weakly with perceptual speed performance. Similarly, a very weak association between TauΔCasp6 and semantic memory was also observed. Aβ levels did not associate with any of the cognitive measures. Previously, we did not observe an association between cognition and Casp6 activity in the ERC when combining NCI, MCI and AD groups, but detected a significant association between episodic and semantic memory and Casp6 activity in the NCI subjects ERC [3, 47]. Larger numbers in each subgroup will be required for a solid conclusion to be drawn. Since the focus of this study was the olfactory bulb, we did not pursue the study in the ERC.

Conclusions

In conclusion, these data support the possibility that the Casp6 activity in the AON of the olfactory bulb reflects degeneration in the ERC and suggest that Casp6 activity in the olfactory bulb could represent degeneration associated with cognitive decline and early AD. Furthermore, Casp6 in the AON could be used as an early marker to detect the initiation of pathways of degeneration in AD.

Abbreviations

AD: Alzheimer disease; AON: Anterior olfactory nucleus; APP: Amyloid precursor protein; CA1: Cornu ammonis 1; Casp3: Caspase-3; Casp6: Caspase-6; Casp7: Caspase-7; ERC: Entorhinal cortex; IQR: Inter-quartile range; MCI: Mild cognitively impaired; MMSE: Mini-mental state examination; NCI: Non-cognitively impaired; NFT: Neurofibrillary tangles; PHF-1: Paired helical filament; PMI: Post-mortem interval; ROS: Religious order study; SD: Standard deviation

Acknowledgements

The authors thank Jenna Ross, Ph. D and Veronica M. Afonso, Ph. D. for their help with the immunostaining of the tissue sections while they were in our laboratories. We are also very grateful to Dr. Peter Davies (The Feinstein Institute for Medical Research, NY) for the gift of the PHF-1 antibody.

Funding

This work was supported by funds from the Canadian Institutes of Health Research (CIHR) MOP-243413-BCA-CGAG-45097, the Canadian Foundation for Innovation, and the JGH Foundation to ALB and National Institutes of Health P30AG10161 and RF1AG15819 to DAB.

Availability of data and materials

The dataset for cognitive measures and subject demographics are available from the ROS at RUSH Medical Centre. The immunohistological staining results are kept electronically and physically in the LeBlanc laboratory. The datasets used and/or analysed and the materials used during the current study may be available upon availability from the corresponding author or Dr D.A. Bennett on reasonable request.

Author's contribution

BF performed and analyzed the immunostainings, prepared the figures and participated in the writing of the manuscript. SA performed the histological examination and reviewed the manuscript. DAB provided the brain materials, the cognitive and demographics data, and reviewed the manuscript. JAC performed the statistical analysis and reviewed the manuscript. ALB conceived, directed the study and wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The study was approved by the Institutional Review Board of Rush University Medical Center. All subjects signed informed consent and an Anatomic Gift Act. The McGill University Institutional Review Board approved the use of these human tissues in research.

Author details

¹Bloomfield Center for Research in Aging, Lady Davis Institute for Medical Research, Jewish General Hospital, 3755 ch. Côte Ste-Catherine, Montreal, QC, Canada. ²Department of Pathology, Montreal Children's Hospital and McGill University, Montreal, QC, Canada. ³Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, USA. ⁴Department of Mathematics and Statistics, McGill University, Montreal, QC, Canada. ⁵Department of Neurology and Neurosurgery, McGill University, Montreal, QC, Canada.

Received: 8 November 2016 Accepted: 1 December 2016

Published online: 08 December 2016

References

1. Abrahama A, Ghoshal N, Gambelin TC, Cryns V, Berry RW, Kuret J, Binder LI (2000) C-terminal inhibition of tau assembly in vitro and in Alzheimer's disease. *J Cell Sci* 113(Pt 21):3737–3745
2. Albrecht S, Bogdanovic N, Ghetti B, Winblad B, LeBlanc AC (2009) Caspase-6 activation in familial Alzheimer disease brains carrying amyloid precursor protein or presenilin I or presenilin II mutations. *J Neuropathol Exp Neurol* 68:1282–1293
3. Albrecht S, Bourdeau M, Bennett D, Mufson EJ, Bhattacharjee M, LeBlanc AC (2007) Activation of caspase-6 in aging and mild cognitive impairment. *Am J Pathol* 170:1200–1209
4. Attems J, Jellinger KA (2006) Olfactory tau pathology in Alzheimer disease and mild cognitive impairment. *Clin Neuropathol* 25:265–271
5. Attems J, Lintner F, Jellinger KA (2005) Olfactory involvement in aging and Alzheimer's disease: an autopsy study. *J Alzheimers Dis* 7:149–157, discussion 173–180
6. Attems J, Walker L, Jellinger KA (2014) Olfactory bulb involvement in neurodegenerative diseases. *Acta Neuropathol* 127:459–475

7. Barnes N, Li L, Yoshikawa K, Schwartz L, Oppenheim R, Milligan C (1998) Increased production of amyloid precursor protein provides a substrate for caspase-3 in dying motoneurons. *J Neurosci* 18:5869–5880
8. Bennett DA, Schneider JA, Aggarwal NT, Arvanitakis Z, Shah RC, Kelly JF, Fox JH, Cochran EJ, Arends D, Treinkman AD, Wilson RS (2006) Decision rules guiding the clinical diagnosis of Alzheimer's disease in two community-based cohort studies compared to standard practice in a clinic-based cohort study. *Neuroepidemiology* 27:169–176
9. Bennett DA, Schneider JA, Buchman AS, Barnes LL, Boyle PA, Wilson RS (2012) Overview and findings from the rush memory and aging project. *Curr Alzheimer Res* 9:646–663
10. Berry RW, Abraha A, Lagalwar S, LaPointe N, Gamblin TC, Cryns VL, Binder LI (2003) Inhibition of tau polymerization by its carboxy-terminal caspase cleavage fragment. *Biochemistry* 42:8325–8331
11. Bertram L, Lill CM, Tanzi RE (2010) The genetics of Alzheimer disease: back to the future. *Neuron* 68:270–281
12. Braak H, Alafuzoff I, Arzberger T, Kretzschmar H, Del Tredici K (2006) Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol (Berl)* 112:389–404
13. Branblett GT, Goedert M, Jakes R, Merrick S, Trojanowski J, Lee V (1993) Abnormal tau phosphorylation at Ser396 in Alzheimer's disease recapitulates development and contributed to reduced microtubule binding. *Neuron* 10:1089–1099
14. Chan SL, Griffin WS, Mattson MP (1999) Evidence for caspase-mediated cleavage of AMPA receptor subunits in neuronal apoptosis and Alzheimer's disease. *J Neurosci Res* 57:315–323
15. Chan SL, Mattson MP (1999) Caspase and calpain substrates: roles in synaptic plasticity and cell death. *J Neurosci Res* 58:167–190
16. Christen-Zaech S, Kraftsik R, Pillevuit O, Kiraly M, Martins R, Khalili K, Miklossy J (2003) Early olfactory involvement in Alzheimer's disease. *Can J Neurol Sci* 30:20–25
17. Cule E, Vineis P, De Iorio M (2011) Significance testing in ridge regression for genetic data. *BMC Bioinf* 12:372
18. Cusack CL, Swahari V, Hampton Henley W, Michael Ramsey J, Deshmukh M (2013) Distinct pathways mediate axon degeneration during apoptosis and axon-specific pruning. *Nat Commun* 4:1876
19. Davies DC, Brooks JW, Lewis DA (1993) Axonal loss from the olfactory tracts in Alzheimer's disease. *Neurobiol Aging* 14:353–357
20. Dunn OJ (1964) Multiple comparisons using rank sums. *Technometrics* 6:241–252
21. Elliott AC, Hynan LS (2011) A SAS (®) macro implementation of a multiple comparison post hoc test for a Kruskal-Wallis analysis. *Comput Methods Programs Biomed* 102:75–80
22. Esiri MM, Wilcock GK (1984) The olfactory bulbs in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 47:56–60
23. Fujishiro H, Tsuboi Y, Lin WL, Uchikado H, Dickson DW (2008) Co-localization of tau and alpha-synuclein in the olfactory bulb in Alzheimer's disease with amygdala Lewy bodies. *Acta Neuropathol* 116:17–24
24. Garman RH (2011) Histology of the central nervous system. *Toxicol Pathol* 39:22–35
25. Godefroy N, Foveau B, Albrecht S, Goodyer CG, LeBlanc AC (2013) Expression and activation of caspase-6 in human fetal and adult tissues. *PLoS One* 8:e79313
26. Gray DC, Mahrus S, Wells JA (2010) Activation of specific apoptotic caspases with an engineered small-molecule-activated protease. *Cell* 142:637–646
27. Guo H, Albrecht S, Bourdeau M, Petzke T, Bergeron C, LeBlanc AC (2004) Active Caspase-6 and Caspase-6 cleaved Tau in neuropil threads, neuritic plaques and neurofibrillary tangles of Alzheimer's Disease. *Am J Pathol* 165:523–531
28. Hoogland PV, van den Berg R, Huisman E (2003) Misrouted olfactory fibres and ectopic olfactory glomeruli in normal humans and in Parkinson and Alzheimer patients. *Neuropathol Appl Neurobiol* 29:303–311
29. Julien O, Zhuang M, Wiita AP, O'Donoghue AJ, Knudsen GM, Craik CS, Wells JA (2016) Quantitative MS-based enzymology of caspases reveals distinct protein substrate specificities, hierarchies, and cellular roles. *Proc Natl Acad Sci U S A* 113(14):E2001–10
30. Kasa P, Rakonczay Z, Gulya K (1997) The cholinergic system in Alzheimer's disease. *Prog Neurobiol* 52:511–535
31. Katzman R, Terry R, DeTeresa R, Brown T, Davies P, Fuld P, Renbing X, Peck A (1988) Clinical, pathological, and neurochemical changes in dementia: a subgroup with preserved mental status and numerous neocortical plaques. *Ann Neurol* 23:138–144
32. Kaushal V, Dye R, Pakavathkumar P, Foveau B, Flores J, Hyman B, Ghetti B, Koller BH, LeBlanc AC (2015) Neuronal NLRP1 inflammasome activation of Caspase-1 coordinately regulates inflammatory interleukin-1-beta production and axonal degeneration-associated Caspase-6 activation. *Cell Death Differ* 22:1676–1686
33. Klaiman G, Champagne N, LeBlanc AC (2009) Self-activation of Caspase-6 in vitro and in vivo: Caspase-6 activation does not induce cell death in HEK293T cells. *Biochim Biophys Acta* 1793:592–601
34. Kovacs T, Cairns NJ (1999) Lantos PL: beta-amyloid deposition and neurofibrillary tangle formation in the olfactory bulb in ageing and Alzheimer's disease. *Neuropathol Appl Neurobiol* 25:481–491
35. Kovacs I, Torok I, Zombori J, Kasa P (1998) Cholinergic structures and neuropathologic alterations in the olfactory bulb of Alzheimer's disease brain samples. *Brain Res* 789:167–170
36. LeBlanc A (1995) Increased production of 4 kDa amyloid beta peptide in serum deprived human primary neuron cultures: possible involvement of apoptosis. *J Neurosci* 15:7837–7846
37. LeBlanc AC (2013) Caspase-6 as a novel early target in the treatment of Alzheimer's disease. *Eur J Neurosci* 37:2005–2018
38. LeBlanc A, Liu H, Goodyer C, Bergeron C, Hammond J (1999) Caspase-6 role in apoptosis of human neurons, amyloidogenesis, and Alzheimer's disease. *J Biol Chem* 274:23426–23436
39. LeBlanc AC, Ramcharitar J, Afonso V, Hamel E, Bennett DA, Pakavathkumar P, Albrecht S (2014) Caspase-6 activity in the CA1 region of the hippocampus induces age-dependent memory impairment. *Cell Death Differ* 21:696–706
40. Mai JK, Paxinos G, Voss T (2008) Atlas of the Human Brain. Third Edition edn. Elsevier, Amsterdam
41. Mann DM, Tucker CM, Yates PO (1988) Alzheimer's disease: an olfactory connection? *Mech Ageing Dev* 42:1–15
42. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R et al (2011) The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7:263–269
43. Mortamais M, Ash JA, Harrison J, Kaye J, Kramer J, Randolph C, Pose C, Albala B, Ropacki M, Ritchie CW, Ritchie K (2016) Detecting cognitive changes in preclinical Alzheimer's disease: A review of its feasibility. *Alzheimers Dement*
44. Mundinano IC, Caballero MC, Ordóñez C, Hernández M, DiCauda C, Marcilla I, Erro ME, Tunon MT, Luquin MR (2011) Increased dopaminergic cells and protein aggregates in the olfactory bulb of patients with neurodegenerative disorders. *Acta Neuropathol* 122:61–74
45. Nikolaev A, McLaughlin T, O'Leary DD, Tessier-Lavigne M (2009) APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature* 457:981–989
46. Ohm TG, Braak H (1987) Olfactory bulb changes in Alzheimer's disease. *Acta Neuropathol* 73:365–369
47. Ramcharitar J, Afonso VM, Albrecht S, Bennett DA, LeBlanc AC (2013) Caspase-6 activity predicts lower episodic memory ability in aged individuals. *Neurobiol Aging* 34:1815–1824
48. Rovelet-Lecrux A, Hannequin D, Raux G, Le Meur N, Laquerriere A, Vital A, Dumanchin C, Feuillette S, Brice A, Vercelletto M et al (2006) APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet* 38:24–26
49. Saiz-Sanchez D, Flores-Cuadrado A, Ubeda-Banon I, de la Rosa-Prieto C, Martínez-Marcos A (2016) Interneurons in the human olfactory system in Alzheimer's disease. *Exp Neurol* 276:13–21
50. Selznick L, Holtzman D, Han B, Gokden M, Srinivasan A, Johnson E, Roth K (1999) In situ immunodetection of neuronal caspase-3 activation in Alzheimer's disease. *J Neuropathol and Exp Neurol* 58:1020–1026
51. Shipley M, Reyes P (1991) Anatomy of the human olfactory bulb and central olfactory pathways. In: Laing DG, Doty R, Breipohl W (eds) *In The sense of smell*. Springer, Heidelberg
52. Simon DJ, Pitts J, Hertz NT, Yang J, Yamagishi Y, Olsen O, Tesic Mark M, Molina H, Tessier-Lavigne M (2016) Axon Degeneration Gated by Retrograde Activation of Somatic Pro-apoptotic Signaling. *Cell* 164:1031–1045
53. Simon DJ, Weimer RM, McLaughlin T, Kallop D, Stanger K, Yang J, O'Leary DD, Hannoush RN, Tessier-Lavigne M (2012) A caspase cascade regulating developmental axon degeneration. *J Neurosci* 32:17540–17553
54. Simpson J, Yates CM, Gordon A, St Clair DM (1984) Olfactory tubercle choline acetyltransferase activity in Alzheimer-type dementia, Down's syndrome and Huntington's chorea. *J Neurol Neurosurg Psychiatry* 47:1138–1139
55. Sivananthan S, Lee A, Goodyer CG, LeBlanc AC (2010) Familial amyloid precursor protein mutants cause caspase-6-dependent but amyloid

- β -peptide-independent neuronal degeneration in primary human neuron cultures. *Cell Death Dis* 1(1):e100. doi:10.1038/cddis.2010.74
56. Sokolowski JD, Gamage KK, Heffron DS, LeBlanc AC, Deppmann CD, Mandell JW (2014) Caspase-mediated cleavage of actin and tubulin is a common feature and sensitive marker of axonal degeneration in neural development and injury. *Acta Neuropathol Commun* 2:16
 57. Stadelmann C, Deckwerth T, Srinivasan A, Bancher C, Bruck W, Jellinger K, Lassmann H (1999) Activation of caspase-3 in single neurons and autophagic granules of granulovacuolar degeneration in Alzheimer's disease. *Am J Pathol* 155:1459–1466
 58. Struble RG, Clark HB (1992) Olfactory bulb lesions in Alzheimer's disease. *Neurobiol Aging* 13:469–473
 59. ter Laak HJ, Renkawek K, van Workum FP (1994) The olfactory bulb in Alzheimer disease: a morphologic study of neuron loss, tangles, and senile plaques in relation to olfaction. *Alzheimer Dis Assoc Disord* 8:38–48
 60. Thomann PA, Dos Santos V, Seidl U, Toro P, Essig M, Schroder J (2009) MRI-derived atrophy of the olfactory bulb and tract in mild cognitive impairment and Alzheimer's disease. *J Alzheimers Dis* 17:213–221
 61. Thomann PA, Dos Santos V, Toro P, Schonknecht P, Essig M, Schroder J (2009) Reduced olfactory bulb and tract volume in early Alzheimer's disease—a MRI study. *Neurobiol Aging* 30:838–841
 62. Tounekti O, Zhang Y, Klaiman G, Goodyer CG, LeBlanc A (2004) Proteasomal degradation of caspase-6 in 17 β -estradiol-treated neurons. *J Neurochem* 89:561–568
 63. Tsuboi Y, Wszolek ZK, Graff-Radford NR, Cookson N, Dickson DW (2003) Tau pathology in the olfactory bulb correlates with Braak stage, Lewy body pathology and apolipoprotein epsilon4. *Neuropathol Appl Neurobiol* 29:503–510
 64. Uribe V, Wong BK, Graham RK, Cusack CL, Skotte NH, Pouladi MA, Xie Y, Feinberg K, Ou Y, Ouyang Y et al (2012) Rescue from excitotoxicity and axonal degeneration accompanied by age-dependent behavioral and neuroanatomical alterations in caspase-6-deficient mice. *Human Mol Gen* 21:1954–1967
 65. Venables WN, Ripley BD (2002) *Modern Applied Statistics with S*, 4th edn. Springer, New York
 66. Wilson RS, Arnold SE, Schneider JA, Tang Y, Bennett DA (2007) The relationship between cerebral Alzheimer's disease pathology and odour identification in old age. *J Neurol Neurosurg Psychiatry* 78:30–35
 67. Wilson RS, Arnold SE, Tang Y, Bennett DA (2006) Odor identification and decline in different cognitive domains in old age. *Neuroepidemiology* 26:61–67
 68. Wilson RS, Schneider JA, Arnold SE, Tang Y, Boyle PA, Bennett DA (2007) Olfactory identification and incidence of mild cognitive impairment in older age. *Arch Gen Psychiatry* 64:802–808

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

