



CMKLR1 activation *ex vivo* does not increase proportionally to serum total chemerin in obese humans

Jay Toulany¹, Sebastian D Parlee², Christopher J Sinal³, Kathryn Slayter⁴, Shelly McNeil⁴ and Kerry B Goralski^{1,3}

¹College of Pharmacy, Dalhousie University, Halifax, Nova Scotia, Canada

²Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, Michigan, USA

³Department of Pharmacology, Dalhousie University, Halifax, Nova Scotia, Canada

⁴Canadian Centre for Vaccinology, IWK Health Centre, Nova Scotia Health Authority and Dalhousie University, Halifax, Nova Scotia, Canada

Correspondence should be addressed to K Goralski

Email
kerry.goralski@dal.ca

Abstract

Prochemerin is the inactive precursor of the adipokine chemerin. Proteolytic processing is obligatory for the conversion of prochemerin into active chemerin and subsequent regulation of cellular processes via the chemokine-like receptor 1 (CMKLR1). Elevated plasma or serum chemerin concentrations and differential processing of prochemerin have been reported in obese humans. The impact of these changes on CMKLR1 signalling in humans is unknown. The objective of this pilot study was to develop a cellular bioassay to measure CMKLR1 activation by chemerin present in human serum and to characterise how obesity modifies serum activation of CMKLR1 under fasted and fed conditions. Blood samples were collected from control ($N=4$, BMI 20–25) and obese ($N=4$, BMI >30) female subjects after an overnight fast ($n=2$) and at regular intervals ($n=7$) following consumption of breakfast over a period of 6 h. A cellular CMKLR1-luminescent reporter assay and a pan-chemerin ELISA were used to determine CMKLR1 activation and total chemerin concentrations, respectively. Serum total chemerin concentration (averaged across all samples) was higher in obese vs control subjects (17.9 ± 1.8 vs 10.9 ± 0.5 nM, $P < 0.05$), but serum activation of CMKLR1 was similar in both groups. The CMKLR1 activation/total chemerin ratio was lower in obese vs control subjects (0.33 ± 0.04 vs 0.58 ± 0.05 , $P < 0.05$). After breakfast, serum total chemerin or CMKLR1 activation did not differ from baseline values. In conclusion, the unexpected observation that obese serum activation of CMKLR1 did not match increased total chemerin concentrations suggests impaired processing to and/or enhanced degradation of active chemerin in serum of obese humans.

Key Words

- ▶ adipokines
- ▶ bioassay
- ▶ chemerin
- ▶ chemokine-like receptor 1 (CMKLR1)
- ▶ obesity

Endocrine Connections
(2016) 5, 70–81

Introduction

After characterisation as an adipokine by our group in 2007, research concerning chemerin and its relationship to obesity has risen steadily. Chemerin is secreted as an inactive precursor prochemerin, which undergoes processing at distinct sites on the C-terminus by proteases of the inflammatory, coagulation and fibrinolytic cascades to become biologically active

chemerin (herein active chemerin) (1, 2, 3, 4). Several isoforms of active chemerin can be produced that vary in their C-terminal amino acid, their ability to bind and activate the chemokine-like receptor 1 (CMKLR1) as well as their capacity to facilitate their distinctive inflammatory, immune and metabolic processes (1, 2, 3, 4, 5, 6, 7, 8, 9, 10).



The established biological functions of chemerin including the recruitment of immune cells to sites of inflammation (3, 9, 11) and the regulation of adipogenesis, lipolysis and glucose metabolism (12, 13, 14, 15, 16, 17) have implicated chemerin as a target of interest in the pathophysiology of obesity and the metabolic syndrome. In line with a reputed role for chemerin in obesity, total chemerin concentrations are elevated in murine and human obesity, are correlated to markers of adiposity including body mass index (BMI), waist-to-hip ratio (WHR) and fat mass and decline after surgical, diet and activity-based weight loss (13, 14, 18, 19, 20, 21, 22). Independent observations in humans, animals and cell models further support adipose tissue/adipocytes as a significant and modifiable source of circulating total chemerin (pro and active chemerin) (14, 18, 19, 23, 24, 25, 26, 27). The exact role, beneficial or detrimental, that chemerin plays in obesity is complicated by the fact that both pro- (28, 29, 30) and anti-inflammatory effects (5, 31, 32, 33) have been described for chemerin/CMKLR1 signalling. Furthermore, the biological effects of chemerin are not exclusively dependent upon total circulating concentrations. Rather, the mechanisms by which prochemerin is activated, the relative abundance of chemerin isoforms and their activity at CMKLR1 are important determinants of biological function (1, 2, 4, 6, 8, 19). Accordingly, a significant limitation of the majority of previous studies has been a reliance on quantifying total but not active chemerin. As such, much of the clinical chemerin research to date has inferred a role for elevated chemerin in obesity and metabolic disorders based on the assumption that parity exists between total and active chemerin concentrations. However, this assumption is not supported by *in vitro* or *in vivo* studies that demonstrated a large discrepancy between total and active chemerin under a TNF-mediated acute inflammatory response and in synovial fluid of arthritic patients (18, 19, 34). A disparity between total and active chemerin is further supported by a recent human study that discovered elevated concentrations of shorter and previously uncharacterised chemerin isoforms in plasma of obese humans compared to lean controls (23). How these novel degraded chemerin isoforms influence CMKLR1 activation and the biological processes that ensue in obesity remains unknown. The objective of our study was to utilise a cellular bioassay to measure CMKLR1 activation by chemerin present within human serum samples to address whether CMKLR1 activation parallels total chemerin concentrations in obese and normal weight humans under fed and fasted conditions.

Materials and methods

Subjects

The study was open to men and women, ages 18–55 years, with a BMI of 20–25 (control weight, $n=4$) or BMI >30 (obese, $n=4$). Subjects were recruited through the Izaak Walton Killam (IWK) Health Centre. Exclusion criteria included subjects taking medications associated with enhanced chemerin production (insulin and metformin) (26) or inhibition of degradation (ACE inhibitors) (6, 35). Additionally, subjects with renal impairment (creatinine clearance <60 mL/min), type 2 diabetes mellitus or who have previously undergone gastric bypass surgery were excluded as these conditions may influence chemerin concentrations independent of obesity (36). The IWK Research Ethics Board approved the experimental protocol (Project # 1013395), and all participants gave their written informed consent before their inclusion in the study.

Clinical procedures

Subjects fasted overnight (10–12 h) prior to their arrival to the Clinical Research Unit of the IWK Health Centre at approximately 06:00 h on the study day. All subjects remained in the unit for the duration of the study. Upon admission, subjects were assessed for anthropometric measures of body fat including BMI, WHR and waist circumference in addition to their systolic (SBP) and diastolic (DBP) blood pressure and heart rate. A saline lock was subsequently placed, and fasted blood samples were collected at 07:00 h (baseline 1) and 08:00 h (baseline 2). Immediately after collection of the second sample at 08:00 h, subjects were offered a selection of breakfast foods (fruit cups, fresh fruit, assorted bagels, muffins and cereals, yogurt, cheese and hard-boiled eggs), beverages (fruit juices, milk, tea and coffee) and condiments (cream cheese, peanut butter and sugar). Subjects chose from these breakfast items without any restrictions and were allowed to consume breakfast over a 30-min period. Blood samples were subsequently taken at 08:30, 09:00, 09:30, 10:00, 11:00, 12:00 and 13:00 h. Two samples (4 mL each) were collected at each time point and placed into a 3.2% sodium citrate tube for plasma preparation and an uncoated tube for serum. Based on reported time for maximal activation of chemerin in serum (9), blood samples taken for serum were left to clot at room temperature for 2 h after collection. All samples

were centrifuged at 2500g for 5 min at 4°C, aliquoted and stored at –70°C until the appropriate analysis was performed.

Quantifying CMKLR1 activation using the CMKLR1 bioassay

The CMKLR1 bioassay methods were based on our previously reported protocol for measuring CMKLR1

activation by chemerin in serum and adipocyte media samples (18, 19). Briefly, a genetically modified HEK293 cell line (HTLA) that constitutively expresses a fusion protein comprised a tobacco etch virus (TEV) protease linked to human β -arrestin2, and a transcriptional-transactivator (tTA)-dependent luciferase reporter gene was transiently transfected with a plasmid-expressing human CMKLR1 fused to a tTa via a TEV protease recognition sequence (37). A batch transfection

Table 1 Baseline subject characteristics.

Parameter	Normal weight (BMI 20–25)	Obese (BMI >30)	Ref. range
Physical characteristics			
Age (years)	30.8±11.6	39±8.0	NA
Weight (kg)	66.8±12.1	106.0±18.6*	NA
Height (cm)	170.2±9.8	163.5±6.0	NA
Waist-to-hip ratio (WHR)	0.81±0.03	0.87±0.05	≥0.85 ^a
Body mass index (BMI) (kg/m ²)	23.0±1.3	38.8±7.5*	≥30 ^b
Cardiovascular and renal			
Heart rate (beats/min)	74±3	72±20	NA
Systolic BP (mm/Hg)	116.8±9.8	125.3±14.6	NA
Diastolic PB (mm/Hg)	72.0±8.5	78.5±14.3	NA
Creatinine (μmol/L)	59.0±8.5	58.8±7.5	<98
eGFR (mL/min)	117.7±21.1	99.4±19.8	≥90
Metabolism/inflammation			
Insulin (pmol/L)	69±22	118±118	≤209
Glucose (mmol/L)	4.65±0.48	5.53±0.96	3.3–5.6
Triglycerides (mmol/L)	0.78±0.17	1.85±1.08	0.5–1.6
Free fatty acids (μmol/L)	573±168	490±239	100–900
Cholesterol (mmol/L)	3.93±0.43	4.55±0.87	2.9–5.7
Hs-CRP (mg/L)	8.25±11.6	8.23±6.17	0–5.0
PAI (ng/mL)	4.48±0.44	9.49±2.26*	NA
Immunological			
White blood cells (×10 ⁹ /L)	6.67±1.77	5.65±1.05	4.0–11
Abs neutrophils (×10 ⁹ /L)	3.90±1.11	3.33±1.24	1.5–8.0
Abs lymphocytes (×10 ⁹ /L)	1.80±0.71	1.80±0.29	1.5–5.0
Abs monocytes (×10 ⁹ /L)	0.63±0.14	0.38±0.10*	0.0–1.0
Abs eosinophils (×10 ⁹ /L)	0.30±0.13	0.15±0.06	0.0–0.5
% Neutrophils	59.2±4.0	57.2±10.9	44–79
% Lymphocytes	26.5±5.3	33.3±10.5	28–48
% Monocytes	9.4±1.2	6.7±1.2*	1.0–9.0
% Eosinophils	4.5±2.3	2.3±0.83	0.0–5.0
% Basophils	0.37±0.06	0.43±0.29	0.0–1.0
Haematological			
Red blood cells (×10 ¹² /L)	4.17±0.39	4.27±0.31	4.1–5.1
Haemoglobin (g/L)	128.3±9.1	112.3±10.6*	120–160
Haematocrit (L/L)	0.39±0.03	0.35±0.03	0.36–0.46
Mean corpuscular volume (fL)	92.4±3.6	82.3±1.9*	77–102
Mean corpuscular haemoglobin (pg)	30.8±1.0	26.3±0.9*	26.0–35.0
Mean corpuscular haemoglobin concentration (g/L)	334±10	320±15	320–370
Red cell distribution width (%)	12.6±0.3	15.3±0.2*	12.2–14.3
Platelet count (×10 ⁹)	225±61	258±45	186–353
Mean platelet volume (fL)	10.9±0.4	10.6±1.2	6.8–11.2

All of the above parameters were measured in the 1st morning baseline plasma or serum samples after an overnight fast. All values are listed as the mean±s.d. mean ($n=4$ per group). The typical reference values for various parameters are shown in the right hand column.

^aA WHR ≥0.85 was used as the cutoff to define central obesity. ^bA BMI of ≥30 kg/m² to define obesity. The estimated glomerular filtration rate (eGFR) was determined from the measured serum creatinine concentration using the Cockcroft–Gault equation with correction for ideal body weight. *The values were significantly different ($P<0.05$) compared to the control group by both unpaired t -test. NA, not applicable; PAI-1, plasminogen activator inhibitor 1.

mixture was prepared containing (per well) 40 μ L Opti-MEM, 200 ng human *CMKLR1*-tTA plasmid, 200 ng β -galactosidase reference plasmid and 2×10^4 HTLA cells (at passage 5–20) in 160 μ L DMEM supplemented with 0.4 μ L of 1 mg/mL polyethylenimine (PEI) and added to a 96-well plate. After 24 h, the transfection mix was aspirated and replaced with 50 μ L of diluted (1:10 with optimum) human serum. To match the sample compositions, recombinant human chemerin₁₅₇ (R&D Systems) 50 μ L standards were prepared in heat-inactivated bovine serum diluted 1:10 in Opti-MEM (0.1, 0.3, 0.6, 1, 3, 6, 10 and 30 nM final concentrations). In this assay, activation of CMKLR1 leads to the recruitment of the β -arrestin2-TEV protease to the C-terminus of the CMKLR1-tTA fusion protein, resulting in tTa cleavage and migration of free tTa to the nucleus where it transcribes the luciferase reporter gene. After 16 h, the serum or standards were aspirated and the cells were incubated for 5 min with shaking (1000 rev min⁻¹) in 100 μ L reporter lysis buffer (RLT; Promega) followed by a single rapid freeze/thaw cycle. To quantify the luciferase activity, 10 μ L of lysate or RLT blank was transferred to a 96-well white luminometer plate. Eighty microlitres of luciferase assay reagent (Promega) was injected into each well, and the luminescence was monitored for 10 ms using a Biotek synergy HT plate reader (Biotek). For the β -galactosidase assay, 30 μ L of lysate or blank was transferred to a clear 96-well plate and incubated with 30 μ L of 2 \times β -galactosidase assay buffer (200 mM NaPO₄ pH 7.3, 2 mM MgCl₂, 100 mM 2-mercaptoethanol and 1.33 mg/mL *ortho*-nitrophenyl- β -galactoside (ONPG)) for 15 min at 37°C. The reaction was stopped by the addition of 100 μ L of 1 M Na₂CO₃, and the absorbance at 420 nm was measured. The luciferase and β -galactosidase measurements were corrected for the respective blanks.

Table 2 Serum CMKLR1 bioassay performance.

Chemerin ₁₅₇ standards (nM)	Serum chemerin standards		
	Predicted chemerin concentrations		
	Accuracy	Inter-assay % CV	Intra-assay % CV
0.10	142.8%	51.4%	26.8%
0.30	102.8%	13.6%	12.6%
0.60	97.7%	13.8%	5.6%
1.00	97.8%	4.5%	12.8%
3.00	102.2%	35.9%	14.5%
6.00	79.2%	47.2%	21.2%
10.00	64.6%	55.2%	46.8%
30.00	56.6%	98.5%	61%

The assay performance values were based on 4 independent experimental replicates with each standard measured in duplicate.

Activation of the receptor by each standard and sample was expressed as the fold-change in luciferase/ β -galactosidase activity relative to the 1:10 diluted blank serum sample. Each standard was assayed in duplicate, and each sample was assayed in triplicate.

Using GraphPad Prism (GraphPad Software), the concentration of each chemerin₁₅₇ standard was plotted against the corresponding fold-change in luciferase/ β -galactosidase activity. The standards were fit to a 4-phase logarithmic nonlinear regression curve (log (agonist) vs normalised response), and the level of CMKLR1 receptor activation (expressed as chemerin₁₅₇




Figure 1 CMKLR1 bioassay performance. Panels A and B are representative of two independent CMKLR1 bioassay standard curves of chemerin₁₅₇ in bovine serum diluted 1:10 in optimum media. Representative serum samples diluted 1:10 from 2 subjects with BMI 20–25 and BMI >30 show that the luciferase/ β -galactosidase values fall within the most accurate and reproducible portion of the curve.

equivalents) in each sample was interpolated based on its luciferase/ β -galactosidase activity.

Total chemerin measurements

Total chemerin was quantified using a human pan-chemerin ELISA (LOT # 1316723, R&D Systems) according to the manufacturer's instructions and our previously reported methods (38). A dose-response curve using a best fit of a 4-parameter logistic equation was used to calculate the total chemerin concentration.

Quantification of other factors in plasma and serum

The 1st baseline sample was used for the baseline complete blood cell count, metabolic/inflammatory and serum

creatinine analysis. These assays were performed in the IWK clinical chemistry facility using standard methods. In addition, we used commercially available analytical kits to quantify glucose, free fatty acids (FFA), high-sensitivity C-reactive protein (hs-CRP), cholesterol, triglycerides (Cayman Chemical) and insulin (Millipore) in all samples in order to compare changes in these parameters in the post-prandial vs baseline period. A commercially available ELISA assay was used for measures of plasminogen activator inhibitor-1 (PAI-1) in the baseline samples (Abcam). The manufacturer's instructions were followed for all assays.

Data and statistical analysis

The primary outcome was to compare the average CMKLR1 activation, total chemerin and CMKLR1




Figure 2

The effect of obesity on serum total chemerin and serum activation of CMKLR1. Serum total chemerin (A), serum activation of CMKLR1 expressed as chemerin₁₅₇ equivalents (C) and CMKLR1 activation/total chemerin (E) were measured in 4 obese (BMI >30) and 4 normal weight females (BMI 20–25) after an overnight fast (baseline, BL), which was followed by breakfast (BF) consumed over 30 min and then intermittent sample measurements over a 5.5-h duration in the post-breakfast period. The between-group statistical analysis was performed on the average data for all time points combined (B, D, F). ** $P < 0.01$ and *** $P < 0.001$ compared to the BMI >30 group, unpaired t -test.

activation/total chemerin ratios between the obese and control groups. For these variables, the average of 9 samples over time in plasma and serum in each patient was calculated. The average values were compared using an unpaired 2-tailed *t*-test. The secondary outcome was to evaluate for differences in average CMKLR1 activation, total chemerin and CMKLR1 activation/total chemerin ratios in the baseline vs post-prandial period within each group. For this analysis, the average of the two baseline measures were compared to the maximum and minimum values for CMKLR1 activation, total chemerin and CMKLR1 activation/total chemerin that were observed in the post-prandial period. A one-way analysis of variance with Bonferroni *post hoc* test was used. Differences between groups were considered significant when $P < 0.05$. SPSS, version 23 was used for statistical analysis.

Results

Subject characteristics

The 1st eight subjects that applied for entry into the study met the inclusion criteria and were enrolled after their initial screening visit. No subjects were excluded based on the predetermined exclusion criteria. Individual subject characteristics are summarised in Table 1. All subjects were female and were categorised as either normal weight ($n=4$) with BMIs between 20 and 25 or obese ($n=4$) with BMIs >30 . As per the study inclusion criteria, weight and BMI were significantly higher in the obese vs control group. Age, height, WHR, cardiovascular and renal parameters

were similar between groups. High-sensitivity C-reactive protein was more than two-fold higher than the upper normal values for one normal weight and 2 obese subjects, but did not differ between groups. Plasminogen activator inhibitor-1 (PAI-1) was significantly higher in the BMI >30 group. Immune cell counts fell with the normal reference range for both groups, although absolute and percent monocyte counts were significantly lower in the BMI >30 group. Haemoglobin, mean corpuscular volume and mean corpuscular haemoglobin were lower and red cell distribution width was higher in the obese vs normal weight groups and was suggestive of mild iron-deficiency anaemia. After breakfast, both groups displayed a significant rise in insulin concentration (after 1.5–2 h) and a later reduction in free fatty acids (2.5–5 h) (Supplementary Fig. 1 and Supplementary Table 1, see section on supplementary data given at the end of this article). In the post-prandial period, significant increases in glucose, triglycerides and cholesterol were observed in the normal weight but not obese group.

CMKLR1 bioassay performance

The CMKLR1 bioassay accuracy and inter- and intra-assay coefficients of variation for each standard curve chemerin concentration are shown in Table 2. The lower limit for the quantification of CMKLR1 activation occurred at approximately 100 pM chemerin₁₅₇, but the standard curve typically overestimated the actual chemerin₁₅₇ concentrations in this range, and the inter-assay and intra-assay CVs were unacceptably high. Between the

Table 3 Post-prandial changes in serum total chemerin and CMKLR1 activation.

Parameter	Baseline	Maximum	Minimum	t_{\max} (h)	t_{\min} (h)
Serum total chemerin (nmol/L)					
Normal weight	10.4 ± 1.1	13.5 ± 1.4	8.69 ± 0.61	1.75 (1.5–3.0)	1.75 (1.5–5.0)
Obese	17.0 ± 3.1	22.9 ± 3.4	13.2 ± 2.2	4.5 (3.0–6.0)	4.0 (1.5–6.0)
Serum CMKLR1 activation (expressed as nmol/L chemerin ₁₅₇ equivalents)					
Normal weight	6.03 ± 1.28	7.45 ± 0.97	5.24 ± 0.98	4.0 (2.5–6.0)	2.0 (1.5–5.0)
Obese	5.70 ± 1.74	6.51 ± 1.99	5.34 ± 1.82	3.0 (2.0–4.0)	2.5 (2.0–6.0)
Serum CMKLR1 activation/total chemerin ratio					
Normal weight	0.583 ± 0.136	0.710 ± 0.154	0.436 ± 0.091*	4.0 (3.0–6.0)	1.75 (1.5–2.0)
Obese	0.333 ± 0.087	0.423 ± 0.145	0.259 ± 0.056	4.0 (2.0–6.0)	2.25 (1.5–2.5)
Plasma total chemerin (nmol/L)					
Normal weight	10.3 ± 1.2	11.5 ± 1.2	7.56 ± 0.42	4.25 (1.5–6)	4.0 (2.5–5.0)
Obese	14.0 ± 3.6	17.5 ± 2.0	12.2 ± 3.0	3.0 (2.5–6.0)	3.0 (1.5–6.0)

t_{\max} and t_{\min} = the time at which the maximum and minimum values occurred, respectively. Baseline, maximum and minimum values are expressed as mean ± s.d. and t_{\max} and t_{\min} as median with lower and upper values in brackets. For each parameter, the average of the two baseline samples was compared to the maximum (or minimum) value obtained in the post-feeding period using a one-way repeated-measures ANOVA, followed by Bonferroni *post hoc* comparison.

* $P < 0.05$ compared to baseline.

concentrations of 0.3–1.0 nM, the assay was highly accurate and the inter- and intra-assay coefficients were in the acceptable range (<15%). The assay performance progressively deteriorated with increasing chemerin concentrations at above 3.0 nM. Two representative standard curves along with samples from two normal weight and two obese subjects demonstrated that a 1:10 dilution of serum samples provided predicted chemerin values within the most accurate and reproducible range (0.3–1.0 nM) of the standard curve (Fig. 1).




Figure 3

The effect of obesity on plasma total chemerin concentration. Plasma total chemerin (A) was measured in 4 obese (BMI >30) and 4 normal weight females (BMI 20–25) after an overnight fast (baseline, BL), which was followed by breakfast (BF) consumed over 30 min and then intermittent sample measurements over a 5.5-h duration in the post-breakfast period. The between-group statistical analysis was performed on the average data for all time points combined (B). ** $P < 0.01$ compared to the BMI >30 group, unpaired t -test.

Total chemerin was elevated in the serum of obese females but did not lead to greater CMKLR1 activation

Serum total chemerin concentrations (measured by a pan-chemerin ELISA) were consistently higher in the obese vs normal weight group, and this was reflected by a significant 1.6-fold higher average total chemerin over the entire sampling duration (Fig. 2A and B). In comparison, CMKLR1 activation (expressed as chemerin₁₅₇ equivalents) in the bioassay was similar between groups (Fig. 2C and D). In matched samples, the average apparent chemerin₁₅₇ concentration determined by the CMKLR1 bioassay ranged between 43% and 66% (normal weight group) and 25% and 45% (obese group) of serum total chemerin concentrations. Consistent with this, the average ratio of CMKLR1 activation to total chemerin concentrations was significantly higher in the normal weight vs obese groups (Fig. 2E and F). Reflecting a small degree of fluctuation in serum total chemerin and CMKLR1 activation over time, the maximum and minimum values for total chemerin and CMKLR1 activation after feeding did not differ from baseline in the normal weight and obese groups (Table 3). The times at which the maximum (t_{max}) and minimum (t_{min}) values were observed were also highly variable (Table 3). With respect to the CMKLR1 activation/total chemerin ratio, a significant nadir was observed in the normal weight but not obese group at between 1.5 and 2 h after breakfast.

Assessment of plasma total chemerin and plasma activation of CMKLR1

Similar to serum, plasma total chemerin was significantly increased in obese vs normal weight subjects (Fig. 3A and B) and did not differ in the post-prandial period vs baseline in either group (Table 3). When data from all eight subjects were combined, the mean serum total chemerin concentration (14.4 ± 3.9) was significantly higher than the mean plasma total chemerin concentration (11.9 ± 3.3), $P = 0.01$ paired t -test. We attempted to measure CMKLR1 activation by plasma samples using the CMKLR1 bioassay. During assay development, it was determined that a 1:20 dilution of bovine plasma with 3.2% Na citrate provided optimal signal to background and generated chemerin standard curves with similar assay performance characteristics as observed for 1:10 diluted serum standards. However, when the subject plasma samples were diluted to 1:20, the level of CMKLR1 activation was at or below the lower limit of quantification

in the assay. Therefore, these data are not presented, and firm conclusions regarding the activation of CMKLR1 by plasma chemerin were not possible.

Discussion

Previous studies examining total chemerin concentrations in humans and animals have resulted in the hypothesis that chemerin is relevant to obesity and obesity-associated comorbidities (12, 39, 40, 41, 42). However, by focusing solely on total chemerin, the majority of these studies have not accounted for chemerin processing to active forms and subsequent activation of CMKLR1. Thus, the biological significance of elevated chemerin in the context of human obesity remains unclear. Our proof-of-concept study has advanced understanding in this area by adopting a CMKLR1 cellular-reporter gene assay (18, 19, 37) to determine if CMKLR1 activation by human serum paralleled total chemerin concentrations and if this relationship differed in lean and obese subjects.

As expected, obese subjects had significantly increased serum total chemerin. Unexpectedly, this did not result in proportionally greater activation of CMKLR1 in the bioassay. The importance of this result is that functional alterations in chemerin/CMKLR1 signalling in obesity, whether pro- or anti-inflammatory may be less than expected based on measures of total chemerin alone. A possible explanation for the observed disparity between total chemerin concentrations and CMKLR1 activation by serum is that prochemerin is degraded to shorter chemerin products that were detected by the pan-chemerin ELISA but did not activate CMKLR1. Supporting the latter argument, Chang and coworkers demonstrated increased plasma total chemerin in obesity using the pan-chemerin ELISA assay, but using isoform-specific ELISAs did not demonstrate significant changes in high-activity chemerin₁₅₇, low-activity chemerin₁₅₅ and inactive prochemerin (23). Furthermore, the combined concentrations of chemerin₁₅₅, chemerin₁₅₇ and prochemerin only accounted for 57% of total chemerin detected by the pan-chemerin ELISA. The remaining 43% was attributed to chemerin isoforms that underwent further C-terminal processing, resulting in shorter proteins. As the ability of chemerin to activate CMKLR1 depends primarily on the 9 C-terminal (149–157) residues (43), these shorter chemerin proteins would be predicted to be inactive. Additionally, it is plausible that these peptides may act as competitive antagonists for the active

chemerin forms. However, both of these possibilities remain to be tested experimentally.

In addition to increased degradation, there could be reduced conversion of prochemerin to active chemerin isoforms. In one pathway, prochemerin is converted to chemerin₁₅₇ via a two-step process involving cleavage of amino acids 159–163 by plasmin or tryptase to form chemerin₁₅₈, an isoform with low activity following removal of the C-terminal lysine by plasma carboxypeptidase B or N to form chemerin₁₅₇ (1, 4). In obese humans, increased circulating concentrations of PAI-1, a serine protease inhibitor that blocks the enzymatic conversion of plasminogen to plasmin, have been well documented (44). Consistent with this, baseline PAI-1 was elevated in our obese subjects, suggesting that the plasmin/carboxypeptidase pathway of chemerin₁₅₇ formation could be less active.

We report the novel observation that within subjects, total chemerin was elevated (21%) in serum compared to plasma. There are at least two possible explanations for this. Platelets store prochemerin and secrete it upon activation (1). Therefore, it is possible that platelet activation during blood clotting increases prochemerin release and total chemerin concentrations in serum vs plasma. The chemerin ELISA demonstrates lower reactivity towards prochemerin, chemerin₁₅₈ and chemerin₁₅₅ compared to chemerin₁₅₇ (23). Thus, it is also possible that the ELISA assay underestimates total chemerin concentrations in plasma, where prochemerin predominates while providing more accurate determinations in serum where chemerin₁₅₇ concentrations are higher.

We previously observed that serum chemerin concentrations in mice oscillated with time of day, and the pattern of oscillation was altered with obesity (18). Other studies have demonstrated that short-term or long-term changes in nutritional status (fasting and re-feeding) affected chemerin mRNA expression and chemerin protein secretion from adipocytes and/or serum chemerin concentrations (22, 45). These latter effects may have been related to insulin stimulation of prochemerin secretion by adipocytes (26, 45). Despite these observations, the majority of clinical experimental studies have assessed plasma or serum chemerin concentrations in a single sample after an overnight fast and have thus not accounted for potential effects of time of day or nutritional effects on chemerin. To begin to address this gap in understanding, the clinical sampling procedures of our pilot study were designed to assess if serum total chemerin and CMKLR1 activation change acutely in association with insulin secretion

after breakfast. The observation of similar values for serum total chemerin and CMKLR1 activation in the post-prandial vs baseline periods suggested that the post-prandial insulin response did not appreciably affect serum chemerin concentrations and CMKLR1 activation. Furthermore, if insulin stimulation of adipose tissue chemerin production was linked to serum total chemerin, the rise and fall of chemerin should have been delayed relative to the insulin response and occurred over a narrow time range such as that observed for free fatty acid concentrations. This was not observed. Our results agreed with the findings of Bauer and coworkers in which a similar acute insulin increase produced by a fasting oral glucose tolerance test did not increase serum chemerin over a period of 1–2 h (46). Our results also expanded on these earlier findings by evaluating a typical breakfast, evaluating more time points over a longer duration and showing that immunodetectable chemerin and CMKLR1 activation are similarly unaffected. Our findings differed from those of Tan and coworkers, which demonstrated an approximate 2-fold increase in plasma chemerin 4 h after initiating a hyperinsulinaemic normoglycemic clamp (26). As the peak insulin concentrations observed in our study were similar to those in the study by Tan and coworkers, our data reaffirm that insulin regulation of chemerin is most likely to occur in conditions when there are sustained rather than acutely elevated insulin concentrations.

There are a number of limitations to our study. The bioassay provided a measure of CMKLR1 activation expressed as chemerin₁₅₇ equivalents but did not measure the 'true' chemerin₁₅₇ concentration and we were unable to reference the CMKLR1 bioassay results to purified prochemerin or other chemerin isoforms. Based on the generation of multiple chemerin isoforms by adipocytes and the presence of multiple chemerin isoforms in plasma (5, 23), there are also likely to be multiple chemerin forms in serum, which may be active or inactive to varying degrees and/or activate the CMKLR1 bioassay with a differing stoichiometry than chemerin₁₅₇. Furthermore, it is unknown if there could be interactions between multiple chemerin isoforms in human serum that could influence the activation of the receptor compared to chemerin₁₅₇ alone.

A second limitation is that the CMKLR1 bioassay only assesses one aspect of CMKLR1 activation and signalling, that being the β -arrestin2-mediated pathway. Emerging research supports that chemerin/CMKLR1 signalling is complex and not completely defined (47). The pleiotropic functions of chemerin involves classical CMKLR1

signalling via $G\alpha_{i/o}$ proteins as well as β -arrestin2 with the involvement of several intracellular effector pathways including intracellular calcium, p38 mitogen-activated protein kinases (MAPKs), extracellular signal-related kinases (ERK1/2) and phosphoinositide-3-kinase/protein kinase B (PI3K/AKT) and the RhoA/Rock pathway (3, 47, 48, 49). Although we can conclude that the CMKLR1 β -arrestin2 pathway is activated by human serum, we cannot currently make inferences regarding biased ligand signalling, activation of specific downstream effector molecules and the resulting impacts on cellular functions and/or CMKLR1 internalisation and desensitisation.

A third limitation was that we assessed the activation of CMKLR1 and not other chemerin receptors. In addition to CMKLR1, chemerin binds G-protein receptor 1 (GPR1) and the non-signalling chemokine C–C motif receptor-like 2 (CCRL2) (30, 37). The majority of chemerin functions have been attributed to CMKLR1 signalling (50). However, recent studies have confirmed functional human GPR1 signalling, which mediates chemotaxis of human gastric adenocarcinoma cells (47). Future studies that compare β -arrestin2 and $G\alpha_{i/o}$ protein-based CMKLR1 and GPR1 bioassays along with signalling reporter assays for the downstream pathways (as described by Rourke and coworkers (47)) will be important to further delineate the mechanisms of chemerin signalling and determine whether β -arrestin2 vs $G\alpha_{i/o}$ biased signalling can be influenced by the variation in chemerin isoforms produced in normal weight vs obese subjects. Our results also cannot account for potential effects of CCRL2, which binds and increases local levels of active chemerin facilitating chemerin interactions with CMKLR1 on neighbouring cells (30).

Although the small number of subjects was a limitation, this was offset by the repeated sampling procedures used for the primary analysis. Furthermore, the mean differences in plasma total chemerin between obese and normal weight subjects were similar to historical data, providing confidence in our primary analyses (38, 46, 51). Given the small sample size, we conducted a conservative comparison of post-feeding minimum and maximum values to baseline for our secondary analysis. The relatively stable serum total chemerin concentrations and CMKLR1 activation during early daytime hours observed in our study is consistent with the results of Tan and coworkers in which plasma chemerin concentrations measured at 30 min to 4-h intervals over 24 h in a small group of control subjects demonstrated little fluctuation (26). However, it is possible that we missed significant changes, given the 30- to 60-min frequency of sampling and that we were

only able to sample for part of the day. Nonetheless, the pilot data provide the basis for sample size determinations for future studies aimed at conducting a more robust two-way mixed analysis of variance to assess the effects of time and obesity on chemerin concentrations and CMKLR1 and GPR1 activation.

The inability to measure CMKLR1 activation by plasma was a limitation. Plasma should better represent circulating active chemerin isoforms, whereas serum activation of CMKLR1 accounts for active chemerin that is formed during the coagulation process. It is possible that the relationships between time and obesity on CMKLR1 activation by plasma could differ from those observed for serum. The ability to detect CMKLR1 activation by serum but not plasma was likely the result of lower concentrations of active chemerin₁₅₇ in plasma (10, 23) and is consistent with previous studies demonstrating that serum- but not plasma-mediated chemotaxis of CMKLR1-expressing β -lymphoma cells (9).

The literature, although mixed, suggests that gender has small or insignificant effects on plasma or serum total chemerin (52, 53, 54, 55, 56, 57). Furthermore, Chang and coworkers, did not observe an effect of gender on the plasma concentration of specific chemerin isoforms (23). However, as only women enrolled in the study, we cannot identify or rule out gender differences in serum activation of CMKLR1.

Our study included healthy obese subjects only. The assessment of whether the relationship between total chemerin and CMKLR1 activation changes with more severe obesity coupled with comorbidities remains to be determined.

In summary, our study is the first to identify a disparity between serum total chemerin and CMKLR1 activation in obesity, in that obese subjects displayed increased immunodetectable chemerin that did not lead to a comparable increase in *ex vivo* CMKLR1 activation. These results affirm our position that inferences regarding chemerin signalling or function based solely on measures of serum total chemerin concentrations should be made with extreme caution. Ideally, future clinical studies should include measures of total chemerin or specific chemerin isoforms (using isoform-specific antibodies or mass spectrometry) combined with bioassay measures of CMKLR1 and/or GPR1 activation.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/EC-16-0065>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

The work was supported by the Dalhousie Pharmacy Endowment Fund Operating Grant and a Dalhousie University, Faculty of Health Professions Research Development Grant.

Author contribution statement

J T performed the research, generated experimental data, analysed the data and co-wrote the manuscript. S D P conceived and designed the research and co-wrote the manuscript. K S served as the clinical study Co-ordinator/Pharmacist and recruited and assessed study subjects and edited the manuscript. C J S conceived and designed the research and edited the manuscript. S M supervised the clinical procedures as the study physician and edited the manuscript. K B G was the study's Principal Investigator, conceived and designed the research, analysed data and co-wrote the manuscript.

Acknowledgements

The authors express their sincere thanks to the subjects who volunteered their time to participate in the clinical study. They would like to sincerely thank Catherine Brown (Clinical Operations Coordinator, Canadian Centre for Vaccinology) and her staff for organising and conducting the clinical sampling procedures. They are grateful to Nichole McMullen, Dr Alexandra Roman and Yan Wang for their assistance with technical aspects of the project and Dr Gilad Barnea (Brown University) who provided the HTLA cells and the CMKLR1-TA expression plasmids that were used in the CMKLR1 bioassay.

References

- Du XY, Zabel BA, Myles T, Allen SJ, Handel TM, Lee PP, Butcher EC & Leung LL. Regulation of chemerin bioactivity by plasma carboxypeptidase N, carboxypeptidase B (activated thrombin-activable fibrinolysis inhibitor), and platelets. *Journal of Biological Chemistry* 2009 **284** 751–758. (doi:10.1074/jbc.M805000200)
- Wittamer V, Bondue B, Guillabert A, Vassart G, Parmentier M & Communi D. Neutrophil-mediated maturation of chemerin: a link between innate and adaptive immunity. *Journal of Immunology* 2005 **175** 487–493. (doi:10.4049/jimmunol.175.1.487)
- Wittamer V, Franssen JD, Vulcano M, Mirjole JF, Le Poul E, Migeotte I, Brezillon S, Tyldesley R, Blanpain C, Detheux M, *et al.* Specific recruitment of antigen-presenting cells by chemerin, a novel processed ligand from human inflammatory fluids. *Journal of Experimental Medicine* 2003 **198** 977–985. (doi:10.1084/jem.20030382)
- Zabel BA, Allen SJ, Kulig P, Allen JA, Cichy J, Handel TM & Butcher EC. Chemerin activation by serine proteases of the coagulation, fibrinolytic, and inflammatory cascades. *Journal of Biological Chemistry* 2005 **280** 34661–34666. (doi:10.1074/jbc.M504868200)
- Dranse HJ, Muruganandan S, Fawcett JP & Sinal CJ. Adipocyte-secreted chemerin is processed to a variety of isoforms and influences MMP3 and chemokine secretion through an NF κ B-dependent mechanism. *Molecular and Cellular Endocrinology* 2016 **436** 114–129. (doi:10.1016/j.mce.2016.07.017)
- Guillabert A, Wittamer V, Bondue B, Godot V, Imbault V, Parmentier M & Communi D. Role of neutrophil proteinase 3 and

- mast cell chymase in chemerin proteolytic regulation. *Journal of Leukocyte Biology* 2008 **84** 1530–1538. (doi:10.1189/jlb.0508322)
- 7 Kulig P, Kantyka T, Zabel BA, Banas M, Chyra A, Stefanska A, Tu H, Allen SJ, Handel TM, Kozik A, et al. Regulation of chemerin chemoattractant and antibacterial activity by human cysteine cathepsins. *Journal of Immunology* 2011 **187** 1403–1410. (doi:10.4049/jimmunol.1002352)
 - 8 Yamaguchi Y, Du XY, Zhao L, Morser J & Leung LL. Proteolytic cleavage of chemerin protein is necessary for activation to the active form, Chem157S, which functions as a signaling molecule in glioblastoma. *Journal of Biological Chemistry* 2011 **286** 39510–39519. (doi:10.1074/jbc.M111.258921)
 - 9 Zabel BA, Silverio AM & Butcher EC. Chemokine-like receptor 1 expression and chemerin-directed chemotaxis distinguish plasmacytoid from myeloid dendritic cells in human blood. *Journal of Immunology* 2005 **174** 244–251. (doi:10.4049/jimmunol.174.1.244)
 - 10 Zhao L, Yamaguchi Y, Sharif S, Du XY, Song JJ, Lee DM, Recht LD, Robinson WH, Morser J & Leung LL. Chemerin158K protein is the dominant chemerin isoform in synovial and cerebrospinal fluids but not in plasma. *Journal of Biological Chemistry* 2011 **286** 39520–39527. (doi:10.1074/jbc.M111.258954)
 - 11 Parolini S, Santoro A, Marcenaro E, Luini W, Massardi L, Facchetti F, Communi D, Parmentier M, Majorana A, Sironi M, et al. The role of chemerin in the colocalization of NK and dendritic cell subsets into inflamed tissues. *Blood* 2007 **109** 3625–3632. (doi:10.1182/blood-2006-08-038844)
 - 12 Bozaoglu K, Bolton K, McMillan J, Zimmet P, Jowett J, Collier G, Walder K & Segal D. Chemerin is a novel adipokine associated with obesity and metabolic syndrome. *Endocrinology* 2007 **148** 4687–4694. (doi:10.1210/en.2007-0175)
 - 13 Ernst MC, Haidl ID, Zuniga LA, Dranse HJ, Rourke JL, Zabel BA, Butcher EC & Sinal CJ. Disruption of the chemokine-like receptor-1 (CMKLR1) gene is associated with reduced adiposity and glucose intolerance. *Endocrinology* 2012 **153** 672–682. (doi:10.1210/en.2011-1490)
 - 14 Ernst MC, Issa M, Goralski KB & Sinal CJ. Chemerin exacerbates glucose intolerance in mouse models of obesity and diabetes. *Endocrinology* 2010 **151** 1998–2007. (doi:10.1210/en.2009-1098)
 - 15 Goralski KB, McCarthy TC, Hanniman EA, Zabel BA, Butcher EC, Parlee SD, Muruganandan S & Sinal CJ. Chemerin, a novel adipokine that regulates adipogenesis and adipocyte metabolism. *Journal of Biological Chemistry* 2007 **282** 28175–28188. (doi:10.1074/jbc.M700793200)
 - 16 Roh SG, Song SH, Choi KC, Katoh K, Wittamer V, Parmentier M & Sasaki SI. Chemerin-A new adipokine that modulates adipogenesis via its own receptor. *Biochemical and Biophysical Research Communications* 2007 **362** 1013–1018. (doi:10.1016/j.bbrc.2007.08.104)
 - 17 Sell H, Laurencikiene J, Taube A, Eckardt K, Cramer A, Horrichs A, Arner P & Eckel J. Chemerin is a novel adipocyte-derived factor inducing insulin resistance in primary human skeletal muscle cells. *Diabetes* 2009 **58** 2731–2740. (doi:10.2337/db09-0277)
 - 18 Parlee SD, Ernst MC, Muruganandan S, Sinal CJ & Goralski KB. Serum chemerin levels vary with time of day and are modified by obesity and tumor necrosis factor- α . *Endocrinology* 2010 **151** 2590–2602. (doi:10.1210/en.2009-0794)
 - 19 Parlee SD, McNeil JO, Muruganandan S, Sinal CJ & Goralski KB. Elastase and trypsin govern TNF α -mediated production of active chemerin by adipocytes. *PLoS ONE* 2012 **7** e51072. (doi:10.1371/journal.pone.0051072)
 - 20 Röss C, Tschoner A, Engl J, Klaus A, Tilg H, Ebenbichler CF, Patsch JR & Kaser S. Effect of bariatric surgery on circulating chemerin levels. *European Journal of Clinical Investigation* 2010 **40** 277–280. (doi:10.1111/j.1365-2362.2010.02255.x)
 - 21 Sell H, Divoux A, Poitou C, Basdevant A, Bouillot JL, Bedossa P, Tordjman J, Eckel J & Clement K. Chemerin correlates with markers for fatty liver in morbidly obese patients and strongly decreases after weight loss induced by bariatric surgery. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 2892–2896. (doi:10.1210/jc.2009-2374)
 - 22 van Herpen NA, Sell H, Eckel J, Schrauwen P & Mensink RP. Prolonged fasting and the effects on biomarkers of inflammation and on adipokines in healthy lean men. *Hormone and Metabolic Research* 2013 **45** 378–382. (doi:10.1055/s-0032-1330015)
 - 23 Chang SS, Eisenberg D, Zhao L, Adams C, Leib R, Morser J & Leung L. Chemerin activation in human obesity. *Obesity* 2016 **24** 1522–1529. (doi:10.1002/oby.21534)
 - 24 Maghsoudi Z, Kelishadi R & Hosseinzadeh-Attar MJ. Association of chemerin levels with anthropometric indexes and C-reactive protein in obese and non-obese adolescents. *ARYA Atherosclerosis* 2015 **11** 102–108.
 - 25 Sell H & Eckel J. Chemotactic cytokines, obesity and type 2 diabetes: in vivo and in vitro evidence for a possible causal correlation? *Proceedings of the Nutrition Society* 2009 **68** 378–384. (doi:10.1017/s0029665109990218)
 - 26 Tan BK, Chen J, Farhatullah S, Adya R, Kaur J, Heutling D, Lewandowski KC, O'Hare JP, Lehnert H & Randeve HS. Insulin and metformin regulate circulating and adipose tissue chemerin. *Diabetes* 2009 **58** 1971–1977. (doi:10.2337/db08-1528)
 - 27 Yang S, Wang Q, Huang W, Song Y, Feng G, Zhou L & Tan J. Are serum chemerin levels different between obese and non-obese polycystic ovary syndrome women? *Gynecological Endocrinology* 2016 **32** 38–41. (doi:10.3109/09513590.2015.1075501)
 - 28 Berg V, Sveinbjornsson B, Bendiksen S, Brox J, Meknas K & Figenschau Y. Human articular chondrocytes express ChemR23 and chemerin; ChemR23 promotes inflammatory signalling upon binding the ligand chemerin (21–157). *Arthritis Research and Therapy* 2010 **12** R228. (doi:10.1186/ar3215)
 - 29 Graham KL, Zabel BA, Loghavi S, Zuniga LA, Ho PP, Sobel RA & Butcher EC. Chemokine-like receptor-1 expression by central nervous system-infiltrating leukocytes and involvement in a model of autoimmune demyelinating disease. *Journal of Immunology* 2009 **183** 6717–6723. (doi:10.4049/jimmunol.0803435)
 - 30 Zabel BA, Nakae S, Zuniga L, Kim JY, Ohyama T, Alt C, Pan J, Suto H, Soler D, Allen SJ, et al. Mast cell-expressed orphan receptor CCRL2 binds chemerin and is required for optimal induction of IgE-mediated passive cutaneous anaphylaxis. *Journal of Experimental Medicine* 2008 **205** 2207–2220. (doi:10.1084/jem.20080300)
 - 31 Bondue B, Vosters O, de Nadai P, Glineur S, De Henau O, Luangsay S, Van Gool F, Communi D, De Vuyst P, Desmecht D, et al. ChemR23 dampens lung inflammation and enhances anti-viral immunity in a mouse model of acute viral pneumonia. *PLoS Pathogens* 2011 **7** e1002358. (doi:10.1371/journal.ppat.1002358)
 - 32 Cash JL, Hart R, Russ A, Dixon JP, Colledge WH, Doran J, Hendrick AG, Carlton MB & Greaves DR. Synthetic chemerin-derived peptides suppress inflammation through ChemR23. *Journal of Experimental Medicine* 2008 **205** 767–775. (doi:10.1084/jem.20071601)
 - 33 Luangsay S, Wittamer V, Bondue B, De Henau O, Rouger L, Brait M, Franssen JD, de Nadai P, Huaux F & Parmentier M. Mouse ChemR23 is expressed in dendritic cell subsets and macrophages, and mediates an anti-inflammatory activity of chemerin in a lung disease model. *Journal of Immunology* 2009 **183** 6489–6499. (doi:10.4049/jimmunol.0901037)
 - 34 Zhao RJ & Wang H. Chemerin/ChemR23 signaling axis is involved in the endothelial protection by K(ATP) channel opener iptakalim. *Acta Pharmacologica Sinica* 2011 **32** 573–580. (doi:10.1038/aps.2011.19)
 - 35 John H, Hierer J, Haas O & Forssmann WG. Quantification of angiotensin-converting-enzyme-mediated degradation of human chemerin 145–154 in plasma by matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry. *Analytical Biochemistry* 2007 **362** 117–125. (doi:10.1016/j.ab.2006.12.001)

- 36 Rutkowski P, Sledzinski T, Zielinska H, Lizakowski S, Goyke E, Szrok-Wojtkiewicz S, Swierczynski J & Rutkowski B. Decrease of serum chemerin concentration in patients with end stage renal disease after successful kidney transplantation. *Regulatory Peptides* 2012 **173** 55–59. (doi:10.1016/j.regpep.2011.09.005)
- 37 Barnea G, Strapps W, Herrada G, Berman Y, Ong J, Kloss B, Axel R & Lee KJ. The genetic design of signaling cascades to record receptor activation. *PNAS* 2008 **105** 64–69. (doi:10.1073/pnas.0710487105)
- 38 Parlee SD, Wang Y, Poirier P, Lapointe M, Martin J, Bastien M, Cianflone K & Goralski KB. Biliopancreatic diversion with duodenal switch modifies plasma chemerin in early and late post-operative periods. *Obesity* 2015 **23** 1201–1208. (doi:10.1002/oby.21084)
- 39 Dong B, Ji W & Zhang Y. Elevated serum chemerin levels are associated with the presence of coronary artery disease in patients with metabolic syndrome. *Internal Medicine* 2011 **50** 1093–1097. (doi:10.2169/internalmedicine.50.5025)
- 40 Lehrke M, Becker A, Greif M, Stark R, Laubender R, von Ziegler F, Lebherz C, Tittus J, Reiser M, Becker C, *et al.* Chemerin is associated with markers of inflammation and components of the metabolic syndrome but does not predict coronary atherosclerosis. *European Journal of Endocrinology* 2009 **161** 339–344. (doi:10.1530/EJE-09-0380)
- 41 Tonjes A, Fasshauer M, Kratzsch J, Stumvoll M & Bluher M. Adipokine pattern in subjects with impaired fasting glucose and impaired glucose tolerance in comparison to normal glucose tolerance and diabetes. *PLoS ONE* 2010 **5** e13911. (doi:10.1371/journal.pone.0013911)
- 42 Yang M, Yang G, Dong J, Liu Y, Zong H, Liu H, Boden G & Li L. Elevated plasma levels of chemerin in newly diagnosed type 2 diabetes mellitus with hypertension. *Journal of Investigative Medicine* 2010 **58** 883–886. (doi:10.2310/JIM.0b013e3181ec5db2)
- 43 Wittamer V, Gregoire F, Robberecht P, Vassart G, Communi D & Parmentier M. The C-terminal nonapeptide of mature chemerin activates the chemerin receptor with low nanomolar potency. *Journal of Biological Chemistry* 2004 **279** 9956–9962. (doi:10.1074/jbc.M313016200)
- 44 Barnard SA, Pieters M & De Lange Z. The contribution of different adipose tissue depots to plasma plasminogen activator inhibitor-1 (PAI-1) levels. *Blood Reviews* 2016 **30** 421–429. (doi:10.1016/j.blre.2016.05.002)
- 45 Stelmanska E, Sledzinski T, Turyn J, Presler M, Korczynska J & Swierczynski J. Chemerin gene expression is regulated by food restriction and food restriction-refeeding in rat adipose tissue but not in liver. *Regulatory Peptides* 2013 **181** 22–29. (doi:10.1016/j.regpep.2012.12.001)
- 46 Bauer S, Bala M, Kopp A, Eisinger K, Schmid A, Schneider S, Neumeier M & Buechler C. Adipocyte chemerin release is induced by insulin without being translated to higher levels in vivo. *European Journal of Clinical Investigation* 2012 **42** 1213–1220. (doi:10.1111/j.1365-2362.2012.02713.x)
- 47 Rourke JL, Dranse HJ & Sinal CJ. CMKLR1 and GPR1 mediate chemerin signaling through the RhoA/ROCK pathway. *Molecular and Cellular Endocrinology* 2015 **417** 36–51. (doi:10.1016/j.mce.2015.09.002)
- 48 Graham KL, Zhang JV, Lewen S, Burke TM, Dang T, Zoudilova M, Sobel RA, Butcher EC & Zabel BA. A novel CMKLR1 small molecule antagonist suppresses CNS autoimmune inflammatory disease. *PLoS ONE* 2014 **9** e112925. (doi:10.1371/journal.pone.0112925)
- 49 Yoshimura T & Oppenheim JJ. Chemokine-like receptor 1 (CMKLR1) and chemokine (C-C motif) receptor-like 2 (CCRL2); two multifunctional receptors with unusual properties. *Experimental Cell Research* 2011 **317** 674–684. (doi:10.1016/j.yexcr.2010.10.023)
- 50 Rourke JL, Dranse HJ & Sinal CJ. Towards an integrative approach to understanding the role of chemerin in human health and disease. *Obesity Reviews* 2013 **14** 245–262. (doi:10.1111/obr.12009)
- 51 Catalan V, Gomez-Ambrosi J, Rodriguez A, Ramirez B, Rotellar F, Valenti V, Silva C, Gil MJ, Salvador J & Fruhbeck G. Increased levels of chemerin and its receptor, chemokine-like receptor-1, in obesity are related to inflammation: tumor necrosis factor-alpha stimulates mRNA levels of chemerin in visceral adipocytes from obese patients. *Surgery for Obesity and Related Diseases* 2013 **9** 306–314. (doi:10.1016/j.soard.2011.11.001).
- 52 Bozaoglu K, Segal D, Shields KA, Cummings N, Curran JE, Comuzzie AG, Mahaney MC, Rainwater DL, Vandenberg JL, Maccluer JW, *et al.* Chemerin is associated with metabolic syndrome phenotypes in a Mexican American Population. *Journal of Clinical Endocrinology and Metabolism* 2009 **94** 3085–3088. (doi:10.1210/jc.2008-1833)
- 53 Chakaroun R, Raschpichler M, Kloting N, Oberbach A, Flehmig G, Kern M, Schon MR, Shang E, Lohmann T, Dressler M, *et al.* Effects of weight loss and exercise on chemerin serum concentrations and adipose tissue expression in human obesity. *Metabolism* 2012 **61** 706–714. (doi:10.1016/j.metabol.2011.10.008).
- 54 El-Mesallamy HO, El-Derany MO & Hamdy NM. Serum omentin-1 and chemerin levels are interrelated in patients with Type 2 diabetes mellitus with or without ischaemic heart disease. *Diabetic Medicine* 2011 **28** 1194–1200. (doi:10.1111/j.1464-5491.2011.03353.x)
- 55 Hu W & Feng P. Elevated serum chemerin concentrations are associated with renal dysfunction in type 2 diabetic patients. *Diabetes Research and Clinical Practice* 2011 **91** 159–163. (doi:10.1016/j.diabres.2010.11.016)
- 56 Takahashi M, Inomata S, Okimura Y, Iguchi G, Fukuoka H, Miyake K, Koga D, Akamatsu S, Kasuga M & Takahashi Y. Decreased serum chemerin levels in male Japanese patients with type 2 diabetes: sex dimorphism. *Endocrine Journal* 2013 **60** 37–44. (doi:10.1507/endocrj.EJ12-0201)
- 57 Weigert J, Obermeier F, Neumeier M, Wanninger J, Filarsky M, Bauer S, Aslanidis C, Rogler G, Ott C, Schaffler A, *et al.* Circulating levels of chemerin and adiponectin are higher in ulcerative colitis and chemerin is elevated in Crohn's disease. *Inflammatory Bowel Diseases* 2010 **16** 630–637. (doi:10.1002/ibd.21091)

Received in final form 18 October 2016

Accepted 8 November 2016

Accepted Preprint published online 8 November 2016

