

The innate immune receptor Dectin-2 mediates the phagocytosis of cancer cells by Kupffer cells for the suppression of liver metastasis

Yoshitaka Kimura^a, Asuka Inoue^{a,b}, Sho Hangai^{a,c}, Shinobu Saijo^d, Hideo Negishi^a, Junko Nishio^a, Sho Yamasaki^e, Yoichiro Iwakura^f, Hideyuki Yanai^{a,c}, and Tadatsugu Taniguchi^{a,c,1}

^aDepartment of Molecular Immunology, Institute of Industrial Science, The University of Tokyo, Tokyo 153-8505, Japan; ^bJapan Research and Open Innovation, Sanofi K.K., Tokyo 163-1488, Japan; ^cMax Planck–The University of Tokyo Center for Integrative Inflammation, Tokyo 153-8505, Japan; ^dDepartment of Molecular Immunology, Medical Mycology Research Center, Chiba University, Chiba 260-8673, Japan; ^eDivision of Molecular Immunology, Medical Institute of Bioregulation, Kyushu University, Fukuoka 812-8582, Japan; and ^fCenter for Animal Disease Models, Research Institute for Biomedical Sciences, Tokyo University of Science, Chiba 278-0022, Japan

Contributed by Tadatsugu Taniguchi, October 30, 2016 (sent for review October 23, 2016; reviewed by Ruslan Medzhitov and Nobuyuki Tanaka)

Tumor metastasis is the cause of most cancer deaths. Although metastases can form in multiple end organs, the liver is recognized as a highly permissive organ. Nevertheless, there is evidence for immune cell-mediated mechanisms that function to suppress liver metastasis by certain tumors, although the underlying mechanisms for the suppression of metastasis remain elusive. Here, we show that Dectin-2, a C-type lectin receptor (CLR) family of innate receptors, is critical for the suppression of liver metastasis of cancer cells. We provide evidence that Dectin-2 functions in resident macrophages in the liver, known as Kupffer cells, to mediate the uptake and clearance of cancer cells. Interestingly, Kupffer cells are selectively endowed with Dectin-2-dependent phagocytotic activity, with neither bone marrow-derived macrophages nor alveolar macrophages showing this potential. Concordantly, subcutaneous primary tumor growth and lung metastasis are not affected by the absence of Dectin-2. In addition, macrophage C-type lectin, a CLR known to be complex with Dectin-2, also contributes to the suppression of liver metastasis. Collectively, these results highlight the hitherto poorly understood mechanism of Kupffer cell-mediated control of metastasis that is mediated by the CLR innate receptor family, with implications for the development of anticancer therapy targeting CLRs.

liver metastasis | C-type lectin receptor | Dectin-2 | Kupffer cell | phagocytosis

Metastasis to distal organs is a critical pathological feature of cancer malignancies. Among the types of metastatic disease, liver metastasis occurs in many cancer types and is strongly correlated with poor prognosis (1). Colon cancer is notable, in that ~80% of metastasis is confined to the liver (2). Therefore, the understanding on how liver metastasis is controlled is of general interest to both basic and clinical tumor immunology.

Innate immune cells, such as Kupffer cells and natural killer (NK) cells, are known to play critical roles in the regulation of liver metastasis (3). Numerous highly specialized cell types are distributed within the sinusoidal structure of the liver, with hepatocytes composing a major proportion of the total number (4). The cells of the innate immune system, including Kupffer cells, NK cells, NKT cells, and dendritic cells (DCs), also reside within the sinusoid, where they play a role in immunity (5). Kupffer cells are particularly critical to the maintenance of homeostasis, with their absence resulting in pathogen invasion and/or systemic inflammation (6). On the other hand, Kupffer cells also can contribute to pathogenesis, as has been reported in such conditions as nonalcoholic fatty liver disease, in which the innate receptor Toll-like receptor (TLR) 4 plays a critical role (7). Thus, like many other components of the innate immune system, the appropriate functional activity of Kupffer cells is critical to the maintenance of a healthy organism.

Compared with other tissue macrophages, Kupffer cells have some unique features, such as their high phagocytotic ability (8). It

also has been reported that Kupffer cells can directly kill cancer cells through the secretion of cytotoxic molecules, such as tumor necrosis factor (TNF)- α and reactive oxygen species, and Kupffer cells enhance antitumor responses mediated by other immune cells, such as NK cells (9). On the other hand, several reports have argued that Kupffer cells also have a protumorigenic effect through the production of inflammatory cytokines and chemokines, which contribute to extracellular matrix remodeling and angiogenesis (9). Thus, the actual role of Kupffer cells in liver metastasis has warranted further investigation.

Previously we showed that Dectin-1, a C-type lectin receptor (CLR) family member, plays a critical role in the suppression of tumor growth and metastasis by NK cells. This suppression is indirect, in that cancer cell recognition by Dectin-1 results in the receptor activation in DCs and macrophages, which in turn can enhance the tumoricidal activity of NK cells (10). That study prompted us to study whether other CLR family receptors manifest a similar or distinct antitumor function in the innate antitumor responses, particularly in the context of the regulation of metastasis. Dectin-2 was of particular interest because of its high sequence homology to Dectin-1 (11) and because, similarly if not identical to Dectin-1, it recognizes high-mannose carbohydrate structures presented on bacteria and fungi (12). Also of interest

Significance

The liver is a common site for metastatic disease, and liver metastasis is strongly correlated with poor prognosis. Therefore, an understanding of how liver metastasis is regulated by the immune system is one of the most important issues in cancer immunology. Liver-resident immune cells may either suppress or promote liver metastasis. In this study, we show that Dectin-2 and macrophage C-type lectin, both of which belong to the C-type lectin family of innate receptors, is expressed on resident liver macrophages known as Kupffer cells and play critical roles in the suppression of liver metastasis by enhancing the cells' phagocytotic activity against cancer cells. Our study sheds light on the protective role of Kupffer cells in liver metastasis with therapeutic implications.

Author contributions: Y.K., H.Y., and T.T. designed research; Y.K., A.I., S.H., and H.Y. performed research; S.S., S.Y., and Y.I. contributed new reagents/analytic tools; Y.K., A.I., S.H., H.N., J.N., and H.Y. analyzed data; and Y.K., H.Y., and T.T. wrote the paper.

Reviewers: R.M., Yale University School of Medicine; and N.T., Institute of Gerontology, Nippon Medical School.

The authors declare no conflict of interest.

Data deposition: The microarray data reported in this paper have been deposited in the Gene Expression Omnibus (GEO) database, www.ncbi.nlm.nih.gov/geo (accession no. GSE88809).

¹To whom correspondence should be addressed. Email: tada@m.u-tokyo.ac.jp.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1617903113/-DCSupplemental.

was the fact that stimulation of Dectin-2 induces phagocytosis through signaling by Fc receptor γ chain (FcR γ) (13).

Here we provide evidence for the antimetastatic function of Dectin-2 in Kupffer cells in the liver. We first show the enhancement of cancer cell metastasis in the livers of mice deficient in the Dectin-2 gene. We also report that among the liver-resident cells, Dectin-2 is dominantly expressed in Kupffer cells, and that the removal of these cells also results in enhanced metastasis. Interestingly, Kupffer cells engulf cancer cells, a process that is impaired by the absence of Dectin-2. Such Dectin-2-mediated activity is specific to Kupffer cells, because neither bone marrow-derived macrophages (BMDMs) nor alveolar macrophages engulf the same cancer cells in Dectin-2-dependent manner.

Furthermore, we also present evidence that macrophage C-type lectin (MCL; also known as Dectin-3), which is known to form a heterodimer with Dectin-2 (14), contributes to the suppression of liver metastasis by enhancing the phagocytotic activity of Kupffer cells, indicating that Dectin-2 and MCL cooperatively suppress liver metastasis. These findings shed light on the hitherto poorly understood mechanism of Kupffer cell-mediated control of liver metastasis and suggest the promising prospect of the manipulation of CLR-mediated antitumor responses for controlling liver metastasis.

Results

Selective Contribution of Dectin-2 to the Suppression of Liver Metastasis. We first examined the contribution of Dectin-2 to antitumor immunity by evaluating wild-type (WT) and Dectin-2-deficient (Dectin-2 KO) mice for subcutaneous tumor growth, lung metastasis, and liver metastasis of colon carcinoma cell line SL4, Lewis lung carcinoma cell line 3LL, and melanoma cell line B16F1 and B16F10, all of which have been demonstrated to undergo metastasis in mice (15–18). We found that Dectin-2 KO mice showed more metastatic nodules in the liver compared with WT mice at 14 d after the intrasplenic inoculation with SL4, B16F1, or B16F10 cells, but not after inoculation with 3LL cells (Fig. 1A). Consistently, liver weight was significantly increased in the mice inoculated with these three cell lines (Fig. 1B). In the case of SL4 cell liver metastasis, the tumor-replaced area was approximately 10-fold larger in Dectin-2 KO mice compared with WT mice (Fig. S1).

Interestingly, notable differences between WT and Dectin-2 KO mice were not observed for subcutaneous tumor growth and lung metastasis of SL4, 3LL, B16F1, or B16F10 cancer cell lines (Fig. 1C and D). These observations are quite different from what we previously reported for another CLR family member, Dectin-1, the absence of which affects the subcutaneous growth and lung metastasis of some cancer cells (10), and indicate that Dectin-2 is selectively involved in the suppression of liver metastasis.

Essential Role of Kupffer Cells in Dectin-2-Mediated Suppression of Liver Metastasis. Cancer cell metastasis, which is initiated by cells leaving the primary site and entering the circulation, proceeds in a stepwise manner that includes, for example, cell adhesion to the endothelial wall of the target organ, extravasation, establishment of micrometastatic colonies, and subsequent tumor growth (1). Previous studies have shown that when cancer cells are intrasplenically inoculated, approximately one-half of the cells are extravasated by around 24 h after inoculation and undergo micrometastasis 4 d later (19, 20). To determine the stage of the metastatic event that is suppressed by Dectin-2, we inoculated SL4 cells expressing green fluorescent protein (GFP) into mice and monitored GFP mRNA in the liver at various time points thereafter. We found a marked increase in GFP mRNA levels in the livers of Dectin-2 KO mice compared with the livers of WT mice as early as 12 h after cancer cell inoculation (Fig. S2A). These results suggest that Dectin-2 mediates antitumor responses during an early phase of liver metastasis, perhaps before or during the extravasation step.




Fig. 1. Selective contribution of Dectin-2 to the suppression of liver metastasis. (A and B) SL4 cells (2×10^5 cells), 3LL cells (3×10^5 cells), B16F1 cells (1×10^6 cells), or B16F10 cells (2×10^5 cells) were inoculated into the spleens of WT and Dectin-2 KO mice. Fourteen days later, the livers were observed macroscopically (A) and liver weights were measured (B). (Scale bar: 1 cm.) (C) SL4 cells (2×10^5 cells), 3LL cells (5×10^5 cells), B16F1 cells (5×10^5 cells), or B16F10 cells (1×10^5 cells) were inoculated s.c. into WT and Dectin-2 KO mice, and tumor volumes were measured every 3 or 4 d. (D) SL4-GFP cells (3×10^5 cells), 3LL-GFP cells (3×10^5 cells), B16F1 cells (1×10^6 cells), or B16F10 cells (5×10^5 cells) were inoculated i.v. into WT and Dectin-2 KO mice, and the metastatic levels of SL4-GFP and 3LL-GFP cells were evaluated by quantifying GFP mRNA in the lung on day 12. The numbers of B16F1 and B16F10 colonies in the lung were counted at 14 d after inoculation. Data are shown as mean \pm SEM. * $P < 0.05$. N.S., not significant.

We next asked which cell types use Dectin-2 for the antitumor response. To address this question, we assessed cells residing in the liver for Dectin-2 expression by flow cytometry analysis of the cellular populations. As shown in Fig. 2A, CD11b⁺ F4/80⁺ cells expressed Dectin-2 at high levels, whereas neither CD11c⁺ cells nor CD11b⁺ Gr1⁺ cells expressed Dectin-2. In addition, Dectin-2 expression was not observed on NK cells, CD4⁺ T cells, CD8⁺ T cells, or CD45⁻ T cells (Fig. S2B). Perhaps expectedly, hepatocytes showed little if any expression of Dectin-2 mRNA (Fig. S2C). These observations suggest that CD11b⁺ F4/80⁺ Kupffer cells are critically involved in the Dectin-2-mediated suppression of liver metastasis.

To further address the role of Dectin-2 in Kupffer cells in the suppression of metastasis, we treated WT and Dectin-2 KO mice with clodronate liposomes to deplete macrophages during the early phase of liver metastasis. We found that the administration of clodronate liposomes markedly enhanced SL4 metastasis in both WT and Dectin-2 KO mice, with no significant difference in tumor burden between them (Fig. 2B and C and Fig. S2D and E). This result indicates that Kupffer cells play a central role in Dectin-2-triggered antitumor immunity against liver metastasis.

It has been reported that gut commensal microbiota can influence tumor development in tissues distal from the intestine (21, 22). Given that Dectin-2 expression in intestinal tissues has been




Fig. 2. Requirement of Kupffer cells for the Dectin-2-mediated antitumor system against liver metastasis. (A) Dectin-2 expression on liver-residing cells was analyzed with flow cytometry. Plots gated on CD45⁺ cells are shown. (B and C) WT and Dectin-2 KO mice were treated with control (ctrl) liposomes or clodronate liposomes 2 d before and after intrasplenic inoculation of SL4 cells (2×10^5 cells). On day 10, the livers were collected. Macroscopic images of the liver (B) and liver weights (C) are shown. (Scale bar: 1 cm.) Data are displayed as mean \pm SEM. ** $P < 0.01$. N.S., not significant.

reported (23), and Dectin-2 is known to recognize high-mannose carbohydrate structures present on bacteria and fungi (12), we next examined whether commensal bacteria and fungi are involved in the suppression of liver metastasis by Dectin-2. We treated WT mice with antibacterial or antifungal antibiotics and then inoculated the mice with SL4 cells to examine liver metastasis. We found no significant increase in SL4 cell metastatic levels in the livers of mice treated with these antibiotics (Fig. S2 F and G), further supporting the idea that Dectin-2 in Kupffer cells acts directly on cancer cells.

Dectin-2-Dependent Phagocytotic Activity of Kupffer Cells Against Cancer Cells. Flow cytometry analysis of the cell composition in livers revealed similar proportions and numbers of Kupffer cells in WT and Dectin-2 KO mice (Fig. S3 A–C). These data implicate Dectin-2 in the regulation of Kupffer cell function, but not in Kupffer cell expansion, in the suppression of liver metastasis.

How does Dectin-2 function in Kupffer cells? Because Dectin-2 can trigger the engulfment of its ligand into cells (24, 25), we first asked whether Kupffer cells engulf cancer cells in a Dectin-2-dependent manner. Kupffer cells sorted from WT or Dectin-2 KO mice were cocultured with carboxyfluorescein diacetate succinimidyl ester (CFSE)-labeled SL4 cells, and then subjected to flow cytometry analysis. As shown in Fig. 3A, this coculturing of SL4 and Kupffer cells resulted in a significant increase in CFSE intensity in WT Kupffer cells, indicating engulfment of the SL4 cancer cells by Kupffer cells. Consistent with this, we observed SL4 cells engulfed by Kupffer cells by confocal microscopy and time-lapse imaging (Fig. 3B and Movie S1). Notably, the intensity of CFSE was significantly lower in the Dectin-2-deficient Kupffer cells (Fig. 3A). These results indicate that Dectin-2 enhances the

phagocytotic activity of Kupffer cells against SL4 cancer cells. However, the fact that Dectin-2 deficiency did not completely abrogate this phagocytotic activity suggests the involvement of additional molecule(s) in this process (Fig. 3A). Interestingly, 3LL cells, which underwent liver metastasis independently of Dectin-2 (Fig. 1 A and B), showed more marked resistance to the engulfment by Kupffer cells compared with SL4 cells (Fig. S3D).

We also analyzed the phagocytotic potential of BMDMs and alveolar macrophages, and found that neither of these cells showed any significant dependence on Dectin-2 for phagocytotic activity against SL4 cells (Fig. S3 E and F), even though both exhibited substantial levels of Dectin-2 expression (Fig. S3G). This observation is consistent with the foregoing data showing that subcutaneous tumor growth and lung metastasis are not affected by Dectin-2 deficiency (Fig. 1 C and D).

Interestingly, after SL4 cells were cocultured with Kupffer cells, we found a decreased proportion of cells negative for DAPI, an indicator of dead cells, among the unphagocytosed SL4 cells (Fig. S3H). This finding suggests that Kupffer cells may actively engulf living cancer cells. Indeed, the number of SL4 cells was reduced after coculturing with Kupffer cells in vitro, which was inhibited by the treatment of Kupffer cells with cytochalasin D, a phagocytosis inhibitor (Fig. 3C). Furthermore, a significantly higher number of SL4 cells was observed after cancer cells were cocultured with Dectin-2-deficient Kupffer cells (Fig. 3D),




Fig. 3. Dectin-2-dependent engulfment and clearance of cancer cells by Kupffer cells. (A) Kupffer cells (1×10^5 cells) isolated from WT and Dectin-2 KO mice were cocultured with or without CFSE-labeled SL4 cells (0.25×10^5 cells) for 2 h. The CFSE intensity in CD45⁺ CD11b⁺ F4/80⁺ cells was analyzed by flow cytometry. (Left) Representative histograms of CFSE level. The cells with a CFSE level exceeding the red line were identified as CFSE⁺ cells. (Right) Proportion of CFSE⁺ cells. (B) Kupffer cells (1×10^5 cells) were cocultured with CFSE-labeled SL4 cells (0.25×10^5 cells) and after 2 h, the cells were observed by confocal microscopy. (C) CFSE-labeled SL4 cells (0.25×10^5 cells) were cultured in the presence or absence of Kupffer cells (1×10^5 cells) pretreated with DMSO or cytochalasin D, and the number of PI⁻ CD45⁻ CFSE⁺ cells was determined at 24 h after culturing. (D) CFSE-labeled SL4 cells (0.25×10^5 cells) were cultured in the presence or absence of Kupffer cells (1×10^5 cells) derived from WT and Dectin-2 KO mice, and the number of PI⁻ CD45⁻ CFSE⁺ cells was determined at 24 h after culturing. Data are shown as mean \pm SEM. * $P < 0.05$. N.S., not significant.

further supporting the role of Dectin-2 in the phagocytosis of cancer cells by Kupffer cells (26).

We next asked whether Dectin-2 can recognize a particular molecular structure on cancer cells. We previously showed that soluble Dectin-1 conjugated to human IgG1 Fc (Dectin-1-Fc) can bind to the surface of cancer cells (10). We then generated Dectin-2-Fc (soluble Dectin-2; sDectin-2) and evaluated its binding to SL4 cells. We were unable to detect any significant binding of sDectin-2 to SL4 cells, however (Fig. S4A). This finding may indicate either that the affinity of sDectin-2 is too low to allow detection of its binding or that Dectin-2 does not directly bind to the cells. In addition, because Dectin-2 recognizes carbohydrate structures on bacteria and fungi (12), we also examined the effects of *N*-glycosidase or *O*-glycosidase treatment of SL4 cells on the differential phagocytotic activity between WT and Dectin-2-deficient Kupffer cells. We found that the activity remained essentially unchanged after treatment with either of these glycosidases (Fig. S4B). Therefore, unlike the previous study showing the Dectin-1 recognition of *N*-glycan structures on cancer cells (10), *N/O*-glycan structures on SL4 cells are dispensable for the Dectin-2-mediated engulfment by Kupffer cells. As such, the nature of the Dectin-2-mediated recognition of cancer cells requires future investigation.

The foregoing observations raise the interesting question of whether Dectin-2 signaling is involved in the phagocytotic activity of Kupffer cells. Because Dectin-2 signaling is known to induce the expression of inflammatory cytokines (24, 25), we examined whether the interaction between Kupffer cells and cancer cells results in Dectin-2-dependent gene expression for various cytokine mRNAs by coculturing these cells. Although the expression levels of interleukin 6 (*Il6*), *Il23a*, chemokine (C-X-C motif) ligand 1 (*Cxcl1*), and chemokine (C-C motif) ligand 2 (*Ccl2*) mRNA were up-regulated by the coculture, Dectin-2 deficiency did not affect the mRNA expression levels (Fig. S4C and D). We also examined the mRNA expression profile by microarray analysis (GEO accession no. GSE88809), and found no Dectin-2-dependent induction of any mRNA measured. These results lend support to the notion that Dectin-2 participates in the engulfment of cancer cells by Kupffer cells independently of gene induction.

Of final note, we also found no evidence of Dectin-2-dependent skewing of Kupffer cell polarization toward an M1 or M2 type in this experimental setting. In WT and Dectin-2-deficient Kupffer cells cocultured with SL4 cells, the mRNA expression signatures for M1 polarization (*Il6*, *Il23a*, and *Tnf*) and those for M2 polarization [arginases, liver (*Arg1*) and mannose receptor, C-type 1 (*Mrc1*; *Cd206*)] remained unchanged (Fig. S4D and E). Although further clarification may be necessary, these observations suggest that Kupffer cells exert their antimetastatic function *in vivo* without their polarization.

Contribution of CLR Family Member MCL (Dectin-3) to the Suppression of Liver Metastasis. Previous studies have shown that Dectin-2's ability to protect the host from fungal infection requires that it form a complex with another CLR member, MCL (or Dectin-3) (14). MCL regulates another CLR, Mincle, by enhancing its cell surface expression (27). Mincle signaling inhibits signal transduction downstream of Dectin-1 (28). Given our finding of mRNA expression for these three CLRs in Kupffer cells (Fig. 4A), we examined whether Dectin-1, MCL, and Mincle play roles in liver metastasis. Intrasplenic inoculation of SL4 cells into mice deficient in any of these CLR members revealed that a deficiency of MCL or Dectin-1, but not of Mincle, aggravated liver metastasis, indicating that MCL and Dectin-1 are also involved in the antimetastatic immune response in the liver (Fig. 4B and C).

Consistent with the foregoing *in vivo* data, MCL-deficient Kupffer cells showed weaker engulfing activity for SL4 cells compared with WT cells (Fig. 4D). The decrease in SL4 cell uptake in MCL-deficient Kupffer cells was not due to the down-regulation of Dectin-2, given that Dectin-2 expression on MCL KO mouse-derived Kupffer cells was similar to that on WT Kupffer cells (Fig. S5A). It is unlikely that the impaired phagocytotic activity of




Fig. 4. MCL-mediated uptake of cancer cells by Kupffer cells and suppression of liver metastasis. (A) Expression levels of *Mcl*, *Dectin-1*, and *Mincle* mRNAs in hepatocytes and Kupffer cells were analyzed by qRT-PCR. (B and C) SL4 cells (2×10^5 cells) were inoculated into the spleens of WT, MCL KO, Dectin-1 KO, and Mincle KO mice. On day 14, macroscopic images of livers were obtained (B), and livers were weighed (C). (Scale bar: 1 cm.) (D) Kupffer cells collected from WT and MCL KO mice were cocultured with or without CFSE-labeled SL4 cells (0.25×10^5 cells) for 2 h. The CFSE intensity in CD45⁺ CD11b⁺ F4/80⁺ cells was analyzed by flow cytometry. (Left) Representative histograms of CFSE levels. Cells with a CFSE level exceeding the red line were identified as CFSE⁺ cells. (Right) Proportion of CFSE⁺ cells. Data are displayed as mean \pm SEM. * $P < 0.05$. N.S., not significant; N.D., not detected.

Dectin-2-deficient Kupffer cells is caused by MCL down-regulation, because *Mcl* mRNA expression remained unaffected in Dectin-2-deficient Kupffer cells (Fig. S5B). Taken together, these results suggest that, similar to the antifungal innate response, MCL contributes to the engulfment of cancer cells by Kupffer cells by pairing with Dectin-2.

In contrast to MCL, although Kupffer cells expressed *Dectin-1* mRNA at high levels (Fig. 4A), the phagocytotic activity of Kupffer cells against SL4 cells was not affected by Dectin-1 deficiency (Fig. S5C). Instead, Dectin-1 induced antitumor killing mediated by liver nonparenchymal cells (NPCs) (Fig. S5D). Indeed, NK cells are mainly responsible for the cytotoxic activity of liver NPCs (29), and this observation is consistent with our previous report showing that Dectin-1 signaling in DCs and macrophages enhances NK cell-mediated tumoricidal activity (10). Of note, neither Dectin-2 nor MCL contributed to this cytotoxic response of the NPCs (Fig. S5D and E), further supporting the idea that, unlike Dectin-1-mediated antitumor responses, Dectin-2- and MCL-mediated antimetastatic responses are mediated by enhanced phagocytotic activity of Kupffer cells.

Discussion

Liver metastasis is a feature of many types of malignant cancers and is correlated with poor prognosis. Although innate immune cells have been identified as key players in the control of liver metastasis, innate immune receptors, such as TLRs, can promote

tumor metastasis (30, 31). The role of CLRs in mediating anti-metastatic responses has remained largely elusive. Here we provide evidence that Dectin-2, a CLR family member, promotes the engulfment and clearance of cancer cells by Kupffer cells and suppresses liver metastasis. This suppression mechanism is not operational for all cancer cells, however; the liver metastasis level for the 3LL cancer cell line remained unaffected in the absence of Dectin-2 (Fig. 1 *A* and *B*). It is worth recalling that Dectin-1 plays a critical role in controlling the metastasis of 3LL cancer cells through activation of NK cells (10). Thus, these results indicate the differential contribution of CLR members in the control of metastasis; that is, Dectin-2, together with MCL, contributes to the phagocytosis of cancer cells by Kupffer cells, whereas Dectin-1 contributes to the NK cell-mediated killing of cancer cells.

Interestingly, the Dectin-2-mediated antitumor response is selective for Kupffer cells, as demonstrated by our finding that Dectin-2 deficiency did not affect cancer cell engulfment by BMDMs and alveolar macrophages. Nor is Dectin-2 involved in the suppression of lung metastasis or subcutaneous tumor growth. Moreover, our results show that MCL, a heterodimeric counterpart of Dectin-2, is also critical for the suppression of liver metastasis, consistent with the idea that Dectin-2 and MCL function through heterodimeric complex formation (14). These findings provide insights into the mechanism of the antimetastatic response mediated by the CLR innate receptor family, i.e., the enhanced phagocytosis of cancer cells by Kupffer cells.

Analysis of the mechanism of Dectin-2-mediated suppression of metastasis revealed that Dectin-2 facilitates the engulfment of SL4 cells by Kupffer cells, but not by BMDMs or alveolar macrophages (Fig. 3*A* and Fig. S3*E* and *F*). These results suggest that Dectin-2 functions as a phagocytotic receptor against cancer cells selectively in Kupffer cells. Kupffer cells exhibit greater phagocytotic ability than other types of macrophages, such as alveolar and peritoneal macrophages (8); thus, Kupffer cells may have some specific molecular features that enhance phagocytotic activity, with Dectin-2 involved in their unique phagocytotic function against cancer cells. Previous studies have shown that several CLR family members, including CLEC4G, CD207, and CLEC4F, are more highly expressed in Kupffer cells compared with other macrophages (32); therefore, it is possible that the Dectin-2–MCL complex cooperates with such CLRs, leading to the selective antitumor function in Kupffer cells. This is an interesting issue that will be addressed in future studies.

We further examined the antitumor potential of Kupffer cells and found data suggesting that the ability of living cell-engulfing Kupffer cells to eliminate cancer cells depends on Dectin-2 (Fig. 3 *C* and *D* and Fig. S3*H*). Previous studies have shown that macrophages engulf dead cells to trigger adaptive immune responses through antigen presentation to T cells, and regulate the tumor microenvironment by producing immune mediators in response to cancer (33). It was recently suggested that macrophages phagocytose living cancer cells and suppress tumor development (26, 34), although the molecular mechanism for this remains largely unknown. In view of our present study, it will be interesting to examine the contributions of CLR family members other than Dectin-2 and MCL to these antitumor responses.

Dectin-2-mediated phagocytosis against fungi is associated with the up-regulation of inflammatory cytokines, such as IL-6 and CXCL1 (24, 25). Nevertheless, our analysis of Dectin-2-dependent gene induction in response to cancer cells revealed that Dectin-2 did not regulate the mRNA expression of any tested genes, even *Il6* and *Cxcl1* (Fig. S4*D*). These observations suggest the interesting idea that the Dectin-2-triggered response in Kupffer cells against cancer cells is distinct from the responses reported previously. Supporting this notion, CARD9, a signaling molecule downstream of Dectin-2, was found to promote the liver metastasis of SL4 cells by manipulating the tumor microenvironment (35), which is inconsistent with our findings. Moreover, NLRP3, which is known to

be activated dependently on Dectin-2 (13), enhances NK cell-mediated antitumor killing to suppress liver metastasis (36), although we found that Dectin-2 was dispensable for NK cell-mediated cytotoxicity against SL4 cells (Fig. S5*D*). Therefore, Dectin-2-triggered phagocytosis of cancer cells may be mediated by a unique signaling pathway.

The molecular nature of how Kupffer cells recognize some, but not all, cancer cells merits further examination. We could not identify the binding of sDectin-2 to cancer cells, and it is possible that, similar to the antifungal response (14), Dectin-2 may need to associate with MCL for the recognition of structure(s) associated with cancer cells. Alternatively, ligand recognition by the Dectin-2–MCL complex might not be required, in that the complex may augment the phagocytotic activity induced by other receptor molecules that recognize cancer cell-associated ligands. Clearly, this is an interesting issue for future studies.

At present, however, we cannot exclude the possibility that conventional signal transduction mediated by FcR γ downstream of Dectin-2 contributes to the Kupffer cell-mediated phagocytosis. FcR γ activation induces endocytosis through the phosphorylation of its own ITAM motif (37). Such posttranslational modifications of the FcR γ are also observed when an anti-Dectin-2 agonistic antibody is taken up by macrophages (25). Previous studies have shown that Mincle transduces FcR γ -mediated signaling pathways (13); however, our data indicate that Mincle is not involved in the control of liver metastasis (Fig. 4 *B* and *C*). Therefore, Dectin-2 may possess some selectivity in sensing cancer cells to induce antitumor responses.

Of note, it has been reported that treatment with an anti-CD47 antibody effectively suppresses *in vivo* tumor growth without co-administration of other chemotherapeutic agents (38, 39). Given that CD47 is a well-known “don’t eat me” signal for inhibiting phagocytosis (40), the uptake of cancer cells by phagocytes is a promising target for anticancer therapy, particularly when combined with a method that accelerates the cancer cell phagocytosis. As such, our findings may reveal a way to enhance the phagocytotic activity of macrophages by developing an agonist for Dectin-2 and/or other CLR family members for controlling metastasis in the liver and other organs.

Materials and Methods

Mice. C57BL/6 mice were purchased from CLEA Japan. *Clec7a*^{−/−} mice (Dectin-1 KO mice), *Clec4n*^{−/−} mice (Dectin-2 KO mice), *Clec4e*^{−/−} mice (Mincle KO mice), and *Clec4d*^{−/−} mice (MCL KO mice) on a C57BL/6 background were generated as described previously (41–44). All animal experiments were approved and performed in accordance with guidelines of The University of Tokyo’s Animal Research Committee.

Cells. Mouse colon carcinoma cell line SL4 was kindly provided by Dr. T. Irimura (The University of Tokyo). Mouse melanoma cell lines B16F1 and B16F10 and Lewis lung carcinoma cell line 3LL were maintained as described previously (10). GFP-transduced SL4 cells (SL4-GFP) and 3LL cells (3LL-GFP) were prepared as described previously (10). Mouse embryonic fibroblasts (MEFs) were retrovirally transfected with pmCherry-N1 vector (Clontech) and used as MEF-mCherry cells after selection with puromycin.

Liver Metastasis Model. The liver metastasis model has been described previously (36). In brief, after the mouse was anesthetized, the spleen was exposed from a small incision in the left flank, and 2×10^5 SL4 cells, 3×10^5 3LL cells, 1×10^6 B16F1 cells, or 2×10^5 B16F10 cells were inoculated into the spleen. Five minutes later, the spleen was excised, and the incision was closed by clip. The mouse was killed on day 14, followed by the macroscopic observation of the liver and the measurement of liver weight. For the evaluation of tumor burden at early stage of liver metastasis, 2×10^5 SL4-GFP cells were inoculated, and liver specimens were collected 4, 8, and 12 h later. The GFP mRNA level in the liver was measured by quantitative RT-PCR (qRT-PCR).

Additional information is provided in [SI Materials and Methods](#).

ACKNOWLEDGMENTS. We thank Y. Miyake, M. Oh-hora, and K. Shibata for helpful comments and advice, and M. Sugahara, M. Taniguchi, T. Mizutani,

S. Chiba, and members of the FACS core laboratory of the Institute of Medical Science, The University of Tokyo for technical assistance. This work is supported in part by Grant-In-Aid for Scientific Research (S) 15638461 from the Ministry of Education, Culture, Sports, Science, and Technology

and by Grant 15656877 from the Japan Agency for Medical Research and Development. The Department of Molecular Immunology at The University of Tokyo is supported by BONAC Corporation and Kyowa Hakko Kirin Co., Ltd.

1. Valastyan S, Weinberg RA (2011) Tumor metastasis: Molecular insights and evolving paradigms. *Cell* 147(2):275–292.
2. Manfredi S, et al. (2006) Epidemiology and management of liver metastases from colorectal cancer. *Ann Surg* 244(2):254–259.
3. Van den Eynden GG, et al. (2013) The multifaceted role of the microenvironment in liver metastasis: Biology and clinical implications. *Cancer Res* 73(7):2031–2043.
4. Protzer U, Maini MK, Knolle PA (2012) Living in the liver: Hepatic infections. *Nat Rev Immunol* 12(3):201–213.
5. Heymann F, Tacke F (2016) Immunology in the liver—from homeostasis to disease. *Nat Rev Gastroenterol Hepatol* 13(2):88–110.
6. Bilzer M, Roggel F, Gerbes AL (2006) Role of Kupffer cells in host defense and liver disease. *Liver Int* 26(10):1175–1186.
7. Bieghs V, Trautwein C (2013) The innate immune response during liver inflammation and metabolic disease. *Trends Immunol* 34(9):446–452.
8. Laskin DL, Weinberger B, Laskin JD (2001) Functional heterogeneity in liver and lung macrophages. *J Leukoc Biol* 70(2):163–170.
9. Paschos KA, Majeed AW, Bird NC (2010) Role of Kupffer cells in the outgrowth of colorectal cancer liver metastases. *Hepatol Res* 40(1):83–94.
10. Chiba S, et al. (2014) Recognition of tumor cells by Dectin-1 orchestrates innate immune cells for anti-tumor responses. *eLife* 3:e04177.
11. Arizumi K, et al. (2000) Cloning of a second dendritic cell-associated C-type lectin (dectin-2) and its alternatively spliced isoforms. *J Biol Chem* 275(16):11957–11963.
12. McGreal EP, et al. (2006) The carbohydrate-recognition domain of Dectin-2 is a C-type lectin with specificity for high mannose. *Glycobiology* 16(5):422–430.
13. Sancho D, Reis e Sousa C (2012) Signaling by myeloid C-type lectin receptors in immunity and homeostasis. *Annu Rev Immunol* 30:491–529.
14. Zhu LL, et al. (2013) C-type lectin receptors Dectin-3 and Dectin-2 form a heterodimeric pattern-recognition receptor for host defense against fungal infection. *Immunity* 39(2):324–334.
15. Morimoto-Tomita M, Ohashi Y, Matsubara A, Tsuiji M, Irimura T (2005) Mouse colon carcinoma cells established for high incidence of experimental hepatic metastasis exhibit accelerated and anchorage-independent growth. *Clin Exp Metastasis* 22(6):513–521.
16. Cullen R, Germanov E, Shimaoka T, Johnston B (2009) Enhanced tumor metastasis in response to blockade of the chemokine receptor CXCR6 is overcome by NKT cell activation. *J Immunol* 183(9):5807–5815.
17. Qi K, et al. (2004) Impact of cirrhosis on the development of experimental hepatic metastases by B16F1 melanoma cells in C57BL/6 mice. *Hepatology* 40(5):1144–1150.
18. Bezuhly M, et al. (2009) Role of activated protein C and its receptor in inhibition of tumor metastasis. *Blood* 113(14):3371–3374.
19. Martin MD, et al. (2010) Rapid extravasation and establishment of breast cancer micrometastases in the liver microenvironment. *Mol Cancer Res* 8(10):1319–1327.
20. Ritsma L, et al. (2012) Intravital microscopy through an abdominal imaging window reveals a pre-micrometastasis stage during liver metastasis. *Sci Transl Med* 4(158):158ra145.
21. Rutkowski MR, et al. (2015) Microbially driven TLR5-dependent signaling governs distal malignant progression through tumor-promoting inflammation. *Cancer Cell* 27(1):27–40.
22. Iida N, et al. (2013) Commensal bacteria control cancer response to therapy by modulating the tumor microenvironment. *Science* 342(6161):967–970.
23. Taylor PR, et al. (2005) Dectin-2 is predominantly myeloid restricted and exhibits unique activation-dependent expression on maturing inflammatory monocytes elicited in vivo. *Eur J Immunol* 35(7):2163–2174.
24. Ifrim DC, et al. (2014) Role of Dectin-2 for host defense against systemic infection with *Candida glabrata*. *Infect Immun* 82(3):1064–1073.
25. Sato K, et al. (2006) Dectin-2 is a pattern recognition receptor for fungi that couples with the Fc receptor γ chain to induce innate immune responses. *J Biol Chem* 281(50):38854–38866.
26. Feng M, et al. (2015) Macrophages eat cancer cells using their own calreticulin as a guide: Roles of TLR and Btk. *Proc Natl Acad Sci USA* 112(7):2145–2150.
27. Miyake Y, Matsutugu OH, Yamasaki S (2015) C-type lectin receptor MCL facilitates Mincle expression and signaling through complex formation. *J Immunol* 194(11):5366–5374.
28. Wevers BA, et al. (2014) Fungal engagement of the C-type lectin mincle suppresses dectin-1-induced antifungal immunity. *Cell Host Microbe* 15(4):494–505.
29. Cohen SA, Tzung SP, Doerr RJ, Goldrosen MH (1990) Role of asialo-GM1-positive liver cells from athymic nude or polyinosinic-polycytidylic acid-treated mice in suppressing colon-derived experimental hepatic metastasis. *Cancer Res* 50(6):1834–1840.
30. Kim S, et al. (2009) Carcinoma-produced factors activate myeloid cells through TLR2 to stimulate metastasis. *Nature* 457(7225):102–106.
31. Yu LX, et al. (2014) Platelets promote tumour metastasis via interaction between TLR4 and tumour cell-released high-mobility group box1 protein. *Nat Commun* 5:5256.
32. Okabe Y, Medzhitov R (2014) Tissue-specific signals control reversible program of localization and functional polarization of macrophages. *Cell* 157(4):832–844.
33. Clarke C, Smyth MJ (2007) Calreticulin exposure increases cancer immunogenicity. *Nat Biotechnol* 25(2):192–193.
34. Kopatz J, et al. (2013) Siglec-h on activated microglia for recognition and engulfment of glioma cells. *Glia* 61(7):1122–1133.
35. Yang M, et al. (2014) Tumor cell-activated CARD9 signaling contributes to metastasis-associated macrophage polarization. *Cell Death Differ* 21(8):1290–1302.
36. Dupaul-Chicoine J, et al. (2015) The Nlrp3 inflammasome suppresses colorectal cancer metastatic growth in the liver by promoting natural killer cell tumoricidal activity. *Immunity* 43(4):751–763.
37. Williams M, Bruhns P, Saeys Y, Hammad H, Lambrecht BN (2014) The function of Fc γ receptors in dendritic cells and macrophages. *Nat Rev Immunol* 14(2):94–108.
38. Weiskopf K, et al. (2016) CD47-blocking immunotherapies stimulate macrophage-mediated destruction of small-cell lung cancer. *J Clin Invest* 126(7):2610–2620.
39. Liu X, et al. (2015) CD47 blockade triggers T cell-mediated destruction of immunogenic tumors. *Nat Med* 21(10):1209–1215.
40. Brown GC, Neher JJ (2012) Eaten alive! Cell death by primary phagocytosis: “Phagoptosis”. *Trends Biochem Sci* 37(8):325–332.
41. Saijo S, et al. (2007) Dectin-1 is required for host defense against *Pneumocystis carinii* but not against *Candida albicans*. *Nat Immunol* 8(1):39–46.
42. Saijo S, et al. (2010) Dectin-2 recognition of α -mannans and induction of Th17 cell differentiation is essential for host defense against *Candida albicans*. *Immunity* 32(5):681–691.
43. Yamasaki S, et al. (2009) C-type lectin Mincle is an activating receptor for pathogenic fungus, *Malassezia*. *Proc Natl Acad Sci USA* 106(6):1897–1902.
44. Miyake Y, et al. (2013) C-type lectin MCL is an Fc γ -coupled receptor that mediates the adjuvanticity of mycobacterial cord factor. *Immunity* 38(5):1050–1062.
45. Yanai H, et al. (2013) Conditional ablation of HMGB1 in mice reveals its protective function against endotoxemia and bacterial infection. *Proc Natl Acad Sci USA* 110(51):20699–20704.