





Draft Genome Sequence of Extended-Spectrum- β -Lactamase-Producing *Escherichia coli* Strain CCUG 62462, Isolated from a Urine Sample

Anna Johnning,^{a,b} Hedvig E. Jakobsson,^{b,c} Fredrik Boulund,^{a,b} Francisco Salvà-Serra,^{b,c,d} © Edward R. B. Moore,^{b,c} Christina Åhrén,^{b,c,e} Nahid Karami,^{b,c} Erik Kristiansson^{a,b}

Department of Mathematical Sciences, Chalmers University of Technology, Gothenburg, Sweden^a; Centre for Antibiotic Resistance Research (CARe) at University of Gothenburg, Gothenburg, Gothenburg, Sweden^b; Department of Infectious Diseases, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden^c; Department of Biology, University of the Balearic Islands, Palma de Mallorca, Spain^d; Swedish Strategic Programme Against Antibiotic Resistance, Västra Götaland, Sweden^e

The draft genome sequence has been determined for an extended-spectrum- β -lactamase (ESBL)-producing ($bla_{CTX-M-15}$) Escherichia coli strain (CCUG 62462), composed of 119 contigs and a total size of 5.27 Mb. This E. coli is serotype O25b and sequence type 131, a pandemic clonal group, causing worldwide antimicrobial-resistant infections.

Received 14 October 2016 Accepted 21 October 2016 Published 15 December 2016

Citation Johnning A, Jakobsson HE, Boulund F, Salvà-Serra F, Moore ERB, Åhrén C, Karami N, Kristiansson E. 2016. Draft genome sequence of extended-spectrum-β-lactamase-producing Escherichia coli strain CCUG 62462, isolated from a urine sample. Genome Announc 4(6):e01382-16. doi:10.1128/genomeA.01382-16.

Copyright © 2016 Johnning et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license. Address correspondence to Nahid Karami, nahid.karami@microbio.gu.se.

he Escherichia coli sequence type 131 (ST131) has emerged as a globally dispersed pathogen, the predominant lineage causing extraintestinal infections, and is often resistant to antibiotic treatment (1). The E. coli ST131 clonal group carries a multitude of resistance and virulence genes, including extended-spectrum β-lactamases (ESBLs). The ESBL-producing *E. coli* strain E10394 (= CCUG 62462) was isolated from a urinary tract infection (UTI) sample as part of an outbreak of ESBL-producing Enterobacteriaceae in a neonatal postsurgery ward at Sahlgrenska University Hospital, Gothenburg, Sweden, in 2008 (2). The strain was characterized as resistant to cefotaxime, tobramycin, and trimethoprim but susceptible to ciprofloxacin, and it was defined as an ESBL-producing strain. It was further characterized as sequence type 131 and serotype O25b. The strain was cultured on blood agar medium, and genomic DNA was isolated from bacterial biomass using a PureLink genomic DNA minikit (Invitrogen, Carlsbad, CA, USA). The genome was sequenced with an Illumina MiSeq instrument (Genomics Core Facility at University of Gothenburg), generating 1,427,585 paired-end reads of 250 bp, yielding a total of 357 Mb. The sequence reads were trimmed using Trim Galore! version 0.3.7 (http://www.bioinformatics.babraham.ac.uk/projects/trim _galore/) with command line arguments "—stringency 3—retain_unpaired." After quality filtering, 1,424,760 paired reads and 2,356/75 unpaired forward/reverse reads were retained for analysis. The high-quality reads were assembled *de novo* with SPAdes version 3.7.0 (3), with command line arguments "-carefulcov_cutoff 5." The resulting assembly consisted of 119 contigs, with a total assembly length of 5.27 Mb. The longest contig measured 515,034 bp, the N_{50} was 328,690 bp, and the average genome coverage of the assembly was $64\times$. The G+C content of the genome sequence was estimated at 50.8%. The genome sequence was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (4); 5,104 coding sequences and 80 tRNAs were identified. Using ResFinder (5), a number of antibiotic resistance

genes were found: bla_{CTX-M-15} on contig LZET01000074, bla_{TEM-1} on contig LZET01000061, aac(3)-IIa on contig LZET01000076, and aadA5, mph(A), sul1, and dfrA17 on contig LZET01000047. PlasmidFinder (6) identified the following plasmid marker genes: IncFIA (nucleotide accession no. AP001918) on contig LZET01000040, IncFIB (accession no. AP001918) on contig LZET01000042, and *IncFII* (accession no. AY458016) on contigs LZET01000050, LZET01000068, and LZET01000072. In addition, genes involved in type IV secretion systems (T4SS) were identified, using CONJscan (7), included genes belonging to the mating-pair formation family F (MPF_F), relaxases (MOB_F, MOB_P, and MOB_O), and type IV coupling proteins (T4CP). Three small plasmids were completely assembled. First, a 5,627-bp plasmid (contig LZET01000060), which was 99% identical to the pJJ1886_ 3 plasmid previously described in E. coli strain JJ1886 (accession no. CP006787.1), was found. Second, a 5,166-bp plasmid (contig LZET01000063), and last, a 1,718-bp plasmid (contig LZET01000087), which was 99% identical to plasmid pEC648 previously described in E. coli strain ST648 (accession no. CP008716.1), were found.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. LZET00000000. The version described in this paper is version LZET00000000.1.

ACKNOWLEDGMENTS

The Culture Collection University of Gothenburg (CCUG; http://www.ccug.se) is supported by the Department of Clinical Microbiology, Sahlgrenska University Hospital. *Escherichia coli* strain CCUG 62462 was isolated in the Clinical Bacteriology Laboratory of Sahlgrenska University Hospital. We acknowledge the efforts of the Genomics Core Facility (http://cf.gu.se/Genomics), Sahlgrenska Academy, University of Gothenburg.

F.S.-S. was supported by a stipend for Basic and Advanced Research from the CCUG, through the Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg.

FUNDING INFORMATION

This work, including the efforts of Nahid Karami, Edward R. B. Moore, and Erik Kristiansson, was funded by Swedish Västra Götaland (437221). This work, including the efforts of Hedvig E. Jakobsson, Anna Johnning, Fredrik Boulund, Francisco Salvà Serra, Edward R. B. Moore, and Erik Kristiansson, was funded by European Commission (EC) (602860).

REFERENCES

- Nicolas-Chanoine M-H, Bertrand X, Madec JY. 2014. Escherichia coli ST131, an intriguing clonal group. Clin Microbiol Rev 27:543–574. http:// dx.doi.org/10.1128/CMR.00125-13.
- Karami N, Helldal L, Welinder-Olsson C, Ahrén C, Moore ER. 2013. Sub-typing of extended-spectrum-beta-lactamase-producing isolates from a nosocomial outbreak: application of a 10-loci generic *Escherichia coli* multi-locus variable number tandem repeat analysis. PLoS One 8:e83030. http://dx.doi.org/10.1371/journal.pone.0083030.
- 3. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV,

- Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. http://dx.doi.org/10.1089/cmb.2012.0021.
- Tatusova T, Ciufo S, Federhen S, Fedorov B, McVeigh R, O'Neill K, Tolstoy I, Zaslavsky L. 2015. Update on RefSeq microbial genomes resources. Nucleic Acids Res 43:D599–D605. http://dx.doi.org/10.1093/nar/ gku1062.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother 67:2640–2644. http:// dx.doi.org/10.1093/jac/dks261.
- Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. Antimicrob Agents Chemother 58:3895–3903. http://dx.doi.org/ 10.1128/AAC.02412-14.
- Guglielmini J, de la Cruz F, Rocha EP. 2013. Evolution of conjugation and type IV secretion systems. Mol Biol Evol 30:315–331. http://dx.doi.org/ 10.1093/molbev/mss221.