

THE PROBLEM OF "DORMANCY" IN BACTERIAL SPORES¹

JACKSON W. FOSTER AND E. STATEN WYNNE²

Department of Bacteriology, The University of Texas, Austin, Texas

Received for publication January 21, 1948

Our experience with the germination of spores of various species of clostridia, especially *Clostridium botulinum*, has led to conclusions that we believe warrant the consideration of investigators of this group of organisms and of the aerobic sporeformers as well. Pertinent observations have been made by previous authors, but it appears that all the recent available evidence on, and various aspects of, the dormancy problem have not hitherto been crystallized into one comprehensive concept.

The essential points already established (Morrison and Rettger, 1930*a,b*, for review of previous literature; Curran and Evans, 1937; Olsen and Scott, 1946; Wynne and Foster, 1948*a*; Foster and Wynne, 1948) are these:

(1) Colony counts from a spore suspension may be submaximum in a few days' incubation, and increase with time over prolonged periods. This indicates delayed germination and is a well-known phenomenon for both aerobic and anaerobic spores. This is the characteristic feature of classical dormancy.

(2) The effect depends on the medium used, being marked in some and non-existent in others.

(3) The final spore counts of a given suspension depend greatly upon the medium employed, though in each medium counts are at the maximum obtainable in it.

(4) Certain substances, notably soluble starch (0.1 per cent), eliminate the delayed germination in some media. In others the starch increases the final maximum count, whereas in still others it has no apparent effect. In general, all complex media preferred for counting spores show the starch effect.

Data exemplifying all these points may be found in table 1. From the information in the literature, as well as in table 1, the following points may be made:

(a) The so-called "dormancy" in these organisms is not an inherent function of the spores, but rather of the medium in which the spores are placed to germinate.

(b) Even when "dormancy" occurred, or in media with low final counts, the large size of colonies speaks against a nutritional inadequacy.

¹ The research which this paper reports was undertaken in co-operation with the Committee on Food Research of the Quartermaster Food and Container Institute for the Armed Forces. The views or conclusions contained in this report are those of the authors. They are not to be construed as necessarily reflecting the views or having the endorsement of the War Department.

² Present address: Department of Plant Sciences, University of Oklahoma, Norman, Oklahoma.

(c) Accordingly, a plausible conclusion is that "dormancy," i.e., delayed germination, and also submaximal counts are due to substances in normal media inhibitory to spore germination. "Dormancy" would then be only a manifestation of antigermination substances in the medium and seemingly is, therefore, merely an artifact.

(d) In the case of the characteristic delayed rate of spore germination during "dormancy," one may consider this to be a function of subinhibitory concentrations of certain substances in the normal medium.

TABLE 1
C. botulinum 62A spore recoveries in various agar media*

BASAL MEDIUM	1 PER CENT SOLUBLE STARCH	INCUBATION, DAYS			
		3	6	17	33
Yesair's plus 0.1% starch.....	—	45	41	36†	—
Difco nutrient.....	—	28	30	22	—
Difco nutrient.....	+	35	42	35	—
Difco brain heart infusion.....	—	4	6	6	—
Difco brain heart infusion.....	+	12	23	30	30
BBL anaerobic.....	—	0	0	2	—
BBL anaerobic.....	+	0	1	6	8
Difco yeast extract, 1.0%.....	—	19	17	15	—
Difco yeast extract, 1.0%.....	+	15	12	11	—
Corn steep liquor solids, 1.0%.....	—	8	10	8	—
Corn steep liquor solids, 1.0%.....	+	7	9	11	14
Fresh beef infusion‡.....	—	43	50	47	—
Fresh beef infusion‡.....	+	60	60	53	—
Fresh liver infusion‡.....	—	24	24	23	—
Fresh liver infusion‡.....	+	46	52	49	—

* All media contained 0.2 per cent glucose and 0.2 per cent thioglycolate; initial pH 7.2 to 7.4.

† The somewhat lower counts in this column as compared to the 6-day values are due to the obscuring of small colonies by the development of gas bubbles and cloudiness in the media.

‡ Prepared exactly like Yesair's medium (cf Wynne and Foster, 1948).

(e) In the case of submaximal counts, the inhibitory substances are of such a nature and concentration that the germination rate of a portion of the spore population is zero, i.e., they never germinate, for all practical purposes.

(f) The inhibitors evidently are of different types, but one is the C₁₈ unsaturated fatty acids (oleic, linoleic, etc.), which are present in almost all complex natural media. The addition of these substances to media suppresses the spore germination rate of clostridia in a manner reminiscent of "dormancy."

(g) The appreciable increase in germination rate, occasioned by the soluble starch in some media, is due to absorption of certain inhibitory substances, including the C₁₈ fatty acids. In some media starch has little or no effect in enhancing counts known to be low.

(h) The different degrees of inactivation of germination inhibitors by starch indicate that different media contain varying amounts of different germination inhibitors. Solvent extraction of some media indicates that some of these naturally occurring inhibitors are nonlipoid.

(i) The significance of these substances inhibitory to germination is evidenced by the fact that the fresh infusions of beef, pork, and liver, ordinarily considered ideal nutrients, contain significant amounts of germination inhibitors, as demonstrated by the considerable enhancement of germination caused by the addition of starch. Incidentally, fresh beef or liver infusion with starch is just as suitable as the widely used (for anaerobes) pork infusion medium of Yesair, with added starch.

(j) Effects similar to that of starch have been observed in the past when supplements containing native proteins and polysaccharides were added to the usual bacteriological media. Though this action has been arbitrarily ascribed to unspecified nutritional factors, an equally plausible idea is that such substances function physically in the manner of starch, through colloid absorption of inhibitory substances, and in some cases this has been proved definitely.

(k) Unless it is shown that no effect is obtained by starch, the foregoing data and ideas make it imperative that this substance be an invariable component of all media in work on anaerobic as well as aerobic spores when maximum counts in the shortest time are desired. Even though starch apparently does not counteract all known cases of retarded germination, it is the simplest over-all treatment available without altering the nutritional status of the basal medium.

(l) The term "dormancy" in the foregoing connection no longer has the degree of validity it formerly enjoyed, and there now appears to be some question as to the justification of its continued use, unless it occurs in cases in which the effect is shown to be independent of the germination medium.

ACKNOWLEDGMENT

We wish to thank Dr. O. B. Williams for his stimulating interest in this project.

REFERENCES

- CURRAN, H. R., AND EVANS, F. R. 1937 The importance of enrichments in the cultivation of bacterial spores previously exposed to lethal agencies. *J. Bact.*, **34**, 179-189.
- FOSTER, J. W., AND WYNNE, E. S. 1948 Physiological studies on spore germination, with special reference to *Clostridium botulinum*. IV. Inhibition of germination by unsaturated C₁₈ fatty acids. *J. Bact.*, **55**, 495-501.
- MORRISON, E. W., AND RETTGER, L. W. 1930a Bacterial spores. I. A study in heat resistance and dormancy. *J. Bact.*, **20**, 299-311.
- MORRISON, E. W., AND RETTGER, L. W. 1930b Bacterial spores. II. A study of bacterial spore germination in relation to environment. *J. Bact.*, **20**, 313-342.
- OLSEN, A. M., AND SCOTT, W. J. 1946 Influence of starch in media used for the detection of heated bacterial spores. *Nature*, **157**, 337.
- WYNNE, E. S., AND FOSTER, J. W. 1948a Physiological studies on spore germination, with special reference to *Clostridium botulinum*. I. Development of a quantitative method. *J. Bact.*, **55**, 61-68.