

Systemic therapy of brain metastases: non–small cell lung cancer, breast cancer, and melanoma

Marc C. Chamberlain, Christina S. Baik, Vijayakrishna K. Gadi, Shailender Bhatia, and Laura Q.M. Chow

Seattle Cancer Center Alliance, Seattle, Washington (M.C.C., C.S.B., V.K.G., S.B., L.Q.M.C.); Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington (C.S.B., V.K.G., L.Q.M.C.); Departments of Neurology and Neurological Surgery, University of Washington, Seattle, Washington (M.C.C.); Division of Medical Oncology, University of Washington, Seattle, Washington (C.S.B., V.K.G., S.B., L.Q.M.C)

Corresponding Author: Marc C. Chamberlain, MD, Seattle Cancer Center Alliance, 825 Eastlake Avenue East, Box 358081 (G6-800), Seattle, Washington 98109-1023; (marccchamberlain@gmail.com)

Abstract

Brain metastases (BM) occur frequently in many cancers, particularly non–small cell lung cancer (NSCLC), breast cancer, and melanoma. The development of BM is associated with poor prognosis and has an adverse impact on survival and quality of life. Commonly used therapies for BM such as surgery or radiotherapy are associated with only modest benefits. However, recent advances in systemic therapy of many cancers have generated considerable interest in exploration of those therapies for treatment of intracranial metastases.

This review discusses the epidemiology of BM from the aforementioned primary tumors and the challenges of using systemic therapies for metastatic disease located within the central nervous system. Cumulative data from several retrospective and small prospective studies suggest that molecularly targeted systemic therapies may be an effective option for the treatment of BM from NSCLC, breast cancer, and melanoma, either as monotherapy or in conjunction with other therapies. Larger prospective studies are warranted to further characterize the efficacy and safety profiles of these targeted agents for the treatment of BM.

Keywords:

Brain metastases | breast cancer | blood-brain barrier | melanoma | non–small cell lung cancer

Involvement of the central nervous system (CNS) is a complication of many cancers. Brain metastases (BM) from systemic malignancies account for the majority of intracranial cancers, with an estimated incidence rate of 8.3 to 11.0 per 100000 as compared with an incidence rate of 6.6 per 100000 for all primary malignant CNS tumors.¹ The development of BM is usually associated with poor prognosis and significant adverse effects on survival and quality of life. Overall, BM are associated with a low 2-year survival rate (8%) and a high burden of neurologic symptoms including headaches, nausea and vomiting, focal motor deficits, cognitive decline, delirium, and seizures.²

BM affect 8%–10% of all cancer patients^{1,3} and 40% of patients with metastatic cancer.^{4,5} The majority of BM originate from lung cancer (40%–50%), breast cancer (15%–25%), and melanoma (5%–20%).^{1,3} BM predominate in the cerebral

hemispheres (80%) followed by the cerebellum (15%) and brainstem (5%), a pattern of distribution reflective of proportional blood flow to these respective regions.⁶ The incidence of BM is believed to be increasing, likely resulting from longer patient survival due to more effective systemic therapies for the primary cancer and the increased use of neuroimaging in neurologically asymptomatic patients.^{7,8}

Treatment options for BM are limited and suboptimal. Historically, the mainstay of therapy has been local treatments such as surgery or radiation therapy (RT) (ie, whole brain radiation therapy (WBRT), stereotactic radiosurgery (SRS), or stereotactic radiation therapy (SRT)). The choice of local therapy is generally guided by the number and location of BM, the extent and prognosis of systemic disease, and the performance status of the patient. Patients with minimal systemic disease, good performance status, and solitary brain

metastasis in a noneloquent location of the brain are often treated with surgical resection followed by RT.⁹ Patients with minimal systemic disease, good performance, and oligometastatic disease in the brain are generally treated with SRT and deferred WBRT.¹⁰ In contrast, patients with greater metastatic burden are usually treated with WBRT despite a lack of randomized trials showing effectiveness of WBRT compared with best standard care; there is an emerging perspective, however, that these patients can be treated with SRS, especially when effective systemic therapy may be available.^{11,12} In a select group of patients (<15% of all patients with BM), aggressive local treatment can prolong survival to ≥ 12 months.¹³ The evolving classification of specific molecular subtypes within most cancers (eg, hormone receptor status in breast cancer or anaplastic lymphoma kinase [*ALK*] rearrangement in non-small cell lung cancer [NSCLC]) will likely alter treatment of BM based on improved prognostication linked to specific tumor subtypes.

Historically, the role of systemic therapy in the treatment of BM has been limited by concerns regarding limited penetration across the blood-brain barrier (BBB), rapid efflux from brain, or intrinsic chemotherapy resistance resulting from multiple prior lines of therapy.¹⁴ Furthermore, patients with symptomatic or uncontrolled BM have generally been excluded from randomized controlled trials of systemic pharmacotherapies. Additionally, CNS outcome measures were often not reported separately from systemic efficacy outcomes. However, novel targeted therapies have substantially improved systemic disease control and survival in molecularly defined cancer populations, which have generated considerable interest in the investigation of these therapies to complement or even replace local therapies for treatment of BM. Although the constraints on parenchymal brain drug delivery also apply to targeted therapies, these agents are increasingly being selected for further study in patients with BM based, in part, on their CNS penetration.

The objective of this review is to summarize the efficacy of emerging systemic therapies, especially targeted therapies and immunotherapies, for the treatment of BM, focusing on the 3 cancers (NSCLC, breast cancer, and melanoma) that most often metastasize to the CNS.

Targeting Brain Metastases: Challenges

When employed in treating BM, a systemic therapy must traverse the BBB, the blood-cerebrospinal fluid (CSF) barrier, and the blood-tumor barrier in sufficient concentration to enable its therapeutic effect (Fig. 1). These various barriers render the brain and CSF inaccessible to most chemotherapeutic drugs because of large size (>150kDa), ionization, hydrophilicity, and/or protein-binding.^{7,14} Additionally, many of the small lipophilic molecules that cross the BBB and blood-tumor barriers can be exported from the brain by highly regulated transmembrane efflux pumps located in the endothelial vasculature of the CNS.^{7,14} Although BM may increase permeability to systemic drugs by disrupting the structural integrity of the BBB and blood-tumor barrier, this is usually not sufficient

to achieve therapeutic levels of most drugs in the CNS.^{7,14} Dose escalation of systemically administered drugs (eg, high-dose methotrexate) is generally limited by systemic toxicity. Furthermore, subtherapeutic concentrations of anticancer drugs achieved in the brain may potentially contribute to acquired treatment resistance.¹⁵

Brain Metastases from Non-small Cell Lung Cancer

Lung cancer, the most common cancer overall, has the highest incidence of BM among all cancers. Approximately 40%–50% of all CNS metastases arise from lung cancer (Table 1), and approximately half of all patients with NSCLC develop BM during the course of their disease. The median overall survival (OS) for patients with BM from NSCLC is 7 months per the Grade Prognostic Assessment (GPA) index.¹⁶ Fortunately, emerging systemic therapies may improve outcomes for this challenging subgroup of patients.

Non-small cell lung cancer is heterogeneous and composed of several molecular subtypes associated with specific driver oncogenes (Fig. 2)^{17,18}; these molecular subtypes are characterized by different prognoses and responses to therapy. Recent advances in the treatment of NSCLC can be credited to improved understanding of the pathogenesis of these molecular subtypes. For example, 10%–35% of NSCLC tumors harbor a somatic activating mutation in the epidermal growth factor receptor (*EGFR*) gene.¹⁹ Patients with activating mutations in *EGFR* have overall response rates up to 85% and progression-free survival (PFS) as long as 13 months with the *EGFR* tyrosine kinase inhibitors (TKIs) gefitinib, erlotinib, or afatinib compared with cytotoxic chemotherapy (up to 38% and 7 mo, respectively).^{20–25} Rearrangement of *ALK* is observed in approximately 5% of the adenocarcinoma population and predicts a good overall response rate to ALKTKIs as with crizotinib (65%) versus cytotoxic chemotherapy (20%).²⁶ Immune checkpoint inhibitors have also led to impressive durable responses in a subgroup of patients.^{27–30} All of these emerging therapies are increasingly proving to be effective in treating patients with BM.

Targeted Therapy with *EGFR* Tyrosine Kinase Inhibitors

Patients with *EGFR*-mutant NSCLC often develop multiple small BM with little peritumoral edema.^{31,32} Efficacy of *EGFR* TKIs in patients with *EGFR*-mutant NSCLC is well established; however, efficacy in patients with BM is not as clear because patients with symptomatic or uncontrolled BM were excluded from pivotal, randomized controlled trials. Consequently, data on the efficacy of TKI therapy in the CNS have been gleaned mostly from retrospective studies of those trials that enrolled patients with BM.

While the low molecular weight and nonpolar nature of TKIs permit passive diffusion across the BBB, many TKIs are substrates for efflux transporter proteins (Table 2).




Fig. 1. Pathways across the blood-brain barrier, blood-cerebrospinal fluid barrier, and blood-tumor barrier. The blood-brain barrier (BBB) is a continuous layer of epithelial cells joined by high-resistance tight junctions surrounded by pericytes and sealed by astrocytic perivascular endfeet.^{7,14} The blood-cerebrospinal fluid (CSF) barrier demarcates the space between the choroid plexus and CSF. Unlike the BBB capillaries, the functional unit of the choroid plexus is fenestrated, has no tight junctions, and thus facilitates molecular transport such as CSF bulk flow, metabolic inactivation, and transcapillary exchange. The blood-tumor barrier refers to a tumor's vasculature that is a variously permeable barrier. Similar to the BBB, it is a major factor limiting the access of many therapeutic agents to brain tumors.¹⁸² The blood-tumor barrier consists of continuous, nonfenestrated capillaries (like those of normal brain), fenestrated capillaries, and interendothelial gaps.¹⁸² Expression of efflux transporters located at the blood-tumor barrier represents an additional mechanism that prevents intracellular penetration of anticancer drugs.¹⁸³ (Brain with tumor adapted with permission under a Creative Commons Attribution License from OpenStax College, <http://legacy.cnx.org/content/m45981/1.4/>; blood-brain barrier reused with permission under a Creative Commons Attribution License from Kübelbeck A, https://commons.wikimedia.org/wiki/File:Blood-brain_barrier_transport_en.png; anatomy of tight junction adapted with permission under a Creative Commons Attribution License from Polakis P. *J Cell Biol.* 2008;183(3):371–373; blood CSF barrier adapted with permission under a Creative Commons Attribution License from Bhaskar S, et al. *Part Fibre Toxicol.* 2010;7:3; blood-tumor barrier adapted from Bredel M. Anticancer drug resistance in primary human brain tumors. *Brain Res Brain Res Rev.* 2001;35(2):161–204, with permission from Elsevier.)

Nevertheless, CSF concentrations of erlotinib, gefitinib, and afatinib exceed those required to inhibit growth of cells harboring *EGFR* mutations in vitro (Table 2). An early prospective study of 41 patients with unselected NSCLC BM treated with gefitinib resulted in 4 (10%) intracranial partial responses with a median duration of response of 13.5 months (Table 3).^{33,34} In another study of unselected patients, a 70% CNS response was observed with first-line erlotinib or gefitinib.³⁵ These patients were mostly Asian, female never-smokers who had a high incidence of *EGFR* mutations; thus, the prevalence of mutation in this study was likely higher than the general NSCLC population.¹⁹

Data from other retrospective studies also suggest that patients with BM from *EGFR*-mutant NSCLC have better outcomes with either WBRT or TKI therapy than patients with BM from *EGFR*-wild-type NSCLC (Table 3).^{31,36,37} A retrospective analysis of 69 cases previously treated with erlotinib reported time to progression within the

brain of 11.7 months for patients with *EGFR* mutations compared with 5.8 months for those with *EGFR*-wild-type or unassessed tumors, despite the fact that only 16% of patients with *EGFR* mutations had received WBRT versus 85% of those without the mutation (Table 3).³⁸ Subsequent prospective phase 2 studies of patients with *EGFR*-mutant NSCLC indicated that TKI therapy provided intracranial responses of $\geq 75\%$.^{39,40}

Despite an initial favorable intracranial response, patients often have CNS progression while maintaining systemic disease control on TKI therapy.^{41,42} A linear correlation is seen between plasma and CSF concentrations of *EGFR* TKIs, suggesting that a higher dose may lead to higher CSF concentration and thereby potentially improve CNS disease response.^{41,43} For instance, in case reports and early phase clinical trials, a response as high as 81% has been reported with an erlotinib twice weekly pulse dose level of 600–1350mg.⁴⁴ However, an independent,

Table 1. Prevalence of brain metastases by subtype of lung, breast, and skin cancers

Primary Tumor Site/Type	Frequency/Prevalence of Brain Metastases
Lung	40%–50% of all BM ^{1,3}
NSCLC	≈50% of NSCLC cases ¹⁸⁵
Somatic <i>EGFR</i> mutant	44% of NSCLC BM cases (despite a genotype prevalence of only 10% in nonsquamous NSCLC) ³¹
<i>ALK</i> positive	35%–50% of NSCLC BM cases ^{26,32,64}
SCLC	40%–50% of SCLC cases (10% of cases at diagnosis) ¹⁸⁶
Breast	15%–25% of all BM ^{1,3}
Triple negative	25%–46% of TNBC cases ^{89,126}
HER2 positive	38% of HER2-positive breast cancer cases ¹⁸⁷
Luminal	5% of luminal breast cancer cases ¹²⁸
Melanoma	5%–20% of all BM ^{1,3}
<i>BRAF</i> WT	≤50% of melanoma cases ^{148,188,189} ; 55%–75% of melanoma cases based on autopsy report ^{1,148,149}
<i>BRAF</i> mutant	Similar frequency of BM as <i>BRAF</i> -WT melanoma ¹⁹⁰

Abbreviations: *ALK*, anaplastic lymphoma kinase; BM, brain metastases; *BRAF*, v-Raf murine sarcoma viral oncogene homolog B; *EGFR*, epidermal growth factor receptor; HER2, human epidermal growth factor receptor 2; NSCLC, non-small cell lung cancer; SCLC, small cell lung cancer; TNBC, triple-negative breast cancer; WT, wild-type.

retrospective study of 10 patients showed minimal efficacy with this strategy.⁴⁵ Similarly equivocal data have been reported regarding use of high-dose afatinib.^{46,47} A novel TKI in development, AZD3759, shows significant CNS penetration in preclinical models and promising early clinical efficacy in *EGFR*-mutant NSCLC patients with BM.⁴⁸

Later-generation *EGFR* inhibitors osimertinib and rociletinib specifically target the T790M resistance mutation in *EGFR* and have demonstrated efficacy (objective response rates, 61% and 59%, and PFS, 9.6 and 13.1 mo, respectively) in patients with the T790M resistance mutation who have progressed on prior TKI therapy.^{49,50} Osimertinib was recently approved for this patient population.⁵¹ Patients with stable treated BM were included in studies for both drugs, with reports of systemic and possible CNS responses in these studies indicating that these novel agents may be active in the CNS.^{52,53}

The use of *EGFR* inhibitors concurrently with RT has been proposed to improve intracranial efficacy based on preclinical studies in murine models.^{54,55} A retrospective analysis of 63 patients reported improved intracranial response rates in patients who received an *EGFR*TKI during WBRT, especially patients with *EGFR* mutations (Table 3).³⁷ In a phase 2 trial of erlotinib in combination with WBRT in 40 patients with BM, overall response rate was 86%, and median survival time was 11.8 months.⁵⁶ Despite the fact that the Radiation Therapy Oncology Group (RTOG) phase 3 trial of WBRT plus SRS with either temozolomide or erlotinib in unselected NSCLC patients with 1–3 BM did not meet its accrual goals and closed prematurely, there was a trend suggesting that adjunctive temozolomide or erlotinib may increase toxicity with a deleterious effect on time to progression and survival.⁵⁷ Similarly, no significant benefit was observed by the addition of gefitinib to WBRT in a phase 2 randomized trial in unselected patients with BM.⁵⁸ Data are currently inadequate to indicate whether TKI therapy administered concurrently with CNS-directed

RT in patients with *EGFR* mutations is beneficial. Given the possible safety concerns raised in the RTOG study, concurrent TKI therapy is not routinely recommended in patients receiving WBRT.

Targeted Therapy with *ALK* Tyrosine Kinase Inhibitors

Although NSCLC with *ALK* rearrangement comprises a small subset of all NSCLC patients (4%–8%), this is an important subpopulation with distinct epidemiology and biology. Patients with *ALK* rearrangement (*ALK*-positive disease) are younger and usually have no or light smoking history. *ALK*-positive tumors are sensitive to *ALK*TKIs, with excellent systemic disease control. Initial findings from clinical trials of patients with *ALK*-positive NSCLC treated with *ALK* TKIs have shown promising CNS responses (Table 3). Crizotinib was the first *ALK* inhibitor approved for treatment of patients with metastatic *ALK*-positive NSCLC. While this drug has demonstrated clinically meaningful disease control, the brain is the most common or only site for disease progression.^{59,60} This observation can likely be attributed to subtherapeutic crizotinib concentrations in the brain (Table 2). Pooled analysis of a phase 3 randomized trial (PROFILE 1007) with a single-arm phase 2 trial (PROFILE 1005) reported a 12-week intracranial disease control rate with crizotinib of 56% among 109 patients with untreated asymptomatic BM compared with 62% in 166 patients with previously treated BM.⁶¹ In the small subset of patients with CNS target lesions at baseline, the confirmed intracranial response rate was 18% in patients who did not receive brain RT and 33% in patients with previously treated BM. Thus, CNS disease control may be achievable with crizotinib initially but is not durable.

Second-generation *ALK* TKIs ceritinib, alectinib, and brigatinib have shown efficacy in crizotinib-resistant patients




Fig. 2. Non-small cell lung cancer mutations. Although more than one-third of driver mutations in non-small cell lung cancer (NSCLC) are still unknown, approximately one-half of primary single mutations have been identified. The approval of targeted therapies for epidermal growth factor receptor (*EGFR*)-mutated and anaplastic lymphoma kinase (*ALK*)-rearranged lung adenocarcinomas has led to a better understanding of individual driver and resistance mutations upon tumor progression.^{17,18} (Large pie chart adapted with permission. © 2016 American Society of Clinical Oncology. All rights reserved.^{17,18})

including activity against BM, in part because of improved BBB penetration (Tables 2 and 3). Ceritinib was approved for treatment of patients with metastatic *ALK*-positive NSCLC whose disease progressed while taking crizotinib or who are intolerant to crizotinib.⁶² Ceritinib is selective for *ALK* at low concentrations in vitro and exhibits activity against crizotinib-resistant tumors in *ALK*-positive NSCLC xenograft models.⁶³ In a phase 1 trial of patients with *ALK*-positive advanced solid tumors (ASCEND-1), approximately half had BM at baseline, and single-agent ceritinib produced an objective response rate of 59% with responses observed in both crizotinib-pretreated and crizotinib-naïve patients.⁶⁴ Median PFS was 6.9 months in crizotinib-pretreated patients and 10.4 months in crizotinib-naïve patients.⁶⁴ In a 1-year follow-up, objective response rates were 56.4% (92/163) and 72% (60/83), respectively, with median duration of response of 8.3 and 17.0 months, respectively.⁶⁵ Half of patients had BM, and the intracranial disease control rate was 65.3% (median time to intracranial response, 6.1wk) in crizotinib-pretreated patients and 79% (median time to intracranial response, 9.9wk) in crizotinib-naïve patients. An ongoing phase 2 study (NCT02336451) is specifically looking at the activity of ceritinib as first-line therapy in crizotinib-treated and crizotinib-naïve patients with untreated, asymptomatic and measurable BM and leptomeningeal disease.⁶⁶

Results from dose-finding phase 1 and phase 2 studies indicate that alectinib has promising antitumor activity in patients with *ALK*-rearranged NSCLC after progression on crizotinib, including those with CNS metastases (Table 3).^{67–69} Intracranial response rates >60% have been reported with alectinib.^{67–69} Based on these data, alectinib was recently approved for treatment of patients with *ALK*-rearranged NSCLC who had progressed post-crizotinib.⁷⁰ In recent phase 2 studies of alectinib in crizotinib-refractory, *ALK*-rearranged NSCLC, 51%–61% of patients

had CNS metastases.^{71,72} Among patients with BM at baseline, the disease control rate was 83%, with a duration of response of 10.3 months. Alectinib is also being investigated specifically in patients with BM in several studies (NCT02075840, NCT02521051, and NCT02604342).⁶⁶

Brigatinib is another second-generation *ALK* inhibitor. In a phase 1 study of 79 patients with *ALK*-positive NSCLC, the response rate and disease control rate were 53% and 87%, respectively, in the 15 patients with measurable BM CNS disease.⁷³ Lorlatinib (PF-06463922) has demonstrated clinical activity in patients with *ALK*-rearranged and *ROS1*-rearranged NSCLC with promising CNS efficacy.^{74,75}

Recent studies have demonstrated a survival benefit with the combination of TKI and RT in patients with BM from *ALK*-rearranged NSCLC. In a retrospective, multi-institution study examining OS and intracranial PFS in 90 patients, treatment with SRS or WBRT and TKIs prolonged survival.⁷⁶

Other potential molecular drivers and signaling transduction pathways—such as *ROS1*, *RET* proto-oncogene (*RET*), mesenchymal-epithelial transition factor receptor tyrosine kinase gene (*MET*), v-Raf murine sarcoma viral oncogene homolog B (*BRAF*), and tyrosine kinase receptor (*TRK*)/tyrosine kinase receptor B (*TRKB*)—are being explored in early-phase clinical trials as therapeutic targets in NSCLC and other cancers. The ability of these agents to penetrate the CNS and elicit intracranial responses will be a major factor in developing these novel targeted therapies and improving the survival of patients with NSCLC.

Immunotherapy

Immune checkpoint inhibitors, particularly those targeting the programmed death 1 (PD-1) pathway, result in

Table 2. Physicochemical and pharmacokinetic properties of new targeted therapies that may be useful for treatment of brain metastases in non-small cell lung cancer, breast cancer, and melanoma

Therapeutic Agent	MW, Da ^a	Log P ^b	BCRP and P-gp Substrate	Inhibitory Concentration, ng/ml	CSF Concentration in Patients with BM, ng/ml	CSF Penetration Rate in Patients with BM, % ^c	Drug Efflux Transporters Restricted CNS Penetration, BCRP/P-gp ^d	Clinical Development Status	
Erlotinib	429.90	2.7	Yes ¹⁸³	Yes ¹⁹¹	7.9 (EGFR-WT NSCLC) ¹⁹²	24–54 (N = 25) ^{192–194}	2.8–5.1 ^{192–194}	Yes ^{183,191}	FDA approved for locally advanced or metastatic EGFR-mutant NSCLC and locally advanced, unresectable or metastatic pancreatic cancer, in combination with gemcitabine
Gefitinib	446.9	3.2	Yes ¹⁹⁵	Yes ¹⁹⁶	0.13 (EGFR-WT NSCLC) ¹⁹⁷	3.7–6.2 (N = 30) ^{194,198}	1.1–1.4 ^{194,198}	Yes ¹⁹⁵	FDA approved for locally advanced or metastatic EGFR-mutant NSCLC
Afatinib	485.9	3.7	Yes ¹⁹⁹	Yes ¹⁹⁹	0.5 ²⁰⁰	0.5 (N = 1) ²⁰⁰	<1 ²⁰⁰	Unknown	FDA approved for metastatic EGFR-mutant NSCLC
Crizotinib	450.3	1.8	Yes ²⁰¹	Yes ¹⁹⁸	108 (EML4-ALK E13;A20 translocation; NCI-H3122) ²⁰² ; 48 (H228) ⁶³	0.62 (N = 1) ²⁰²	0.26 ²⁰²	Yes ²⁰¹	FDA approved for locally advanced or metastatic ALK-positive NSCLC
Ceritinib	558.1	5.0	Unknown	Unknown	11 (Ba/F3 NPM-ALK WT) ²⁰³ ; 2 (H228); 0.08 (ALK enzymatic assay) ⁶³	Unknown	Unknown	Yes ²⁰⁴	FDA approved for patients with ALK-positive NSCLC who have progressed on or are intolerant to crizotinib
Alectinib	482.6	5.5	No ^{6,205}	Unknown	0.92 (ALK cell-free assay) ⁶⁷	1.3 ⁶⁷	86 ⁶⁷	No ²⁰⁵	FDA breakthrough therapy designation for patients with ALK-positive NSCLC who have progressed on crizotinib
Nivolumab	146000	N/A	N/A	N/A	Not expected to cross intact BBB. Mechanism of action is in the periphery on T cells, which then cross the BBB	Not expected to cross intact BBB	Not expected to cross the BBB		FDA approved for: (1) advanced squamous NSCLC after platinum-based chemotherapy; and (2) unresectable or metastatic melanoma and disease progression following ipilimumab and, if BRAF-V600-mutation positive, a BRAF inhibitor
Trastuzumab	145000	N/A	N/A	N/A	43.5 (p185 ^{HER2} extra-cellular domain) ²⁰⁶	Not expected to cross intact BBB			FDA approved for HER2-overexpressing breast cancer, and HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma
Lapatinib	580.5	5.1 ²⁰⁷	Yes ²⁰⁷	Yes ²⁰⁷	6 (EGFR and HER2) ²⁰⁷	1.3–4.5 (N = 2) ²⁰⁸	0.9–1.3 ²⁰⁸	Yes ²⁰⁷	FDA approved as part of combination treatment for advanced or metastatic (with an anthracycline, a taxane, and trastuzumab) and postmenopausal (with letrozole) HER2-overexpressing breast cancer

Therapeutic Agent	MW, Da ^a	Log <i>p</i> ^b	BCRP and P-gp Substrate	Inhibitor	Median Inhibitory Concentration, ng/ml	CSF Concentration in Patients with BM, ng/ml	CSF Penetration Rate in Patients with BM, % ^c	Drug Efflux Transporters Restricted CNS Penetration, BCRP/P-gp ^d	Clinical Development Status
Vemurafenib	489.9	5.1	Yes ^{209,210}	Yes ²¹⁰		470 (N = 6) ²¹¹	0.98 ²¹¹	Yes ^{209,212}	FDA approved for unresectable or metastatic melanoma with BRAF-V600E mutation
Dabrafenib	615.7	5.4	Yes ^{162,213}	Unknown			Unknown	Yes ¹⁶²	FDA approved for unresectable or metastatic melanoma with BRAF-V600E mutation and in combination with trametinib for BRAF-V600E and BRAF-V600K mutations
Ipilimumab	148000	N/A	N/A	N/A	<148 (CTLA-4) ²¹⁴		Not expected to cross intact BBB. Mechanism of action is in the periphery on T cells, which then cross the BBB		FDA approved for unresectable or metastatic melanoma
Pembrolizumab	149000	N/A	N/A	N/A			Not expected to cross intact BBB. Mechanism of action is in the periphery on T cells, which then cross the BBB		FDA approved for unresectable or metastatic melanoma and disease progression following ipilimumab and, if BRAF-V600 mutation positive, a BRAF inhibitor

Abbreviations: ALK, anaplastic lymphoma kinase; BBB, blood-brain barrier; BCRP, breast cancer resistance protein; BM, brain metastases; BRAF, v-Raf murine sarcoma viral oncogene homolog B; CNS, central nervous system; CSF, cerebrospinal fluid; CTLA-4, cytotoxic T-lymphocyte antigen; EGFR, epidermal growth factor receptor; EML4, echinoderm microtubule associated protein like 4; FDA, U.S. Food and Drug Administration; HER2, human epidermal growth factor receptor 2; MW, molecular weight; N/A, not applicable; NCI, National Cancer Institute; NPM, nucleophosmin; NSCLC, non-small cell lung cancer; P-gp, P-glycoprotein; WT, wild type.

^aObtained from the PubChem Compound Database (<http://www.ncbi.nlm.nih.gov/pcccompound>).

^bObtained from the Drugbank database (<http://www.drugbank.ca/>) and ChemsSpider database (<http://www.chemspider.com>).

^cAssessed using human blood and CSF samples.

^dBased on nonclinical studies using mouse models.

^eAlectinib was not transported by P-gp in cell transport assay, suggesting it is a poor or non-P-gp substrate.

Table 3. Efficacy of whole brain radiation therapy and targeted therapies for brain metastases from non-small cell lung cancer, stratified by driver mutation status

Publication and Study Design	No. and Type of Patients	Any Prior Therapy for BM	Targeted Therapy Dosage Regimen	Intracranial ORR, % (CR + PR)	Intracranial Disease Control, % (CR + PR + SD)	Median Time to CNS Progression, mo	PFS, mo	OS, Incidence or Median Duration
Studies of unselected patients with NSCLC								
Ceresoli et al., ³³ prospective, noncomparative	41 consecutive (80% platinum pretreated)	WBRT, 44%	Gefitinib 250 mg/d	10	27	NR	3	27% at 11 mo
Wu et al., ³⁴ phase 2	40 chemotherapy pretreated		Gefitinib 250 mg/d	32 (0+32)	77	NR	9	15 mo
Kim et al., ³⁵ prospective, noncomparative	23 Korean never-smokers; chemotherapy-naïve	No	Gefitinib 250 mg/d or Erlotinib 150 mg/d	70 (0+70)	74	NR	7.1	18.8 mo
Pesce et al., ⁵⁸ multicenter, randomized phase 2	59	Only prior chemotherapy allowed	WBRT + temozolomide 75 mg/m ² × 21/28 d (N = 43) WBRT + gefitinib 250 mg/d (N = 16)	NR	NR	1.8	1.8	4.9 mo
Sperduto et al., ⁵⁷ multicenter, randomized phase 3	125 with 1–3 BM	Prior brain resection allowed	WBRT + SRS (N = 44) WBRT + SRS + temozolomide 75 mg/m ² × 21 d (N = 40) WBRT + SRS + erlotinib 150 mg/d (N = 41)	NR	NR	8.1	NR	13.4 mo
Studies in EGFR-mutant NSCLC								
Gow et al., ³⁷ retrospective	63 Taiwanese with BM from adenocarcinomas (73% mutant)	No	WBRT WBRT + TKI	All, 46 ^a EGFRWT, 24 ^a EGFR mutant, 54 ^{b,*} All, 67 ^a EGFR mutant, 84 ^a	NR	NR	NR	14.7 mo
Eicher et al., ³¹ retrospective	93 (44% mutant)	83% WBRT (alone, 53%; + craniotomy, 22%; + SRS, 8%); 5% erlotinib	Treatment with TKI after BM diagnosis (EGFRWT, 19%; EGFR mutant, 78%)	NR	NR	EGFRWT, 8.4 EGFR mutant, 12.4	NR	EGFRWT, 7.6 mo EGFR mutant, 14.5 mo All patients, TKI vs no TKI: 19.1 vs 7.3 mo (P = .001)
Lee et al., ³⁶ retrospective	43 (70% mutant)	No	WBRT (31% of EGFR-WT 50% of EGFR-mutant patients also received a TKI)	All, 70 (12+58) EGFRWT, 46 EGFR mutant, 80*	84	18 12 21**	NR	15 mo 11 mo 15 mo*

Publication and Study Design	No. and Type of Patients	Any Prior Therapy for BM	Targeted Therapy Dosage Regimen	Intracranial ORR, % (CR + PR)	Intracranial Disease Control, % (CR + PR + SD)	Median Time to CNS Progression, mo	PFS, mo	OS, Incidence or Median Duration
Porta et al., ³⁸ retrospective	69 (25% mutant)	80% WBRT	Erlotinib 150 mg/d ^b	All, 26 (15+11) EGFR mutant, 82 (47+35) ^{***}	NR	2.9 11.7	NR	4.3 mo 12.9 mo
Park et al., ³⁹ open-label, single-center phase 2	28 Korean with mutant EGFR	No	Gefitinib 250 mg/d (N=22) or erlotinib 150 mg/d (N=6)	NR	93	6.6	NR	15.9 mo
Wu et al., ⁴⁰ open-label, multicenter phase 2	48 Chinese with asymptomatic BM; chemotherapy pretreated (17% mutant EGFR)	No	Erlotinib 150 mg/d	All, 58 (4+54) EGFRWT, 33 (0+33) EGFR mutant, 75 (12+62)	NR	All, 10.1 EGFRWT, 4.4 ^d EGFR mutant, 15.2 ^{d,*}	NR	All, 18.9 EGFRWT, 18.4 mo EGFR mutant, 37.5 mo
Studies in ALK-positive NSCLC								
Costa et al., ⁶¹ retrospective	275 asymptomatic with BM	60% WBRT 40% Untreated	Crizotinib 250 mg BID	33 ^e 18 ^e	62 ^f 56 ^f	13.2 7.0	6.0 5.9	74% at 6 mo 77% at 6 mo
Shaw et al., ²¹⁵ phase 1	124 (74 with BM)	79% ALK inhibitor	Ceritinib 750 mg/d	All, 35 Pretreated, 29 Treatment-naive, 60	NR	NR	8.3 7.0 NE	NR
Mok et al., ²¹⁶ open-label multicenter phase 2	20 ^g	Chemotherapy and crizotinib	Ceritinib 750 mg/d	NR	80	NR	5.4	NR
Felip et al., ²¹⁷ open-label multicenter phase 2	10 ALK inhibitor naive ^g	Prior chemotherapy allowed	Ceritinib 750 mg/d	NR	80	NR	NR	NR
Gadgeel et al., ⁶⁷ open-label multicenter phase 1/2	21 with resistance or intolerance to crizotinib	81% WBRT; 95% chemotherapy	Alectinib 300–900 mg BID	52 (29+24)	62	NR	NR	NR
Gandhi et al., ⁶⁸ open-label multicenter phase 2	16 with BM after crizotinib ^g	Prior chemotherapy allowed	Alectinib 600 mg BID	69 (13+56)	100	NR	NR	NR
Ou et al., ⁶⁹ open-label multicenter phase 2	34 with BM after crizotinib ^g	Prior chemotherapy allowed	Alectinib 600 mg BID	56 (15+41)	NR	NR	NR	NR
Camidge et al., ²¹⁸ open-label multicenter phase 1/2	12	All other therapies allowed	Brigatinib 30–300 mg/d	50	NR	22	NR	NR

Publication and Study Design	No. and Type of Patients	Any Prior Therapy for BM	Targeted Therapy Dosage Regimen	Intracranial ORR, % (CR + PR)	Intracranial Disease Control, % (CR + PR + SD)	Median Time to CNS Progression, mo	PFS, mo	OS, Incidence or Median Duration
Bauer et al., ⁷⁴ and Shaw et al., ⁷⁵ phase 1/2	17 with BM after ≥1TKI	Treatment naïve or disease progression after previous ALK inhibitor	PF-06463922 10–200 mg/d	29	NR	NR	NR	NR
Gettinger et al., ²¹⁹ multicenter phase 2	10	No	AP26113 180 mg/d	40	60	NR	NR	NR

Abbreviations: ALK, anaplastic lymphoma kinase; BID, twice daily; BM, brain metastases; CNS, central nervous system; CR, complete response; d, day; EGFR, epidermal growth factor receptor; mo, month; NE, not estimable; NR, not reported; NSCLC, non-small cell lung cancer; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; PR, partial response; SD, stable disease; SRS, stereotactic radiosurgery; TKI, tyrosine kinase inhibitor; WBRT, whole brain radiotherapy; WT, wild type.

^aA combination of major and good responses.

^bEight of 17 patients with brain metastases from EGFR-mutant NSCLC received erlotinib as sole therapy.

^cThe remaining patients underwent radiosurgery or SRS with or without WBRT, or no localized therapy.

^dPFS definition was a composite of occurrence of symptomatic BM, confirmed morphologically proven intracranial progressive disease, and extracranial progressive disease.

^eBased on confirmed CNS responses.

^fAt week 12.

^gMeasurable BM at baseline.

* $P < .05$ vs EGFR WT; ** $P < .01$ vs EGFR WT; *** $P < .001$ vs unselected control patients.

impressive disease control in a subset of patients with NSCLC. Tumors evade the immune system using multiple mechanisms including the expression of PD-1 ligands (PD-L1 or PD-L2) by cells in the tumor microenvironment; binding of PD-L1 or PD-L2 to PD-1 receptors leads to inhibition of cytotoxic T cells. Pharmacologic inhibition of the PD-1 receptor/ligand interaction reverses such immune evasion and restores T-cell immunity against the tumor.^{77,78} Notably, PD-1 inhibitors lead to systemic activation of T cells, which can cross the BBB.^{79,80} Two monoclonal antibodies to PD-1, nivolumab and pembrolizumab, are approved for the treatment of NSCLC. Although the objective response rate is generally low with these agents, responses are impressively durable, and treatment leads to a meaningful survival improvement in responders while preserving quality of life.

Phase 3 studies confirming the efficacy of nivolumab as second-line therapy for NSCLC included patients with treated stable BM^{27,28}; there was no indication of increased neurological complications or toxicities in these patients. A large phase 1 study has demonstrated the efficacy of pembrolizumab in the treatment of advanced NSCLC (objective response rate, 19%; median duration of response, 12.5 mo); however, only 10% of patients included in the study had BM.²⁹ Preliminary results from an ongoing phase 2 trial of pembrolizumab showed an intracranial response rate of 45% in 11 patients with untreated BM; there were no serious neurologic complications.³⁰

These early data show that immune checkpoint inhibitors may be an effective treatment for patients with BM, although whether the responses are durable remains to be determined. There are ongoing trials specifically looking at responses of BM from NSCLC and/or melanoma to nivolumab (NCT02621515, NCT02374242, and NCT02320058) and pembrolizumab (NCT02085070).⁶⁶

Chemotherapy

In patients who are refractory to targeted TKI treatments or refractory to or not candidates for immunotherapy, cytotoxic therapy remains an option for suitable patients with NSCLC and will continue to play a role in providing modest systemic responses and improving survival. Intracranial responses with cytotoxic chemotherapy usually correlate with systemic responses in patients with NSCLC. Intracranial response rates as high as 68% have been reported with chemotherapy in asymptomatic patients⁸¹; however, PFS is usually limited to several months, and OS ranges from 5 to 16 months (Table 4). The best intracranial outcomes were achieved using regimens containing cisplatin and pemetrexed^{82,83} or bevacizumab, carboplatin, and paclitaxel.⁸¹

Brain Metastases from Breast Cancer


Autopsy reports indicate that the incidence of BM in women with metastatic breast cancer is as high as 30%.^{84–86} There is also evidence of an increasing incidence of BM from breast cancer.⁸ Notably, according to the GPA, prognostic

Table 4. Efficacy of cytotoxic chemotherapies for brain metastases from non–small cell lung cancer

Publication and Study Design	No. and Type of Patients	Any Prior Local Therapy for BM	Dosage Regimen	ORR, % (CR+ PR)	Intracranial ORR, % (CR + PR)	Intracranial Disease Control, % (CR + PR + SD)	Median Overall PFS, mo	Median Time to CNS Progression, mo	Median OS, mo
Robin et al., ²⁰ multicenter, randomized phase 3	176 inoperable single BM	No	Cisplatin 100 mg/m ² on d 1 and vinorelbine 30 mg/m ² on d 1, 8, 15, 22 Q4W + WBRT 30 Gy/10 fx/12 on progression (arm A) or on d 1 (arm B)	Arm A: 21 Arm B: 20	Arm A, 27 (1+26) Arm B, 33 (8+25)	NR NR	Arm A, 4 Arm B, 3	NR NR	Arm A, 6 Arm B, 5
Barlesi et al., ⁸² multicenter phase 2	43 asymptomatic inoperable BM	No	Cisplatin 75 mg/m ² + pemetrexed 500 mg/m ² Q3W for 4 cycles ^a	35	42 (2+40)	84	4.0	NR	7.4
Galetta et al., ²²¹ multicenter phase 2	25 asymptomatic inoperable BM	No	2 cycles of fotemustine 80 mg/m ² d 1, 8 and cisplatin 80 mg/m ² d 1, Q3W	12 (0+12)	NR	60	2.6	NR	4.7
Dingjin et al., ⁸³ single-center phase 2	41 newly diagnosed inoperable BM	No	Cisplatin 75 mg/m ² + pemetrexed 500 mg/m ² Q3W for 6 cycles + WBRT 30 Gy/10 fx/12 d on d 1–12 of cycle 1	37 (0+37)	68 (2+66)	97.5	8.9	10.6	12.6
Besse et al., ⁸¹ multicenter phase 2	67 asymptomatic BM	No	Bevacizumab 15 mg/kg + carboplatin AUC x 6 + paclitaxel 200 mg/m ² Q3W	63	61	NR	6.7	NR	16.0
Brosnan et al., ²²² retrospective	8 progressive BM	WBRT + SRS	Bevacizumab + irinotecan	NR	>50%	NR	4.0	NR	5.2

Abbreviations: AUC, area under the concentration-time curve; BM, brain metastases; CNS, central nervous system; CR, complete response; d, day; NR, not reported; NSCLC, non–small cell lung cancer; mo, month; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; PR, partial response; Q3W, every 3 weeks; Q4W, every 4 weeks; SD, stable disease; SRS, stereotactic radiosurgery; WBRT, whole-brain radiotherapy.

^aResponding patients were eligible for 2 additional cycles.



index patients with BM from breast cancer have the longest median OS (13.8 mo).¹⁶ Several oncogenes have been identified as drivers of breast cancer (Fig. 3),^{87,88} and the propensity for BM from breast cancer is dependent on tumor subtype, with the highest frequency of BM observed in patients with triple-negative breast cancer (TNBC), followed by human epidermal growth factor receptor 2 (HER2)-positive and luminal breast cancers (Table 1).⁸⁹ Patients with HER2-positive metastatic breast cancer are 2–4 times more likely to develop BM than patients with HER2-negative disease.^{90–92} The standard of care for this common complication of breast cancer includes WBRT, SRS, and surgery, which yield a median OS of 1–2 years in patients with HER2-positive breast cancer and ≤7 months in patients with TNBC.^{89,93,94}

Therapies for Brain Metastases from HER2-positive Breast Cancer

Overview

Until recently, development of targeted treatments in patients with BM and breast cancer was a low priority as BM presents at an advanced stage with little appreciable effect on OS. Currently, the only nonendocrine therapeutic target in breast cancer is HER2, which is overexpressed in 25%–30% of patients.⁹⁵ The development of HER2-targeting therapies has been associated with an improvement in OS.⁹⁶ Consequently, controlling or preventing BM in patients with HER2-positive breast cancer has increasingly become an important treatment consideration.

In patients with HER2-positive metastatic breast cancer, the recommended first-line systemic treatment is a combination of pertuzumab, trastuzumab, and taxane.^{97–99} Patients with active BM from HER2-positive breast cancer were excluded from participation in the

pivotal clinical trials supporting this combination, and treatment regimens containing trastuzumab or lapatinib failed to prevent CNS relapse (MA.31 trial: lapatinib arm 18%, trastuzumab arm 24%; CLEOPATRA: 13% in both treatment arms; CEREBEL: lapatinib arm 3%, trastuzumab arm 5%).^{100–102} In a phase 3 randomized study of capecitabine plus lapatinib versus capecitabine alone for advanced-stage trastuzumab-refractory breast cancer, fewer patients in the combination arm had symptomatic CNS progression as part of the first progression event compared with those not receiving lapatinib.¹⁰³ There is evidence that treatment regimens containing capecitabine may afford greater protective efficacy against BM from HER2-positive breast cancer,¹⁰⁴ and capecitabine alone is known to have activity within the CNS, even in the absence of concurrent HER2-directed therapy.^{105,106} While fewer BM tend to occur in treatment arms containing small molecules (ie, lapatinib and/or capecitabine) than in those containing large biologics such as trastuzumab, pertuzumab, or ado-trastuzumab emtansine (T-DM1), OS in patients with BM is substantially improved in patients randomized to better-performing treatment arms.^{104,107} Finally, neratinib, an irreversible HER2-targeting small molecule TKI, plus paclitaxel may be more effective than trastuzumab plus paclitaxel in reducing CNS progression from HER2-positive metastatic breast cancer.¹⁰⁸

Targeted Therapies

The failure of trastuzumab to prevent CNS relapse among patients with HER2-positive breast cancer can be ascribed to a lack of BBB penetration due to its large size (Table 2). Although retrospective data associated trastuzumab with extending OS in patients with BM from HER2-positive breast cancer, it is likely that this effect resulted from improved systemic disease control rather than any direct intracranial effect.^{109–111} Retrospective analysis of the EMILIA trial revealed that the more potent antibody-cytotoxic chemotherapy conjugate T-DM1 was associated with a protective efficacy similar to that of lapatinib plus capecitabine against BM from HER2-positive breast cancer: CNS progression in those without CNS metastases at baseline occurred in 2.0% of patients who received T-DM1 and 0.7% who received lapatinib plus capecitabine.¹⁰⁴ In patients with asymptomatic BM at baseline, T-DM1 appeared to increase duration of OS compared with lapatinib plus capecitabine (Table 5).¹⁰⁴

Lapatinib has low-level antitumor activity as a single agent for HER2-positive breast cancer patients with progressive BM or BM refractory to trastuzumab and cranial radiotherapy.^{112,113} Positron emission tomography scanning shows that lapatinib distributes into HER2-positive breast cancer BM but not within normal brain tissue. This finding is consistent with clinical data indicating that lapatinib is suitable for treating rather than preventing BM in this population.¹¹⁴ Neratinib was also tested as a single agent in a phase 2 trial of women with HER2-positive BM from breast cancer who had CNS progression after resection, WBRT, or SRS.¹¹⁵ In this trial, the CNS objective response rate (all partial) was 8%, and the median PFS was 1.9 months.

Table 5. Efficacy of targeted therapies for brain metastases from HER2-amplified breast cancer

Publication and Study Design	No. and Type of Patients	Any Prior Therapy for BM	Targeted Therapy Dosage Regimen	Intracranial ORR, % (CR + PR)	Intracranial Disease Control, % (CR + PR + SD)	Median Time to CNS Progression, mo	PFS, mo	OS, Incidence or Median Duration
Bartsch et al., ¹⁰⁹ case-control retrospective	17	WBRT	Trastuzumab 8 mg/kg STAT then 6 mg/kg Q3W	NR	NR	6	NR	7 mo
Park et al., ¹¹⁰ retrospective	78 with symptomatic BM	WBRT, SRS, intrathecal chemotherapy	Trastuzumab before BM Trastuzumab after BM	42 44	68 72	3.9 7.8	NR	4.0 mo 13.6 mo
Church et al., ¹¹¹ retrospective	26 chemotherapy pretreated	Neurosurgery, SRS, WBRT	No trastuzumab Trastuzumab	NR	NR	2.9 NR	NR	HER2 amplified, 11.9 mo* Not HER2 amplified, 3.8 mo
Krop et al., ¹⁰⁴ phase 3 subgroup analysis	95 with BM after trastuzumab and a taxane	69% WBRT and/or local treatment 70% WBRT and/or local treatment	T-DM1 (N = 45) Capecitabine-lapatinib (n = 50)	NR	NR	NR	5.9 5.7	26.8 mo* 12.9 mo
Lin et al., ¹¹² phase 2	39 with BM after trastuzumab	95% WBRT or SRS or both	Lapatinib 750 mg BID	2.6 (0+2.6)	15.4 ^a	NR	3.0	
Lin et al., ¹¹³ open-label, multicenter phase 3	242 with BM after trastuzumab 50 phase 2 completers	95% WBRT; 26% SRS	Lapatinib 750 mg BID Lapatinib + capecitabine	6 (0+6) ^b 20 (0+20)	43 NR		2.4 3.7	6.4 mo
Freedman et al., ¹¹⁵ open-label, multicenter phase 2	40 with BM after CNS-directed therapy	78% WBRT	Neratinib 240 mg QD	8 (0+8)	NR	NR	1.9	8.7 mo
Bartsch et al., ¹¹⁸ retrospective	43 with KPS >70	Local therapy	Trastuzumab ± chemotherapy (N = 28) Trastuzumab ± lapatinib (N = 15)	NR	NR	NR	NR	13 mo NYR
Bachelot et al., ¹²¹ open-label, multicenter phase 2	44 with untreated BM	No	Lapatinib 1250 mg/d + capecitabine (2 g/m ² d-14) x 21 d	66 (0+66)	NR	5.5	5.5	17.0

Abbreviations: BID, twice daily; BM, brain metastases; CNS, central nervous system; CR, complete response; d, day; HER2, human epidermal growth factor receptor 2; KPS, Karnofsky Performance Score; mo, month; NR, not reported; NYR, not yet reached; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; PR, partial response; Q3W, every 3 weeks; QD, once daily; SD, stable disease; SRS, stereotactic radiosurgery; STAT, loading dose; T-DM1, ado-trastuzumab emtansine; WBRT, whole brain radiotherapy.

^aIn both CNS and non-CNS sites.

^bCompared with an ORR of 15% in the 130 patients with measurable extra-CNS disease at baseline.

**P* < .01 vs comparator arm.

Combinational Therapies

There is also interest in a combination of trastuzumab and nanoparticle albumin-bound paclitaxel following impressive objective response rates and PFS as first-line treatment for metastatic breast cancer (excluding BM).^{116,117} Case-controlled data indicate that median OS was improved in patients with BM from HER2-positive breast cancer when lapatinib was added to trastuzumab (not yet reached) compared with trastuzumab alone (13 mo).¹¹⁸

The efficacy of lapatinib alone in the treatment of active BM from HER2-positive breast cancer was increased when coadministered with capecitabine (objective response rate, 38% with lapatinib plus capecitabine vs 0% with lapatinib plus topotecan).^{113,119} Furthermore, the addition of lapatinib to capecitabine significantly prolonged time to tumor progression relative to capecitabine alone (8.4 vs 4.4 mo) in patients with metastatic HER2-positive breast cancer, although progressive BM was an exclusion criterion.¹²⁰ Subsequent findings from the phase 2 LANDSCAPE trial established primary systemic therapy with lapatinib plus capecitabine as an effective and safe alternative to WBRT in patients with asymptomatic to oligosymptomatic BM from HER2-positive breast cancer. The intracranial response rate was 66% in this selected population, thereby delaying the need for WBRT by 8.3 months.¹²¹ A prospective study showed significant uptake of both lapatinib and capecitabine into BM following surgical resection in patients with metastatic breast cancer.¹²²

Dual anti-HER2 therapy with trastuzumab and lapatinib plus capecitabine was an effective and well-tolerated regimen in patients with metastatic HER2-positive breast cancer.¹²³ One case report describes how trastuzumab and lapatinib plus capecitabine as second-line therapy after T-DM1 resulted in partial remission of BM without systemic disease progression and a >14-month delay in time to WBRT.¹²⁴

Finally, in a case series, ONT-380, a HER2-targeting covalently binding TKI, demonstrated promising activity against BM from HER2-positive breast cancer in combination with other systemic agents.¹²⁵

Therapies for Brain Metastases from Other Breast Cancer Subtypes

Targeted Therapies and Hormonal Manipulation

Up to 40% of patients with TNBC develop symptomatic BM during the course of their disease,¹²⁶ yet no targeted therapies for this disease subtype have been developed. Notably, the high expression of PD-L1 in TNBC suggests that this pathway is a potential therapeutic target. In preliminary results of a phase 1 study (KEYNOTE 012) of pembrolizumab, patients with TNBC (13% with BM) had an overall response rate of 19%.¹²⁷ Furthermore, several ongoing clinical trials are investigating pembrolizumab in TNBC (NCT02447003, NCT02555657, NCT02622074, NCT02513472, and NCT02648477).⁶⁶

About 5% of patients with metastatic luminal-type, estrogen-receptor-overexpressing breast cancer develop BM during their course of disease.¹²⁸ Of note, estrogen-receptor expression changes from initial overexpression in the primary tumor to absence of expression in the

corresponding BM in up to 50% of cases.^{129,130} Moreover, in cases where the estrogen receptor continues to be expressed, the estrogen receptor 1 gene (*ESR1*) often contains mutations that result in a constitutively active protein.¹³¹ Concentrations of the estrogen receptor antagonist tamoxifen and its metabolites have been shown to be up to 46-fold higher in brain tissue and BM than in serum.¹³² Furthermore, case reports of patients with BM from breast cancer have demonstrated prolonged survival and prolonged remissions with tamoxifen endocrine therapy, the aromatase inhibitor letrozole, and the progestin megestrol acetate.^{133–135} With more clinically approved drugs for estrogen-receptor-expressing breast cancer, including the mechanistic target of rapamycin (mTOR) inhibitor everolimus, it will be of interest to see if a further extension of survival is possible for TNBC BM. Despite an ability to cross the BBB, palbociclib, an inhibitor of cyclin-dependent kinase (CDK)4/6 recently approved for first- and subsequent-line management of hormone-receptor-positive metastatic breast cancer, is unlikely to be effective against BM from breast cancer as it is a substrate for CNS drug efflux pumps.¹³⁶

Chemotherapy

When treating BM from breast cancer with cytotoxic chemotherapy, agents that exert antitumor activity against breast cancer in the extraneural compartment are selected as opposed to agents with extensive penetration of the BBB but limited systemic activity (eg, temozolomide).^{137–140} In patients naïve to cyclophosphamide and anthracyclines, FEC (5-FU, epirubicin, cyclophosphamide) and CMF (cyclophosphamide, methotrexate, 5-FU) have purported activity in patients with BM.¹⁴¹ Of the few cytotoxic chemotherapeutic agents that can penetrate the BBB when disrupted by BM or radiation, cisplatin demonstrated clinical activity in patients with BM from breast cancer, particularly TNBC, as a single agent and in combination with other chemotherapies or with vinorelbine plus WBRT.^{142–145} Phase 2 trials have also been completed evaluating capecitabine monotherapy in patients with CNS progression after WBRT alone or with SRS and no prior systemic therapy for BM (NCT01077726, NCT00977379, and NCT00570908).⁶⁶

The novel cytotoxic agent sagopilone, a microtubule stabilizer that penetrates the BBB and is not a substrate for CNS efflux transporters, has been evaluated in a single-arm, phase 2 study of 15 breast cancer patients with BM. A CNS partial response was seen in 13% of patients, with a median PFS and OS of 1.4 and 5.3 months, respectively.¹⁴⁶ Additionally, a peptide-facilitated, brain-penetrating formulation of paclitaxel (GRN1005) was well tolerated and decreased tumor size in heavily pretreated patients with advanced solid tumors, including those who had BM and/or failed prior taxane therapy.¹⁴⁷

Brain Metastases from Melanoma

An estimated 50% of patients with stage IV melanoma develop BM, but the prevalence may be as high as 75% based on autopsy reports.^{1,148,149} According to the GPA for

BM, patients with BM from melanoma have a median OS of 6.74 months.¹⁶ Patients with multiple BM and extensive extracranial disease have extremely poor survival outcomes (as short as 1–2 mo in neurologically symptomatic patients).^{150,151} The prognosis may be somewhat better for patients with brain involvement at initial diagnosis of stage IV melanoma than for those who develop BM later.¹⁵² Some patients with solitary BM without known extracranial disease may survive for several years after local treatment.^{152,153}

Melanoma is generally not considered as sensitive to RT or traditional cytotoxic chemotherapy as many other primary malignancies, and these treatment shortcomings are accentuated when a patient develops BM. While surgery and RT may lead to prolonged survival and symptom palliation in patients with oligometastatic CNS involvement, these therapies do not protect against development of new BM.¹⁵⁴ Systemic therapy, although underinvestigated in melanoma patients with CNS involvement, represents a more viable treatment approach for what is essentially a systemic disease with subclinical metastases. Temozolomide and fotemustine were considered promising treatments for BM because of their CNS penetration; however, response rates with these agents were poor, and responses were transient. Fortunately, recent advances in systemic therapy for melanoma, both in molecularly targeted therapy and immunotherapy, render new hope for effective use of systemic agents for BM.

Targeted Therapy with BRAF and/or MEK Inhibitors

Recent discoveries have identified numerous driver genetic mutations in melanoma, particularly in *BRAF* and neuroblastoma RAS viral (v-Ras) oncogene homolog (*NRAS*; Fig. 4). Approximately 40%–60% of melanoma patients harbor a *BRAF* driver mutation, which results in the substitution of valine at codon 600 of the BRAF serine-threonine kinase (BRAF V600). The small molecule BRAF inhibitors dabrafenib and vemurafenib target the RAF/MEK/ERK (MAPK) pathway,^{155,156} and are associated with high response rates and improved survival in metastatic melanoma patients with BRAF V600-mutant tumors.^{157,158} The efficacy is improved further when BRAF inhibitors are used in combination with MEK inhibitor therapy to counter reactivation of the MAPK pathway.^{159–161} BRAF inhibitors are associated with quick-onset regressions in the vast majority of patients and represent a rational option for control and palliation of BM from melanoma.

BRAF inhibitors have been associated with intracranial responses despite limited intracranial bioavailability (Table 2).¹⁶² Dabrafenib therapy led to regression of BM in a phase 1 trial of patients with untreated BM from BRAF V600-mutant melanoma.¹⁶³ In a phase 2 trial of patients with BRAF V600E-mutant melanoma, dabrafenib resulted in comparable intracranial and overall response rates (39% and 38%, respectively) in those with previously untreated BM and similar response rates (31% and 31%, respectively) in those with progressive BM despite prior treatment, indicating that the central and peripheral activity of this agent is concordant (Table 6).¹⁶⁴ Duration of response in this




Fig. 4. Melanoma mutations. Mutations in v-Raf murine sarcoma viral oncogene homolog B (*BRAF*) account for almost half of the driving mutations in melanoma, with another 16% driven by mutations in neuroblastoma RAS viral (v-Ras) oncogene homolog (*NRAS*).¹⁸⁴ Therapies designed to target these mutations have improved outcomes in patients with melanoma.

study ranged from 20 to 28 months. Similarly, in an open-label trial of patients with BRAF V600-mutant melanoma and symptomatic BM, vemurafenib demonstrated anti-tumor activity at both intracranial and extracranial sites, with a duration of response in the brain of 4.4 months (Table 6).¹⁶⁵ MEK inhibition with MEK162 was also promising in a phase 2 study that included patients with treated and stable BM harboring *NRAS* or *BRAF* mutations.¹⁶⁶ Dual BRAF and MEK inhibition (eg, with dabrafenib plus trametinib or vemurafenib plus cobimetinib) is now established as the standard of care for appropriately selected patients with advanced melanoma and tumors that harbor a BRAF V600 mutation, although the efficacy in patients with BM has not yet been determined.^{159,161}

Immunotherapy

Another major drug development in melanoma has been the approval of immune checkpoint inhibitors targeting the PD-1 and cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4) pathways. The anti-CTLA-4 monoclonal antibody ipilimumab was approved after demonstration of improved OS in previously treated patients with metastatic melanoma.¹⁶⁷ Ipilimumab was investigated in a phase 2 trial of patients with advanced melanoma and BM.¹⁶⁸ In the cohort of patients with neurologically asymptomatic disease who were not receiving corticosteroid treatment at study entry ($N = 51$), ipilimumab elicited a CNS response of 16% and CNS disease control rate of 24%, with intracranial responses generally concordant with extracranial responses. The 2-year survival rate in this cohort was 26%, suggesting the possibility of long-term survival in a sizable proportion of patients with an otherwise poor prognosis.

Nivolumab and pembrolizumab have also been approved for treatment of advanced melanoma.^{169,170} Nivolumab was associated with significant improvements in OS

Table 6. Efficacy of targeted therapies for brain metastases from metastatic melanoma

Publication and Study Design	No. and Type of Patients	Any Prior Local Therapy for BM	Targeted Therapy Dosage Regimen	ORR, % (CR + PR)	Intracranial ORR, % (CR + PR)	Disease Control, % (CR + PR + SD)	Intracranial Disease Control, % (CR + PR + SD)	Median Duration of Intracranial Response, mo	Median Overall PFS, mo	Median OS, mo
Long et al., ¹⁶⁴ multicenter phase 2	BRAF-V600E mutant BM (N = 139)	47%	Dabrafenib 150 mg BID	37.8	39.2	79.7	81.1	4.6	3.7	7.6
				30.8	30.8	83.1	89.2	6.5	3.8	7.2
Dummer et al., ¹⁶⁵ pilot study	BRAF-V600K mutant BM (n = 33)	54%	Vemurafenib 960 mg BID	0	6.7	46.7	33.3	2.9	1.9	3.7
				27.8	22.2	50.0	50.0	3.8	3.7	5.0
Margolin et al., ¹⁶⁸ multicenter phase 2	BRAF-V600 mutation with symptomatic BM (N = 24)	Yes	Ipilimumab 10 mg/kg + corticosteroids	42 (0 + 42) ^a	16 (0 + 16) ^a	80	84 ^a	4.4 ^b	3.9	5.3
				10 (0 + 10)	16 (0 + 16)	18	24	NR	1.4 ^c	7.0
	Symptomatic BM (N = 21)	48%		5 (0 + 5)	5 (5 + 0)	5	10		1.2 ^d	3.7

Abbreviations: BID, twice daily; BM, brain metastases; BRAF, v-Raf murine sarcoma viral oncogene homolog B; CR, complete response; mo, month; NR, not reported; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; PR, partial response; SD, stable disease.

^aAmong 19 patients with measurable intracranial disease at baseline.

^bCompared with 3.8 months at extracranial sites.

^cCompared with a median of 1.5 months for disease progression in the brain.

^dCompared with a median of 1.2 months for disease progression in the brain.

and PFS relative to dacarbazine in previously untreated patients without a *BRAF* mutation.¹⁷¹ Pembrolizumab was associated with prolonged PFS and OS and a more favorable toxicity profile than ipilimumab in patients with advanced melanoma who had not received previous therapy with immune checkpoint inhibitors.¹⁷² Additionally, pembrolizumab resulted in a high rate of sustained tumor regression among patients with advanced melanoma, including those with disease progression despite receiving ipilimumab.¹⁷³

To date, there are no published data on the safety and efficacy of pembrolizumab or nivolumab for BM from melanoma because patients with active BM were excluded from entering these studies.¹⁷¹⁻¹⁷³ However, several trials of nivolumab and pembrolizumab are specifically investigating response in BM from melanoma (NCT02621515, NCT02374242, NCT02320058, and NCT02085070).⁶⁶ An interim analysis of the ongoing clinical trial of pembrolizumab (NCT02085070) reported durable partial responses in melanoma patients with untreated BM.¹⁷⁴ Another ongoing trial is investigating the safety and efficacy of the combination of nivolumab plus ipilimumab in this setting (NCT02320058).⁶⁶

Chemotherapy

Cytotoxic therapies have had only modest activity against melanoma in general and therefore have limited utility in the treatment of BM from melanoma. Dacarbazine, the long-standing standard of care for metastatic melanoma, has not reliably demonstrated response in CNS metastasis across numerous trials.^{175,176} Although the oral dacarbazine analog temozolomide has improved capacity to cross the BBB, most studies report response rates of <10% in patients with BM.^{176,177} Fotemustine, a nitrosourea that crosses the BBB, showed promise in early trials that included patients with BM, but intracranial response rates were low (6%) in later trials.^{178,179}

Local and Systemic Combinatorial Approaches

The synergistic potential of RT in facilitating an immune response from immune checkpoint blockade is under evaluation. In a single-institution study of patients with BM from melanoma, univariate analysis revealed that SRS before or during treatment with ipilimumab was associated with higher rates of OS at 1 year than SRS after ipilimumab (65% or 56% vs 40%) and fewer instances of regional brain recurrences at 1 year (64% or 69% vs 92%).¹⁸⁰ There was a trend toward improved local control in those who received SRS concomitantly with ipilimumab (1-y local recurrence, 0%) compared with those who received SRS before (13%) or after (11%) ipilimumab.

Outcomes in patients with BM from *BRAF* V600E-mutant melanoma when combining SRS or WBRT prior to or concomitantly with vemurafenib were also positive in a retrospective analysis.¹⁸¹ Most evaluable patients had an improvement in neurological symptoms (7/11; 64%) and a radiographic response of index lesions (36/48; 75%), of which 23 (48%) were complete responses and 13 (27%) were partial responses. The CNS local control rate, freedom

from new BM, and OS at 6 months were 75%, 57%, and 92%, respectively.

Discussion

Systemic therapy of advanced cancer has been revolutionized by the advent of novel targeted therapeutics, which are associated with prolongation of survival and improvement in quality of life. The application of these therapies to patients with BM requires an understanding of their clinical pharmacology, efficacy, and safety as it relates to the CNS. Unfortunately, patients with active BM have generally been excluded from clinical trials of novel targeted therapies because of the concern for unexpected toxicities and the likelihood that the poor survival of patients with BM will reduce the effect size between comparator agents. Although knowledge about the efficacy and safety of targeted therapies for BM had previously been limited to retrospective observations and small prospective studies, that trend appears to be changing with increasing investigation of these novel therapies in prospective studies specifically for patients with BM.

The few single-arm phase 2 studies that have focused on targeted therapy in patients with BM from NSCLC, breast cancer, or melanoma highlight the potential of targeted systemic therapy to address intracranial disease. There are encouraging findings regarding the utility of the EGFR TKIs gefitinib and erlotinib in BM from *EGFR*-mutant tumors and the second-generation ALK TKIs ceritinib, alectinib, and brigatinib in BM from *ALK*-positive disease. Similar findings are evident with use of anti-HER2 targeted therapies trastuzumab, lapatinib, and T-DM1 for BM from HER2-overexpressing breast cancer, although efficacy is improved when capecitabine is added to either trastuzumab or lapatinib. The *BRAF* and MEK inhibitors, either alone or in combination, appear active in patients with *BRAF*-mutant melanoma. Various studies have reported intracranial response rates ranging from 31% (with dabrafenib in *BRAF* V600E-mutant melanoma) to as high as 75% (with erlotinib in *EGFR*-mutant NSCLC).^{35,40,67,121,164} Unfortunately intracranial responses with many of these drugs may not be as robust as extracranial responses, perhaps reflecting insufficient drug concentrations in brain tissue. Responses with these therapies are generally of quick onset but may not always be durable. Monotherapy with a TKI and close monitoring for progressive disease is a reasonable strategy for patients with small volume, asymptomatic BM associated with appropriate molecular subgroups of these diseases.

Emerging data suggest that immunotherapy with monoclonal antibodies that target the PD-1 or CTLA-4 pathways may lead to substantial and durable intracranial responses concordant with their systemic activity. This may be because immune checkpoint inhibitors do not require direct access to brain parenchyma, as their effects are mediated by proxy on peripheral T cells, which in turn penetrate into the CNS. The hallmark of successful immunotherapy is the potential for durable responses and long-term survival in responding patients, which appears to be preserved in patients with BM. Immunotherapy often has

a delayed onset of response and may lead to inflammatory treatment effects, which are highly relevant to the patients with BM at risk of neurologic complications due to increased mass effects.

Given the aforementioned advances, there is increasing recognition of the need to include patients with untreated BM in clinical trials and to perform trials of systemic therapy specifically in patients with BM. This strategy is not without its challenges because such trials are intensive, expensive, and come with a risk of neurologic complications. However, such studies are required to determine the optimal combination of targeted treatments and traditional therapies used to treat patients with BM.

In conclusion, the emerging evidence for the potential of novel targeted therapies that successfully target BM from NSCLC, breast cancer, and melanoma represents a paradigm shift in the management of BM. Patients with BM increasingly have a realistic hope that current targeted therapies and further research into new targeted therapies will permit an improvement in both their survival and quality of life.

Funding

Writing and editorial assistance for this manuscript, detailed in the acknowledgments, was provided by QXV, a UDG Healthcare Company, and funded by Novartis.

Acknowledgments

Writing and editorial assistance for this manuscript were provided by Nathan Yardley, PhD, and Shannon Davis of QXV, a UDG Healthcare Company, and funded by Novartis.

Conflict of interest statement. CB has received honoraria for advisory board participation from Novartis and Clovis, and works as a co-investigator on clinical trials in the Division of Medical Oncology of the University of Washington, which receives grant funding on an ongoing basis from Clovis, Merck Novartis, Pfizer, Genentech, Inc., and AstraZeneca. VG has received honoraria for advisory board participation from Genentech and works as a principal or co-investigator on clinical trials in the Breast Program in the Division of Medical Oncology at the University of Washington, which receives grant funding on an ongoing basis from Novartis, Pfizer, Genentech, and AstraZeneca. LC has received honoraria for advisory board participation from Novartis, Merck, and Seattle Genetics, for participating as the steering committee chair of a clinical trial from Novartis, and for consulting work from Amgen. LC also works on clinical trials at the University of Washington, which receives grant funding on an ongoing basis from Novartis, Merck, Pfizer, Genentech, and AstraZeneca/Medimmune. SB works at the University of Washington, which receives grant funding from Bristol-Myers-Squibb, EMD-Serono, Merck, Immune Design, Oncosec, NantKwest, and Amgen. MC has no conflicts of interest.

References

- Barnholtz-Sloan J, Sloan A, Davis F, Vignea F, Lai P, Sawaya R. Incidence proportions of brain metastases in patients diagnosed (1973 to 2001) in the metropolitan Detroit cancer surveillance system. *J Clin Oncol.* 2004;22(14):2865–2872.
- Hall WAD, Jalilian HR, Nussbaum ES, Cho KH. Long-term survival with metastatic cancer to the brain. *Med Oncol.* 2000;17(4):279–286.
- Schouten LJ, Rutten JH, Huveneers HA, Wijnstra A. Incidence of brain metastases in a cohort of patients with carcinoma of the breast, colon, kidney, and lung and melanoma. *Cancer.* 2002;94(10):2698–2705.
- Gavrilovic I, Posner JB. Brain metastases: epidemiology and pathophysiology. *J Neurooncol.* 2005;75(1):5–14.
- Tabouret E, Chinot O, Metellus P, Tallet A, Viens P, Gonçalves A. Recent trends in epidemiology of brain metastases: an overview. *Anticancer Res.* 2012;32(11):4655–4662.
- Delattre JY, Krol G, Thaler H, Posner JB. Distribution of brain metastases. *Arch Neurol.* 1988;45(7):741–744.
- Eichler AF, Chung EK, Kodack D, Loeffler JS, Fukumura D, Jain RK. The biology of brain metastases—translation to new therapies. *Nat Rev Clin Oncol.* 2011;8(6):344–356.
- Frisk GS, Svensson TB, Backlund L, Lidbrink E, Blomqvist P, Smedby KE. Incidence and time trends of brain metastases admissions among breast cancer patients in Sweden. *Br J Cancer.* 2012;106(11):1850–1853.
- Patchell RA, Tibbs PA, Walsh JW. A randomized trial of surgery in the treatment of single metastases to the brain. *N Engl J Med.* 1990;322(8):494–500.
- Brown PD, Asher AL, Ballman KV. A phase III randomized trial of whole brain radiation therapy (WBRT) in addition to radiosurgery (SRS) in patients with 1 to 3 brain metastases. *J Clin Oncol.* 2015;33(suppl):abstr LBA4.
- Yamamoto M, Serizawa T, Higuchi Y. A multi-institutional prospective observational study of stereotactic radiosurgery (SRS) for patients with multiple brain metastases (BMs): updated results of the JLGK0901 study—long-term results of irradiation-related complications and neurocognitive function (NCF). *J Clin Oncol.* 2015;33(suppl):abstr 2020.
- Tsao MN, Sultanem K, Chiu D. Supportive care management of brain metastases: what is known and what we need to know. Conference proceedings of the National Cancer Institute of Canada (NCIC) Workshop on Symptom Control in Radiation Oncology. *Clin Oncol (R Coll Radiol).* 2003;15(7):429–434.
- Mehta MP, Rodrigus P, Terhaard CH. Survival and neurologic outcomes in a randomized trial of motexafin gadolinium and whole-brain radiation therapy in brain metastases. *J Clin Oncol.* 2003;21(13):2529–2536.
- Fortin D. The blood-brain barrier: its influence in the treatment of brain tumors metastases. *Curr Cancer Drug Targets.* 2012;12(3):247–259.
- Fu F, Nowak M, Bonhoeffer S. Spatial heterogeneity in drug concentrations can facilitate the emergence of resistance to cancer therapy. *PLoS Comput Biol.* 2015;11(3):e1004142.
- Sperduto P, Wkased N, Roberge D. Summary report on the graded prognostic assessment: an accurate and facile diagnosis-specific tool to estimate survival for patients with brain metastases. *J Clin Oncol.* 2012;30(4):419–425.
- Kris MG, Johnson BE, Berry LD. Using multiplexed assays of oncogenic drivers in lung cancers to select targeted drugs. *JAMA.* 2014;311(19):1998–2006.
- Awad MM, Oxnard GR, Jackman DM. MET exon 14 mutations in non-small-cell lung cancer are associated with advanced age and stage-dependent MET genomic amplification and c-Met overexpression. *J Clin Oncol.* 2016;34(7):721–730.

19. Herbst RSHeymach JVLippman SM. Lung cancer. *N Engl J Med*. 2008;359(13):1367–1380.
20. Han JYPark KKim SW. First-SIGNAL: first-line single-agent icressa versus gemcitabine and cisplatin trial in never-smokers with adenocarcinoma of the lung. *J Clin Oncol*. 2012;30(10):1122–1128.
21. Mitsudomi TMorita SYatabe Y. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol*. 2010;11(2):121–128.
22. Maemondo MInoue AKobayashi K. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med*. 2010;362(25):2380–2388.
23. Zhou CWu YLChen G. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol*. 2011;12(8):735–742.
24. Rosell RCarcerey EGervais R. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol*. 2012;13(3):239–246.
25. Sequist LVYang JCYamamoto N. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol*. 2013;31(27):3327–3334.
26. Shaw ATKim DWNakagawa K. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med*. 2013;368(25):2385–2394.
27. Brahmer JReckamp KLBaas P. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med*. 2015;373(2):123–135.
28. Borghaei HPaz-Ares LHorn L. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med*. 2015;373(17):1627–1639.
29. Garon EBRizvi NAHui R. Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med*. 2015;372(21):2018–2028.
30. Goldberg SBGettinger SNMahajan A. Activity and safety of pembrolizumab in patients with metastatic non-small cell lung cancer with untreated brain metastases. *J Clin Oncol*. 2015;33(suppl):abstr 8035.
31. Eichler AFKahle KTWang DL. EGFR mutation status and survival after diagnosis of brain metastasis in nonsmall cell lung cancer. *Neuro Oncol*. 2010;12(11):1193–1199.
32. Gainor JFOu SHLogan JBorges LFSaw AT. The central nervous system as a sanctuary site in ALK-positive non-small-cell lung cancer. *J Thorac Oncol*. 2013;8(12):1570–1573.
33. Ceresoli GLCappuzzo FGregorc VBartolini SCrinò LVilla E. Gefitinib in patients with brain metastases from non-small-cell lung cancer: a prospective trial. *Ann Oncol*. 2004;15(7):1042–1047.
34. Wu CLi YLWang ZMLi ZZhang TXWei Z. Gefitinib as palliative therapy for lung adenocarcinoma metastatic to the brain. *Lung Cancer*. 2007;57(3):359–364.
35. Kim JELee DHChoi Y. Epidermal growth factor receptor tyrosine kinase inhibitors as a first-line therapy for never-smokers with adenocarcinoma of the lung having asymptomatic synchronous brain metastasis. *Lung Cancer*. 2009;65(3):351–354.
36. Lee HLCchung TSTing LL. EGFR mutations are associated with favorable intracranial response and progression-free survival following brain irradiation in non-small cell lung cancer patients with brain metastases. *Radiat Oncol*. 2012;7:181.
37. Gow CHChien CRChang YL. Radiotherapy in lung adenocarcinoma with brain metastases: effects of activating epidermal growth factor receptor mutations on clinical response. *Clin Cancer Res*. 2008;14(1):162–168.
38. Porta RSánchez-Torres JMPaz-Ares L. Brain metastases from lung cancer responding to erlotinib: the importance of EGFR mutation. *Eur Respir J*. 2011;37(3):624–631.
39. Park SJKim HTLee DH. Efficacy of epidermal growth factor receptor tyrosine kinase inhibitors for brain metastasis in non-small cell lung cancer patients harboring either exon 19 or 21 mutation. *Lung Cancer*. 2012;77(3):556–560.
40. Wu YLZhou CCheng Y. Erlotinib as second-line treatment in patients with advanced non-small-cell lung cancer and asymptomatic brain metastases: a phase II study (CTONG-0803). *Ann Oncol*. 2013;24(4):993–999.
41. Jackman DMHolmes AJLindeman N. Response and resistance in a non-small-cell lung cancer patient with an epidermal growth factor receptor mutation and leptomeningeal metastases treated with high-dose gefitinib. *J Clin Oncol*. 2006;24(27):4517–4520.
42. Katayama TShimizu JSuda K. Efficacy of erlotinib for brain and leptomeningeal metastases in patients with lung adenocarcinoma who showed initial good response to gefitinib. *J Thorac Oncol*. 2009;4(11):1415–1419.
43. Togashi YMasago KFukudo M. Efficacy of increased-dose erlotinib for central nervous system metastases in non-small cell lung cancer patients with epidermal growth factor receptor mutation. *Cancer Chemother Pharmacol*. 2011;68(4):1089–1092.
44. Yu HASima CSReales D. A phase I study of twice weekly pulse dose and daily low dose erlotinib as initial treatment for patients (pts) with EGFR-mutant lung cancers. *J Clin Oncol*. 2015;33(suppl):abstr 8017.
45. Jackman DMMach SLHeng JC. Pulsed dosing of erlotinib for central nervous system progression in EGFR-mutant non-small cell lung cancer [abstr 8116]. *J Clin Oncol*. 2013;31(suppl).
46. Yap TAVidal LAdam J. Phase I trial of the irreversible EGFR and HER2 kinase inhibitor BIBW 2992 in patients with advanced solid tumors. *J Clin Oncol*. 2010;28(25):3965–3972.
47. Bordi PTiseo MBortesi BNaldi NButi SARDizzoni A. Overcoming T790M-driven acquired resistance to EGFR-TKIs in NSCLC with afatinib: a case report. *Tumori*. 2014;100(1):e20–e23.
48. Kim D-WYang JC-HChen K. AZD3759, an EGFR inhibitor with blood brain barrier (BBB) penetration for the treatment of non-small cell lung cancer (NSCLC) with brain metastasis (BM): Preclinical evidence and clinical cases. *J Clin Oncol*. 2015;33(suppl):abstr 8016.
49. Jänne PAYang JCKim DW. AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer. *N Engl J Med*. 2015;372(18):1689–1699.
50. Sequist LVSoria JCGoldman JW. Rociletinib in EGFR-mutated non-small-cell lung cancer. *N Engl J Med*. 2015;372(18):1700–1709.
51. Tagrisso (osimertinib) tablet, for oral use. Available at http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Label_ApprovalHistory#labelinfo. Accessed January 11, 2016.
52. Varga ACamidge DRSequist LV. Activity of rociletinib in EGFR mutant NSCLC patients with a history of CNS involvement. *Eur J Cancer*. 2015;51(suppl 3):S598.
53. Ahn MJTsai CMYang JCH. AZD9291 activity in patients with EGFR-mutant advanced non-small cell lung cancer (NSCLC) and brain metastases: data from phase II studies. *Eur J Cancer*. 2015;51(suppl 3):S625.
54. Akimoto THunter NRBuchmiller LMason KAng KKMilas L. Inverse relationship between epidermal growth factor receptor expression and radio-curability of murine carcinomas. *Clin Cancer Res*. 1999;5(10):2884–2890.
55. Chinnaiyan PHuang SVallabhaneni G. Mechanisms of enhanced radiation response following epidermal growth factor receptor signaling inhibition by erlotinib (Tarceva). *Cancer Res*. 2005;65(8):3328–3335.
56. Welsh JWKomaki RAmiri A. Phase II trial of erlotinib plus concurrent whole-brain radiation therapy for patients with brain metastases from non-small-cell lung cancer. *J Clin Oncol*. 2013;31(7):895–902.
57. Sperduto PWWang MRobins HI. A phase 3 trial of whole brain radiation therapy and stereotactic radiosurgery alone versus WBRT and SRS with temozolomide or erlotinib for non-small cell lung cancer and 1 to 3 brain metastases: Radiation Therapy Oncology Group 0320. *Int J Radiat Oncol Biol Phys*. 2013;85(5):1312–1318.

58. Pesce GAKlingbiel DRibi K. Outcome, quality of life and cognitive function of patients with brain metastases from non-small cell lung cancer treated with whole brain radiotherapy combined with gefitinib or temozolomide. A randomised phase II trial of the Swiss Group for Clinical Cancer Research (SAKK 70/03). *Eur J Cancer*. 2012;48(3):377–384.
59. Camidge DRBang YJKwak EL. Activity and safety of crizotinib in patients with ALK-positive non-small-cell lung cancer: updated results from a phase 1 study. *Lancet Oncol*. 2012;13(10):1011–1019.
60. Weickhardt AJScheier BBurke JM. Local ablative therapy of oligoprogressive disease prolongs disease control by tyrosine kinase inhibitors in oncogene-addicted non-small-cell lung cancer. *J Thorac Oncol*. 2012;7(12):1807–1814.
61. Costa DBShaw ATOu SH. Clinical experience with crizotinib in patients with advanced ALK-rearranged non-small-cell lung cancer and brain metastases. *J Clin Oncol*. 2015;33(17):1881–1888.
62. Zykadia (ceritinib) capsules for oral use. Prescribing Information. Available at <http://www.pharma.us.novartis.com/product/pi/pdf/zykadia.pdf> Accessed August 31, 2015.
63. Friboulet LLI NKatayama R. The ALK inhibitor ceritinib overcomes crizotinib resistance in non-small cell lung cancer. *Cancer Discov*. 2014;4(6):662–673.
64. Shaw ATKim DWMehra R. Ceritinib in ALK-rearranged non-small-cell lung cancer. *N Engl J Med*. 2014;370(13):1189–1197.
65. Kim DWMehra RTan DS. Activity and safety of ceritinib in patients with ALK-rearranged non-small-cell lung cancer (ASCEND-1): updated results from the multicentre, open-label, phase 1 trial. *Lancet Oncol*. 2016;17(4):452–463.
66. ClinicalTrials.gov. Available at <https://clinicaltrials.gov/>. Accessed May 28, 2016.
67. Gadgeel SMGandhi LRIely GJ. Safety and activity of alectinib against systemic disease and brain metastases in patients with crizotinib-resistant ALK-rearranged non-small-cell lung cancer (AF-002JG): results from the dose-finding portion of a phase 1/2 study. *Lancet Oncol*. 2014;15(10):1119–1128.
68. Gandhi LShaw AGadgeel SM. A phase II, open-label, multicenter study of the ALK inhibitor alectinib in an ALK+ non-small-cell lung cancer (NSCLC) U.S./Canadian population who had progressed on crizotinib (NP28761). *J Clin Oncol*. 2015;33(suppl):abstr 8019.
69. Ou S-HIAhn JSPetris LD. Efficacy and safety of the ALK inhibitor alectinib in ALK+ non-small-cell lung cancer (NSCLC) patients who have failed prior crizotinib: an open-label, single-arm, global phase 2 study (NP28673). *J Clin Oncol*. 2015;33(suppl):abstr 8008.
70. Alecensa (alectinib) capsules, for oral use. Prescribing Information. Available at http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Label_ApprovalHistory#labelinfo. Accessed January 8, 2016.
71. Ou S-HIshaw AGandhi L. BMET-18 assessing central nervous system (CNS) response to alectinib in two phase II studies of pre-treated ALK+ non-small cell lung cancer (NSCLC): RECIST versus RANO criteria. *Neuro Oncol*. 2015;17(suppl 5):v48–v49.
72. Ou SHAhn JSDe Petris L. Alectinib in crizotinib-refractory ALK-rearranged non-small-cell lung cancer: a phase II global study. *J Clin Oncol*. 2016;34(7):661–668.
73. Gettinger SNBazhenova LSalgia R. Brigatinib (AP26113) efficacy and safety in ALK+ NSCLC: phase 1/2 trial results. Presented at 16th World Conference on Lung Cancer; September 9, 2015; Denver, CO.
74. Bauer TMShaw ATSolomon B. Phase I/II study of PF-06463922, an ALK/ROS1 tyrosine kinase inhibitor, in patients with advanced non-small-cell lung cancer harboring specific molecular alterations. *J Clin Oncol*. 2015;33(suppl):abstr TPS2620.
75. Shaw ATBauer TMFelip E. Clinical activity and safety of PF-06463922 from a dose escalation study in patients with advanced ALK+ or ROS1+ NSCLC. *J Clin Oncol*. 2015;33(suppl):abstr 8018.
76. Johung KLYeh NDesai NB. Extended survival and prognostic factors for patients with ALK-rearranged non-small-cell lung cancer and brain metastasis. *J Clin Oncol*. 2016;34(2):123–129.
77. Freeman GJLong AJIwai Y. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med*. 2000;192(7):1027–1034.
78. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer*. 2012;12(4):252–264.
79. Bechmann IGalea IPerry VH. What is the blood-brain barrier (not)? *Trends Immunol*. 2007;28(1):5–11.
80. Platten MOchs Klemke DOpitz CWick W. Microenvironmental clues for glioma immunotherapy. *Curr Neurol Neurosci Rep*. 2014;14(4):440.
81. Besse BLe Moulec SMazières J. Bevacizumab in patients with non-squamous non-small cell lung cancer and asymptomatic, untreated brain metastases (BRAIN): a nonrandomized, phase II study. *Clin Cancer Res*. 2015;21(8):1896–1903.
82. Barlesi FGervais RLena H. Pemetrexed and cisplatin as first-line chemotherapy for advanced non-small-cell lung cancer (NSCLC) with asymptomatic inoperable brain metastases: a multicenter phase II trial (GFPC 07-01). *Ann Oncol*. 2011;22(11):2466–2470.
83. Dinglin XXHuang YLiu HZeng YDHou XChen LK. Pemetrexed and cisplatin combination with concurrent whole brain radiotherapy in patients with brain metastases of lung adenocarcinoma: a single-arm phase II clinical trial. *J Neurooncol*. 2013;112(3):461–466.
84. Tsukada YFouad APickren JWLane WW. Central nervous system metastasis from breast carcinoma. Autopsy study. *Cancer*. 1983;52(12):2349–2354.
85. Lee YT. Breast carcinoma: pattern of metastasis at autopsy. *J Surg Oncol*. 1983;23(3):175–180.
86. Cho SYChoi HY. Causes of death and metastatic patterns in patients with mammary cancer. Ten-year autopsy study. *Am J Clin Pathol*. 1980;73(2):232–234.
87. Kumar RSharma ATiwari RK. Application of microarray in breast cancer: an overview. *J Pharm Bioallied Sci*. 2012;4(1):21–26.
88. COSMIC, catalogue of somatic mutations in cancer. Available at <http://cancer.sanger.ac.uk/cosmic/browse/tissue#sn=breast&ss=NS&hn=carcinoma>. Accessed January 15, 2016.
89. Lin NUAmiri-Kordestani LPalmieri DLiewehr DJSteege PS. CNS metastases in breast cancer: old challenge, new frontiers. *Clin Cancer Res*. 2013;19(23):6404–6418.
90. Leyland-Jones B. Human epidermal growth factor receptor 2-positive breast cancer and central nervous system metastases. *J Clin Oncol*. 2009;27(31):5278–5286.
91. Kaal ECVeitch CJ. CNS complications of breast cancer: current and emerging treatment options. *CNS Drugs*. 2007;21(7):559–579.
92. Gabos ZSinha RHanson J. Prognostic significance of human epidermal growth factor receptor positivity for the development of brain metastasis after newly diagnosed breast cancer. *J Clin Oncol*. 2006;24(36):5658–5663.
93. Cho ERubinstein LStevenson P. The use of stereotactic radiosurgery for brain metastases from breast cancer: who benefits most? *Breast Cancer Res Treat*. 2015;149(3):743–749.
94. Niwińska AMurawska MPogoda K. Breast cancer brain metastases: differences in survival depending on biological subtype, RPA RTOG prognostic class and systemic treatment after whole-brain radiotherapy (WBRT). *Ann Oncol*. 2010;21(5):942–948.
95. Slamon DJLeyland-Jones BShak S. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med*. 2001;344(11):783–792.
96. Lin NU. Weighing the options for human epidermal growth factor receptor 2-directed therapy in metastatic breast cancer. *J Clin Oncol*. 2015;33(14):1530–1533.

97. Gradishar WJ, Anderson BO, Blair SL. Breast cancer version 3.2014. *J Natl Compr Canc Netw*. 2014;12(4):542–590.
98. Giordano SH, Temin SK, Irshner JJ. Systemic therapy for patients with advanced human epidermal growth factor receptor 2-positive breast cancer: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2014;32(19):2078–2099.
99. Cardoso F, Costa A, Norton L. ESO-ESMO 2nd international consensus guidelines for advanced breast cancer (ABC2). *Breast*. 2014;23(5):489–502.
100. Swain SM, Kim SBC, Cortés J. Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA study): overall survival results from a randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Oncol*. 2013;14(6):461–471.
101. Swain SM, Baselga J, Kim SB. Pertuzumab, trastuzumab, and docetaxel in HER2-positive metastatic breast cancer. *N Engl J Med*. 2015;372(8):724–734.
102. Pivot X, Manikhas A, Żurawski B. CEREBEL (EGF111438): a phase III, randomized, open-label study of lapatinib plus capecitabine versus trastuzumab plus capecitabine in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer. *J Clin Oncol*. 2015;33(14):1564–1573.
103. Cameron D, Casey M, Press M. A phase III randomized comparison of lapatinib plus capecitabine versus capecitabine alone in women with advanced breast cancer that has progressed on trastuzumab: updated efficacy and biomarker analyses. *Breast Cancer Res Treat*. 2008;112(3):533–543.
104. Krop I, Lin NU, Blackwell K. Trastuzumab emtansine (T-DM1) versus lapatinib plus capecitabine in patients with HER2-positive metastatic breast cancer and central nervous system metastases: a retrospective, exploratory analysis in EMILIA. *Ann Oncol*. 2015;26(1):113–119.
105. Wang M, Yung WK, Royce M, Schomer D, Theriault RL. Capecitabine for 5-fluorouracil-resistant brain metastases from breast cancer. *Am J Clin Oncol*. 2001;24(4):421–424.
106. Siegelmann-Danieli N, Stein M, Bar-Ziv J. Complete response of brain metastases originating in breast cancer to capecitabine therapy. *Isr Med Assoc J*. 2003;5(11):833–834.
107. Swain SM, Baselga J, Miles D. Incidence of central nervous system metastases in patients with HER2-positive metastatic breast cancer treated with pertuzumab, trastuzumab, and docetaxel: results from the randomized phase III study CLEOPATRA. *Ann Oncol*. 2014;25(6):1116–1121.
108. Awada A, Colomer R, Bondarenko I. Efficacy and CNS progression analysis from the randomized phase 2 trial of neratinib + paclitaxel vs trastuzumab + paclitaxel as first-line treatment for HER2+ metastatic breast cancer (NEfERTT). *J Clin Oncol*. 2015;33(suppl):abstr 610.
109. Bartsch R, Rottenfusser A, Wenzel C. Trastuzumab prolongs overall survival in patients with brain metastases from Her2 positive breast cancer. *J Neurooncol*. 2007;85(3):311–317.
110. Park I, Ho J, Lee K, Nam B, Kwon Y, Shin KH. Trastuzumab treatment beyond brain progression in HER2-positive metastatic breast cancer. *Ann Oncol*. 2009;20(1):56–62.
111. Church DN, Modgil R, Gugliani S. Extended survival in women with brain metastases from HER2 overexpressing breast cancer. *Am J Clin Oncol*. 2008;31(3):250–254.
112. Lin NU, Carey L, Liu MC. Phase II trial of lapatinib for brain metastases in patients with human epidermal growth factor receptor 2-positive breast cancer. *J Clin Oncol*. 2008;26(12):1993–1999.
113. Lin NU, Diéras V, Paul D. Multicenter phase II study of lapatinib in patients with brain metastases from HER2-positive breast cancer. *Clin Cancer Res*. 2009;15(4):1452–1459.
114. Saleem A, Searle G, Kenny LM. Lapatinib access into normal brain and brain metastases in patients with Her-2 overexpressing breast cancer. *EJNMMI Res*. 2015;5:30.
115. Freedman RA, Gelman R, Wefel JS. Translational breast cancer research consortium (TBCRC) 022: a phase II trial of neratinib for patients with human epidermal growth factor receptor 2-positive breast cancer and brain metastases. *J Clin Oncol*. 2016;34(9):945–952.
116. Mirtschling BC, Cosgriff T, Harker G, Keaton M, Chidiac T, Min M. A phase II study of weekly nanoparticle albumin-bound paclitaxel with or without trastuzumab in metastatic breast cancer. *Clin Breast Cancer*. 2011;11(2):121–128.
117. Conlin AK, Seidman A, Bach A. Phase II trial of weekly nanoparticle albumin-bound paclitaxel with carboplatin and trastuzumab as first-line therapy for women with HER2-overexpressing metastatic breast cancer. *Clin Breast Cancer*. 2010;10(4):281–287.
118. Bartsch R, Berghoff A, Pluschnig U. Impact of anti-HER2 therapy on overall survival in HER2-overexpressing breast cancer patients with brain metastases. *Br J Cancer*. 2012;106(1):25–31.
119. Lin NU, Eierman W, Greil R. Randomized phase II study of lapatinib plus capecitabine or lapatinib plus topotecan for patients with HER2-positive breast cancer brain metastases. *J Neurooncol*. 2011;105(3):613–620.
120. Geyer CE, Forster J, Lindquist D. Lapatinib plus capecitabine for HER2-positive advanced breast cancer. *N Engl J Med*. 2006;355(26):2733–2743.
121. Bachelot T, Romieu G, Campone M. Lapatinib plus capecitabine in patients with previously untreated brain metastases from HER2-positive metastatic breast cancer (LANDSCAPE): a single-group phase 2 study. *Lancet Oncol*. 2013;14(1):64–71.
122. Morikawa A, Peereboom DM, Thorsheim HR. Capecitabine and lapatinib uptake in surgically resected brain metastases from metastatic breast cancer patients: a prospective study. *Neuro Oncol*. 2015;17(2):289–295.
123. Alés-Martínez J, Filipovich E, Llano J, LJViro J, Sánchez-Escribano R. Dual anti-HER2 therapy (lapatinib and trastuzumab) plus capecitabine is a very effective and well-tolerated regimen (CLT) in metastatic HER2-positive breast cancer patients. *J Clin Oncol*. 2014;32(suppl):abstr e11513.
124. Bergen E, Berghoff A, Rudas M, Preusser M, Bartsch R. Breast cancer brain metastases responding to lapatinib plus capecitabine as second-line primary systemic therapy. *Anticancer Drugs*. 2015;26(5):579–581.
125. Ferrario C, Welch S, Chaves JM. ONT-380 in the treatment of HER2+ breast cancer central nervous system (CNS) metastases (mets). *J Clin Oncol*. 2015;33(suppl):abstr 612.
126. Weil R, Palmieri D, Bronder J, Stark A, MSteege PS. Breast cancer metastasis to the central nervous system. *Am J Pathol*. 2005;167(4):913–920.
127. Nanda R, Chow LQ, Dees EC. A phase Ib study of pembrolizumab (MK-3475) in patients with advanced triple-negative breast cancer [abstract S1-09]. Available at <https://www.sabcs.org/About/Past-Symposia>. Accessed March 11, 2016.
128. Arvold N, O'H KS, Niemierko A. Brain metastases after breast-conserving therapy and systemic therapy: incidence and characteristics by biologic subtype. *Breast Cancer Res Treat*. 2012;136(1):153–160.
129. Duchnowska R, Ziadziusko R, Trojanowski T. Conversion of epidermal growth factor receptor 2 and hormone receptor expression in breast cancer metastases to the brain. *Breast Cancer Res*. 2012;14(4):R119.
130. van Kruchten M, de Vries EGB, Brown M. PET imaging of oestrogen receptors in patients with breast cancer. *Lancet Oncol*. 2013;14(11):e465–e475.
131. Jeselsohn R, Yelensky R, Buchwalter G. Emergence of constitutively active estrogen receptor- α mutations in pretreated advanced estrogen receptor-positive breast cancer. *Clin Cancer Res*. 2014;20(7):1757–1767.
132. Lien EA, Wester KL,ønning P, Solheim E, Ueland PM. Distribution of tamoxifen and metabolites into brain tissue and brain metastases in breast cancer patients. *Br J Cancer*. 1991;63(4):641–645.
133. Salvati MC, Cervoni L, Innocenzi G, Bardella L. Prolonged stabilization of multiple and single brain metastases from breast cancer with tamoxifen. Report of three cases. *Tumori*. 1993;79(5):359–362.

134. Stewart DJ, Dahrour S. Response of brain metastases from breast cancer to megestrol acetate: a case report. *J Neurooncol.* 1995;24(3):299–301.
135. Madhup RK, Kirti SB, Bhatt ML, Srivastava PK, Srivastava MK, Kumar S. Letrozole for brain and scalp metastases from breast cancer—a case report. *Breast.* 2006;15(3):440–442.
136. Raub TJ, Wishart GN, Kulanthaivel P. Brain exposure of two selective dual CDK4 and CDK6 inhibitors and the antitumor activity of CDK4 and CDK6 inhibition in combination with temozolomide in an intracranial glioblastoma xenograft. *Drug Metab Dispos.* 2015;43(9):1360–1371.
137. Christodoulou CB, Fafaloukos DL, Linardou H. Temozolomide (TMZ) combined with cisplatin (DDP) in patients with brain metastases from solid tumors: a Hellenic Cooperative Oncology Group (HeCOG) phase II study. *J Neurooncol.* 2005;71(1):61–65.
138. Kouvaris JR, Miliadou AK, Kouloulas VE. Phase II study of temozolomide and concomitant whole-brain radiotherapy in patients with brain metastases from solid tumors. *Onkologie.* 2007;30(7):361–366.
139. Addeo R, De Rosa CF, Aiola V. Phase 2 trial of temozolomide using protracted low-dose and whole-brain radiotherapy for non-small cell lung cancer and breast cancer patients with brain metastases. *Cancer.* 2008;113(9):2524–2531.
140. Siena SC, Rindò LD, Novoa M. Dose-dense temozolomide regimen for the treatment of brain metastases from melanoma, breast cancer, or lung cancer not amenable to surgery or radiosurgery: a multicenter phase II study. *Ann Oncol.* 2010;21(3):655–661.
141. Boogerd WG, Groeneweld FL, Linn SB, Baars JW, Brandsma D, van Tinteren H. Chemotherapy as primary treatment for brain metastases from breast cancer: analysis of 115 one-year survivors. *J Cancer Res Clin Oncol.* 2012;138(8):1395–1403.
142. Kolaric KR, Roth AJ, Jelic IM, Matkovic A. Phase II clinical trial of cisplatin-diammine platinum (Cis DDP) in metastatic brain tumors. *J Cancer Res Clin Oncol.* 1982;104(3):287–293.
143. Viñolas NG, Graus FM, Mellado BC, Aral L, Estapé J. Phase II trial of cisplatin and etoposide in brain metastases of solid tumors. *J Neurooncol.* 1997;35(2):145–148.
144. Franciosi VC, Cocconi GM, Michiara M. Front-line chemotherapy with cisplatin and etoposide for patients with brain metastases from breast carcinoma, non-small cell lung carcinoma, or malignant melanoma: a prospective study. *Cancer.* 1999;85(7):1599–1605.
145. Cassier PA, Ray-Coquard IS, Sunyach MP. A phase 2 trial of whole-brain radiotherapy combined with intravenous chemotherapy in patients with brain metastases from breast cancer. *Cancer.* 2008;113(9):2532–2538.
146. Freedman RA, Bullitt ES, Sun L. A phase II study of sagopilone (ZK 219477; ZK-EPO) in patients with breast cancer and brain metastases. *Clin Breast Cancer.* 2011;11(6):376–383.
147. Kurzrock RG, Gabrail N, Chandhasin C. Safety, pharmacokinetics, and activity of GRN1005, a novel conjugate of angiopep-2, a peptide facilitating brain penetration, and paclitaxel, in patients with advanced solid tumors. *Mol Cancer Ther.* 2012;11(2):308–316.
148. Amer MH, Al-Sarraf MB, Baker LH, Vaitkevicius VK. Malignant melanoma and central nervous system metastases: incidence, diagnosis, treatment and survival. *Cancer.* 1978;42(2):660–668.
149. Budman DR, Camacho EW, Wittes RE. The current causes of death in patients with malignant melanoma. *Eur J Cancer.* 1978;14(4):327–330.
150. Gupta GR, Robertson AG, Mackie RM. Cerebral metastases of cutaneous melanoma. *Br J Cancer.* 1997;76(2):256–259.
151. Fife KM, Colman MH, Stevens GN. Determinants of outcome in melanoma patients with cerebral metastases. *J Clin Oncol.* 2004;22(7):1293–1300.
152. Sampson JH, Carter JH, Jr, Friedman AH, Seigler HF. Demographics, prognosis, and therapy in 702 patients with brain metastases from malignant melanoma. *J Neurosurg.* 1998;88(1):11–20.
153. Zacest AC, Besser M, Stevens GT, Thompson JF, McCarthy WH, Culjak G. Surgical management of cerebral metastases from melanoma: outcome in 147 patients treated at a single institution over two decades. *J Neurosurg.* 2002;96(3):552–558.
154. Gorantla VK, Kirkwood J, Tawbi HA. Melanoma brain metastases: an unmet challenge in the era of active therapy. *Curr Oncol Rep.* 2013;15(5):483–491.
155. Davies H, Bignell G, Cox C. Mutations of the BRAF gene in human cancer. *Nature.* 2002;417(6892):949–954.
156. Flaherty KT, Infante JR, Daud A. Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N Engl J Med.* 2012;367(18):1694–1703.
157. Chapman PB, Hauschild A, Robert C. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med.* 2011;364(26):2507–2516.
158. Hauschild AG, Grob JJ, Demidov LV. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet.* 2012;380(9839):358–365.
159. Larkin J, Ascierto P, Dréno B. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med.* 2014;371(20):1867–1876.
160. Robert C, Karaszewska B, Schachter J. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med.* 2015;372(1):30–39.
161. Long GV, Stroyakovskiy DG, Gogas H. Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma. *N Engl J Med.* 2014;371(20):1877–1888.
162. Mittapalli RK, Vaidyanathan SD, Dudek AZ, Elmquist WF. Mechanisms limiting distribution of the threonine-protein kinase B-Raf(V600E) inhibitor dabrafenib to the brain: implications for the treatment of melanoma brain metastases. *J Pharmacol Exp Ther.* 2013;344(3):655–664.
163. Falchook GS, Long GV, Kurzrock R. Dabrafenib in patients with melanoma, untreated brain metastases, and other solid tumours: a phase 1 dose-escalation trial. *Lancet.* 2012;379(9829):1893–1901.
164. Long GV, Trefzger U, Davies MA. Dabrafenib in patients with Val600Glu or Val600Lys BRAF-mutant melanoma metastatic to the brain (BREAK-MB): a multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2012;13(11):1087–1095.
165. Dummer RG, Goldinger S, Turttschi CP. Vemurafenib in patients with BRAF(V600) mutation-positive melanoma with symptomatic brain metastases: final results of an open-label pilot study. *Eur J Cancer.* 2014;50(3):611–621.
166. Ascierto P, Schadendorf D, Berking C. MEK162 for patients with advanced melanoma harbouring NRAS or Val600 BRAF mutations: a non-randomised, open-label phase 2 study. *Lancet Oncol.* 2013;14(3):249–256.
167. Hodi FS, O'Day SJ, McDermott DF. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med.* 2010;363(8):711–723.
168. Margolin K, Ernstoff M, Shami O. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol.* 2012;13(5):459–465.
169. Weber JS, D'Angelo S, Minor D. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol.* 2015;16(4):375–384.
170. Topalian SL, Sznol M, McDermott DF. Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab. *J Clin Oncol.* 2014;32(10):1020–1030.
171. Robert C, Long GV, Brady B. Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med.* 2015;372(4):320–330.
172. Robert C, Schachter J, Long GV. Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med.* 2015;372(26):2521–2532.
173. Hamid O, Robert C, Daud A. Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *N Engl J Med.* 2013;369(2):134–144.
174. Kluger HM, Goldberg S, Sznol M. Safety and activity of pembrolizumab in melanoma patients with untreated brain metastases. *J Clin Oncol.* 2015;33(suppl):abstr 9009.

175. Sloan AENock CJEinstein DB. Diagnosis and treatment of melanoma brain metastasis: a literature review. *Cancer Control*. 2009;16(3):248–255.
176. Long GVMargolin KA. Multidisciplinary approach to brain metastasis from melanoma: the emerging role of systemic therapies. *Am Soc Clin Oncol Educ Book*. 2013:393–398.
177. Agarwala SSKirkwood JMGore M. Temozolomide for the treatment of brain metastases associated with metastatic melanoma: a phase II study. *J Clin Oncol*. 2004;22(11):2101–2107.
178. Jacquillat CKhayat DBanzet P. Final report of the French multicenter phase II study of the nitrosourea fotemustine in 153 evaluable patients with disseminated malignant melanoma including patients with cerebral metastases. *Cancer*. 1990;66(9):1873–1878.
179. Avril MFAamdal SGrob JJ. Fotemustine compared with dacarbazine in patients with disseminated malignant melanoma: a phase III study. *J Clin Oncol*. 2004;22(6):1118–1125.
180. Kiess APWolchok JDBarker CA. Stereotactic radiosurgery for melanoma brain metastases in patients receiving ipilimumab: safety profile and efficacy of combined treatment. *Int J Radiat Oncol Biol Phys*. 2015;92(2):368–375.
181. Narayana AMathew MTam M. Vemurafenib and radiation therapy in melanoma brain metastases. *J Neurooncol*. 2013;113(3):411–416.
182. Groothuis DR. The blood-brain and blood-tumor barriers: a review of strategies for increasing drug delivery. *Neuro Oncol*. 2000;2(1):45–59.
183. Elmeliyeg MACarcaboso AMTagen MBai FStewart CF. Role of ATP-binding cassette and solute carrier transporters in erlotinib CNS penetration and intracellular accumulation. *Clin Cancer Res*. 2011;17(1):89–99.
184. Beadling CHEinrich MCWarrick A. Multiplex mutation screening by mass spectrometry evaluation of 820 cases from a personalized cancer medicine registry. *J Mol Diagn*. 2011;13(5):504–513.
185. Sørensen JBHansen HHHansen MDombernowsky P. Brain metastases in adenocarcinoma of the lung: frequency, risk groups, and prognosis. *J Clin Oncol*. 1988;6(9):1474–1480.
186. Quan ALVidetic GMSuh JH. Brain metastases in small cell lung cancer. *Oncology*. 2004;18(8):961–972; discussion 974, 979–980, 987.
187. Brufsky AMMayer MRugo HS. Central nervous system metastases in patients with HER2-positive metastatic breast cancer: incidence, treatment, and survival in patients from registHER. *Clin Cancer Res*. 2011;17(14):4834–4843.
188. Davies MALiu PMcIntyre S. Prognostic factors for survival in melanoma patients with brain metastases. *Cancer*. 2011;117(8):1687–1696.
189. Peuvrel LSaint-Jean MQuéreux G. Incidence and characteristics of melanoma brain metastases developing during treatment with vemurafenib. *J Neurooncol*. 2014;120(1):147–154.
190. Gummadi TZhang BYValpione S. Impact of BRAF mutation and BRAF inhibition on melanoma brain metastases. *Melanoma Res*. 2015;25(1):75–79.
191. de Vries NABuckle TZhao JBeijnen JHSchellens JHvan Tellingen O. Restricted brain penetration of the tyrosine kinase inhibitor erlotinib due to the drug transporters P-gp and BCRP. *Invest New Drugs*. 2012;30(2):443–449.
192. Deng YFeng WWu J. The concentration of erlotinib in the cerebrospinal fluid of patients with brain metastasis from non-small-cell lung cancer. *Mol Clin Oncol*. 2014;2(1):116–120.
193. Togashi YMasago KFukudo M. Cerebrospinal fluid concentration of erlotinib and its active metabolite OSI-420 in patients with central nervous system metastases of non-small cell lung cancer. *J Thorac Oncol*. 2010;5(7):950–955.
194. Togashi YMasago KMasuda S. Cerebrospinal fluid concentration of gefitinib and erlotinib in patients with non-small cell lung cancer. *Cancer Chemother Pharmacol*. 2012;70(3):399–405.
195. Agarwal SSane RGallardo JLOhlfest JREImquist WF. Distribution of gefitinib to the brain is limited by P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2)-mediated active efflux. *J Pharmacol Exp Ther*. 2010;334(1):147–155.
196. Leggas MPanetta JCZhuang Y. Gefitinib modulates the function of multiple ATP-binding cassette transporters in vivo. *Cancer Res*. 2006;66(9):4802–4807.
197. Coldren CDHelfrich BAWitta SE. Baseline gene expression predicts sensitivity to gefitinib in non-small cell lung cancer cell lines. *Mol Cancer Res*. 2006;4(8):521–528.
198. Zhao JChen MZhong W. Cerebrospinal fluid concentrations of gefitinib in patients with lung adenocarcinoma. *Clin Lung Cancer*. 2013;14(2):188–193.
199. Wind SGiessmann TJungnik A. Pharmacokinetic drug interactions of afatinib with rifampicin and ritonavir. *Clin Drug Investig*. 2014;34(3):173–182.
200. Hoffknecht PTufman AWehler T. Efficacy of the irreversible ErbB family blocker afatinib in epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI)-pretreated non-small-cell lung cancer patients with brain metastases or leptomeningeal disease. *J Thorac Oncol*. 2015;10(1):156–163.
201. Tang SCNguyen LNSparidans RWWagenaar EBeijnen JHSchinkel AH. Increased oral availability and brain accumulation of the ALK inhibitor crizotinib by coadministration of the P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) inhibitor elacridar. *Int J Cancer*. 2014;134:1484–1494.
202. Costa DBKobayashi SPandya SS. CSF concentration of the anaplastic lymphoma kinase inhibitor crizotinib. *J Clin Oncol*. 2011;29(15):e443–e445.
203. Fontana DCecon MGambacorti-Passerini CMogni L. Activity of second-generation ALK inhibitors against crizotinib-resistant mutants in an NPM-ALK model compared to EML4-ALK. *Cancer Med*. 2015;4(7):953–965.
204. Kort ASparidans RWWagenaar EBeijnen JHSchinkel AH. Brain accumulation of the EML4-ALK inhibitor ceritinib is restricted by P-glycoprotein (P-GP/ABCB1) and breast cancer resistance protein (BCRP/ABCG2). *Pharmacol Res*. 2015;102:200–207.
205. Kodama TTsukaguchi TYoshida MKondoh OSakamoto H. Selective ALK inhibitor alectinib with potent antitumor activity in models of crizotinib resistance. *Cancer Lett*. 2014;351(2):215–221.
206. Carter PPresta LGorman CM. Humanization of an anti-p185HER2 antibody for human cancer therapy. *Proc Natl Acad Sci U S A*. 1992;89(10):4285–4289.
207. Polli JWHumphreys JEHarmon KA. The role of efflux and uptake transporters in [N-{3-chloro-4-[(3-fluorobenzyl)oxy]phenyl}-6-[5-([2-(methylsulfonyl)ethyl]amino)methyl]-2-furyl]-4-quinazolinamine (GW572016, lapatinib) disposition and drug interactions. *Drug Metab Dispos*. 2008;36(4):695–701.
208. Gori SLunardi Ginno A. Lapatinib concentration in cerebrospinal fluid in two patients with HER2-positive metastatic breast cancer and brain metastases. *Ann Oncol*. 2014;25(4):912–913.
209. Mittapalli RKVaidyanathan SSane REImquist WF. Impact of P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) on the brain distribution of a novel BRAF inhibitor: vemurafenib (PLX4032). *J Pharmacol Exp Ther*. 2012;342(1):33–40.
210. *Zelboraf (vemurafenib) [US package insert]*. South San Francisco, CA: Genentech;2014.
211. Sakji-Dupré LLe Rhun ETemplier CDesmedt EBlanchet BMortier L. Cerebrospinal fluid concentrations of vemurafenib in patients treated for brain metastatic BRAF-V600 mutated melanoma. *Melanoma Res*. 2015;25(4):302–305.
212. Durmus SSparidans RWWagenaar EBeijnen JHSchinkel AH. Oral availability and brain penetration of the B-RAFV600E inhibitor vemurafenib can be enhanced by the P-GLYCOPROTEIN (ABCB1) and breast

- cancer resistance protein (ABCG2) inhibitor elacridar. *Mol Pharm.* 2012;9(11):3236–3245.
213. *Tafinlar (dabrafenib) [US package insert]*. Research Triangle Park, NC: GlaxoSmithKline;2014.
214. O'Day SJ, Hamid O, Urba WJ. Targeting cytotoxic T-lymphocyte antigen-4 (CTLA-4): a novel strategy for the treatment of melanoma and other malignancies. *Cancer.* 2007;110(12):2614–2627.
215. Shaw AT, Mehra RT, Tan DSW. *Evaluation of ceritinib-treated patients with anaplastic lymphoma kinase rearranged (ALK+) non-small cell lung cancer (NSCLC) and brain metastases in the ASCEND-1 study [abstract 1293P]*. Presented at the European Society For Medical Oncology Annual Meeting; September 26–30, 2014; Madrid, Spain.
216. Mok T, Spigel D, Felip E. ASCEND-2: a single-arm, open-label, multicenter phase II study of ceritinib in adult patients (pts) with ALK-rearranged (ALK+) non-small cell lung cancer (NSCLC) previously treated with chemotherapy and crizotinib (CRZ). *J Clin Oncol.* 2015;33(suppl):abstr 8059.
217. Felip E, Orlov S, Park K. ASCEND-3: a single-arm, open-label, multicenter phase II study of ceritinib in ALK-naïve adult patients (pts) with ALK-rearranged (ALK+) non-small cell lung cancer (NSCLC). *J Clin Oncol.* 2015;33(suppl):abstr 8060.
218. Camidge DR, Bazhenova L, Salgia R. Safety and efficacy of brigatinib (AP26113) in advanced malignancies, including ALK+ non-small cell lung cancer (NSCLC). *J Clin Oncol.* 2015;33(suppl):abstr 8062.
219. Gettinger SN, Bazhenova L, Salgia R. Updated efficacy and safety of the ALK inhibitor AP26113 in patients (pts) with advanced malignancies, including ALK+ non-small cell lung cancer (NSCLC). *J Clin Oncol.* 2014;32(suppl):abstr 8047.
220. Robinet G, Thomas P, Breton JL. Results of a phase III study of early versus delayed whole brain radiotherapy with concurrent cisplatin and vinorelbine combination in inoperable brain metastasis of non-small-cell lung cancer: Groupe Français de Pneumo-Cancérologie (GFPC) Protocol 95-1. *Ann Oncol.* 2001;12(1):59–67.
221. Galetta D, Gebbia V, Silvestris N. Cisplatin, fotevustine and whole-brain radiotherapy in non-small cell lung cancer patients with asymptomatic brain metastases: a multicenter phase II study of the Gruppo Oncologico Italia Meridionale (GOIM 2603). *Lung Cancer.* 2011;72(1):59–63.
222. Brosnan EM, Fadul C, Dragnev K, Davis M, Shah K, Eskey CJ. Outcome of combined bevacizumab and irinotecan for patients with progressive brain metastases from non-small cell lung cancer (NSCLC). *J Clin Oncol.* 2015;33(suppl):abstr e19059.