

ERRATUM

Open Access



Erratum to: Incidence and risk factor prevalence of community-acquired pneumonia in adults in primary care in Spain (NEUMO-ES-RISK project)

I. Rivero-Calle^{1,2†}, J. Pardo-Seco², P. Aldaz³, D. A. Vargas⁴, E. Mascarós⁵, E. Redondo⁶, J. L. Díaz-Maroto⁷, M. Linares-Rufo⁸, M. J. Fierro-Alacio⁹, A. Gil¹⁰, J. Molina¹¹, D. Ocaña¹², Federico Martínón-Torres^{1,2*†} and on behalf of NEUMOEXPERTOS group

Erratum

In the original publication of this article [1], the figure files provided were incorrect. The correct figures can be found at the end of this text.

Furthermore, the acknowledgments section omitted the following information in the original article [1]:

"The data for the realization of this publication are part of the database BIFAP managed by the Spanish Agency of Medicines and Sanitary Products (AEMPS). The results, discussion and conclusions of this study are those perceived by the authors only and do not represent in any way the position of the AEMPS on this subject.

The authors are grateful for the excellent collaboration of family physicians and primary care pediatricians who participated in BIFAP".

Author details

¹Translational Pediatrics and Infectious Diseases Section, Pediatrics Department, Hospital Clínico Universitario de Santiago de Compostela, Travesía da Choupana, s/n, 15706 Santiago de Compostela, Spain. ²Genetics, Vaccines, Infections and Pediatrics Research Group (GENVIP), Healthcare Research Institute of Santiago de Compostela, Santiago de Compostela, Spain. ³Member of the Infectious Diseases Prevention Group PAPPs-SEMIFYC, Primary Health Care Center San Juan, Pamplona, Spain. ⁴Versatile Hospitalization Unit, Hospital de Alta Resolución El Toyo, Agencia Pública Sanitaria, Hospital de Poniente, Almería, Spain. ⁵Health Department, Hospital Dr Peset, Primary Care Center Fuente de San Luís, Valencia, Spain. ⁶Preventive and Public Health Activities Group SEMERGEN, International Health Center, Madrid, Spain. ⁷Primary Care Health Center Guadalajara, Infectious Diseases Group SEMERGEN, Guadalajara, Spain. ⁸Primary Care and Clinical Microbiology, Infectious Diseases Group SEMERGEN, Fundación io, Spain. ⁹Primary Care Health Center El Olivillo, Cádiz, Spain. ¹⁰Preventive Medicine and Public Health, Rey Juan Carlos University, Madrid, Spain. ¹¹Primary Care Respiratory Group, Health Care Center Francia, Fuenlabrada, Madrid, Spain. ¹²Primary Care Respiratory Group, Health Care Center Algeciras, Algeciras, Spain.

Received: 29 November 2016 Accepted: 14 December 2016

Published online: 12 January 2017

Reference

1. Rivero-Calle, et al. Incidence and risk factor prevalence of community-acquired pneumonia in adults in primary care in Spain (NEUMO-ES-RISK project). *BMC Infect Dis* 2016;16:645. doi:10.1186/s12879-016-1974-4.

* Correspondence: federico.martinon.torres@sergas.es

†Equal contributors

¹Translational Pediatrics and Infectious Diseases Section, Pediatrics Department, Hospital Clínico Universitario de Santiago de Compostela, Travesía da Choupana, s/n, 15706 Santiago de Compostela, Spain

²Genetics, Vaccines, Infections and Pediatrics Research Group (GENVIP), Healthcare Research Institute of Santiago de Compostela, Santiago de Compostela, Spain




Fig. 1 Distribution of the different lifestyle risk factors and comorbidities in patients seeking medical assistance because of pneumonia at primary care, according to age




Fig. 2 Distribution of the different lifestyle risk factors and comorbidities in patients seeking medical assistance because of pneumonia at primary care, according to gender




Fig. 3 Global incidence of CAP stratified by age




Fig. 4 Global prevalence of the different lifestyle risk factors and comorbidities stratified by age