



Evaluation of Veterinary-Specific Interpretive Criteria for Susceptibility Testing of *Streptococcus equi* Subspecies with Trimethoprim-Sulfamethoxazole and Trimethoprim-Sulfadiazine

Carmen Sadaka,^{a,b,c} Theo Kanellos,^b Luca Guardabassi,^{c,d} Joseph Boucher,^a Jeffrey L. Watts^a

Zoetis Global Therapeutics Research, Kalamazoo, Michigan, USA^a; Zoetis Inc., Paris, France^b; Department of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen, Frederiksberg, Denmark^c; Department of Biomedical Sciences, Ross University School of Veterinary Medicine, Basseterre, St. Kitts and Nevis, West Indies^d

ABSTRACT Antimicrobial susceptibility test results for trimethoprim-sulfadiazine with *Streptococcus equi* subspecies are interpreted based on human data for trimethoprim-sulfamethoxazole. The veterinary-specific data generated in this study support a single breakpoint for testing trimethoprim-sulfamethoxazole and/or trimethoprim-sulfadiazine with *S. equi*. This study indicates trimethoprim-sulfamethoxazole as an acceptable surrogate for trimethoprim-sulfadiazine with *S. equi*.

KEYWORDS breakpoints, horses, interpretive criteria, potentiated sulfonamides, strangles, veterinary microbiology

Trimethoprim-sulfadiazine (SXD) is one of the antimicrobial agents prescribed for the treatment of strangles and other equine infections caused by *Streptococcus equi* subsp. *equi* and *Streptococcus equi* subsp. *zoepidemicus* (1). The CLSI veterinary standard (VET01-A4) lists breakpoints for trimethoprim-sulfamethoxazole (SXT) for *Streptococcus pneumoniae* species, but not for SXD and beta-hemolytic streptococci (2). Additionally, these breakpoints are based on human-derived data (3). Therefore, the interpretation of antimicrobial susceptibility test (AST) results in compliance with CLSI for *S. equi* is based on the clinical breakpoint for *Streptococcus pneumoniae* with SXT used in human medicine (susceptible [S] ≤ 0.5/9.5 µg/ml, intermediate [I] = 1/19 to 2/38 µg/ml, resistant [R] ≥ 4/76 µg/ml and S ≥ 19 mm, I = 16 to 18 mm, R ≤ 15 mm) (2, 3). This breakpoint may not be appropriate for equine isolates due to the pharmacokinetic/pharmacodynamic (PK/PD) differences of the trimethoprim-sulfonamide combinations between horses and humans and because of the PK/PD variability of different potentiated sulfonamide combinations on different bacterial species, sometimes even within the same genus (4). There is limited information on the susceptibility of *S. equi* subspecies to both SXT and SXD. Moreover, SXT is used as a class representative for susceptibility testing of SXD by veterinary diagnostic laboratories, but this use has not been validated specifically for equine isolates (2). This study had four goals: (i) determine if SXT is an acceptable surrogate drug for SXD when testing equine streptococcal isolates; (ii) assess the *in vitro* activity of SXD and SXT against clinical isolates of *S. equi*; (iii) evaluate the synergistic ratios for SXD and SXT with equine *S. equi*; and (iv) evaluate whether the human breakpoints are appropriate for use with equine isolates.

(This study was presented in part at the 57th Annual Conference of the American Association of Veterinary Laboratory Diagnosticians, Kansas City, MO, 16 to 22 October

Received 27 July 2016 **Returned for modification** 19 August 2016 **Accepted** 3 November 2016

Accepted manuscript posted online 9 November 2016

Citation Sadaka C, Kanellos T, Guardabassi L, Boucher J, Watts JL. 2017. Evaluation of veterinary-specific interpretive criteria for susceptibility testing of *Streptococcus equi* subspecies with trimethoprim-sulfamethoxazole and trimethoprim-sulfadiazine. *J Clin Microbiol* 55:326–330. <https://doi.org/10.1128/JCM.01610-16>.

Editor B. W. Fenwick, University of Tennessee

Copyright © 2016 Sadaka et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Carmen Sadaka, carmen.sdk@hotmail.com, or Jeffrey L. Watts, jeffrey.l.watts@zoetis.com.

2014 [5] and the International Conference on One Health Antimicrobial Resistance, Copenhagen, Denmark, 30 September to 2 October 2015 [6]).

A total of 270 *S. equi* isolates divided equally between *S. equi* subsp. *equi* ($n = 135$) and *S. equi* subsp. *zoepidemicus* ($n = 135$) isolated from clinical cases of strangles or equine infectious respiratory disease were used in the study. All isolates were collected from laboratories participating in the Zoetis Global Therapeutics Research surveillance program (2002 to 2009) from different geographic locations (USA [$n = 156$], Canada [$n = 94$], and Europe [$n = 20$]).

Minimal inhibitory concentration (MIC) as determined by the broth microdilution procedure and agar disk diffusion tests were conducted ($n = 270$) as previously described (2, 7). SXT disks (1.25/23.75 μg) (Oxoid Ltd.) were used in the disk diffusion assay. MICs were determined for sulfadiazine (SDZ), sulfamethoxazole (SMX), trimethoprim (TMP), and their corresponding TMP-sulfa combinations (1:19 ratio) (Sigma-Aldrich). Quality control was conducted using CLSI-recommended quality control strains (for MIC assay, *Enterococcus faecalis* American Type Culture Collection [ATCC] 29212, *Streptococcus pneumoniae* ATCC 49619, and *Staphylococcus aureus* ATCC 29213; for disk diffusion assay, *E. faecalis* ATCC 29212, *S. pneumoniae* ATCC 49619, *S. aureus* ATCC 25923, and *Escherichia coli* ATCC 25922) and antimicrobial agents (SXT disks, amoxicillin-clavulanic acid combination [2:1], SXT combination [1:19], and ampicillin). Cation-adjusted Mueller-Hinton broth (CA-MHB) and cation-adjusted Mueller-Hinton agar (CA-MHA) (Fisher Scientific) were prepared in-house as per the instructions of the manufacturer. Thymidine phosphorylase (200 U/liter) (Sigma-Aldrich) was added to all media prior to use to make sure that the media were as thymidine-free and thymine-free as possible. Lysed horse blood (2 to 5%) (Hema Resource) was added to CA-MHB, and sheep blood (5%) (Cleveland Scientific) was added to CA-MHA used for *Streptococcus* species only.

Fractional inhibitory concentration (FIC) indices were determined using the checkerboard technique to evaluate optimal drug ratios for synergy (FIC index of ≤ 0.5) on a selection of isolates (10 *S. equi* subsp. *zoepidemicus* isolates and 10 *S. equi* subsp. *equi* isolates) (8). These isolates were chosen on the basis of very poor activity of individual sulfonamides (MIC $> 2,084 \mu\text{g/ml}$). Quality control included testing for synergy (amoxicillin and clavulanic acid) and antagonism (sparfloxacin and chloramphenicol) repro-




FIG 1 Histograms of MIC distributions for TMP, SMX, SDZ, SXT, and SXD for 270 *S. equi* isolates. MIC values for SXT and SXD are plotted based on the TMP concentration in each combination antimicrobial (TMP-to-sulfonamide combination ratio 1:19). The concentration ranges for SXT and SXD tested are 0.03/0.6 to 32/608 $\mu\text{g/ml}$.

ducibility in the reference *S. aureus* strain ATCC 29213 (9, 10). FIC indices were determined by using the method described by Krogstad and Moellering (8).

The MIC results are summarized in Fig. 1. Poor activity was recorded against individual drugs (MIC₉₀ values of 8, 256, and >2,048 μg/ml for TMP, SMX, and SDZ, respectively). MICs of SXT (MIC₉₀ of 0.25/4.8 μg/ml) and SXD (MIC₉₀ of 0.25/4.8 μg/ml) at a ratio 1:19 were recorded for all tested isolates. A bimodal distribution was noticed with an epidemiological cutoff value of ≤1/19 μg/ml for wild type for both SXT and SXD (Fig. 1). Test error rates were determined by the standard error rate-bounded method (11, 12) using the R statistical computing program (Frisbee Sailing 2013-09-25 version 3.0.2) to plot scattergrams.

The disk diffusion method remains one of the most frequently employed AST method (13). One of the challenges of this method with potentiated sulfonamides is testing for SXD susceptibility using SXT disks without knowing whether the latter is representative (2). Veterinary-specific epidemiological cutoffs were set for both subspecies. Based on the MIC (*S* ≤ 0.5/9.5 μg/ml; *I* = 1/19 to 2/38 μg/ml, and *R* ≥ 4/76 μg/ml) and test zone diameter (*S* ≥ 19 mm; *I* = 16 to 18 mm, and *R* ≤ 15 mm) cutoffs, the discrepancy rates were within the acceptable range (very major errors ≤ 1%, major errors ≤ 3.5%, and minor errors ≤ 5%) (Fig. 2). SXT disks are therefore good surrogates for AST of potentiated sulfonamides. Moreover, the scattergram for SXD MICs versus SXT MICs showed that SXT is a suitable representative for SXD (*R* = 0.85) (Fig. 3), with the consideration that SXT may be more active by at least one twofold dilution (40.74% [110/270]) (data not shown).

The FIC indices (for *S. equi* subsp. *equi* and *S. equi* subsp. *zooepidemicus*, SXT FIC_{index average} values of 0.333 and 0.335, respectively, and SXD FIC_{index average} values of 0.321 and 0.381, respectively) showed synergy at a wide range of TMP-to-sulfonamide ratios and concentrations (1:1 to 1:256) (see Table S1 in the supplemental material). The current findings are in accordance with those from a similar study conducted on 59 different equine pathogens (5). Additionally, the results comply with those from a susceptibility study by the agar dilution method where TMP and nine sulfonamide combinations exhibited synergy against equine *Salmonella enterica* serovars (*n* = 62) at all tested TMP-to-sulfonamide ratios (1:1 to 1:160) for all tested sulfonamides, including SDZ but not SMX (synergy recorded at all ratios except at the 1:1 ratio) (14). In human medicine, potentiated sulfonamides are administered at a 1:5 TMP-to-sulfa ratio to




FIG 2 Scattergrams of (a) SXT MIC and (b) SXD MIC versus SXT (1.25/23.75 μg) disk diffusion test zone diameter (in millimeters) for 270 *S. equi* isolates. Numbers represent the numbers of isolates at each MIC/test zone diameter pair. The horizontal and vertical lines represent MIC and disk diffusion cutoffs, respectively.




FIG 3 Scattergram of SXD versus SXT MICs (in micrograms per milliliter) for the 270 *S. equi* isolates tested. The scattergram also shows the linear regression with $R = 0.85$. Numbers represent the numbers of isolates at each SXD/SXT MIC pair.

achieve the 1:20 plasma ratio required for clinical efficacy. It has been shown in equine practice that potentiated sulfonamides achieve the 1:20 plasma ratio and can be regarded to as therapeutically effective for susceptible *S. equi* when administered at 5 mg of TMP/kg of body weight and 25 mg of SDZ/kg (1:5 ratio) with a dosing interval of 12 h (15–19). The results support the available literature, since the FIC indices showed synergy at a wide range of TMP-to-sulfonamide ratios and concentrations.

Overall, the data indicated low resistance figures ($\leq 2.2\%$) for both subspecies regardless of the AST method used (data not shown). Similarly, Schwarz et al.(20) have also reported low resistance figures (0%). However, high resistance values have been reported in the literature (25.9% and 42% for *S. equi* subsp. *equi* and 45.5% and 61.1% for *S. equi* subsp. *zooepidemicus* [21, 22]). While the reasons for the discrepancies in resistance levels are not known, test methodologies could account for some of these differences, as it is well established that thymidine levels in the test media can diminish the activity of sulfas resulting in false resistance (23). We recommend following the CLSI guidelines to determine whether the conditions of the growth medium are satisfactory and ensure that excessive thymidine levels are not present in the test medium (2).

On the basis of the data generated in this study, we propose that the current CLSI interpretive criteria for *S. pneumoniae* (MIC cutoffs of $S \leq 0.5/9.5 \mu\text{g/ml}$, $I = 1/19$ to $2/38 \mu\text{g/ml}$, and $R \geq 4/76 \mu\text{g/ml}$ and test zone diameter cutoffs of $S \geq 19 \text{ mm}$, $I = 16$ to 18 mm , and $R \leq 15 \text{ mm}$) are suitable for testing equine *S. equi*. In the absence of resistant strains, we recommend veterinary-specific interpretive criteria for susceptible strains only for testing SXT and SXD with equine streptococcal isolates (MIC cutoff of $S \leq 0.5/9.5 \mu\text{g/ml}$ and test zone diameter cutoff of $S \geq 19 \text{ mm}$).

In summary, the data generated in this study indicate that SXT is an appropriate class representative for AST of SXD in *S. equi* subsp. The generated data are consistent with the human-derived interpretative criteria for SXT with *S. pneu-*

moniae. On the basis of the generated data, we propose a single susceptible breakpoint for testing SXT and SXD in *S. equi* strains: a MIC cutoff of $\leq 0.5/9.5 \mu\text{g/ml}$ and a zone diameter cutoff of ≥ 19 mm.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/JCM.01610-16>.

TEXT S1, PDF file, 0.6 MB.

ACKNOWLEDGMENTS

This work was supported by the EU Initial Training Network TRAIN-ASAP (grant agreement 289285).

We thank Lacie Johansen, Victoria Dow, and Michael Sweeney for providing the isolates and for their reviews.

Zoetis, as an animal health company, sells a veterinary antibiotic product containing trimethoprim-sulfadiazine (Tucoprim). The authors involved in the research group that are employees of Zoetis are not directly involved in the sales and marketing of any Zoetis products.

REFERENCES

- Wilson DW. 2001. Rational selection of antimicrobials for use in horses, p 75–93. *In* Proceedings of the 47th Annual Convention of the American Association of Equine Practitioners (AAEP), vol 47. American Association of Equine Practitioners, Lexington, KY.
- Clinical and Laboratory Standards Institute. 2013. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; approved standard—fourth edition. CLSI document VET01-A4. Clinical and Laboratory Standards Institute, Wayne, PA.
- Clinical and Laboratory Standards Institute. 2011. Performance standards for antimicrobial susceptibility testing: twenty-first informational supplement M100-S2. Clinical and Laboratory Standards Institute, Wayne, PA.
- Lees P, Shojaee Aliabadi F. 2002. Rational dosing of antimicrobial drugs: animals versus humans. *Int J Antimicrob Agents* 19:269–284. [https://doi.org/10.1016/S0924-8579\(02\)00025-0](https://doi.org/10.1016/S0924-8579(02)00025-0).
- Sadaka C, Guardabassi L, Kanellos T, Sweeney MT, Watts JL. 2014. Determination of optimal *in vitro* drug ratios of trimethoprim/sulfamethoxazole and trimethoprim/sulfadiazine against equine pathogens, poster 08, p 160. *In* Proceedings of the 57th Annual Conference of the American Association of Veterinary Laboratory Diagnosticians, Kansas City, MO, 16 to 22 October 2014. American Association of Veterinary Laboratory Diagnosticians, Visalia, CA.
- Sadaka C, Kanellos T, Guardabassi L, Boucher J, Watts JL. 2015. Determination of *in vitro* activity of trimethoprim/sulfamethoxazole and trimethoprim/sulfadiazine against *Streptococcus equi* subsp. *equi* and *Streptococcus equi* subsp. *zooepidemicus*, p 52–53. Proceedings of the International Conference on One Health Antimicrobial Resistance, 30 September to 2 October 2015. International Conference on One Health Antimicrobial Resistance, Copenhagen, Denmark.
- Clinical and Laboratory Standards Institute. 2008. Development of *in vitro* susceptibility testing criteria and quality control parameters for veterinary antimicrobial agents; approved guideline—3rd ed. CLSI document VET02-A3. Vol 28, no 7. Clinical and Laboratory Standards Institute, Wayne, PA.
- Krogstad DJ, Moellering RC, Jr. 1986. Antimicrobial combinations, p 537–595. *In* Lorian V (ed), *Antibiotics in laboratory medicine*, 2nd ed. Williams and Wilkins, Baltimore, MD.
- Eliopoulos GM, Moellering RC, Jr. 1991. Laboratory methods used to assess the activity of antimicrobial combinations, p 432–492. *In* Lorian V (ed), *Antibiotics in laboratory medicine*, 3rd ed. The Williams & Wilkins Co., Baltimore, MD.
- Neu HC. 1989. Synergy of fluoroquinolones with other antimicrobial agents. *Rev Infect Dis* 11(Suppl 5):S1025–S1035. https://doi.org/10.1093/clindis/11.Supplement_5.S1025.
- Metzler CM, DeHaan RM. 1974. Susceptibility tests of anaerobic bacteria: statistical and clinical considerations. *J Infect Dis* 30:588–594.
- Brunden MN, Zurenko GE, Kapik B. 1992. Modification of the error-rate bounded classification scheme for use with two MIC break points. *Diagn Microbiol Infect Dis* 15:135–140. [https://doi.org/10.1016/0732-8893\(92\)90037-T](https://doi.org/10.1016/0732-8893(92)90037-T).
- Brooks MB, Morley PS, Dargatz DA, Hyatt DR, Salman MD, Akey BL. 2002. Survey of antimicrobial susceptibility testing practices of veterinary diagnostic laboratories in the United States. *J Am Vet Med Assoc* 222:169–173.
- van Duijkeren E, van Klingeren B, Vulto AG, Sloet van Oldruitenborgh-Oosterbaan MM, Breukink HJ, van Miert AS. 1994. *In vitro* susceptibility of equine *Salmonella* strains to trimethoprim and sulfonamide alone or in combination. *Am J Vet Res* 55:1386–1390.
- McClure SR, Koenig R, Hawkins PA. 2015. A randomized controlled field trial of a novel trimethoprim-sulfadiazine oral suspension for treatment of *Streptococcus equi* subsp. *zooepidemicus* infection of the lower respiratory tract in horses. *J Am Vet Med Assoc* 246:1345–1353. <https://doi.org/10.2460/javma.246.12.1345>.
- Gustafsson A, Baverud V, Franklin A, Gunnarsson A, Ogren G, Ingvast-Larsson C. 1999. Repeated administration of trimethoprim/sulfadiazine in the horse—pharmacokinetics, plasma protein binding and influence on intestinal microflora. *J Vet Pharmacol Ther* 22:20–26. <https://doi.org/10.1046/j.1365-2885.1999.00183.x>.
- Bushby SRM. 1980. Sulfonamide and trimethoprim combinations. *J Am Vet Med Assoc* 176(10):1049–1053.
- van Duijkeren E, Ensink JM, Meijer LA. 2002. Distribution of orally administered trimethoprim and sulfadiazine into noninfected subcutaneous tissue chambers in adult ponies. *J Vet Pharmacol Ther* 25: 273–277. <https://doi.org/10.1046/j.1365-2885.2002.00418.x>.
- van Duijkeren E, Vulto AG, Sloet van Oldruitenborghoosterbaan MM, Mevius DJ, Kessels BG, Breukink HJ, van Miert AS. 1994. A comparative study of the pharmacokinetics of intravenous and oral trimethoprim/sulfadiazine formulations in the horse. *J Vet Pharmacol Ther* 17: 440–446. <https://doi.org/10.1111/j.1365-2885.1994.tb00275.x>.
- Schwarz S, Alesik E, Grobbel M, Lübke-Becker A, Werckenthin C, Wieler LH, Wallmann J. 2007. Antimicrobial susceptibility of streptococci from various indications of swine, horses, dogs and cats as determined in the BFT-GermVet monitoring program 2004–2006. *Berl Munch Tierarztl Wochenschr* 120:380–390.
- Johns IC, Adams EL. 2015. Trends in antimicrobial resistance in equine bacterial isolates: 1999–2012. *Vet Rec* 176:334. <https://doi.org/10.1136/vr.102708>.
- Erol E, Locke SJ, Donahoe JK, Mackin MA, Carter CN. 2012. Beta-hemolytic *Streptococcus* spp. from horses: a retrospective study (2000–2010). *J Vet Diagn Invest* 24:142–147. <https://doi.org/10.1177/1040638711434138>.
- Feary DJ, Hyatt D, Traub-Dargatz J, Roach S, Jones RL, Wu CC, Morley PS. 2005. Investigation of falsely reported resistance of *Streptococcus equi* subsp. *zooepidemicus* isolates from horses to trimethoprim-sulfamethoxazole. *J Vet Diagn* 17:483–486. <https://doi.org/10.1177/104063870501700515>.