

Published in final edited form as:

Eur J Protistol. 2016 August ; 55(Pt A): 4–11. doi:10.1016/j.ejop.2016.02.002.

Ecological and evolutionary significance of novel protist lineages

Javier del Campo^{a,b,*}, Laure Guillou^c, Elisabeth Hehenberger^a, Ramiro Logares^b, Purificación López-García^d, and Ramon Massana^b

^aDepartment of Botany, University of British Columbia, Vancouver, Canada

^bDepartment of Marine Biology and Oceanography, Institut de Ciències del Mar (CSIC), Barcelona, Catalonia, Spain

^cSorbonne Universités, UPMC Univ. Paris 6, CNRS, Adaptation et Diversité en Milieu Marin (UMR 7144), équipe DIPO, Station Biologique de Roscoff, 29688 Roscoff, France

^dEcologie Systématique Evolution, CNRS, Université Paris-Sud, AgroParisTech, Université Paris-Saclay, 91400 Orsay, France

Abstract

Environmental molecular surveys targeting protist diversity have unveiled novel and uncultured lineages in a variety of ecosystems, ranging from completely new high-rank lineages, to new taxa moderately related to previously described organisms. The ecological roles of some of these novel taxa have been studied, showing that in certain habitats they may be responsible for critical environmental processes. Moreover, from an evolutionary perspective they still need to be included in a more accurate and wider understanding of the eukaryotic tree of life. These seminal discoveries promoted the development and use of a wide range of more in-depth culture-independent approaches to access this diversity, from metabarcoding and metagenomics to single cell genomics and FISH. Nonetheless, culturing using classical or innovative approaches is also essential to better characterize this new diversity. Ecologists and evolutionary biologists now face the challenge of apprehending the significance of this new diversity within the eukaryotic tree of life.

Introduction

During the last 15 years, the discovery of multiple microbial eukaryotic lineages has strongly modified general concepts in ecology, revealing unknown groups, some of which appear to be extremely abundant in the environment (Moreira and López-García, 2002). Due to their abundance and key functional role, they are often critical for biogeochemical processes in a given ecosystem (Caron et al., 2012). This new diversity also challenges our understanding of eukaryotic evolutionary relationships. Important changes in the global taxonomy of eukaryotes happened partly because of the discovery of divergent protist lineages occupying pivotal positions in the phylogenetic tree (Burki, 2014). Studying the life

*Corresponding author at: Department of Botany, University of British Columbia, 3529-6270 University Boulevard, Vancouver, BC, Canada. Tel.: +1 7788611714. javier.delcampo@botany.ubc.ca (J. del Campo).

history traits of such novel groups may bring up shared features among lineages that will help us to better understand the evolutionary history of certain groups and, ultimately, the whole eukaryotic tree. The discovery of this new diversity within protists began with the use of culturing-independent molecular methods. From the first diversity studies based on the amplification, cloning and Sanger sequencing of 18S rRNA genes to more recent high-throughput metabarcoding, metagenomics or single cell genomics, these techniques have allowed us to move forward faster, freeing us of the constraints and limitations of culturing (del Campo et al., 2013a), although they have their own biases and pitfalls (Amend et al., 2010; Berney et al., 2004). The use of molecular tools does not mean that culturing became obsolete, but rather asks for innovative culturing strategies targeting the new diversity. The combined use of culture-independent techniques and renovated culturing efforts during the last years has pushed forward our knowledge of the eukaryotes and is filling the gaps on complex ecological puzzles and the still incomplete eukaryotic tree of life.

Approaches to Learn the Biology of Novel Protists

Environmental surveys

Almost 15 years ago three seminal papers revealed that the picture we had of the microbial eukaryotic diversity in marine systems deviated from reality (Diez et al., 2001; López-García et al., 2001; Moon-van der Staay et al., 2001). Using clone libraries of environmental 18S rRNA genes these studies revealed that protist diversity at the picoplanktonic level was dominated by two novel groups of organisms that were named MALV (Marine Alveolates) (Grosillier et al., 2006; Guillou et al., 2008) and MAST (Marine Stramenopiles) (Massana et al., 2004). The MALV include five independent lineages that branch at the base of dinoflagellates in phylogenetic trees. Known representatives of these environmental clusters belong to Syndiniales (Syndinids, Dinophyta) and are all parasites. As such, they play a key regulating role for high-biomass populations like those forming toxic harmful blooms (Chambouvet et al., 2008; Coats et al., 1996; Kim and Park, 2014). They can also impact higher trophic levels and fisheries by infecting radiolarians (Cachon, 1964), cope-pods (Skovgaard et al., 2005), fish larvae (Meneses et al., 2003) and crustaceans such as lobsters and crabs (Shields, 1994). MAST, on the other hand, include a diverse collection of polyphyletic groups of small (2–5 μ m) bacterivorous flagellates branching at the base of the Stramenopiles, each lineage potentially having slightly different ecological niches (Massana et al., 2014). The large clonal abundance of these new clades (both groups combined represent on average nearly half of the clones, Massana and Pedrós-Alió, 2008), suggest that they may also have a strong impact in the food web and in the biogeochemical cycles. The fact that both MALV and MAST cells are heterotrophic protists seems to reflect the higher difficulty of culturing heterotrophs than phototrophs, and therefore the former are better candidates to hide novel diversity. Apart from these two groups a myriad of branches sprouted in different parts of the eukaryotic tree, corresponding to additional novel, potentially uncultured, lineages.

Subsequently, this bonanza of novel diversity was revealed also in freshwater systems (Amaral-Zettler et al., 2002; Richards et al., 2005), hydrothermal vents (Edgcomb et al., 2002; López-García et al., 2003), and soils (Lawley et al., 2004). Often the relevance of

these novel lineages is not obvious to estimate because they may not necessarily be the most abundant organisms and therefore eclipsed by the sequences of dominant ones. Sometimes, it is even hard to differentiate novel diversity from sequence errors or pseudogenes (Thornhill et al., 2007). A way to make this novelty emerge is through meta-analysis and reanalysis of published data, allowing the report of novel clades within a wider range of groups (Richards and Bass, 2005) and populating specific regions of the eukaryotic tree (del Campo and Massana, 2011; del Campo and Ruiz-Trillo, 2013). An obvious challenge in future investigations is to increase the breadth of habitats to explore, focusing on under sampled, uncommon or remote ecosystems that may harbor additional new diversity.

The environmental clone library approach (Sanger sequencing) has been fundamental to retrieve complete 18S rDNA sequences for phylogenetic reconstructions of the novel diversity. While still useful to explore the diversity of novel divergent lineages, this approach is clearly limited for describing community diversity. Direct massive sequencing using new technologies such as 454 pyrosequencing (now obsolete) (Sogin et al., 2006) and Illumina (Amaral-Zettler et al., 2009) have now replaced clone libraries for the study of protist diversity in the environment. These so called High Throughput Environmental Sequencing (HTES or metabarcoding) techniques are cheaper, faster and retrieve several orders of magnitude more sequences from the analyzed samples than clone libraries; for example a single Illumina MiSeq run can produce up to 25 millions reads [2×300 base pairs] (Logares et al., 2012). The HTES techniques are limited by the size of the retrieved reads, currently about 450 bp, which leads to a decrease in the phylogenetic resolution. To partially overcome this issue, two hyper-variable regions of the 18S are mainly used to characterize communities: V4 and V9 (Stoeck et al., 2010). Complete sets of tools have been developed or adopted to process this big data, such as the suites QIIME (Caporaso et al., 2010) and Mothur (Schloss et al., 2009) as well as R packages like vegan (Dixon, 2003) or phyloseq (McMurdie and Holmes, 2013). Proper reference sequence databases are crucial to work with HTES reads, and nowadays there are two available for eukaryotes: Silva (Quast et al., 2013) and PR2 (Guillou et al., 2013). After the discovery of novel diversity based on clone libraries, HTES has completely changed the picture again, since the increase of the sequencing power gives access to new diversity within the so-called “rare” biosphere encompassing organisms at very low abundances (Logares et al., 2014) and allows sampling a wider range of habitats (Heger et al., 2014).

The use of meta-omic techniques, the retrieval of the genomes and transcriptomes of the whole community, represents an alternative strategy to study novelty. This way has not been as deeply explored as the metabarcoding but has shown its potential (Massana et al., 2008; Not et al., 2009; Piganeau et al., 2008). Furthermore meta-omics are not only used to identify but also to describe the ecology and distribution of novel organisms (Cuvelier et al., 2010; Geisen et al., 2015; Worden et al., 2012). In the next years advances in genome recruiting methods (Albertsen et al., 2013; Luo et al., 2012; Spang et al., 2015) will allow assembling individual genomes (Metagenome-Assembled Genomes or MAGs; Garcia et al., 2015) from unknown organisms present in metagenomic and metatranscriptomic datasets and to infer their putative functions and improve or resolve their phylogenetic position.

FISH linking identity with morphology and function

Although environmental clone libraries were a great approach to unveil novel diversity in their time, as are HTES methods today, both share a common problem. There is not a lot that can be inferred about the abundance and biology of the new taxa just by analyzing the reported sequences, apart from their phylogenetic position and their putative habitat. Here is where a technique like FISH (Fluorescence in situ hybridization) can help in linking sequence identity and morphology and even function. The fundament of this technique is to design fluorescent oligonucleotidic probes (18–22 bp) targeting particular environmental 18S rDNA sequences that will hybridize to the rRNA located in ribosome. As hundreds of ribosomes are usually present in a cell, the target can be observed by epifluorescence microscopy (Amann et al., 1995). By visualizing the novel cells embedded in a complex natural assemblage, FISH provides rough pictures of the cell size and shape, and of the presence of several cell components (such as flagella or chloroplasts). In addition, it provides true cell counts, allowing to determine their real contribution in terms of abundance and biomass in the ecosystem. Moreover, combined with other tools such as bacterivory experiments or sophisticated microscopy, FISH provides also information about functional roles of new clades.

Employing FISH, it was shown that the different MAST lineages include cells of different size, abundance, and bacterial grazing behavior, while they collectively account for a very important fraction of heterotrophic flagellates in marine planktonic euphotic regions (Massana et al., 2006, 2009). In the case of MALV protists, the combination of FISH with confocal microscopy has allowed co-localizing MALV cells within putative hosts to confirm their life cycle. Thus, MALV-II cells initiate their life cycle as free-living infective spores (called dinospores) that infect the dinoflagellate host, form large growing endocellular trophonts that leave the host after sporulation as a multicellular vermiform stage which finally disaggregates into new dinospores (Chambouvet et al., 2008). A third example of the use of FISH has been the investigation of the cellular properties of the Cryptomycota or rozellids, an intriguing lineage located at the base of the fungal clade (Lara et al., 2010). Applying FISH on environmental samples it was shown that Cryptomycota cells have flagella in some stages of their life cycle, and that they always lack chitin, once thought to be the primary cell-wall component characterizing fungi (Jones et al., 2011).

Single cell techniques

Single-cell-omic techniques (Stepanuskas, 2012) represent another culture independent approach to access the novel diversity at the genomic level, sometimes even preserving information on cellular identity, morphology and function. There are three main approaches to obtain single cells: (1) manual picking of single cells under the microscope, a slow method largely dependent on the expertise of the manipulator and the size of the targeted organisms; (2) single cell sorting by Flow Cytometry, a high-throughput method that sorts thousands of single cells in less than a minute (Rose et al., 2004) and is currently the most used; (3) single cell sorting using sophisticated microfluidic devices, an emerging approach that does not only sort the cells but eventually allows to experiment with them at a microscale (Leung et al., 2012; Streets et al., 2014). After sorting, three critical steps are required, each one experiencing current technological improvements: whole genome

amplification, genome sequencing, and sequence analysis (assembly, gene prediction, and annotation).

Single Cell Genomics (SCG) are changing the eukaryotic genomic panorama that was clearly dominated, and still is, by the multicellular groups and their parasites (del Campo et al., 2014). SCG gave access to the first genomic data on two novel eukaryotic groups like Picozoa (previously picobiliphytes) (Yoon et al., 2011) and MAST-4 (Roy et al., 2014). SCG can also help us to better understand the ecology and reveal novel diversity en masse, but instead of analyzing the whole community it is done one cell at a time (Heywood et al., 2011; Lepere et al., 2011; Martinez-Garcia et al., 2012; Woyke et al., 2009). Single Cell Transcriptomics (SCT) is a good complement to SCG that has been already tested with large protists (Kolisko et al., 2014) and not only works fine but ends up being relatively cheap and easy to implement in any lab. Advantages of SCT are that it allows doing comparative transcriptomics (Sebé-Pedrós et al., 2013), provides a more direct view of functional genes, and is a necessary alternative of SCG for organisms with large genomes. Single Cell-omics allow accessing the functional gene pool of uncultured organisms widely reported in the environment, thereby contributing to a better understanding of their ecological function. From an evolutionary perspective, this new genome/transcriptome data can be used to improve the eukaryotic phylogenomic framework (Burki, 2014) by enriching the taxon sampling used by researchers working on specific pathways and processes. Single Cell-omics is a game changer like metabarcoding and will boost our knowledge on the genomics of microeukaryotes.

Culturing the uncultured

The three strategies we have presented are forming the spearhead of the discovery of new protist diversity because they are useful, high-throughput and fast. But we should not forget that culturing has been the most important method not only to describe new diversity but to understand organismal biology (which is still elusive from sequencing data). Nowadays, despite being time consuming, sometimes thankless, and often overshadowed by molecular methods, culturing is still having a strong impact on novel diversity. If we have a look at some discoveries that happened after the molecular revolution, we will notice that via culturing we have discovered organisms that pushed the field forward. Some new isolates have shaken the old-established eukaryotic tree, such as the still orphan (*incertae sedis*) clades *Telonema* (Shalchian-Tabrizi et al., 2006), *Collodictyon* (Zhao et al., 2012) or Picozoa (Seenivasan et al., 2013) as these three lineages might represent new deep branches of the tree that are not yet properly placed. Other cultured species have induced substantial changes within supergroup topology and composition, like *Carpodiomonas* in the Excavates (Kolisko et al., 2010), *Palpitomonas* for Cryptista (Yabuki et al., 2014) or the Colponemids in the Alveolates (Janouskovec et al., 2013). Some of these discoveries have a deeper impact on general biology, such as *Chomera velia*, a novel algal isolate related to the apicomplexan parasites (Moore et al., 2008). *C. velia* is important not only because it represents a brand new algal group but also because its phylogenetic placement may help to understand the mechanisms behind the origin of apicomplexan parasitism (Woo et al., 2015). While most of these strains have been retrieved by standard isolation procedures, novel methods have been recently developed to isolate organisms that are abundant in the environment but have been

reluctant to being cultured. This is the case for *Minorisa minuta*, an heterotrophic flagellate at the base of the Chlorarachniophytes that was first detected by environmental surveys and then cultured using single cell sorting and growth in an oligotrophic medium (del Campo et al., 2013b).

Sometimes a more detailed study of well-established cultures produces significant novel biological insights. Such is the case of *Capsaspora owczarzaki* that was originally isolated and deposited at ATCC as *Nuclearia*. After a proper phylogenetic analysis *Capsaspora* revealed itself as a new branch of the holozoans related to choanoflagellates and metazoans (Ruiz-Trillo et al., 2004) and helped to change our view on the origin and mechanisms of metazoan multicellularity (Sebe-Pedros et al., 2011). Another example within the opisthokonts is the basal fungal clade Rozellida or Cryptomycota (Jones et al., 2011; Lara et al., 2010), a widely diverse parasitic group that was defined by combining data retrieved from environmental surveys and from the cultured *Rozella allomyces*. During the last years an even larger group has been defined at the base of the Fungi that also includes the Microsporidians (Capella-Gutiérrez et al., 2012) and the the Aphelids (Karpov et al., 2013). Finally, another interesting example is *Breviata anathema* (Walker et al., 2006), previously known as *Mastigamoeba invertens* (Cavalier-Smith et al., 2004), once considered to belong to the amoebozoans and now placed at the base of the newly defined eukaryotic supergroup Obazoa, which encompasses Opisthokonts, Apusomonads and Breviates (Brown et al., 2013).

Conclusions and Future Perspectives

The journey on the discovery of novel eukaryotic lineages is far from being over, it is just starting, and we should be conscious of the importance of this venture. Increasing the knowledge of the eukaryotic diversity at any level of the tree, from the root to the leaves, is of paramount ecological and evolutionary importance, having even implications in other fields such as biomedicine or conservation. We need to fill all possible gaps to fully understand eukaryotic evolution, the rise of parasites, the generation of symbioses, the establishment of multicellularity or the origin of the eukaryotic cell. At an ecological level, we need to know who are the main players of the different processes and we need to characterize these actors. We have been doing great advances so far but we are still facing several challenges that need to be overcome:

Keep improving our culture independent techniques

Retrieving more and longer reads, processing and analyzing more images, getting better genomes from single-cell-omics, etc. The tools we have been using have provided great results but as explained before they also have their own pitfalls and limitations. We should work to polish them and find more efficient ways to get better data.

Surfing the data tsunami

Current molecular methods to assess microbial diversity allow processing many samples generating an overwhelming amount of sequences that hardly will be fully analyzed. We should make an effort to escalate the analytical tools available to the amount of data

obtained. It is crucial to develop innovative analytical strategies to get the most from the retrieved information.

A picture speaks a thousand words

Imaging was during decades the only way to describe new organisms for protistologists and there is a relevant knowledge corpus. Except for the few studies that combine ultrastructure and sequencing or tools like FISH, most often imaging and molecular data are decoupled. Integrating both approaches, and keeping them integrated in the future, is extremely important. It is advisable to work with images in the same systematic way than with sequences: imaging acquisition by high-throughput methods and creation of comparable and browsable databases. The technology is already available, just needs to be adopted/adapted to protist studies.

Data need to be free

A critical challenge of the tsunami of data is the lack of proper infrastructures for data sharing in a comprehensive way. As a community, we should make an effort to push for the creation of open databases that allow our peers to have an easy access to the results. Furthermore, additional meta-analytical studies can give a new use of the data and bring up some new results.

From the bench to the computer and back to the bench

So far we have been mostly identifying this new diversity but now we need to fully understand their biology and ecological relevance, demanding experimental scenarios. So it is time for isolation and culturing, and to perform original experiments with natural communities. We encourage the use of classical or novel culturing methods to bring to the lab organisms reported by environmental surveys that can be relevant from an ecological or evolutionary point of view. The protist world is vast and hosts most of the eukaryotic diversity, but we are only starting to grasp the extent of such diversity. Exciting years are ahead of us as we deploy all the potential of high-throughput techniques for the study of microbial eukaryotes. The fact of being understudied from an evolutionary and ecological point of view will make the protists, and the protistologists, the main responsible for reshaping the tree of the eukaryotes and for describing novel ecological interactions.

Acknowledgements

This paper was inspired by the symposium of the same title held during the VII ECOP meeting, Seville, Spain, organized by the Federation of European Protistological Societies. JdC was supported by a Marie Curie International Outgoing Fellowship grant (FP7-PEOPLE-2012-IOF – 331450 CAARL) and by an International Society of Protistologists grant to attend to the meeting. LG was supported by the French ANR project HAPAR (Défi 1: Climatic changes) and the European Project MaCuMBa (FP7-KBBE-2012-6-311975). PLG acknowledges the European Research Council under the European Union's Seventh Framework Program ERC Grant Agreement 322669 'ProtistWorld'. RM was supported by the MEFISTO project (CTM2013-43767-P, MINECO).

References

Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple metagenomes. *Nat Biotechnol.* 2013; 31:533–538. [PubMed: 23707974]

- Amann RI, Ludwig W, Schleifer K-HH. Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol Rev.* 1995; 59:143–169. [PubMed: 7535888]
- Amaral-Zettler LA, Gómez F, Zettler ER, Keenan BGB. Eukaryotic diversity in Spain's river of fire. *Nature.* 2002; 417:137. [PubMed: 12000949]
- Amaral-Zettler LA, McCliment EA, Ducklow HW, Huse SM. A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PLoS ONE.* 2009; 4:e6372. [PubMed: 19633714]
- Amend AS, Seifert KA, Bruns TD. Quantifying microbial communities with 454 pyrosequencing: does read abundance count? *Mol Ecol.* 2010; 19:5555–5565. [PubMed: 21050295]
- Berney C, Fahrni JF, Pawlowski J. How many novel eukaryotic 'kingdoms'? Pitfalls and limitations of environmental DNA surveys. *BMC Biol.* 2004; 2:13. [PubMed: 15176975]
- Brown MW, Sharpe SC, Silberman JD, Heiss AA, Lang BF, Simpson AGB, Roger AJ. Phylogenomics demonstrates that breviate flagellates are related to opisthokonts and apusomonads. *Proc R Soc B Biol Sci.* 2013; 280:20131755.
- Burki F. The eukaryotic tree of life from a global phylogenomic perspective. *Cold Spring Harb Perspect Biol.* 2014; 6:a016147. [PubMed: 24789819]
- Cachon, J. Contribution a l'étude des péridiniens parasites. *Cytologie, Cycles évolutifs. Annales Des Sciences Naturelles, Zoologie.* Paris: 1964. p. 1-158. 12e Série, Tome VI, Paris
- Capella-Gutiérrez S, Marcet-Houben M, Gabaldón T. Phylogenomics supports microsporidia as the earliest diverging clade of sequenced fungi. *BMC Biol.* 2012; 10:47. [PubMed: 22651672]
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods.* 2010; 7:335–336. [PubMed: 20383131]
- Caron DA, Countway PD, Jones AC, Kim DY, Schnetzer A. Marine protistan diversity. *Ann Rev Mar Sci.* 2012; 4:467–493.
- Cavalier-Smith T, Chao EE, Oates B. Molecular phylogeny of Amoebozoa and the evolutionary significance of the unikont *Phalansterium*. *Eur J Protistol.* 2004; 40:21–48.
- Chambouvet A, Morin P, Marie D, Guillou L. Control of toxic marine dinoflagellate blooms by serial parasitic killers. *Science.* 2008; 322:1254–1257. (80). [PubMed: 19023082]
- Coats DW, Adam EJ, Gallegos CL, Hedrick S. Parasitism of photosynthetic dinoflagellates in a shallow subestuary of Chesapeake Bay USA. *Aquat Microb Ecol.* 1996; 11:1–9.
- Cuvelier ML, Allen AE, Monier A, McCrow JP, Messié M, Tringe SG, Woyke T, Welsh RM, Ishoey T, Lee J-H, Binder BJ, et al. Targeted metagenomics and ecology of globally important uncultured eukaryotic phytoplankton. *Proc Natl Acad Sci.* 2010; 107:14679–14684. [PubMed: 20668244]
- del Campo J, Balagué V, Forn I, Lekunberri I, Massana R. Culturing bias in marine heterotrophic flagellates analyzed through seawater enrichment incubations. *Microb Ecol.* 2013a; 66:489–499. [PubMed: 23749062]
- del Campo J, Massana R. Emerging diversity within Chrysophytes, Choanoflagellates and Bicosoecids based on molecular surveys. *Protist.* 2011; 162:435–448. [PubMed: 21239227]
- del Campo J, Not F, Forn I, Sieracki ME, Massana R. Taming the smallest predators of the oceans. *ISME J.* 2013b; 7:351–358. [PubMed: 22810060]
- del Campo J, Ruiz-Trillo I. Environmental survey meta-analysis reveals hidden diversity among unicellular opisthokonts. *Mol Biol Evol.* 2013; 30:802–805. [PubMed: 23329685]
- del Campo J, Sieracki ME, Molestina R, Keeling PJ, Massana R, Ruiz-Trillo I. The others: our biased perspective of eukaryotic genomes. *Trends Ecol Evol.* 2014; 29:252–259. [PubMed: 24726347]
- Díez B, Pedrós-Alió C, Massana R. Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. *Appl Environ Microbiol.* 2001; 67:2932–2941. [PubMed: 11425705]
- Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci.* 2003; 14:927–930.
- Edgcomb VP, Kysela DT, Teske A, de Vera Gomez A, Sogin ML. Benthic eukaryotic diversity in the Guaymas Basin hydrothermal vent environment. *Proc Natl Acad Sci.* 2002; 99:7658–7662. [PubMed: 12032339]

- Garcia SL, Buck M, McMahon KD, Grossart H-P, Eiler A, Warnecke F. Auxotrophy and intrapopulation complementary in the “interactome” of a cultivated freshwater model community. *Mol Ecol*. 2015; 24:4449–4459. [PubMed: 26179741]
- Geisen S, Tveit AT, Clark IM, Richter A, Svenning MM, Bonkowski M, Urich T. Metatranscriptomic census of active protists in soils. *ISME J*. 2015; 9:2178–2190. [PubMed: 25822483]
- Groisillier A, Massana R, Valentin K, Vault D, Guillou L. Genetic diversity and habitats of two enigmatic marine alveolate lineages. *Aquat Microb Ecol*. 2006; 42:277–291.
- Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud GG, de Vargas C, Decelle J, del Campo J, et al. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small subunit rRNA sequences with curated taxonomy. *Nucleic Acids Res*. 2013; 41:D597–D604. [PubMed: 23193267]
- Guillou L, Viprey M, Chambouvet A, Welsh RM, Kirkham AR, Massana R, Scanlan DJ, Worden AZ. Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales (Alveolata). *Environ Microbiol*. 2008; 10:3349–3365. [PubMed: 18771501]
- Heger TJ, Edgcomb VP, Kim E, Lukeš J, Leander BS, Yubuki N. A resurgence in field research is essential to better understand the diversity, ecology, and evolution of microbial eukaryotes. *J Eukaryot Microbiol*. 2014; 61:214–223. [PubMed: 24325268]
- Heywood JL, Sieracki ME, Bellows W, Poulton NJ, Stepanauskas R. Capturing diversity of marine heterotrophic protists: one cell at a time. *ISME J*. 2011; 5:674–684. [PubMed: 20962875]
- Janouškovec J, Tikhonenkov DV, Mikhailov KV, Simdyanov TG, Aleoshin VV, Mylnikov AP, Keeling PJ. Colponemids represent multiple ancient alveolate lineages. *Curr Biol*. 2013; 23:2546–2552. [PubMed: 24316202]
- Jones MDM, Forn I, Gadelha C, Egan MJ, Bass D, Massana R, Richards TA. Discovery of novel intermediate forms redefines the fungal tree of life. *Nature*. 2011; 474:200–203. [PubMed: 21562490]
- Karpov SA, Mikhailov KV, Mirzaeva GS, Mirabdullaev IM, Mamkaeva KA, Titova NN, Aleoshin VV. Obligately phagotrophic aphelids turned out to branch with the earliest-diverging fungi. *Protist*. 2013; 164:195–205. [PubMed: 23058793]
- Kim S, Park MG. *Amoebophrya* spp. from the bloom-forming dinoflagellate *Cochlodinium polykrikoides*: parasites not nested in the “amoebophrya ceratii complex.”. *J Eukaryot Microbiol*. 2014; 61:173–181. [PubMed: 24612333]
- Kolisko M, Boscaro V, Burki F, Lynn DH, Keeling PJ. Single-cell transcriptomics for microbial eukaryotes. *Curr Biol*. 2014; 24:R1081–R1082. [PubMed: 25458215]
- Kolisko M, Silberman JD, Cepicka I, Yubuki N, Takishita K, Yabuki A, Leander BS, Inouye I, Inagaki Y, Roger AJ, Simpson AGB. A wide diversity of previously undetected free-living relatives of diplomonads isolated from marine/saline habitats. *Environ Microbiol*. 2010; 12:2700–2710. [PubMed: 20482740]
- Lara E, Moreira D, López-García P. The environmental clade LKM11 and Rozella form the deepest branching clade of fungi. *Protist*. 2010; 161:116–121. [PubMed: 19674933]
- Lawley B, Ripley S, Bridge P, Convey P. Molecular analysis of geographic patterns of eukaryotic diversity in antarctic soils. *Appl Environ Microbiol*. 2004; 70:5963–5972. [PubMed: 15466539]
- Lepère C, Demura M, Kawachi M, Romac S, Probert I, Vault D. Whole-genome amplification (WGA) of marine photosynthetic eukaryote populations. *FEMS Microbiol Ecol*. 2011; 76:513–523. [PubMed: 21348885]
- Leung K, Zahn H, Leaver T, Konwar KM, Hanson NW, Page AP, Lo C-C, Chain PS, Hallam SJ, Hansen CL. A programmable droplet-based microfluidic device applied to multiparameter analysis of single microbes and microbial communities. *Proc Natl Acad Sci*. 2012; 109:7665–7670. [PubMed: 22547789]
- Logares R, Audic S, Bass D, Bittner L, Boutte C, Christen R, Claverie J-MM, Decelle J, Dolan JR, Dunthorn M, Edvardsen B, et al. Patterns of rare and abundant marine microbial eukaryotes. *Curr Biol*. 2014; 24:813–821. [PubMed: 24704080]
- Logares R, Haverkamp THA, Kumar S, Lanzén A, Neder-bragt AJ, Quince C, Kausserud H. Environmental microbiology through the lens of high-throughput DNA sequencing: synopsis of

- current platforms and bioinformatics approaches. *J Microbiol Methods*. 2012; 91:106–113. [PubMed: 22849829]
- López-García P, Philippe H, Gail F, Moreira D. Autochthonous eukaryotic diversity in hydrothermal sediment and experimental microcolonizers at the Mid-Atlantic Ridge. *Proc Natl Acad Sci*. 2003; 100:697–702. [PubMed: 12522264]
- López-García P, Rodríguez-Valera F, Pedrós-Alió C, Moreira D. Unexpected diversity of small eukaryotes in deep-sea Antarctic plankton. *Nature*. 2001; 409:603–607. [PubMed: 11214316]
- Luo C, Tsementzi D, Kyrpides NCC, Konstantinidis KTT. Individual genome assembly from complex community short-read metagenomic datasets. *ISME J*. 2012; 6:898–901. [PubMed: 22030673]
- Martinez-Garcia M, Brazel D, Poulton NJ, Swan BK, Gomez ML, Masland D, Sieracki ME, Stepanauskas R. Unveiling in situ interactions between marine protists and bacteria through single cell sequencing. *ISME J*. 2012; 6:703–707. [PubMed: 21938022]
- Massana R, Castresana J, Balagué V, Guillou L, Romari K, Groisillier A, Valentin K, Pedrós-Alió C. Phylogenetic and ecological analysis of novel marine stramenopiles. *Appl Environ Microbiol*. 2004; 70:3528–3534. [PubMed: 15184153]
- Massana R, del Campo J, Sieracki ME, Audic S, Logares R. Exploring the uncultured microeukaryote majority in the oceans: reevaluation of ribogroups within stramenopiles. *ISMEJ*. 2014; 8:854–866.
- Massana R, Karniol B, Pommier T, Bodaker I, Bèjà O. Metagenomic retrieval of a ribosomal DNA repeat array from an uncultured marine alveolate. *Environ Microbiol*. 2008; 10:1335–1343. [PubMed: 18266757]
- Massana R, Pedrós-Alió C. Unveiling new microbialeukaryotes in the surface ocean. *Curr Opin Microbiol*. 2008; 11:213–218. [PubMed: 18556239]
- Massana R, Terrado R, Forn I, Lovejoy C, Pedrós-Alió C. Distribution and abundance of uncultured heterotrophic flagellates in the world oceans. *Environ Microbiol*. 2006; 8:1515–1522. [PubMed: 16913912]
- Massana R, Unrein F, Rodríguez-Martínez R, Forn I, Lefort T, Pinhassi J, Not F. Grazing rates and functional diversity of uncultured heterotrophic flagellates. *ISME J*. 2009; 3:588–596. [PubMed: 19129862]
- McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*. 2013; 8:e61217. [PubMed: 23630581]
- Meneses I, Vendrell C, Stratoudakis Y. Mackerel (*Scomber scombrus*) eggs parasitized by *Ichthyodinium chabelardi* in the north-east Atlantic: An overlooked source of mortality. *J Plankton Res*. 2003; 25:1177–1181.
- Moon-van der Staay SY, de Wachter R, Vault D. Oceanic 18S rDNA sequences from picoplankton reveal unsuspected eukaryotic diversity. *Nature*. 2001; 409:607–610. [PubMed: 11214317]
- Moore RB, Oborník M, Janoušková J, Chrudimský T, Vancová M, Green DH, Wright SW, Davies NW, Bolch CJS, Heimann K, Šlapeta J, et al. A photosynthetic alveolate closely related to apicomplexan parasites. *Nature*. 2008; 451:959–963. [PubMed: 18288187]
- Moreira D, López-García P. The molecular ecology of microbial eukaryotes unveils a hidden world. *Trends Microbiol*. 2002; 10:31–38. [PubMed: 11755083]
- Not F, del Campo J, Balagué V, de Vargas C, Massana R. New insights into the diversity of marine picoeukaryotes. *PLoS ONE*. 2009; 4:e7143. [PubMed: 19787059]
- Piganeau G, Desdevises Y, Derelle E, Moreau H. Picoeukaryotic sequences in the Sargasso Sea metagenome. *Genome Biol*. 2008; 9:R5. [PubMed: 18179699]
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res*. 2013; 41:D590–D596. [PubMed: 23193283]
- Richards TA, Bass D. Molecular screening of free-living microbial eukaryotes: diversity and distribution using a meta-analysis. *Curr Opin Microbiol*. 2005; 8:240–252. [PubMed: 15939346]
- Richards TA, Vepritskiy AA, Gouliamova DE, Nierzwicki-Bauer SA. The molecular diversity of freshwater picoeukaryotes from an oligotrophic lake reveals diverse, distinctive and globally dispersed lineages. *Environ Microbiol*. 2005; 7:1413–1425. [PubMed: 16104864]
- Rose JM, Caron DA, Sieracki ME, Poulton NJ. Counting heterotrophic nanoplanktonic protists in cultures and aquatic communities by flow cytometry. *Aquat Microb Ecol*. 2004; 34:263–277.

- Roy RS, Price DC, Schliep A, Cai G, Korobeynikov A, Yoon HS, Yang EC, Bhattacharya D. Single cell genome analysis of an uncultured heterotrophic stramenopile. *Sci Rep.* 2014; 4:4780. [PubMed: 24759094]
- Ruiz-Trillo I, Inagaki Y, Davis LLa, Sperstad S, Landfald B, Roger AJ, Sperstad S. *Capsaspora owczarzakii* is an independent opisthokont lineage. *Curr Biol.* 2004; 14:R946–R947. [PubMed: 15556849]
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, et al. Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol.* 2009; 75:7537–7541. [PubMed: 19801464]
- Sebé-Pedrós A, de Mendoza A, Lang BF, Degnan BM, Ruiz-Trillo I. Unexpected repertoire of metazoan transcription factors in the unicellular holozoan *Capsaspora owczarzakii*. *Mol Biol Evol.* 2011; 28:1241–1254. [PubMed: 21087945]
- Sebé-Pedrós A, Irimia M, del Campo J, Parra-Acero H, Russ C, Nusbaum C, Blencowe BJ, Ruiz-Trillo I. Regulated aggregative multicellularity in a close unicellular relative of metazoa. *Elife.* 2013; 2:e01287. [PubMed: 24368732]
- Seenivasan R, Sausen N, Medlin LK, Melkonian M. *Picomonas judraskeda* gen. et sp. nov.: the first identified member of the Picozoa phylum nov., a widespread group of picoeukaryotes, formerly known as “Picobiliphytes”. *PLoS ONE.* 2013; 8:e59565.
- Shalchian-Tabrizi K, Eikrem W, Klaveness D, Vaulot D, Minge MA, Le Gall F, Romari K, Thronsen J, Botnen A, Massana R, Thomsen HA, et al. Telonemia, a new protist phylum with affinity to chromist lineages. *Proc R Soc B: Biol Sci.* 2006; 273:1833–1842.
- Shields JD. The parasitic dinoflagellates of marine crustaceans. *Annu Rev Fish Dis.* 1994; 4:241–271.
- Skovgaard A, Massana R, Balagué V, Saiz E. Phylogenetic position of the copepod-infesting parasite *Syndinium turbo* (Dinoflagellata, Syndinea). *Protist.* 2005; 156:413–423. [PubMed: 16310746]
- Sogin ML, Morrison HG, Huber JA, Welch DM, Huse SM, Neal PR, Arrieta JM, Herndl GJ. Microbial diversity in the deep sea and the underexplored “rare biosphere”. *Proc Natl Acad Sci.* 2006; 103:12115–12120. [PubMed: 16880384]
- Spang A, Saw JH, Jørgensen SL, Zaremba-Niedzwiedzka K, Martijn J, Lind AE, van Eijk R, Schleper C, Guy L, Ettema TJG. Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature.* 2015; 521:173–179. [PubMed: 25945739]
- Stepanuskas R. Single cell genomics: an individual look at microbes. *Curr Opin Microbiol.* 2012; 15:613–620. [PubMed: 23026140]
- Stoeck T, Bass D, Nebel M, Christen R, Jones MDM, Breiner H-W, Richards TA. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol Ecol.* 2010; 19:21–31. [PubMed: 20331767]
- Streets AM, Zhang X, Cao C, Pang Y, Wu X, Xiong L, Yang L, Fu Y, Zhao L, Tang F, Huang Y. Microfluidic single-cell whole-transcriptome sequencing. *Proc Natl Acad Sci.* 2014; 111:7048–7053. [PubMed: 24782542]
- Thornhill DJ, LaJeunesse TC, Santos SR. Measuring rDNA diversity in eukaryotic microbial systems: how intragenomic variation, pseudogenes, and PCR artifacts confound biodiversity estimates. *Mol Ecol.* 2007; 16:5326–5340. [PubMed: 17995924]
- Walker G, Dacks JB, Martin Embley T. Ultrastructural description of *Breviata anathema*, n. gen., n. sp., the organism previously studied as “*Mastigamoeba invertens*”. *J Eukaryot Microbiol.* 2006; 53:65–78. [PubMed: 16579808]
- Woo YH, Ansari H, Otto TD, Klinger CM, Kolisko M, Michálek J, Saxena A, Shanmugam D, Tayyrov A, Veluchamy A, Ali S, et al. Chromerid genomes reveal the evolutionary path from photosynthetic algae to obligate intracellular parasites. *Elife.* 2015; 4:1–41.
- Worden AZ, Janouškovec J, McRose D, Engman A, Welsh RM, Malfatti S, Tringe SG, Keeling PJ. Global distribution of a wild alga revealed by targeted metagenomics. *Curr Biol.* 2012; 22:R675–R677. [PubMed: 22974991]
- Woyke T, Xie G, Copeland A, González JM, Han CS, Kiss H, Saw JH, Senin P, Yang C, Chatterji S, Cheng J-F, et al. Assembling the marine metagenome, one cell at a time. *PLoS ONE.* 2009; 4:e5299. [PubMed: 19390573]

- Yabuki A, Kamikawa R, Ishikawa Sa, Kolisko M, Kim E, Tanabe AS, Kume K, Ishida K, Inagki Y. *Palpitomonas bilix* represents a basal cryptist lineage: insight into the character evolution in Cryptista. *Sci Rep.* 2014; 4:1–6.
- Yoon HS, Price DC, Stepanauskas R, Rajah VD, Sier-acki ME, Wilson WH, Yang EC, Duffy S, Bhattacharya D. Single-cell genomics reveals organismal interactions in uncultivated marine protists. *Science.* 2011; 332:714–717. (80). [PubMed: 21551060]
- Zhao S, Burki F, Brate J, Keeling PJ, Klaveness D, Shalchian-Tabrizi K. Collodictyon - an ancient lineage in the tree of eukaryotes. *Mol Biol Evol.* 2012; 29:1557–1568. [PubMed: 22319147]