

Review



Cite this article: Costa RP, Mizusaki BEP, Sjöström PJ, van Rossum MCW. 2017 Functional consequences of pre- and postsynaptic expression of synaptic plasticity. *Phil. Trans. R. Soc. B* **372**: 20160153. <http://dx.doi.org/10.1098/rstb.2016.0153>

Accepted: 2 November 2016

One contribution of 16 to a discussion meeting issue 'Integrating Hebbian and homeostatic plasticity'.

Subject Areas:

neuroscience, computational biology

Keywords:

synaptic plasticity, Hebbian plasticity, spike-timing-dependent plasticity, homeostatic plasticity, long-term potentiation, synaptic release

Author for correspondence:

Mark C. W. van Rossum
e-mail: mvanross@inf.ed.ac.uk

Functional consequences of pre- and postsynaptic expression of synaptic plasticity

Rui Ponte Costa^{1,2}, Beatriz E. P. Mizusaki^{3,4}, P. Jesper Sjöström⁴
and Mark C. W. van Rossum¹

¹Institute for Adaptive and Neural Computation, School of Informatics University of Edinburgh, Edinburgh, UK

²Centre for Neural Circuits and Behaviour, University of Oxford, Oxford, UK

³Instituto de Física, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

⁴Centre for Research in Neuroscience, Department of Neurology and Neurosurgery, Program for Brain Repair and Integrative Neuroscience, The Research Institute of the McGill University Health Centre, McGill University, Montreal, Quebec, Canada

id RPC, 0000-0003-2595-2027; PJS, 0000-0001-7085-2223; MCWvR, 0000-0001-6525-6814

Growing experimental evidence shows that both homeostatic and Hebbian synaptic plasticity can be expressed presynaptically as well as postsynaptically. In this review, we start by discussing this evidence and methods used to determine expression loci. Next, we discuss the functional consequences of this diversity in pre- and postsynaptic expression of both homeostatic and Hebbian synaptic plasticity. In particular, we explore the functional consequences of a biologically tuned model of pre- and postsynaptically expressed spike-timing-dependent plasticity complemented with postsynaptic homeostatic control. The pre- and postsynaptic expression in this model predicts (i) more reliable receptive fields and sensory perception, (ii) rapid recovery of forgotten information (memory savings), and (iii) reduced response latencies, compared with a model with postsynaptic expression only. Finally, we discuss open questions that will require a considerable research effort to better elucidate how the specific locus of expression of homeostatic and Hebbian plasticity alters synaptic and network computations.

This article is part of the themed issue 'Integrating Hebbian and homeostatic plasticity'.

1. Introduction

Synapses shape the computations of the nervous system. The combination of thousands of excitatory and inhibitory synaptic inputs determine whether a neuron fires or not. Furthermore, the synapse is known to be a key site of information storage in the brain, although not the only one [1]. Changes in the synapses are hypothesized to allow neuronal networks to change function and to adapt through Hebbian and Hebbian-like mechanisms. At the same time, large perturbations in activity levels such as those occurring during synaptogenesis or eye-opening require negative feedback, so that the network can keep its activity level within reasonable bounds and continue performing its computational tasks properly [2,3]. Such homeostatic control of neuronal activity can occur through changes in intrinsic neuronal properties such as control of dendrite excitability [4,5], somatic excitability [1,6] and movement of the axon hillock relative to the soma [7]. However, in this review, we focus on homeostatic processes at the synapse such as synaptic scaling, which provides a form of negative feedback to counter changes in the activity levels, while providing synaptic normalization and competition among inputs [8,9].

As we explain in detail in this review, irrespective of whether synaptic plasticity is Hebbian or homeostatic, the expression locus of plasticity matters. A fundamental distinction is whether the change is pre- or postsynaptic. Changes

in the number of postsynaptic receptors typically only modify the synaptic gain. However, long-term changes in the presynaptic release probability alter the short-term dynamics of the synapse [10–16]. Synaptic dynamics such as short-term depression and facilitation describe how the synaptic efficacy changes during repeated stimulation of the synapse over a time course of hundreds of milliseconds [13,17–19]. These short-term modifications of synaptic efficacy (reviewed in [19]) have been proposed to underlie computations such as gain control [20], redundancy reduction [21] and adaptive filtering [22]. In the context of a recurrent neuronal network, they can affect the activity dynamics and allow the formation and switching among attractor states [23,24], and have been proposed as the basis for working memory [25].

Synaptic plasticity can thus affect network dynamics, but this poses several questions: What are the functional implications of expressing long-term plasticity pre- or postsynaptically? What are the underlying expression mechanisms? Why is there such a large diversity in the expression? Why is there sometimes both pre- and postsynaptic expression? In this review, we begin by discussing pre- and postsynaptic components of Hebbian and homeostatic synaptic plasticity (§2). Then, in §§3 and 4 we examine some of the consequences of the variability of the expression locus of synaptic plasticity, including those that we recently identified using a biologically tuned computational model of neocortical spike-timing-dependent plasticity (STDP) [16].

2. The biological underpinnings of pre- and postsynaptic expression of plasticity

As old as the field of long-term synaptic plasticity itself is the question of how precisely information is stored in neuronal circuits. Historically, Donald Hebb and Jerzy Konorski argued for the strengthening of already existing connections between neurons as a means for information storage, whereas Santiago Ramon y Cajal favoured the growth of new connections [26]. Several relatively recent studies have found evidence that the formation of new synapses is important for long-term information storage in neuronal circuits [27–30]. Indeed, there is strong evidence both in mammals and in the sea slug *Aplysia* that structural plasticity via formation of new afferent inputs is essential for protein-synthesis-dependent long-term memories [31]. The creation of new afferents would correspond to an increase in the number of release sites (box 1: methods), but it should be noted that the number of release sites might be different from the number of anatomical contacts [58].

With already existing connections between neurons, there are essentially two main ways of increasing synaptic strength: either presynaptic release is increased or postsynaptic receptor channels are upregulated [49,59]. Both can be achieved in a number of ways. The presynaptic release probability is controlled by various factors, such as the number and sensitivity of presynaptic calcium channels, as well as other presynaptic ion channels that can modulate neurotransmitter release (such as the epithelial sodium channel EnaC in case of synaptic scaling at the *Drosophila* neuromuscular junction (NMJ) [60,61]), the setpoint of presynaptic calcium sensors involved in eliciting neurotransmitter release, e.g. the synaptotagmins 1, 2 and 9 [62], and the size of the pool of readily releasable vesicles as well as its replenishment rate (in case of homeostasis, see [63,64]) [13,62].

The postsynaptic contribution to the synaptic response is determined by the number and location of postsynaptic receptors, as well as their properties (e.g. conformational state [65] and subunit composition [66,67]). In addition, the geometry of the extracellular space and the apposition of the release sites have also been suggested as important determinants of the response amplitude [68,69].

Experimentally, determination of the expression locus is far from trivial, and a battery of techniques has been applied (box 1). In long-term potentiation (LTP) experiments, evidence for most of the above mechanisms has been found. The historic pre versus post controversy is now typically interpreted as a reflection of the diversity of LTP phenomena, which we now know depends on multiple factors such as age, synapse state, neuromodulation, synapse type and induction protocol [36,49,70–76] (but see [77]). A combination of pre- and postsynaptic expression is also possible [49].

A similar pre- or postsynaptic expression question exists for synaptic homeostasis. While most studies have focused on postsynaptic expression, also here a wide variety in expression, including presynaptic expression [78–80], has been observed, and for instance whether the expression is pre- or postsynaptic appears to depend on developmental stage [81,82]. Sometimes, diversity in mechanisms can even be observed within one system. For instance, in homeostatic plasticity experiments in the hippocampus, both pre- and postsynaptic expression were observed; whereas some CA3–CA3 connections were unexpectedly *reduced* after activity deprivation, other connections strengthened as expected, perhaps to prevent network instability [83]. Also some forms of synaptic scaling at the *Drosophila* and mammalian NMJ are presynaptic: loss of postsynaptic receptors is compensated by increased transmitter release, which restores the mean amplitude of evoked excitatory postsynaptic potentials (EPSPs) [61,84]. A presynaptic locus of expression of homeostatic plasticity at the NMJ is perhaps to be expected, given that the postsynaptic partner—the muscle myotube—does not integrate its inputs such as a neuron does, but rather serves to fire in response to activation at the synaptic input. The pre- and postsynaptic components of the NMJ are therefore tightly co-regulated in synaptogenesis and after damage to ensure proper activation of the muscle [85], so when postsynaptic NMJ sensitivity is reduced, it is in this context not entirely surprising that the presynaptic machinery compensates accordingly by upscaling neurotransmitter release. This example illustrates how the locus of expression must be understood in the context of function of the synapse type at hand.

Further indication that the exact expression locus is functionally important comes from the fact that the expression of both short-term plasticity [86] and long-term plasticity [76] can depend on pre- and postsynaptic cell type. In the case of short-term plasticity, connections from the same presynaptic neurons onto different cells can short-term depress or facilitate depending on the target cell type [87,88], whereas multiple connections between two neurons are often highly similar [45]. Similarly, while STDP exists at both horizontal and vertical excitatory inputs to visual cortex layer 2/3 pyramidal cells, the mechanistic underpinnings as well as the precise temporal requirements for induction are different [89]. Such specificity suggests that the specific locus of expression of long-term plasticity at a given synapse type is meaningful for the proper functioning of microcircuits in the brain, as otherwise tight

Box 1. Methods to determine the locus of plasticity.

The properties of synaptic release can be used to determine the locus of synaptic plasticity by a variety of methods. Among these, there are methods for studying vesicle release, such as FM1-43 dye labelling to explore changes in presynaptic release [32], glutamate uncaging to explore changes in postsynaptic responsiveness or spine size [33,34], measuring *N*-methyl-D-aspartate : α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (NMDA: AMPA) ratio to look for insertion of postsynaptic receptors [35,36], employing the use-dependent NMDA receptor blocker MK-801 to look for changes in glutamate release [37,38], or exploring changes in paired-pulse ratio suggesting a change in probability of release [15,36] (although see [39]).

It is also common to employ spontaneous release as a metric of the locus of expression, as each spontaneously released vesicle gives rise to a well-defined single postsynaptic quantal response known as a miniPSC. This approach is often used in studies of homeostatic plasticity [40], because, here, it is important to measure synaptic changes globally across a majority of inputs to a cell, but this method has also been used to explore Hebbian plasticity [35,41]. An increase in miniPSC frequency in the absence of a change in miniPSC amplitude is typically interpreted as indicating higher release probability or an increase in the number of synaptic contacts, whereas an increased miniPSC amplitude is most often thought to reflect an increase in postsynaptic responsiveness owing to more efficacious postsynaptic receptors. Alternative interpretations of spontaneous release experiments are, however, also possible; for example in the case of AMPAfication of silent synapses, which leads to an apparent change in release probability even though unsilencing is a postsynaptic process [41].

In the scenario where individual synapses are monitored, it is possible to employ methods that rely on the response variability. One such method is non-stationary noise analysis [42], which has been used to determine the effect of homeostasis on inhibitory connections [43], although this method can be unreliable for dendritic synapses [44]. In the related CV analysis, the peak synaptic response is modelled as a binomial process. The process has as parameters the release probability Pr , and the response to each vesicle, the quantal amplitude q . These parameters are assumed identical across the N release sites, and indeed, such coordination has been found [45]. The CV—which is experimentally quantified as the response standard deviation over the mean—is independent of q , namely $CV = \sqrt{(1 - Pr)/PrN}$, and therefore an increase in the mean without an increase in CV can be interpreted as a postsynaptic increase of q [46]. Conversely, if plasticity is presynaptically expressed, then a change in CV is expected, because the CV is a measure of noise, and because the chief source of noise in neurotransmission is the presynaptic stochasticity of vesicle release. The CV analysis method does, however, come with several caveats. In particular, accidental loss or gain of afferent fibres in extracellular stimulation experiments, or unsilencing or growth of new synapses will confuse the results [46]. It is also not obvious that release is independent at different sites, in which case the binomial model is not suitable [46]. By assuming that one of the parameters does not change during the experiment (e.g. fixed N as is reasonable to assume in some plasticity experiments [47,48]), the variance and mean of postsynaptic responses can be used to estimate $Pr = \text{mean}/Nq$ and $q = \text{variance}/\text{mean} + \text{mean}/N$ [16,49,50].

An alternative way to determine whether synaptic changes correspond to alterations of release probability or of quantal response amplitude is to examine the postsynaptic response to a pair or a train of presynaptic stimuli. The idea is that when the release probability is high, the vesicle pool will be depleted more quickly, leading to a more strongly depressing train of postsynaptic responses. When combined with CV analysis, this method can be used to measure all three parameters—English Pr , N and q —of the binomial release model [51]. By fitting these phenomenological models before and after plasticity induction, one can determine which combination of parameters was changed owing to plasticity. It should be noted that experimental results from paired-pulse experiments should also be treated with caution. For example, unsilencing or specific postsynaptic upregulation of release sites with quite different release probability may lead to changes in short-term dynamics that could erroneously be interpreted as presynaptic in origin, even though the actual site of expression is postsynaptic [39]. There are also postsynaptic contributions to synaptic short-term dynamics [52–54] that can complicate the interpretation of experiments. It is therefore better to employ several methods in parallel in the same study—such as CV analysis, paired-pulse ratio, NMDA : AMPA ratio and spontaneous release [15,36]—to independently verify the locus of expression.

Recently, inference methods of short-term plasticity and quantal parameters have been introduced [55–57]. The sampling method of Costa *et al.* [55] is particularly well suited to dealing with the strong correlation and uncertainty in the synapse parameters. Based on this method, we revealed interesting variations between different neuronal connections and proposed more informative experimental protocols based on irregular spike trains, which would be promising to apply in plasticity experiments.

regulation of expression locus would not have arisen during the evolution of the brain.

3. Pre- and postsynaptic expression of spike-timing-dependent plasticity

While the diverse pathways of plasticity induction and expression are increasingly unravelled, their functional roles are still largely an open question. Recently, we have started exploring some of these consequences, using computational

models of STDP. In STDP experiments, where spikes from the presynaptic neuron are paired with millisecond precision with postsynaptic ones, the question of pre- versus postsynaptic expression has also been extensively examined. Depending on factors such as synapse type, brain area and experimental conditions, there is evidence for both pre- and postsynaptic changes [15,36,89–92]. Because of the synapse-type specificity of STDP [76], we used STDP data of connections between visual cortex layer-5 pyramidal cells only [15,36,93]. At this synapse, it has been observed that using STDP induction protocols potentiation has both pre- and postsynaptic components




Figure 1. A schematic of our biologically tuned STDP model with pre- and postsynaptic expression. (a) The synaptic weight is the product of the release probability Pr and the quantal amplitude q . Changes in these parameters owing to STDP are modelled as functions of presynaptic activity trace x_+ and postsynaptic activity traces y_+ and y_- . (b) The fitted model captures the estimated changes in release probability (left) and quantal amplitude (right) for both positive timing (presynaptic spikes 10 ms before postsynaptic ones; blue) and negative timing (presynaptic spikes 10 ms after postsynaptic ones; red), as a function of the frequency of STDP pairings. Symbols indicate data, whereas lines denote the model fit. (c) After LTP, the release probability is enhanced, which leads to stronger short-term depression. The change in short-term synaptic dynamics in the model (bottom) mimics the data (top). Panels (b) and (c) are reproduced with permission from [16].

[36], whereas long-term depression (LTD) is expressed presynaptically only [15]. Presynaptic-only time-dependent LTD has also been found in other synapse types and brain areas [90,92].

Our model of STDP allows for distinct pre- and postsynaptic expression (figure 1a). This phenomenological model relies on three dynamic variables, one which tracks past presynaptic activity $x_+(t)$, and two that track postsynaptic activity, $y_+(t)$ and $y_-(t)$. These traces increase with every spike and decay exponentially between spikes. The plasticity is expressed as a function of the traces, but in contrast to traditional STDP models where just the synaptic weight changes as a function of them [94], here both the release probability and the quantal amplitude are independently modified. In our model, we assume that the number of release sites N is fixed and that it does not change on the timescale of the experiments, consistent with experimental observations [47,48]. However, the model could be straightforwardly generalized to also include changes in N .

Even though we model the observed phenomenology rather than the observed biophysical or mechanistic details, with caution the components of the model can be interpreted to correspond to specific physiological components. The presynaptic trace (x_+), for example, could represent glutamate binding to postsynaptic NMDA receptors, which when depolarized by postsynaptic spikes unblocks NMDA receptors, leading to classical postsynaptic LTP [59]. Similarly, the postsynaptic trace y_+ can be interpreted as retrograde nitric oxide (NO) signalling, which is read out by presynaptic spikes and leads to presynaptically expressed LTP [36]. Finally, the postsynaptic trace y_- can be linked to endocannabinoid (eCB) retrograde release, which triggers presynaptically expressed LTD when coincident with presynaptic spikes [15,90,92].

As mentioned above, we fitted our model to experimental data of one synapse type only (layer-5 pyramidal cells onto layer-5 pyramidal cells in the visual cortex) [15,36,93], across different frequencies and timings. To ensure the biological realism of the model, we further constrained the model fitting by using data from NO and eCB pharmacological blockade experiments in which either presynaptic LTD or LTP expression alone was abolished [36]. Furthermore, we verified that our model captured the expected interaction of short- and long-term plasticity correctly (figure 1c), which permits the exploration of the functional

implications of changes in short dynamics owing to the induction of long-term plasticity.

In the current model, neither LTD nor LTP depends on the state of the synapse—the values of q and Pr . As a result, the current model does not have a (non-trivial) fixed point, and as the fitting to the data only considered the *relative* changes in these parameters, the initial conditions were arbitrarily set to $q = 1$ and $Pr = 0.5$. An improved model could include state dependence in the plasticity to (i) create a fixed point and a realistic weight distribution and (ii) allow fitting to data that takes into account that plasticity might depend on the state (see also §6). Such extensions would, however, require more data. Similarly, it might be possible to model plasticity at the level of voltage [95] or even calcium [96] to capture finer details observed experimentally.

4. Functional consequences of pre- and postsynaptic spike-timing-dependent plasticity expression

The model reveals several functional implications of expressing synaptic plasticity pre- as well as postsynaptically. First, the locus of expression of plasticity will change the trial-to-trial variability of the synaptic response and overall reliability of neurotransmission. Specifically, by increasing the release probability, trial-to-trial reliability from synaptic transmission can be increased. Thus, joint pre- and postsynaptic plasticity can lead to a larger increase in the signal-to-noise ratio (SNR) than postsynaptic modification alone (figure 2a). The functional impact on SNR of this joint modification is consistent with improved sensory perception and its electrophysiological correlates observed in auditory cortex [97].

Second, the pre- and postsynaptic components can differ in stability properties: some changes might be quick to induce, but hard to stabilize and vice versa. This, in turn, can provide neuronal networks with the necessary flexibility to quickly adapt to environmental changes. Using a simple receptive field development simulation, we propose that this might enable a form of memory savings. Memory savings is a concept introduced by Ebbinghaus [98] and means that repeated learning of information is easier, even if the initially learned




Figure 2. Compared with postsynaptic expression alone, STDP with pre- and postsynaptic expression improves sensory perception, enables memory savings and shortens response latencies. (a) Changes in the signal-to-noise ratio (SNR) during receptive field learning in the STDP model. The SNR is represented by the grey scale; the curves represent the various plasticity trajectories starting from the initial condition in the centre. Poisson train inputs that were stimulated at a high rate ('on') obtain high SNR for postsynaptic-only potentiation (dark blue arrows), but combining pre- and postsynaptic potentiation yields considerably better SNR (dark red arrows). Weakly stimulated inputs ('off') obtain lower SNR in either condition (light blue and light red arrows). These modelling results are in keeping with the observed modifications of *in vivo* synaptic responses to a tone from on and off receptive field positions (dark and light green arrows) [97]. (b) Rapid relearning and memory savings with asymmetrically combined pre- and postsynaptic expression of long-term plasticity. Top: response of a neuron to two stimuli, red and blue. The neuron is initially trained with the blue stimulus, and becomes over time selective to it. This initial learning is slow because the changes in q (bottom panel) are slow. After learning, the memory is overwritten with the red stimulus. However, when switching back to the initial blue stimulus, the relearning is more rapid than at first exposure. Middle: presynaptic LTP and LTD can rapidly completely reverse each other. Bottom: LTP has a postsynaptic component that does not reverse quickly, which means a postsynaptic trace is left behind after overwriting with novel information. This hidden trace enables rapid relearning of previously learnt, but overwritten, information. (c) Left: schematic of a firing-rate model with feed-forward and feedback connections as described in [22]. In this network, recurrent synapses are short-term depressing. Changing release probability Pr affects the short-term dynamics, while changing the postsynaptic amplitude q only scales the postsynaptic response. Right: comparison of changes in the response to a 100 ms step stimulus in the recurrent network model when the recurrent synapses are subject to changes in either Pr or q . Increases in the release probability shorten the latency more than increases in the postsynaptic amplitude. Panels (a) and (b) were reproduced with permission from [16].

information appears to have been forgotten. When memories were overwritten, the presynaptic component of the old memory was erased quickly, but the postsynaptic component stayed largely intact. As a result, information that was initially learned but subsequently overwritten could rapidly be recovered upon relearning, provided that the postsynaptic component had not yet decayed completely (figure 2b). This mechanism could thus enable the brain to adapt quickly to different environments or to different tasks without fully forgetting previous learned information. The savings effect mirrors monocular deprivation experiments showing lasting postsynaptic structural effects on spine density that enable more rapid plasticity on repeated monocular deprivation [99,100].

In the STDP data we saw no evidence for any decrease in the postsynaptic component q , perhaps because its decrease may be very slow. Under other protocols, LTD in q has been observed [33]. As it appears unbiological to have no decrease in q , we assumed that a slow homeostatic-like process can decrease q , and so over very long times q decays and the hidden memory trace decays with it. Without this homeostatic process, the hidden trace in q would not decay and memory savings would occur for memories of any age. Our model also suggests that presynaptic boutons should be more dynamic during learning. Recently, Yang *et al.* [101] imaged layer-5 pyramidal cell synapses and found that boutons tend to grow more often than spines after an auditory fear conditioning task.

Finally, while the effects reported in [16] considered feed-forward networks, the changes in release probability under STDP also have consequences for recurrent networks. Excitation-dominated recurrent networks, connected through strong short-term depressing synapses, can have long response latencies that are governed by the synaptic dynamics. We used the model presented in [22] to examine the effect of different expression loci in a recurrent network. Figure 2c illustrates the response of a firing-rate model when the release probability Pr is increased, versus a case in which the quantal amplitude q is increased. The pre- and postsynaptic modifications were set such that the peak responses were identical. In both cases, the response latency was shortened, but when release probability was allowed to increase owing to LTP, response latency shortened about twice as much compared with the case where only postsynaptic plasticity was enabled.

5. Possible other consequences of diversity in locus of plasticity

The 'embarrassment of riches' in the possible expression sites of plasticity [72] is paralleled in many other biological systems. We mention the work of Marder & Goaillard [102] on ion-channel expression, and Turrigiano has emphasized that the multiple ways to achieve homeostasis is puzzling (e.g. review [103]). Considering Hebbian and homeostatic plasticity

together (see Chen *et al.* [104]) complicates this matter even further. It might have a number of consequences beyond the ones discussed above in the STDP model (§4). First, the multiple expression sites provide robustness to the system and multiple ways to maintain the capacity for plasticity, despite internal or external disruption and compensate for genetic defects. Such redundancy can also be advantageous when an abundance of synapses is subject to somewhat diverse learning rules, as it increases the chance that one or some of the synapses correctly adapts to the task at hand. This diversity argument also occurs on the evolutionary level [105], namely a population can be functionally similar but diverse in mechanism, allowing for better adaptation of the population as a whole to novel circumstances. Yet, the publication of yet another pathway often makes one want to exclaim ‘Who ordered that?’, as Rabi did when the subatomic muon particle was discovered.

Second, the multiple expression sites provide flexibility to local circuits, so that, via synapse-type-specific plasticity, different microcircuit components can be independently regulated [76]. For example, LTD at layer 4 to layer 2/3 connections, but not at layer 2/3 to 2/3 synapses, is more readily induced during the critical period [106,107], whereas thalamocortical LTP is already strongly diminished before the critical period has begun [108]. The locus of expression of long-term plasticity at these different synapse types also differs.

Similarly, different plasticity protocols are affected by distinct forms of neuromodulation. The neuromodulators can specifically control forms of STDP that express, for example, postsynaptically [109–111], providing a potential link between behaviourally relevant behaviours and expression loci.

Finally, LTD is not necessarily the opposite of LTP; this becomes even more pressing when considering the diversity of expression mechanisms. In virtually all computational models, LTP induction followed by LTD induction returns the synapse to its original state. Instead, in the abovementioned STDP model, such a protocol might leave the synapse in a different state, even if the apparent synaptic weight is the same, as happens in the case of memory savings. A more direct experimental research of these issues, for instance using learning and subsequent unlearning, would be worthwhile. These considerations also indicate that both the pre- and postsynaptic component need mechanisms to prevent them from saturating and thereby losing the capacity for change. This might be possible by introducing soft bounds for both the pre- and postsynaptic components, or introducing both pre- and postsynaptic normalization [112].

6. Discussion

To model the impact of synaptic plasticity on circuit computations, it is important to know how synapses change during Hebbian and homeostatic plasticity. Here, we have discussed several possible expression sites of synaptic plasticity. We have demonstrated three candidate effects in an STDP model where both pre- and postsynaptic components are modified: (i) a change in the release probability can improve the SNR in the circuit; (ii) the difference in the timescales of modification can lead to the formation of hidden memory traces; and (iii) as a result of changes in synaptic dynamics, the response latency in recurrent networks can be shortened with plasticity. The possible functional impact of combining pre- and postsynaptic plasticity is certainly not restricted to the three findings we

illustrate here. We have rather just scratched the surface of what is likely an emerging field of study.

There is a large range of open issues. For instance, it has long been argued that the stability of memory in spite of continuous molecular turnover is a quite remarkable problem for nature to solve [113,114]. How synapses maintain stable information storage while staying plastic still remains unclear. The diversity of plasticity expression mechanisms could allow for a staged process by which initial changes are presynaptic, but later changes are consolidated structurally [58]. It is, however, not unlikely that multiple expression mechanisms are active in tandem. How these pre- and postsynaptic alterations are coordinated to ensure the long-term fidelity of information storage will require extensive further research. State-based models with a large range of transition rates between states have been explored to resolve this issue [115–118]; see also [119]. As these models are agnostic about expression, the current model could be seen as a biological implementation of such a multistate model. It would, for instance, be of interest to know if the fast resetting of synaptic weights known to occur with exposure to enriched environments [120] is pre- or postsynaptic. It would also be of interest to research if the storage capacity advantages observed in those more theoretical models will also occur in the current phenomenological model. There is also similarity to a recent study in which homeostasis acted as an independent multiplicative mechanism [121].

Another important issue is the weight dependence of long-term plasticity—LTP is hard to induce at synapses that are already strong [93,122–124]—which has important implications for the synaptic weight distribution, memory stability [125] and information capacity [126]. It has been shown that presynaptic modifications strongly depend on the initial release probability [49], which is expected as release probability is bounded between zero and unity. This demonstrates that the weight dependence can stem from presynaptic considerations. However, postsynaptic mechanisms such as compartmentalization of calcium signals may also explain this weight dependence, as it leads to large spines with long necks being ‘write protected’ [127–130]. This finding together with the fact that spine volume is proportional to the expression of AMPA receptors [131] implies that small spines should be more prone to LTP, which is consistent with experimental observations [34]. Such pre- and postsynaptic mechanisms are of course not mutually exclusive and both may contribute to the weight dependence of plasticity [123]. Including these effects would be an obvious next target for the STDP model. Experimentally, it would be of interest to apply protocols [55] that can accurately probe the short-term plasticity parameters before and after STDP induction.

Long-term synaptic plasticity and homeostatic plasticity have been fruitful modelling topics that have clarified the role of plasticity in biological neuronal networks as well as inspired artificial neuronal networks. Yet, despite experimental evidence for presynaptic components in both Hebbian plasticity and synaptic homeostasis, in the overwhelming majority of computational models presynaptic contributions have been ignored (for an exception, see [132,133]), or the models are agnostic about the expression and only adjust the synaptic weight. However, as we have seen, this is not a neutral assumption, and may affect the outcome of the plasticity on network function.

Interestingly, in recurrent networks, short-term plasticity will have an effect on the pre/post activity patterns, and

thereby change STDP induction [134–136]. Theoretically, such mutually interacting systems are extremely challenging [137].

Our discussion has been restricted to the plasticity of excitatory synapses. Inhibitory neurons, in all their diversity [138–140], bring yet another level of complexity as differential short-term dynamics of excitatory and inhibitory synapses yield considerably richer dynamics [55,86,141,142]. We suspect that only a small fraction of the richness and variety of the experimentally observed plasticity phenomena are understood and currently only a few computational models include them. A continued dialogue between theory and experiment should hopefully advance our understanding.

Competing interests. We have no competing interests

References

- Zhang W, Linden DJ. 2003 The other side of the engram: experience-driven changes in neuronal intrinsic excitability. *Nat. Rev. Neurosci.* **4**, 885–900. (doi:10.1038/nrn1248)
- Watt AJ, Desai NS. 2010 Homeostatic plasticity and STDP: keeping a neuron's cool in a fluctuating world. *Front. Synaptic Neurosci.* **2**, 240. (doi:10.3389/fnsyn.2010.00005)
- Keck T *et al.* 2017 Integrating Hebbian and homeostatic plasticity: the current state of the field and future research directions. *Phil. Trans. R. Soc. B* **372**, 20160158. (doi:10.1098/rstb.2016.0158)
- Poolos NP, Migliore M, Johnston D. 2002 Pharmacological upregulation of h-channels reduces the excitability of pyramidal neuron dendrites. *Nat. Neurosci.* **5**, 767–774. (doi:10.1038/nn891)
- Hoffman DA, Magee JC, Colbert CM, Johnston D. 1997 K⁺ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature* **387**, 869–875. (doi:10.1038/42571)
- Desai NS, Rutherford LC, Turrigiano GG. 1999 Plasticity in the intrinsic electrical properties of cortical pyramidal neurons. *Nat. Neurosci.* **2**, 515–520. (doi:10.1038/9165)
- Grubb MS, Burrone J. 2010 Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature* **465**, 1070–1074. (doi:10.1038/nature09160)
- Turrigiano GG. 2011 Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu. Rev. Neurosci.* **34**, 89–103. (doi:10.1146/annurev-neuro-060909-153238)
- Papouin T, Dunphy J, Tolman M, Foley JC, Haydon PG. 2017 Astrocytic control of synaptic function. *Phil. Trans. R. Soc. B* **372**, 20160154. (doi:10.1098/rstb.2016.0154)
- Markram H, Tsodyks M. 1996 Redistribution of synaptic efficacy between neocortical pyramidal neurons. *Nature* **382**, 807–810. (doi:10.1038/382807a0)
- Zakharenko S, Zablow L, Siegelbaum S. 2001 Visualization of changes in presynaptic function during long-term synaptic plasticity. *Nat. Neurosci.* **4**, 711–717. (doi:10.1038/89498)
- Zakharenko SS, Zablow L, Siegelbaum SA. 2002 Altered presynaptic vesicle release and cycling during mGluR-dependent LTD. *Neuron* **35**, 1099–1110. (doi:10.1016/S0896-6273(02)00898-X)
- Zucker RS, Regehr WG. 2002 Short-term synaptic plasticity. *Annu. Rev. Physiol.* **64**, 355–405. (doi:10.1146/annurev.physiol.64.092501.114547)
- Gerdeman GL, Ronesi J, Lovinger DM. 2002 Postsynaptic endocannabinoid release is critical to long-term depression in the striatum. *Nat. Neurosci.* **5**, 446–451. (doi:10.1038/nn832)
- Sjöström PJ, Turrigiano GG, Nelson SB. 2003 Neocortical LTD via coincident activation of presynaptic NMDA and cannabinoid receptors. *Neuron* **39**, 641–654. (doi:10.1016/S0896-6273(03)00476-8)
- Costa RP, Froemke RC, Sjöström PJ, van Rossum MC. 2015 Unified pre- and postsynaptic long-term plasticity enables reliable and flexible learning. *eLife* **4**, e09457. (doi:10.7554/elife.09457)
- Abbott LF, Regehr WG. 2004 Synaptic computation. *Nature* **431**, 796–803. (doi:10.1038/nature03010)
- Hennig MH. 2013 Theoretical models of synaptic short term plasticity. *Front. Comput. Neurosci.* **7**, 45. (doi:10.3389/fncom.2013.00045)
- Tsodyks M, Wu S. 2013 Short-term synaptic plasticity. *Scholarpedia*. revision #136920.
- Abbott LF, Varela JA, Sen K, Nelson SB. 1997 Synaptic depression and cortical gain control. *Science* **275**, 220–224. (doi:10.1126/science.275.5297.221)
- Goldman MS, Maldonado P, Abbott LF. 2002 Redundancy reduction and sustained firing with stochastic depressing synapses. *J. Neurosci.* **22**, 584–591.
- van Rossum MCW, van der Meer MAA, Xiao D, Oram MW. 2008 Adaptive integration in the visual cortex by depressing recurrent cortical circuits. *Neural Comput.* **20**, 1847–1872. (doi:10.1162/neco.2008.06-07-546)
- Barak O, Tsodyks M. 2007 Persistent activity in neural networks with dynamic synapses. *PLoS Comput. Biol.* **3**, e35. (doi:10.1371/journal.pcbi.0030035)
- York LC, van Rossum MCW. 2009 Recurrent networks with short term synaptic depression. *J. Comput. Neurosci.* **27**, 607–620. (doi:10.1007/s10827-009-0172-4)
- Mongillo G, Barak O, Tsodyks M. 2008 Synaptic theory of working memory. *Science* **319**, 1543–1546. (doi:10.1126/science.1150769)
- Markram H, Gerstner W, Sjöström PJ. 2011 A history of spike-timing-dependent plasticity. *Front. Synaptic Neurosci.* **3**, 4. (doi:10.3389/fnsyn.2011.00004)
- Engert F, Bonhoeffer T. 1999 Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* **399**, 66–70. (doi:10.1038/19978)
- Le Bé J-V, Markram H. 2006 Spontaneous and evoked synaptic rewiring in the neonatal neocortex. *Proc. Natl Acad. Sci. USA* **103**, 13 214–13 219. (doi:10.1073/pnas.0604691103)
- Kwon H-B, Sabatini BL. 2011 Glutamate induces de novo growth of functional spines in developing cortex. *Nature* **474**, 100–104. (doi:10.1038/nature09986)
- Nabavi S, Fox R, Proulx CD, Lin JY, Tsien RY, Malinow R. 2014 Engineering a memory with LTD and LTP. *Nature* **511**, 348–352. (doi:10.1038/nature13294)
- Kandel ER. 2001 The molecular biology of memory storage: a dialogue between genes and synapses. *Science* **294**, 1030–1038. (doi:10.1126/science.1067020)
- Stanton PK, Winterer J, Zhang X-I, Müller W. 2005 Imaging LTP of presynaptic release of FM1-43 from the rapidly recycling vesicle pool of Schaffer collateral-CA1 synapses in rat hippocampal slices. *Eur. J. Neurosci.* **22**, 2451–2461. (doi:10.1111/j.1460-9568.2005.04437.x)
- Dotz H-U, Eder M, Frick A, Zieglgänsberger W. 1999 Precisely localized LTD in the neocortex revealed by infrared-guided laser stimulation. *Science* **286**, 110–113. (doi:10.1126/science.286.5437.110)
- Matsuzaki M, Honkura N, Ellis-Davies GCR, Kasai H. 2004 Structural basis of long-term potentiation in single dendritic spines. *Nature* **429**, 761–766. (doi:10.1038/nature02617)

35. Watt AJ, Sjöström PJ, Häusser M, Nelson SB, Turrigiano GG. 2004 A proportional but slower NMDA potentiation follows AMPA potentiation in LTP. *Nat. Neurosci.* **7**, 518–524. (doi:10.1038/nn1220)
36. Sjöström PJ, Turrigiano GG, Nelson S. 2007 Multiple forms of long-term plasticity at unitary neocortical layer 5 synapses. *Neuropharmacology* **52**, 176–184. (doi:10.1016/j.neuropharm.2006.07.021)
37. Kullmann DM, Erdemli G, Asztély F. 1996 LTP of AMPA and NMDA receptor-mediated signals: evidence for presynaptic expression and extrasynaptic glutamate spill-over. *Neuron* **17**, 461–474. (doi:10.1016/S0896-6273(00)80178-6)
38. Hessler NA, Shirke AM, Malinow R. 1993 The probability of transmitter release at a mammalian central synapse. *Nature* **366**, 569–572. (doi:10.1038/366569a0)
39. Poncer J, Malinow R. 2001 Postsynaptic conversion of silent synapses during LTP affects synaptic gain and transmission dynamics. *Nat. Neurosci.* **4**, 989–996. (doi:10.1038/nn719)
40. Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB. 1998 Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature* **391**, 892–896. (doi:10.1038/36103)
41. Isaac J, Oliet S, Hjelmstad G, Nicoll R, Malenka R. 1996 Expression mechanisms of long-term potentiation in the hippocampus. *J. Physiol. Paris* **90**, 299–303. (doi:10.1016/S0928-4257(97)87901-6)
42. Traynelis SF, Silver RA, Cull-Candy SG. 1993 Estimated conductance of glutamate receptor channels activated during EPSCs at the cerebellar mossy fiber-granule cell synapse. *Neuron* **11**, 279–289. (doi:10.1016/0896-6273(93)90184-S)
43. Kilman V, van Rossum MCW, Turrigiano GG. 2002 Activity deprivation reduces mIPSC amplitude by decreasing the number of postsynaptic GABA_A receptors clustered at neocortical synapses. *J. Neurosci.* **22**, 1328–1337.
44. Feldwisch-Drentrup H, Barrett AB, Smith MT, van Rossum MC. 2012 Fluctuations in the open time of synaptic channels: an application to noise analysis based on charge. *J. Neurosci. Methods* **210**, 15–21. (doi:10.1016/j.jneumeth.2011.11.004)
45. Koester HJ, Johnston D. 2005 Target cell-dependent normalization of transmitter release at neocortical synapses. *Science* **308**, 863–866. (doi:10.1126/science.1100815)
46. Faber DS, Korn H. 1991 Applicability of the coefficient of variation method for analyzing synaptic plasticity. *Biophys. J.* **60**, 1288–1294. (doi:10.1016/S0006-3495(91)82162-2)
47. Bolshakov VV, Golan H, Kandel ER, Siegelbaum SA. 1997 Recruitment of new sites of synaptic transmission during the cAMP-dependent late phase of LTP at CA3-CA1 synapses in the hippocampus. *Neuron* **19**, 635–651. (doi:10.1016/S0896-6273(00)80377-3)
48. Saez I, Friedlander MJ. 2009 Plasticity between neuronal pairs in layer 4 of visual cortex varies with synapse state. *J. Neurosci.* **29**, 15 286–15 298. (doi:10.1523/JNEUROSCI.2980-09.2009)
49. Larkman A, Hannay T, Stratford K, Jack J. 1992 Presynaptic release probability influences the locus of long-term potentiation. *Nature* **360**, 70–73. (doi:10.1038/360070a0)
50. Markram H, Lübke J, Frotscher M, Roth A, Sakmann B. 1997 Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex. *J. Physiol.* **500**, 409–440. (doi:10.1113/jphysiol.1997.sp022031)
51. Loebel A, Silberberg G, Helbig D, Markram H, Tsodyks M, Richardson MJE. 2009 Multiquantal release underlies the distribution of synaptic efficacies in the neocortex. *Front. Comput. Neurosci.* **3**, 27. (doi:10.3389/neuro.10.027.2009)
52. Rozov A, Burnashev N. 1999 Polyamine-dependent facilitation of postsynaptic AMPA receptors counteracts paired-pulse depression. *Nature* **401**, 594–598. (doi:10.1038/44151)
53. Rozov A, Jerecic J, Sakmann B, Burnashev N. 2001 AMPA receptor channels with long-lasting desensitization in bipolar interneurons contribute to synaptic depression in a novel feedback circuit in layer 2/3 of rat neocortex. *J. Neurosci.* **21**, 8062–8071.
54. Heine M, Groc L, Frischknecht R, Beique J-C, Lounis B, Rumbaugh G, Huganir RL, Cognet L, Choquet D. 2008 Surface mobility of postsynaptic AMPARs tunes synaptic transmission. *Science* **320**, 201–205. (doi:10.1126/science.1152089)
55. Costa RP, Sjöström PJ, van Rossum MCW. 2013 Probabilistic inference of short-term synaptic plasticity in neocortical microcircuits. *Front. Comput. Neurosci.* **7**, 75. (doi:10.3389/fncom.2013.00075)
56. Bhumbra GS, Beato M. 2013 Reliable evaluation of the quantal determinants of synaptic efficacy using Bayesian analysis. *J. Neurophysiol.* **109**, 603–620. (doi:10.1152/jn.00528.2012)
57. Barri A, Wang Y, Hansel D, Mongillo G. 2016 Quantifying repetitive transmission at chemical synapses: a generative-model approach. *eNeuro* **3**, ENEURO-0113. (doi:10.1523/ENEURO.0113-15.2016)
58. Loebel A, Le Bé J-V, Richardson MJE, Markram H, Herz AV. 2013 Matched pre- and post-synaptic changes underlie synaptic plasticity over long time scales. *J. Neurosci.* **33**, 6257–6266. (doi:10.1523/JNEUROSCI.3740-12.2013)
59. Bliss TV, Collingridge GL. 1993 A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361**, 31–39. (doi:10.1038/361031a0)
60. Younger MA, Müller M, Tong A, Pym EC, Davis GW. 2013 A presynaptic ENaC channel drives homeostatic plasticity. *Neuron* **79**, 1183–1196. (doi:10.1016/j.neuron.2013.06.048)
61. Davis GW, Müller M. 2015 Homeostatic control of presynaptic neurotransmitter release. *Annu. Rev. Physiol.* **77**, 251–270. (doi:10.1146/annurev-physiol-021014-071740)
62. Kaeser PS, Regehr WG. 2014 Molecular mechanisms for synchronous, asynchronous, and spontaneous neurotransmitter release. *Annu. Rev. Physiol.* **76**, 333. (doi:10.1146/annurev-physiol-021113-170338)
63. Müller M, Liu KSY, Sigrist SJ, Davis GW. 2012 RIM controls homeostatic plasticity through modulation of the readily-releasable vesicle pool. *J. Neurosci.* **32**, 16 574–16 585. (doi:10.1523/JNEUROSCI.0981-12.2012)
64. Wang X, Pinter MJ, Rich MM. 2016 Reversible recruitment of a homeostatic reserve pool of synaptic vesicles underlies rapid homeostatic plasticity of quantal content. *J. Neurosci.* **36**, 828–836. (doi:10.1523/JNEUROSCI.3786-15.2016)
65. Lee H-K, Barbarosie M, Kameyama K, Bear MF, Huganir RL. 2000 Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* **405**, 955–959. (doi:10.1038/35016089)
66. Kessels HW, Malinow R. 2009 Synaptic AMPA receptor plasticity and behavior. *Neuron* **61**, 340–350. (doi:10.1016/j.neuron.2009.01.015)
67. Isaac JT, Ashby MC, McBain CJ. 2007 The role of the GluR2 subunit in AMPA receptor function and synaptic plasticity. *Neuron* **54**, 859–871. (doi:10.1016/j.neuron.2007.06.001)
68. Raghavachari S, Lisman JE. 2004 Properties of quantal transmission at CA1 synapses. *J. Neurophysiol.* **92**, 2456–2467. (doi:10.1152/jn.00258.2004)
69. Lisman JE, Raghavachari S, Tsien RW. 2007 The sequence of events that underlie quantal transmission at central glutamatergic synapses. *Nat. Rev. Neurosci.* **8**, 597–609. (doi:10.1038/nrn2191)
70. Larkman AU, Jack JJB. 1995 Synaptic plasticity: hippocampal LTP. *Curr. Opin. Neurobiol.* **5**, 324–334. (doi:10.1016/0959-4388(95)80045-X)
71. Frey U, Morris RG. 1997 Synaptic tagging and long-term potentiation. *Nature* **385**, 533–536. (doi:10.1038/385533a0)
72. Malenka RC, Bear MF. 2004 LTP and LTD: an embarrassment of riches. *Neuron* **44**, 5–21. (doi:10.1016/j.neuron.2004.09.012)
73. Kullmann DM, Moreau AW, Bakiri Y, Nicholson E. 2012 Plasticity of inhibition. *Neuron* **75**, 951–962. (doi:10.1016/j.neuron.2012.07.030)
74. Padamsey Z, Emptage N. 2014 Two sides to long-term potentiation: a view towards reconciliation. *Phil. Trans. R. Soc. B* **369**, 20130154. (doi:10.1098/rsta.2013.0154)
75. Park P, Volianskis A, Sanderson TM, Bortolotto ZA, Jane DE, Zhuo M, Kaang B-K, Collingridge GL. 2014 Nmda receptor-dependent long-term potentiation comprises a family of temporally overlapping forms of synaptic plasticity that are induced by different patterns of stimulation. *Phil. Trans. R. Soc. B* **369**, 20130131. (doi:10.1098/rsta.2013.0131)
76. Larsen RS, Sjöström PJ. 2015 Synapse-type-specific plasticity in local circuits. *Curr. Opin. Neurobiol.* **35**, 127–135. (doi:10.1016/j.conb.2015.08.001)
77. Granger AJ, Nicoll RA. 2014 Expression mechanisms underlying long-term potentiation: a postsynaptic view, 10 years on. *Phil. Trans. R. Soc. B* **369**, 20130136. (doi:10.1098/rsta.2013.0136)
78. Bacci A, Coco S, Pravettoni E, Schenk U, Armano S, Frassoni C, Verderio C, De Camilli P, Matteoli M.

- 2001 Chronic blockade of glutamate receptors enhances presynaptic release and downregulates the interaction between synaptophysin-synaptobrevin-vesicle-associated membrane protein 2. *J. Neurosci.* **21**, 6588–6596.
79. Burrone J, O'Byrne M, Murthy VN. 2002 Multiple forms of synaptic plasticity triggered by selective suppression of activity in individual neurons. *Nature* **420**, 414–418. (doi:10.1038/nature01242)
80. Thiagarajan TC, Lindskog M, Tsien RW. 2005 Adaptation to synaptic inactivity in hippocampal neurons. *Neuron* **47**, 725–737. (doi:10.1016/j.neuron.2005.06.037)
81. Wierenga CJ, Walsh MF, Turrigiano GG. 2006 Temporal regulation of the expression locus of homeostatic plasticity. *J. Neurophysiol.* **96**, 2127–2133. (doi:10.1152/jn.00107.2006)
82. Han EB, Stevens CF. 2009 Development regulates a switch between post- and presynaptic strengthening in response to activity deprivation. *Proc. Natl Acad. Sci. USA* **106**, 10 817–10 822. (doi:10.1073/pnas.0903603106)
83. Kim J, Tsien RW. 2008 Synapse-specific adaptations to inactivity in hippocampal circuits achieve homeostatic gain control while dampening network reverberation. *Neuron* **58**, 925–937. (doi:10.1016/j.neuron.2008.05.009)
84. Ouanounou G, Baux G, Bal T. 2016 A novel synaptic plasticity rule explains homeostasis of neuromuscular transmission. *eLife* **5**, e12190. (doi:10.7554/eLife.12190)
85. Sanes JR, Lichtman JW. 2001 Induction, assembly, maturation and maintenance of a postsynaptic apparatus. *Nat. Rev. Neurosci.* **2**, 791–805. (doi:10.1038/35097557)
86. Blackman AV, Abrahamsson T, Costa RP, Lalanne T, Sjostrom PJ. 2013 Target-cell-specific short-term plasticity in local circuits. *Front. Synaptic Neurosci.* **5**, 11. (doi:10.3389/fnsyn.2013.00011)
87. Markram H, Wang Y, Tsodyks M. 1998 Differential signaling via the same axon of neocortical pyramidal neurons. *Proc. Natl Acad. Sci. USA* **95**, 5323–5328. (doi:10.1073/pnas.95.9.5323)
88. Buchanan KA, Blackman AV, Moreau AW, Elgar D, Costa RP, Lalanne T, Tudor Jones AA, Oyrer J, Sjostrom PJ. 2012 Target-specific expression of presynaptic NMDA receptors in neocortical microcircuits. *Neuron* **75**, 451–466. (doi:10.1016/j.neuron.2012.06.017)
89. Banerjee A, González-Rueda A, Sampaio-Baptista C, Paulsen O, Rodríguez-Moreno A. 2014 Distinct mechanisms of spike timing-dependent LTD at vertical and horizontal inputs onto L2/3 pyramidal neurons in mouse barrel cortex. *Physiol. Rep.* **2**, e00271. (doi:10.1002/phy2.271)
90. Bender VA, Feldman D. 2006 Two coincidence detectors for spike timing-dependent plasticity in somatosensory cortex. *J. Neurosci.* **26**, 4166–4177. (doi:10.1523/JNEUROSCI.0176-06.2006)
91. Yang Y, Calakos N. 2013 Presynaptic long-term plasticity. *Front. Synaptic Neurosci.* **5**, 8. (doi:10.3389/fnsyn.2013.00008)
92. Andrade-Talavera Y, Duque-Feria P, Paulsen O, Rodríguez-Moreno A. 2016 Presynaptic spike timing-dependent long-term depression in the mouse hippocampus. *Cerebr. Cortex* **26**, 3637–3654. (doi:10.1093/cercor/bhw172)
93. Sjostrom PJ, Turrigiano GG, Nelson SB. 2001 Rate, timing, and cooperativity jointly determine cortical synaptic plasticity. *Neuron* **32**, 1149–1164. (doi:10.1016/S0896-6273(01)00542-6)
94. Pfister J-P, Gerstner W. 2006 Triplets of spikes in a model of spike timing-dependent plasticity. *J. Neurosci.* **26**, 9673–9682. (doi:10.1523/JNEUROSCI.1425-06.2006)
95. Clopath C, Büsing L, Vasilaki E, Gerstner W. 2010 Connectivity reflects coding: a model of voltage-based STDP with homeostasis. *Nat. Neurosci.* **13**, 344–352. (doi:10.1038/nn.2479)
96. Graupner M, Brunel N. 2012 Calcium-based plasticity model explains sensitivity of synaptic changes to spike pattern, rate, and dendritic location. *Proc. Natl Acad. Sci. USA* **109**, 3991–3996. (doi:10.1073/pnas.1109359109)
97. Froemke RC *et al.* 2013 Long-term modification of cortical synapses improves sensory perception. *Nat. Neurosci.* **16**, 79–88. (doi:10.1038/nn.3274)
98. Ebbinghaus H. 1913 *Memory: a contribution to experimental psychology*. New York, NY: Teachers College Press, Columbia University.
99. Hofer S, Mrsic-Flogel T, Bonhoeffer T, Hübener M. 2008 Experience leaves a lasting structural trace in cortical circuits. *Nature* **457**, 313–317. (doi:10.1038/nature07487)
100. Hübener M, Bonhoeffer T. 2010 Searching for engrams. *Neuron* **67**, 363–371. (doi:10.1016/j.neuron.2010.06.033)
101. Yang Y, Liu DQ, Huang W, Deng J, Sun Y, Zuo Y, Poo MM. 2016 Selective synaptic remodeling of amygdalocortical connections associated with fear memory. *Nat. Neurosci.* **19**, 1348–1355. (doi:10.1038/nn.4370)
102. Marder E, Goaillard J-M. 2006 Variability, compensation and homeostasis in neuron and network function. *Nat. Rev. Neurosci.* **7**, 563–574. (doi:10.1038/nrn1949)
103. Turrigiano GG. 2017 The dialectic of Hebb and homeostasis. *Phil. Trans. R. Soc. B* **372**, 20160258. (doi:10.1098/rstb.2016.0258)
104. Yee AX, Hsu Y-T, Chen L. 2017 A metaplasticity view of the interaction between homeostatic and Hebbian plasticity. *Phil. Trans. R. Soc. B* **372**, 20160155. (doi:10.1098/rstb.2016.0155)
105. Daniels BC, Chen Y-J, Sethna JP, Gutenkunst RN, Myers CR. 2008 Sloppiness, robustness, and evolvability in systems biology. *Curr. Opin. Biotechnol.* **19**, 389–395. (doi:10.1016/j.copbio.2008.06.008)
106. Corlew R, Wang Y, Ghermazien H, Erisir A, Philpot BD. 2007 Developmental switch in the contribution of presynaptic and postsynaptic NMDA receptors to long-term depression. *J. Neurosci.* **27**, 9835–9845. (doi:10.1523/JNEUROSCI.5494-06.2007)
107. Banerjee A, Meredith RM, Rodríguez-Moreno A, Mierau SB, Auberson YP, Paulsen O. 2009 Double dissociation of spike timing-dependent potentiation and depression by subunit-preferring NMDA receptor antagonists in mouse barrel cortex. *Cerebr. Cortex* **19**, 2959–2969. (doi:10.1093/cercor/bhp067)
108. Crair MC, Malenka RC. 1995 A critical period for long-term potentiation at thalamocortical synapses. *Nature* **375**, 325–328. (doi:10.1038/375325a0)
109. Seol GH, Ziburkus J, Huang SY, Song L, Kim IT, Takamiya K, Huganir RL, Lee H-K, Kirkwood A. 2007 Neuromodulators control the polarity of spike-timing-dependent synaptic plasticity. *Neuron* **55**, 919–929. (doi:10.1016/j.neuron.2007.08.013)
110. Pawlak V, Wickens JR, Kirkwood A, Kerr JND. 2010 Timing is not everything: neuromodulation opens the STDP gate. *Front. Synaptic Neurosci.* **2**, 146. (doi:10.3389/fnsyn.2010.00146)
111. Frémaux N, Gerstner W. 2016 Neuromodulated spike-timing-dependent plasticity, and theory of three-factor learning rules. *Front. Neural Circuits* **9**, 1178. (doi:10.3389/fncir.2015.00085)
112. Zenke F, Gerstner W. 2017 Hebbian plasticity requires compensatory processes on multiple timescales. *Phil. Trans. R. Soc. B* **372**, 20160259. (doi:10.1098/rstb.2016.0259)
113. Crick F. 1984 Memory and molecular turnover. *Nature* **312**, 101. (doi:10.1038/312101a0)
114. Lisman J. 1989 A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory. *Proc. Natl Acad. Sci. USA* **86**, 9574–9578. (doi:10.1073/pnas.86.23.9574)
115. Fusi S, Drew PJ, Abbott L. 2005 Cascade models of synaptically stored memories. *Neuron* **45**, 599–611. (doi:10.1016/j.neuron.2005.02.001)
116. Leibold C, Kempster R. 2008 Sparseness constrains the prolongation of memory lifetime via synaptic metaplasticity. *Cerebr. Cortex* **18**, 67–77. (doi:10.1093/cercor/bhm037)
117. Barrett AB, van Rossum MCW. 2008 Optimal learning rules for discrete synapses. *PLoS Comput. Biol.* **4**, e1000230. (doi:10.1371/journal.pcbi.1000230)
118. Lahiri S, Ganguli S. 2013 A memory frontier for complex synapses. *Adv. Neural Inf. Process. Syst.* 1034–1042.
119. Liu KKL, Hagan MF, Lisman JE. 2017 Gradation (approx. 10 size states) of synaptic strength by quantal addition of structural modules. *Phil. Trans. R. Soc. B* **372**, 20160328. (doi:10.1098/rstb.2016.0328)
120. Abraham WC, Logan B, Greenwood JM, Dragunow M. 2002 Induction and experience-dependent consolidation of stable long-term potentiation lasting months in the hippocampus. *J. Neurosci.* **22**, 9626–9634.
121. Toyozumi T, Kaneko M, Stryker MP, Miller KD. 2014 Modeling the dynamic interaction of Hebbian and homeostatic plasticity. *Neuron* **84**, 497–510. (doi:10.1016/j.neuron.2014.09.036)
122. Bi G-Q, Poo M-M. 1998 Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. *J. Neurosci.* **18**, 10 464–10 472.
123. Liao D, Jones A, Malinow R. 1992 Direct measurement of quantal changes underlying

- long-term potentiation in CA1 hippocampus. *Neuron* **9**, 1089–1097. (doi:10.1016/0896-6273(92)90068-0)
124. Debanne D, Gähwiler BH, Thompson SM. 1999 Heterogeneity of synaptic plasticity at unitary CA1-CA3 and CA3-CA3 connections in rat hippocampal slice cultures. *J. Neurosci.* **19**, 10 664–10 671.
125. Billings G, van Rossum MCW. 2009 Memory retention and spike-timing-dependent plasticity. *J. Neurophysiol.* **101**, 2775–2788. (doi:10.1152/jn.91007.2008)
126. van Rossum M, Shippi M, Barrett A. 2012 Soft-bound synaptic plasticity outperforms hard-bound plasticity. *PLoS Comput. Biol.* **8**, e1002386. (<http://dx.doi.org/10.1371/journal.pcbi.1002386>)
127. Noguchi J, Matsuzaki M, Ellis-Davies GCR, Kasai H. 2005 Spine-neck geometry determines NMDA receptor-dependent Ca²⁺ signaling in dendrites. *Neuron* **46**, 609–622. (doi:10.1016/j.neuron.2005.03.015)
128. Yasumatsu N, Matsuzaki M, Miyazaki T, Noguchi J, Kasai H. 2008 Principles of long-term dynamics of dendritic spines. *J. Neurosci.* **28**, 13 592–13 608. (doi:10.1523/JNEUROSCI.0603-08.2008)
129. Kalantzis G, Shouval HZ. 2009 Structural plasticity can produce metaplasticity. *PLoS ONE* **4**, e8062. (doi:10.1371/journal.pone.0008062)
130. O'Donnell C, Nolan MF, van Rossum MCW. 2011 Dendritic spine dynamics regulate the long-term stability of synaptic plasticity. *J. Neurosci.* **31**, 16 142–16 156. (doi:10.1523/JNEUROSCI.2520-11.2011)
131. Matsuzaki M, Ellis-Davies GCR, Nemoto T, Miyashita Y, Iino M, Kasai H. 2001 Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat. Neurosci.* **4**, 1086–1092. (doi:10.1038/nn736)
132. Senn W, Markram H, Tsodyks M. 2001 An algorithm for modifying neurotransmitter release probability based on pre- and postsynaptic spike timing. *Neural Comput.* **13**, 35–67. (doi:10.1162/089976601300014628)
133. Lisman J, Raghavachari S. 2006 A unified model of the presynaptic and postsynaptic changes during LTP at CA1 synapses. *Sci. STKE* **2006**, re11. (doi:10.1126/stke.3562006re11)
134. Wang Y, Markram H, Goodman PH, Berger TK, Ma J, Goldman-Rakic PS. 2006 Heterogeneity in the pyramidal network of the medial prefrontal cortex. *Nat. Neurosci.* **9**, 534–542. (doi:10.1038/nn1670)
135. Vasilaki E, Giugliano M. 2014 Emergence of connectivity motifs in networks of model neurons with short- and long-term plastic synapses. *PLoS ONE* **9**, e84626. (doi:10.1371/journal.pone.0084626)
136. Froemke R, Dan Y. 2002 Spike-timing-dependent synaptic modification induced by natural spike trains. *Nature* **416**, 433–438. (doi:10.1038/416433a)
137. Gilson M, Burkitt A, Grayden D, Thomas D, van Hemmen J. 2009 Emergence of network structure due to spike-timing-dependent plasticity in recurrent neuronal networks. I. Input selectivity–strengthening correlated input pathways. *Biol. Cybern.* **101**, 81–102. (doi:10.1007/s00422-009-0319-4)
138. Markram H, Toledo-Rodriguez Maria, Wang Y, Gupta A, Silberberg G, Wu C. 2004 Interneurons of the neocortical inhibitory system. *Nat. Rev. Neurosci.* **5**, 793–807. (doi:10.1038/nrn1519)
139. DeFelipe J *et al.* 2013 New insights into the classification and nomenclature of cortical GABAergic interneurons. *Nat. Rev. Neurosci.* **14**, 202–216. (doi:10.1038/nrn3444)
140. Kepecs A, Fishell G. 2014 Interneuron cell types are fit to function. *Nature* **505**, 318–326. (doi:10.1038/nature12983)
141. Galarreta M, Hestrin S. 1998 Frequency-dependent synaptic depression and the balance of excitation and inhibition in the neocortex. *Nat. Neurosci.* **1**, 587–594. (doi:10.1038/2822)
142. Varela JA, Song S, Turrigiano GG, Nelson SB. 1999 Differential depression at excitatory and inhibitory synapses in visual cortex. *J. Neurosci.* **19**, 4293–4304.