



Jose-Andres C. Portillo,¹ Yalitz Lopez Corcino,¹ Yanling Miao,¹ Jie Tang,² Nader Sheibani,³ Timothy S. Kern,^{2,4,5} George R. Dubyak,⁶ and Carlos S. Subauste^{1,4,7}

CD40 in Retinal Müller Cells Induces P2X₇-Dependent Cytokine Expression in Macrophages/Microglia in Diabetic Mice and Development of Early Experimental Diabetic Retinopathy



Diabetes 2017;66:483–493 | DOI: 10.2337/db16-0051

Müller cells and macrophages/microglia are likely important for the development of diabetic retinopathy; however, the interplay between these cells in this disease is not well understood. An inflammatory process is linked to the onset of experimental diabetic retinopathy. CD40 deficiency impairs this process and prevents diabetic retinopathy. Using mice with CD40 expression restricted to Müller cells, we identified a mechanism by which Müller cells trigger proinflammatory cytokine expression in myeloid cells. During diabetes, mice with CD40 expressed in Müller cells upregulated retinal tumor necrosis factor- α (TNF- α), interleukin 1 β (IL-1 β), intracellular adhesion molecule 1 (ICAM-1), and nitric oxide synthase (NOS2), developed leukostasis and capillary degeneration. However, CD40 did not cause TNF- α or IL-1 β secretion in Müller cells. TNF- α was not detected in Müller cells from diabetic mice with CD40⁺ Müller cells. Rather, TNF- α was upregulated in macrophages/microglia. CD40 ligation in Müller cells triggered phospholipase C-dependent ATP release that caused P2X₇-dependent production of TNF- α and IL-1 β by macrophages. P2X₇^{-/-} mice and mice treated with a P2X₇ inhibitor were protected from diabetes-induced TNF- α , IL-1 β , ICAM-1, and NOS2 upregulation. Our studies indicate that CD40 in Müller cells is sufficient to upregulate retinal inflammatory markers and appears

to promote experimental diabetic retinopathy and that Müller cells orchestrate inflammatory responses in myeloid cells through a CD40-ATP-P2X₇ pathway.

Increasing evidence indicates that chronic low-grade inflammation is important for the development of diabetic retinopathy (1,2). Tumor necrosis factor- α (TNF- α) and interleukin 1 β (IL-1 β) are proinflammatory molecules upregulated in this disease (3,4). Macrophages/microglia in the diabetic retina express TNF- α (4). Moreover, both cytokines contribute to diabetes-induced degeneration of retinal capillaries, a hallmark of diabetic retinopathy (5,6). In addition to macrophages/microglia, Müller cells (the major retinal macroglia) become dysfunctional in diabetes and contribute to the development of diabetic retinopathy (7). However, little is known about whether Müller cells enhance proinflammatory responses in macrophages/microglia in diabetes.

CD40 is an important driver of retinal inflammation in experimental diabetic retinopathy (8,9). CD40 is upregulated in retinal Müller cells, endothelial cells, and microglia in diabetic mice (8). CD40 ligation in Müller cells and endothelial cells upregulates intracellular adhesion molecule 1 (ICAM-1) and chemokine (C-C motif) ligand 2 (CCL2)

¹Division of Infectious Diseases and HIV Medicine, Department of Medicine, Case Western Reserve University, Cleveland, OH

²Division of Molecular Endocrinology, Department of Medicine, Case Western Reserve University, Cleveland, OH

³Department of Ophthalmology, University of Wisconsin-Madison, Madison, WI

⁴Department of Ophthalmology and Visual Sciences, Case Western Reserve University, Cleveland, OH

⁵Louis Stokes Cleveland Veterans Administration Medical Center, Research Service 151, Cleveland, OH

⁶Department of Physiology and Biophysics, Case Western Reserve University, Cleveland, OH

⁷Department of Pathology, Case Western Reserve University, Cleveland, OH

Corresponding author: Carlos S. Subauste, carlos.subauste@case.edu.

Received 11 January 2016 and accepted 7 July 2016.

This article contains Supplementary Data online at <http://diabetes.diabetesjournals.org/lookup/suppl/doi:10.2337/db16-0051/-/DC1>.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

See accompanying article, p. 261.

(8,9). CD40 ligation in monocytes/macrophages/microglia upregulates TNF- α , IL-1 β , inducible nitric oxide synthase 2 (NOS2), and CCL2 (10–12). CD40 drives ICAM-1 and CCL2 upregulation, increases protein nitration and the number of leukocytes adherent to blood vessel walls (leukostasis) in the retina of diabetic mice, and is required for the development of capillary degeneration (8,9).

CD40 in hematopoietic cells has been considered central to the development of inflammatory diseases. Although studies using bone marrow chimeras suggest that CD40 expressed in nonhematopoietic cells is also required for inflammation (13), it is not known whether expression of CD40 restricted to the nonhematopoietic compartment is sufficient for development of inflammatory disorders.

Using transgenic mice with expression of CD40 in Müller cells, we report that after induction of diabetes, CD40 expression in these nonhematopoietic cells was sufficient for inflammatory molecule upregulation and development of capillary degeneration. TNF- α was upregulated in macrophages/microglia rather than in Müller cells. CD40 ligation in Müller cells induced macrophages to secrete TNF- α and IL-1 β via an ATP-P2X₇ receptor pathway. Pharmacologic or genetic inhibition of the P2X₇ receptor in diabetic mice impaired not only TNF- α and IL-1 β upregulation but also upregulation of ICAM-1 and NOS2, molecules reported to be driven by TNF- α and/or IL-1 β . Thus, CD40 in Müller cells orchestrates inflammatory responses in macrophages/microglia and promotes the development of experimental diabetic retinopathy.

RESEARCH DESIGN AND METHODS

Transgenic Mice

Mouse CD40 construct was inserted into the *Eco* RI and *Bam* HI sites of the pTet^{OS} plasmid (14). After sequence verification, the transgene was excised by *Sal* I digestion (14) and microinjected into mouse oocytes (B6). Founder Tet^{OS}-CD40 mice were identified by PCR using the following primers: Tet^{OS}CD40 forward: 5'-GCAACGTGCTGGTTATTGTG-3', reverse: 5'-CCGGGACTTTAAACCACAGA-3'. The driver line consisted of transgenic mice that express tetracycline (Tet)-repressible transactivator (tTA) under the control of the glial fibrillary acidic protein (GFAP) promoter *gfa2* consisting of the 2.2 kb of 5'-flanking DNA of human GFAP (15). Homozygous Tet^{OS}-CD40 (responder) and heterozygous GFAP-tTA (driver) transgenic mice (15) (both B6) were backcrossed onto a CD40^{-/-} (B6) background. To confirm that transgenic mice were CD40^{-/-}, animals were genotyped using primers that detect wild-type CD40 and mutant CD40 (neomycin cassette inserted into exon 3 resulting in lack of functional CD40) (16). Both lines of mice were bred and offspring identified by PCR analysis of genomic DNA. PCR primers for CD40 and tTA were obtained from The Jackson Laboratory. Littermates that inherited only one transgene (single transgenic and nonexpressing) served as controls (Trg-Ctr) for double transgenic animals (Trg-CD40; expressing GFAP promoter-specific CD40 expression).

Induction of Diabetes

Mice were made diabetic using streptozotocin (STZ). Fasted mice (20–25 g body weight) received five daily intraperitoneal injections of STZ (55 mg/kg; MP Biomedicals). Development of diabetes (blood glucose >250 mg/mL) was assessed beginning 1 week after the first injection of STZ. Glycated hemoglobin was measured at 2 months (VARIANT Classic; Bio-Rad). Mice were weighed weekly and, if needed, received insulin to prevent weight loss while maintaining chronic hyperglycemia (target range 350–500 mg/mL). The dose given (0–0.2 units of NPH insulin subcutaneously, 0–3 times per week) was determined individually for each animal. The insulin requirement was similar for all groups of diabetic mice. Studies adhered to the institutional guidelines for humane treatment of animals, *Principles of Laboratory Animal Care* (National Institutes of Health) and to the *Statement for the Use of Animals in Ophthalmic and Visual Research* (Association for Research in Vision and Ophthalmology). Studies were approved by the Case Western Reserve University Institutional Animal Care and Use Committee.

Leukostasis

The number of leukocytes adherent to the retinal microvasculature was determined at 2 months of diabetes. After perfusion with PBS, fluorescein-coupled concanavalin A lectin (20% g/mL; Vector Laboratories) was infused (17). Retinal flat mounts were analyzed using fluorescence microscopy, and brightly fluorescent leukocytes were counted.

Vascular Histopathology

Retinal vasculature was isolated and stained with periodic acid Schiff (8). Eight areas in the mid retina were examined blindly under original magnification $\times 400$. Degenerate capillaries were defined as capillary-sized tubes of material positive for periodic acid Schiff without nuclei along the capillary. Samples were processed blindly.

Immunohistochemistry

Paraffin-embedded eyes fixed with formalin-free zinc fixative (24 h; BD Biosciences) were treated with proteinase K or citrate buffer (8–12 sections per mouse) (8). Antibodies are listed in the Supplementary Data. Retinas were analyzed using a Leica DMI 6000B epifluorescence microscope.

Real-time Quantitative PCR

Real-time PCR was performed as described using primers for ICAM-1 (8), CCL2 (9), TNF- α (18), IL-1 β (19), NOS2 (20), P2X₇ receptor (21), and 18S rRNA (8).

Flow Cytometry

Retinal cell suspensions were obtained and permeabilized as described (22). Antibodies are listed in the Supplementary Data. CD40 expression is expressed as the corrected mean fluorescence intensity (22).

Cells

Primary human Müller cells obtained as described (8) and the human Müller cell line MIO M1 (gift from Dr. Gloria Limb, University College London, London, U.K.; both >95% vimentin⁺, cellular retinaldehyde binding protein [CRALBP]⁺ and

GFAP⁺) were transduced with a retroviral vector that encodes human CD40 or an empty vector (9). The Müller cell lines mRAST and rMC1 (mouse and rat, respectively) were transduced with a retroviral vector that encodes hmCD40. Müller cells were incubated with multimeric human CD154 to induce CD40 stimulation (obtained from Dr. Richard Kornbluth, Multimeric Biotherapeutics Inc., La Jolla, CA) or with a non-functional CD154 mutant (T147N) as the control (13). Cells were also incubated with recombinant FLAG-tagged CD154 (CD40L) plus enhancer (anti-FLAG antibody) or enhancer alone (Enzo Life Sciences). In certain experiments, Müller cells were incubated with BAPTA-AM (1,2-Bis[2-aminophenoxy] ethane-N,N,N',N'-tetraacetic acid tetrakis[acetoxymethyl ester]; 25 $\mu\text{mol/L}$; Sigma Aldrich), the phospholipase C inhibitor U73122, or the inactive analog U73343 (1 $\mu\text{mol/L}$; Tocris Bioscience). Human Müller cells were also cultured with the human monocytic cell line MonoMac6 pretreated with interferon- γ (100 IU for 48 h), with or without the P2X₇ receptor inhibitor A-438079 (10 $\mu\text{mol/L}$; Tocris Bioscience). Mouse Müller cells were incubated with bone marrow-derived macrophages (BMM) from B6 or P2X₇^{-/-} mice plus sodium polyoxotungstate (10 $\mu\text{mol/L}$) to inhibit the ectonucleotidase CD39 expressed in BMM. MonoMac6 and BMM were pretreated with lipopolysaccharide (100 ng/mL) for 4 h in studies of IL-1 β secretion. MonoMac6 cells were transfected with human P2X₇ small interfering (si)RNA or control siRNA (Qiagen) using TransIT-X2 (Mirus).

Measurement of Extracellular ATP

The ecto-ATPase inhibitor β,γ -methylene-ATP (300 $\mu\text{mol/L}$; Sigma-Aldrich) was added 15 min before stimulation with CD154. Extracellular ATP was quantified using an ATP bioluminescence assay kit and an ATP standard curve (Sigma-Aldrich) (23).

Immunoblot

Membranes were probed with antibody to P2X₇ receptor, actin (Santa Cruz Biotechnology), total phospholipase C γ 1 (PLC γ 1; Cell Signaling), or phospho-Tyr783 PLC γ 1 (Cell Signaling).

ELISA

Human IL-1 β , human TNF- α , rat TNF- α , mouse IL-1 β , and TNF- α (all from eBioscience), human CCL2, mouse CRP (both R&D Systems), and CD154 (Boster Bio) were measured using ELISA.

Statistical Analysis

Results are expressed as the mean \pm SEM. Data were analyzed by two-tailed Student *t* test or ANOVA. Differences were considered statistically significant at $P < 0.05$.

RESULTS

CD40 Expression in Müller Cells

We examined the effect of diabetes on serum CD154 (CD40 ligand) levels. Serum CD154 concentrations were significantly increased in mice made diabetic using STZ, whereas CRP levels remained unchanged, and TNF- α was undetectable (<3.9 pg/mL) (Supplementary Fig. 1). Diabetic mice are reported to upregulate CD40 in retinal

Müller cells, and diabetic CD40^{-/-} mice are protected from experimental diabetic retinopathy (8). To examine the role of CD40 expressed in Müller cells in experimental diabetic retinopathy, we used a binary transgenic mouse system to obtain CD40^{-/-} mice where CD40 expression is rescued in retinal Müller cells. The driver line consisted of mice expressing the tTA under the control of the human GFAP promoter *gfa2* (GFAP-tTA mice). These mice (B6 background) were backcrossed with CD40^{-/-} mice (also B6 background). The responder line consisted of homozygous CD40^{-/-} mice containing mouse CD40 cloned downstream of the Tet^{OS} promoter (Supplementary Fig. 2A). After mating GFAP-tTA mice with Tet^{OS} CD40 animals, double-transgenic offspring (Trg-CD40) are predicted to exhibit rescued CD40 expression in retinal macroglia. In contrast, no such rescue should occur in single-transgenic mice carrying the GFAP-tTA or Tet^{OS} CD40. PCR analysis of genomic DNA identified Trg-Ctr or Trg-CD40 mice (Supplementary Fig. 2B). Trg-CD40 mice expressed CD40 in vimentin⁺ cells (Müller cells), whereas Trg-Ctr animals lacked CD40 expression (Supplementary Fig. 2C). Flow cytometry analysis of vimentin⁺ cells confirmed that rescue of CD40 occurred in Trg-CD40 but not in Trg-Ctr mice and that levels of CD40 expression in vimentin⁺ cells were similar in B6 and Trg-CD40 mice (Supplementary Fig. 2D). Staining with an antibody against CRALBP, a specific Müller cell marker, confirmed that CD40⁺ cells from Trg-CD40 mice were Müller cells (Supplementary Figs. 2E and 3A). The pattern of GFAP expression was the same in B6 and transgenic mice and was detected in astrocytic processes present in the ganglion cell layer (Supplementary Fig. 2F). These results are in agreement with evidence that the GFAP promoter drives gene expression in Müller cells independently of endogenous GFAP gene activity (24). Moreover, GFAP⁺ astrocytes lacked detectable CD40 (Supplementary Fig. 2G) because expression driven by the GFAP transgene is lost in astrocytes from adult mice (24). Selective localization of CD40 in Trg-CD40 mice was further confirmed using immunohistochemistry and flow cytometry by the lack of detectable CD40 expression in the other retinal cells that express CD40 in wild-type mice: endothelial cells, microglia/macrophages, and ganglion cells, as well as lack of CD40 expression in leukocytes (Supplementary Figs. 2G and 3B and C). Thus, the transgenic system results in mice with CD40 rescue in Müller cells rather than astrocytes.

Müller Cell CD40 Promotes ICAM-1 Upregulation and Vascular Changes of Experimental Diabetic Retinopathy

Male B6, CD40^{-/-}, Trg-Ctr, and Trg-CD40 mice were rendered diabetic using STZ. Blood glucose concentrations, HbA_{1c} levels, and body weights of diabetic mice were similar ($P > 0.5$) (Supplementary Table 1). ICAM-1 expression is elevated in the diabetic retina (17) and promotes leukostasis (25,26). Compared with nondiabetic mice, ICAM-1 mRNA levels were increased in diabetic Trg-CD40 and B6 mice but not in diabetic Trg-Ctr or

CD40^{-/-} mice (Fig. 1A). Diabetic B6 and Trg-CD40 mice showed increased ICAM-1 expression in retinal capillaries compared with nondiabetic controls (Fig. 1B). In contrast, diabetic CD40^{-/-} and Trg-Ctr mice did not exhibit increased ICAM-1 expression (Fig. 1B).

Next, we determined whether diabetic Trg-CD40 mice develop increased retinal leukostasis. Diabetic Trg-CD40 and B6 mice showed a significant increase in the numbers of adherent leukocytes (Fig. 1C). Degenerate capillaries are a central feature of diabetic retinopathy. Diabetic B6 mice exhibited the expected increase in degenerate capillaries compared with nondiabetic animals, whereas diabetic mice lacking CD40 were protected from capillary degeneration (Fig. 1D). Diabetic Trg-CD40 animals showed increased degeneration of retinal capillaries similar to that

seen in diabetic B6 mice (Fig. 1D). Thus, CD40 in Müller cells promotes ICAM-1 upregulation in the diabetic retina and vascular changes of early diabetic retinopathy.

CD40 Expressed in Müller Cells Promotes Upregulation of TNF- α , IL-1 β , NOS2, and CCL2 mRNA Levels in the Diabetic Retina

TNF- α , IL-1 β , NOS2, and likely CCL2 play a pathogenic role in diabetic retinopathy (3,5,6,27–30). We examined the effects of Müller cell CD40 on mRNA levels of these molecules. In contrast to diabetic Trg-Ctr mice, diabetic Trg-CD40 animals upregulated TNF- α , IL-1 β , NOS2, and CCL2 mRNA levels (Fig. 2). Thus, expression of CD40 on Müller cells drives upregulation of TNF- α , IL-1 β , NOS2, and CCL2 mRNA in the retina of diabetic mice.




Figure 1—CD40 expression in Müller cells from diabetic mice promotes upregulation of ICAM-1 in the retina and vascular changes of early diabetic retinopathy. **A**: At 2 months of diabetes, retinas from diabetic B6, CD40^{-/-}, Trg-Ctr, and Trg-CD40 mice (DM) as well as from nondiabetic (ND) control animals were collected and used for mRNA extraction. mRNA levels of ICAM-1 were assessed by real-time quantitative PCR using 18S rRNA as the internal control. One nondiabetic B6 mouse was given an arbitrary value of 1, and data are expressed as the fold-increase compared with this animal. The horizontal bars represent mean \pm SEM ($n = 8$ –13 animals per group). **B**: At 2 months of diabetes, retinal sections were incubated with anti-ICAM-1 monoclonal antibody and tomato lectin ($n = 6$ mice/group). GCL, ganglion cell layer; INL, inner nuclear layer; IPL, inner plexiform layer. Arrowheads show blood vessels with increased ICAM-1 expression. Scale bar, 10 μ m. **C**: At 2 months of diabetes, adherent leukocytes in the retinal vasculature of diabetic and nondiabetic control mice were quantified by labeling with concanavalin A. Representative image shows an adherent leukocyte (arrowhead) within the vasculature of a diabetic B6 mouse. **D**: At 8 months of diabetes, retinal digests were examined for the presence of degenerate capillaries. The horizontal bars represent the mean \pm SEM ($n = 6$ –10 animals per group). Representative image shows an acellular capillary (arrow) in the retinal digest of a diabetic B6 mouse. ** $P < 0.01$, *** $P < 0.001$ by ANOVA.




Figure 2—CD40 expression in Müller cells from diabetic mice (DM) restores upregulation of TNF- α , IL-1 β , NOS2, and CCL2 mRNA in the retina. At 2 months of diabetes, retinas from diabetic B6, CD40^{-/-}, Trg-Ctr, and Trg-CD40 mice and from nondiabetic (ND) control animals were collected and used for mRNA extraction. mRNA levels were assessed by real-time quantitative PCR using 18S rRNA as the internal control. One nondiabetic B6 mouse was given an arbitrary value of 1, and data are expressed as the fold-increase compared with this animal. The horizontal bars represent the mean \pm SEM ($n = 7$ –15 animals per group). * $P < 0.05$, ** $P < 0.01$ by ANOVA.

Müller Cells Do Not Secrete TNF- α or IL-1 β in Response to CD40 Ligation

We examined the effects of CD40 stimulation on TNF- α and IL-1 β production by Müller cells. CD40 expression in Müller cells is upregulated in diabetes (8). However, in agreement with other nonhematopoietic cells, primary Müller cells and Müller cell lines have very low levels of CD40 expression in vitro under basal conditions. To test whether CD40 ligation caused TNF- α and/or IL-1 β secretion, we induced CD40 expression by transducing Müller cells with a CD40-encoding retroviral vector, an approach well suited to study the effects of CD40 signaling (9,11). CD40⁺ primary human Müller cells or the CD40⁺ human Müller cell line MOI-M1 failed to secrete TNF- α in response to CD154 (CD40 ligand) (Supplementary Fig. 4A). Although human Müller cells also failed to secrete IL-1 β in response to CD154, a marked upregulation in CCL2 was detected (Supplementary Fig. 4A). We tested whether similar findings apply to Müller cells from rodents. The rat Müller cell line rMC-1, which expresses a chimera of the extracellular domain of human CD40, and the intracytoplasmic domain mouse CD40 (hmCD40) was incubated with human CD154. These cells upregulated ICAM-1 in response to human CD154, but TNF- α secretion was not detected (Supplementary Fig. 4B). Mouse Müller cells that express hmCD40 also failed to secrete TNF- α and IL-1 β (Fig. 4I). Thus, CD40 stimulation of Müller cells does not increase TNF- α and IL-1 β secretion.

Retinal Microglia/Macrophages but Not Müller Cells From Diabetic Trg-CD40 Mice Express TNF- α

We examined CCL2 and TNF- α expression by immunohistochemistry. CCL2 was detected in the end feet and stalks of Müller cells from diabetic Trg-CD40 mice but not in diabetic Trg-Ctr mice or nondiabetic animals (Fig. 3A and B and Supplementary Fig. 5A). In contrast, no TNF- α was detected in Müller cells from diabetic Trg-CD40 mice or mice from any other group (Fig. 3C and Supplementary Fig. 5B). Interestingly, TNF- α was observed in microglia/macrophages (Iba-1⁺ cells) from diabetic Trg-CD40 mice (Fig. 3D and E). Analysis of multiple retinal sections revealed that 15% of microglia/macrophages in diabetic Trg-CD40 mice were TNF- α ⁺ (Fig. 3D). Thus, the presence of CD40 in Müller cells from diabetic mice drives CCL2 upregulation in Müller cells and upregulation of TNF- α in microglia/macrophages.

CD40-Activated Müller Cells Secrete ATP and Cause P2X₇ Receptor-Dependent Cytokine Production in Monocytes/Macrophages

We tested whether CD40-activated Müller cells induce IL-1 β and TNF- α production in bystander monocytes/macrophages. We used monocytic cells that lack CD40 (MonoMac6) to avoid the effects of direct CD40 ligation on these cells. Similar to Müller cells, monocytic cells failed to secrete IL-1 β and TNF- α in response to CD154 (Fig. 4A). In contrast, the addition of CD154 to coculture of CD40⁺ human Müller cells with monocytic cells enhanced IL-1 β and TNF- α production (Fig. 4A). Cytokine




Figure 3—Diabetic transgenic mice that express CD40 in retinal Müller cells upregulate CCL2 in Müller cells and TNF- α in microglia/macrophages. **A** and **B**: Retinal sections from diabetic Trg-Ctr and Trg-CD40 mice at 2 months of diabetes and from nondiabetic controls were incubated with anti-CCL2 plus anti-CRALBP antibody. CCL2 expression at the level of Müller cell end feet (ganglion cell layer) (**A**) and at the level of Müller cells stalks (outer plexiform layer) (**B**). **C**: Sections were incubated with anti-TNF- α plus anti-CRALBP antibody. Müller cell end feet and stalk areas are shown. **D** and **E**: Retinal sections were incubated with anti-TNF- α monoclonal antibody plus anti-Iba-1 antibody. **D**: Sections were analyzed to determine the percentages of Iba-1⁺ cells in diabetic mice (DM) and nondiabetic (ND) mice that were stained with anti-TNF- α antibody. *** $P < 0.001$. **E**: Images represent microglia/macrophages present in the inner plexiform layer ($n = 4$ mice/group). Scale bars, 10 μ m.

production was driven by CD40 because IL-1 β and TNF- α secretion were not significantly increased in cultures that contained Müller cells that were largely CD40⁻ (Fig. 4A).

Secretion of ATP by astrocytes has been proposed to cause purinergic receptor-driven IL-1 β production by microglia (31). To determine whether purinergic signaling mediates IL-1 β and TNF- α production during the Müller cells/monocytic cells coculture, we tested whether CD40 ligation in Müller cells increases secretion of extracellular ATP. CD40⁺ human Müller cells incubated with CD154 exhibited increased ATP release that was noted after 15 min of stimulation (Fig. 4B and C). Similar results were observed in mouse Müller cells (Fig. 4D).

Extracellular ATP binds purinergic receptors, among which P2X₇ is central to IL-1 β secretion (32). The addition of a specific P2X₇ receptor inhibitor (A-438079) to CD40⁺ human Müller cells and monocytic cells ablated the enhanced IL-1 β production triggered by CD154 (Fig. 4E).

It appeared unlikely that the P2X₇ receptor inhibitor acted on Müller cells because they did not exhibit detectable functional P2X₇ receptor (Supplementary Fig. 6A). To confirm that monocytes but not Müller cells were responsible for cytokine production, cells were incubated with a purinergic receptor ligand. 3'-O-(4-benzoyl)benzoyl (Bz)-ATP (100–300 μ mol/L) enhanced cytokine production by monocytic cells but not by Müller cells (Supplementary Fig. 6B). In contrast to IL-1 β , P2X₇ receptor inhibitor did not decrease TNF- α production triggered by addition of CD154 to CD40⁺ human Müller cells cultured with monocytic cells (Fig. 4F). Similarly, knockdown of the P2X₇ receptor in monocytic cells impaired IL-1 β production but not TNF- α when these cells were incubated with CD40⁺ Müller cells plus CD154 (Fig. 4G and H). Of relevance, although P2X₇ receptor is central to IL-1 β secretion, other P2 receptors in monocytic cell lines induce TNF- α production (33). Next, we examined the role of P2X₇ receptor in cytokine production by primary mouse macrophages. Whereas mouse hmCD40⁺ Müller cells or mouse BMM failed to secrete IL-1 β or TNF- α when incubated with human CD154, incubation of hmCD40⁺ Müller cells plus BMM resulted in increased IL-1 β and TNF- α production after human CD154 stimulation (Fig. 4I). No significant increase in cytokine production was noted when BMM from P2X₇^{-/-} mice were used (Fig. 4J). Thus, CD40-activated Müller cells secrete extracellular ATP and induce P2X₇ receptor-mediated cytokine production in monocytes/macrophages.

CD40 Appears to Induce ATP Release in Müller Cells Through PLC γ 1

ATP can be released after an increase in intracytoplasmic Ca²⁺ (34), and CD40 elevates intracytoplasmic Ca²⁺ levels (35,36). We examined the effects of BAPTA-AM (chelator of intracellular Ca²⁺) on ATP release. Incubation with BAPTA-AM impaired CD40-mediated ATP release in Müller cells (Fig. 5A). Next, we examined whether CD40 ligation in Müller cells causes phosphorylation of PLC γ 1, a signaling molecule that increases intracytoplasmic Ca²⁺ (37). CD40 ligation caused a rapid PLC γ 1 phosphorylation at Tyr783, a marker of PLC γ 1 activation (Fig. 5B). ATP release was impaired when Müller cells were treated with the PLC inhibitor U73122 but not when incubated with the inactive analog U73343 (Fig. 5C). Thus, CD40 ligation phosphorylates PLC γ 1 and appears to cause PLC γ 1-dependent ATP release in Müller cells.

Deficiency of P2X₇ Receptor or Administration of a P2X₇ Receptor Inhibitor Impairs Upregulation of TNF- α , IL-1 β , ICAM-1, and NOS2 in the Retina of Diabetic Mice

We examined P2X₇ receptor expression to explore its in vivo role in the diabetic retina. P2X₇ receptor mRNA levels were enhanced in the retinas of diabetic Trg-CD40 mice (Fig. 6A). Immunohistochemistry indicated that microglia/macrophages from diabetic Trg-CD40 mice exhibited increased P2X₇ receptor expression (Fig. 6B




Figure 4—CD40-activated Müller cells secrete ATP and induce P2X₇ receptor-dependent production of cytokines by monocytic cells and macrophages. **A:** Control human Müller cell (transduced with empty retroviral vector MIEG3) and CD40⁺ human Müller cells (transduced with CD40-encoding retroviral vector MIEG3-CD40) were incubated with CD40⁻ human monocytic cell lines (MonoMac6) with or without CD154. Cytokines were measured in supernatants by ELISA at predetermined optimal time points (24 h for IL-1β and 4 h for TNF-α). **B:** Human Müller cell line was incubated with or without CD154. Concentrations of extracellular ATP at time 0 and at 15 min of incubation are shown. **C:** CD40⁺ Müller cells were incubated with CD154 and concentrations of extracellular ATP were measured at different times. **D:** Mouse Müller cells transduced with the hmCD40-encoding retroviral vector were incubated with or without human CD154. CD40⁺ human Müller cells were incubated with MonoMac6 cells with or without CD154 in the presence or absence of the P2X₇ receptor inhibitor A-438079, and IL-1β (**E**) and TNF-α (**F**) were measured in supernatants by ELISA. CD40⁺ human Müller cells were incubated with MonoMac6 cells transfected with control or P2X₇ receptor siRNA, and IL-1β (**G**) and TNF-α (**H**) were measured in supernatants by ELISA. **I:** Mouse Müller cells that express hmCD40 were incubated with BMM from B6 or P2X₇^{-/-} mice, and IL-1β (**I**) and TNF-α (**J**) were measured by ELISA. Results are presented as mean ± SEM (*n* = 3). **P* < 0.05, ***P* < 0.01, ****P* < 0.001 by Student *t* test.

and C). Next, we examined the role of P2X₇ receptor on TNF-α and IL-1β expression in diabetic mice and also examined ICAM-1 and NOS2 because retinal ICAM-1

upregulation in diabetes is TNF-α dependent (38) and TNF-α and IL-1β promote NOS2 expression (39). Compared with B6 mice, diabetic P2X₇^{-/-} mice exhibited




Figure 5—CD40 ligation in Müller cells causes Tyr783 phosphorylation (p) of PLC γ 1 and PLC-dependent secretion of ATP. **A:** CD40⁺ Müller cells were incubated with or without BAPTA-AM, followed by stimulation with CD154 and measurement of extracellular ATP. Concentrations of extracellular ATP at time 0 and at 15 min of incubation are shown. **B:** CD40⁺ human Müller cells were incubated with CD154. Expression of p-Tyr783 PLC γ 1 and total PLC γ 1 were assessed by immunoblot. The bars represent quantification of relative p-Tyr783 PLC γ 1 from three different experiments. **C:** CD40⁺ Müller cells were incubated with U73122 or U73343, followed by stimulation with CD154 and measurement of extracellular ATP. Concentrations of extracellular ATP at time 0 and at 15 min of incubation are shown. Results are presented as mean \pm SEM ($n = 3$). ** $P < 0.01$, *** $P < 0.001$ by Student t test.

impaired upregulation of TNF- α , IL-1 β , ICAM-1, and NOS2 (Fig. 6D). We treated diabetic Trg-CD40 mice with the P2X₇ receptor inhibitor Brilliant Blue G (BBG). BBG did not affect blood glucose or HbA_{1c} levels (data not shown) but did diminish diabetes-induced TNF- α , IL-1 β , ICAM-1, and NOS2 upregulation (Fig. 6E). Together, the presence of CD40 in Müller cells from diabetic mice drives upregulation of retinal TNF- α , IL-1 β , ICAM-1, and NOS2 through a P2X₇ receptor-dependent mechanism.

DISCUSSION

We report that expression of CD40 in Müller cells is sufficient for upregulation of inflammatory molecules in the diabetic retina and development of early diabetic retinopathy. In diabetes, CD40 on Müller cells drives upregulation of CCL2 in these cells; however, CD40 does not increase TNF- α and IL-1 β secretion by Müller cells. CD40 stimulation in Müller cells causes release of extracellular ATP, an effect that appears dependent on PLC γ 1. In turn, ATP released by CD40-activated Müller cells induces P2X₇ receptor-mediated secretion of TNF- α and

IL-1 β by macrophages. These effects are relevant in vivo because TNF- α is upregulated in microglia/macrophages from diabetic mice that express CD40 in Müller cells and mice treated with BBG are protected from diabetes-induced upregulation of TNF- α and IL-1 β and also upregulation of ICAM-1 and NOS2. The role of P2X₇ is not only applicable to transgenic mice because diabetic P2X₇^{-/-} mice also exhibit diminished inflammatory molecule upregulation. These findings support a model of amplification of inflammation whereby CD40 engagement in Müller cells triggers an inflammatory response not only in these cells but also in bystander myeloid cells in a manner dependent on the ATP-P2X₇ receptor pathway.

Extracellular ATP acts not only as a neurotransmitter but also as a messenger that triggers cytokine production by macrophages/microglia. We uncovered that CD40 induced the release of extracellular ATP. CD40 stimulation caused rapid Tyr783 phosphorylation of PLC γ 1 in Müller cells, and PLC inhibition prevented release of extracellular ATP after CD40 ligation. Indeed, PLC mediates ATP release (40). Work presented here strongly supports




Figure 6—Diabetic transgenic mice that express CD40 in retinal Müller cells upregulate P2X₇ receptor in microglia/macrophages, and administration of the P2X₇ receptor inhibitor BBG impairs upregulation of retinal mRNA levels of TNF- α and IL-1 β . *A*: At 2 months of diabetes, retinas from diabetic (DM) and nondiabetic (ND) Trg-Ctr and Trg-CD40 mice were collected and used for mRNA extraction. mRNA levels of P2X₇ receptor were assessed by real-time quantitative PCR using 18S rRNA as the internal control. One nondiabetic Trg-Ctr mouse was given an arbitrary value of 1, and data are expressed as the fold-increase compared with this animal. The horizontal bars represent the mean \pm SEM ($n = 7$ –11 animals per group). *B* and *C*: Sections from diabetic and nondiabetic Trg-CD40 mice were incubated with anti-P2X₇ receptor plus anti-Iba-1 antibodies. *B*: Sections were analyzed to determine the percentages of Iba-1⁺ cells that stained brightly with anti-P2X₇ receptor antibody. *C*: Images represent microglia/macrophages present in the inner plexiform layer (IPL) ($n = 4$ mice/group). Scale bar, 10 μ m. *D*: At 2 months of diabetes, retinas from diabetic and nondiabetic B6 and P2X₇^{-/-} mice were collected and used to measure TNF- α , IL-1 β , ICAM-1, and NOS2 mRNA levels. One nondiabetic B6 mouse was given an arbitrary value of 1, and data are expressed as the fold-increase compared with this animal ($n = 7$ –12 animals per group). *E*: Diabetic Trg-CD40 mice were treated with BBG or vehicle daily for 28 days beginning at 1 month of diabetes. Retinas from these animals and from diabetic and nondiabetic Trg-Ctr were collected and used to measure TNF- α , IL-1 β , ICAM-1, and NOS2 mRNA levels ($n = 6$ –11 mice per group). ** $P < 0.01$, *** $P < 0.001$ by ANOVA.

that purinergic signaling promotes proinflammatory cytokine expression in microglia/macrophages because 1) administration of the P2X₇ inhibitor BBG impaired upregulation of TNF- α and IL-1 β in diabetic mice that express CD40 restricted to Müller cells, and 2) diabetic P2X₇^{-/-} mice were protected from upregulation of these cytokines. These findings identified a mechanism that enables CD40 expressed in nonhematopoietic cells to induce macrophages/microglia to produce TNF- α and IL-1 β , cytokines pivotal in inflammation. This mechanism would circumvent the inability or poor capacity of CD40 to trigger secretion of these cytokines in nonhematopoietic cells.

The level of P2X₇ receptor expression is functionally relevant because increased receptor expression is sufficient to cause microglia activation even in the absence of pathologic stimuli (41). Thus, the importance of the P2X₇ receptor in diabetic retinopathy is supported not only by the studies performed in P2X₇^{-/-} mice and in mice treated with BBG but also by the upregulation of the P2X₇ receptor noted in retinal microglia/macrophages. Moreover, P2X₇ receptor upregulation also occurs in diabetic B6 mice (J.-A.C.P., C.S.S., unpublished data). P2X₇ receptor is key for IL-1 β and TNF- α secretion by microglia/macrophages (42–45). Both cytokines contribute to diabetes-induced degeneration of retinal capillaries (5,6). The observation that the concentration of Bz-ATP (100–300 μ mol/L) that induced IL-1 β and TNF- α production in monocytic cells appeared to be higher than the ATP concentration detected in ATP assays does not detract from the role of purinergic signaling in IL-1 β and TNF- α upregulation. The assay based on firefly luciferase underestimates ATP concentrations in the intercellular space. Assays based on targeting luciferase to the surface of intact cells revealed higher concentrations of ATP at the cell surface (100–200 μ mol/L) (46). In addition, P2X₇ receptor upregulation in retinal microglia/macrophages may increase their sensitivity to ATP (47).

Serum CD154 levels are elevated in diabetic mice, and soluble CD154 is biologically active (48). Thus, these results together with the prior demonstration of increased CD40 expression in retinal cells indicate that the CD40-CD154 pathway is likely activated in diabetes. Of relevance, plasma CD15 is increased in patients with diabetic retinopathy (49). Retinal CD154 levels are also likely increased because microthrombosis occurs in diabetic retinopathy and activated platelets express CD154 (50). Differences in soluble CD154 concentrations may contribute to the susceptibility to experimental diabetic retinopathy. However, we have not detected differences in serum CD154 concentrations between diabetic B6 and P2X₇^{-/-} mice, it is unlikely that CD40 expression in Müller cells affects circulating CD154, and CD40 upregulation is sufficient to markedly increase proinflammatory responses upon CD40 ligation. Although we did not detect CD40 expression in cells other than Müller glia, we cannot rule out that potential ectopic CD40 expression could contribute to the results observed. Finally, although these

studies revealed the importance of CD40 in Müller cells for the pathogenesis of experimental diabetic retinopathy, CD40 signaling at the level of microglia/macrophages and/or endothelial cells likely also participates in the development of this disease.

In summary, this study uncovered the pivotal role CD40 expressed in a nonhematopoietic compartment and the purinergic-mediated recruitment of proinflammatory responses in microglia/macrophages in the development of inflammatory responses. These findings may also be relevant to other diseases driven by CD40, including inflammatory bowel disease, atherosclerosis, and lupus nephritis. CD40 expressed in nonhematopoietic cells in intestine, blood vessels, and kidney may trigger the release of extracellular ATP and engagement of purinergic receptors present in infiltrating myeloid cells. Finally, these results may have therapeutic implications. Inhibition of CD40 can lead to a therapeutic approach against inflammatory disorders. Signaling pathways downstream of CD40 have different relative roles in inducing proinflammatory responses in hematopoietic and nonhematopoietic cells (11). Targeting a signaling pathway in nonhematopoietic cells may inhibit proinflammatory responses in these cells and in neighboring myeloid cells.

Acknowledgments. The authors thank Alejandro Sosa for performing Ca²⁺ measurement studies, Andrea Boyd Tressler for assistance in ATP release measurements, Scott Howell for image acquisition, Catherine Doller for tissue processing for histopathology (all from Case Western Reserve University, Cleveland, OH), and Richard Kornbluth (Multimeric Biotherapeutics Inc., La Jolla, CA), Gloria Limb (University College London, London, U.K.), and Vijay Sarthy (Northwestern University, Chicago, IL) for providing reagents.

Funding. This work was supported by National Institute of General Medical Sciences grant GM-36387 (G.R.D.), National Eye Institute grants EY-019250 (C.S.S.) and P30-EY-11373, and Juvenile Diabetes Foundation International grant 1-2009-204 (C.S.S.).

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. J.-A.C.P., Y.L.C., Y.M., J.T., T.S.K., G.R.D., and C.S.S. acquired and analyzed data. J.-A.C.P., Y.L.C., Y.M., J.T., and C.S.S. conducted experiments. J.-A.C.P. and C.S.S. designed experiments and wrote the manuscript. N.S. contributed with reagents. C.S.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Tang J, Kern TS. Inflammation in diabetic retinopathy. *Prog Retin Eye Res* 2011;30:343–358
2. Antonetti DA, Klein R, Gardner TW. Diabetic retinopathy. *N Engl J Med* 2012; 366:1227–1239
3. Krady JK, Basu A, Allen CM, et al. Minocycline reduces proinflammatory cytokine expression, microglial activation, and caspase-3 activation in a rodent model of diabetic retinopathy. *Diabetes* 2005;54:1559–1565
4. Yang LP, Sun HL, Wu LM, et al. Baicalein reduces inflammatory process in a rodent model of diabetic retinopathy. *Invest Ophthalmol Vis Sci* 2009;50: 2319–2327
5. Jousseaume AM, Doehmen S, Le ML, et al. TNF- α mediated apoptosis plays an important role in the development of early diabetic retinopathy and long-term histopathological alterations. *Mol Vis* 2009;15:1418–1428

6. Vincent JA, Mohr S. Inhibition of caspase-1/interleukin-1 β signaling prevents degeneration of retinal capillaries in diabetes and galactosemia. *Diabetes* 2007;56:224–230
7. Wang J, Xu X, Elliott MH, Zhu M, Le YZ. Müller cell-derived VEGF is essential for diabetes-induced retinal inflammation and vascular leakage. *Diabetes* 2010;59:2297–2305
8. Portillo JA, Greene JA, Okenka G, et al. CD40 promotes the development of early diabetic retinopathy in mice. *Diabetologia* 2014;57:2222–2231
9. Portillo JA, Schwartz I, Zarini S, et al. Proinflammatory responses induced by CD40 in retinal endothelial and Müller cells are inhibited by blocking CD40-Traf2,3 or CD40-Traf6 signaling. *Invest Ophthalmol Vis Sci* 2014;55:8590–8597
10. Kiener PA, Moran-Davis P, Rankin BM, Wahl AF, Aruffo A, Hollenbaugh D. Stimulation of CD40 with purified soluble gp39 induces proinflammatory responses in human monocytes. *J Immunol* 1995;155:4917–4925
11. Portillo JA, Greene JA, Schwartz I, Subauste MC, Subauste CS. Blockade of CD40-TRAF2,3 or CD40-TRAF6 is sufficient to impair pro-inflammatory responses in non-haematopoietic cells. *Immunology* 2015;144:21–33
12. Portillo JA, Muniz-Feliciano L, Subauste MC, Heinzel FP, Subauste CS. CD40 and tumor necrosis factor- α co-operate to up-regulate inducible nitric oxide synthase expression in macrophages. *Immunology* 2012;135:140–150
13. Portillo JA, Van Grol J, Zheng L, et al. CD40 mediates retinal inflammation and neurovascular degeneration. *J Immunol* 2008;181:8719–8726
14. Sarao R, Dumont DJ. Conditional transgene expression in endothelial cells. *Transgenic Res* 1998;7:421–427
15. Lin W, Kemper A, McCarthy KD, et al. Interferon- γ induced medulloblastoma in the developing cerebellum. *J Neurosci* 2004;24:10074–10083
16. Kawabe T, Naka T, Yoshida K, et al. The immune responses in CD40-deficient mice: impaired immunoglobulin class switching and germinal center formation. *Immunity* 1994;1:167–178
17. Jousseaume AM, Poulaki V, Le ML, et al. A central role for inflammation in the pathogenesis of diabetic retinopathy. *FASEB J* 2004;18:1450–1452
18. Johnson LL, Lanthier P, Hoffman J, Chen W. Vaccination protects B cell-deficient mice against an oral challenge with mildly virulent *Toxoplasma gondii*. *Vaccine* 2004;22:4054–4061
19. Overbergh L, Valckx D, Waer M, Mathieu C. Quantification of murine cytokine mRNAs using real time quantitative reverse transcriptase PCR. *Cytokine* 1999;11:305–312
20. Park EM, Cho S, Frys K, et al. Interaction between inducible nitric oxide synthase and poly(ADP-ribose) polymerase in focal ischemic brain injury. *Stroke* 2004;35:2896–2901
21. Milano PM, Douillet CD, Riesenman PJ, et al. Intestinal ischemia-reperfusion injury alters purinergic receptor expression in clinically relevant extraintestinal organs. *J Surg Res* 2008;145:272–278
22. Portillo JA, Okenka G, Kern TS, Subauste CS. Identification of primary retinal cells and ex vivo detection of proinflammatory molecules using flow cytometry. *Mol Vis* 2009;15:1383–1389
23. Blum AE, Joseph SM, Przybylski RJ, Dubyak GR. Rho-family GTPases modulate Ca(2+)-dependent ATP release from astrocytes. *Am J Physiol Cell Physiol* 2008;295:C231–C241
24. Kuzmanovic M, Dudley VJ, Sarthy VP. GFAP promoter drives Müller cell-specific expression in transgenic mice. *Invest Ophthalmol Vis Sci* 2003;44:3606–3613
25. McLeod DS, Lefer DJ, Merges C, Luty GA. Enhanced expression of intracellular adhesion molecule-1 and P-selectin in the diabetic human retina and choroid. *Am J Pathol* 1995;147:642–653
26. Miyamoto K, Khosrof S, Bursell SE, et al. Prevention of leukostasis and vascular leakage in streptozotocin-induced diabetic retinopathy via intercellular adhesion molecule-1 inhibition. *Proc Natl Acad Sci U S A* 1999;96:10836–10841
27. Mitamura Y, Takeuchi S, Matsuda A, Tagawa Y, Mizue Y, Nishihira J. Monocyte chemoattractant protein-1 in the vitreous of patients with proliferative diabetic retinopathy. *Ophthalmologica* 2001;215:415–418
28. Zhang SX, Wang JJ, Gao G, Shao C, Mott R, Ma JX. Pigment epithelium-derived factor (PEDF) is an endogenous anti-inflammatory factor. *FASEB J* 2006;20:323–325
29. Zheng L, Du Y, Miller C, et al. Critical role of inducible nitric oxide synthase in degeneration of retinal capillaries in mice with streptozotocin-induced diabetes. *Diabetologia* 2007;50:1987–1996
30. Huang H, Gandhi JK, Zhong X, et al. TNF α is required for late BRB breakdown in diabetic retinopathy, and its inhibition prevents leukostasis and protects vessels and neurons from apoptosis. *Invest Ophthalmol Vis Sci* 2011;52:1336–1344
31. Monif M, Burnstock G, Williams DA. Microglia: proliferation and activation driven by the P2X7 receptor. *Int J Biochem Cell Biol* 2010;42:1753–1756
32. Solle M, Labasi J, Perregeux DG, et al. Altered cytokine production in mice lacking P2X(7) receptors. *J Biol Chem* 2001;276:125–132
33. Domercq M, Brambilla L, Pilati E, Marchaland J, Volterra A, Bezzi P. P2Y1 receptor-evoked glutamate exocytosis from astrocytes: control by tumor necrosis factor-alpha and prostaglandins. *J Biol Chem* 2006;281:30684–30696
34. Bal-Price A, Moneer Z, Brown GC. Nitric oxide induces rapid, calcium-dependent release of vesicular glutamate and ATP from cultured rat astrocytes. *Glia* 2002;40:312–323
35. Klaus GG, Choi MS, Holman M. Properties of mouse CD40. Ligation of CD40 activates B cells via a Ca(++)-dependent, FK506-sensitive pathway. *Eur J Immunol* 1994;24:3229–3232
36. Lazaar AL, Amrani Y, Hsu J, et al. CD40-mediated signal transduction in human airway smooth muscle. *J Immunol* 1998;161:3120–3127
37. Yang YR, Follo MY, Cocco L, Suh PG. The physiological roles of primary phospholipase C. *Adv Biol Regul* 2013;53:232–241
38. Jousseaume AM, Poulaki V, Mitsiades N, et al. Nonsteroidal anti-inflammatory drugs prevent early diabetic retinopathy via TNF- α suppression. *FASEB J* 2002;16:438–440
39. Liu J, Zhao ML, Brosnan CF, Lee SC. Expression of type II nitric oxide synthase in primary human astrocytes and microglia: role of IL-1beta and IL-1 receptor antagonist. *J Immunol* 1996;157:3569–3576
40. Liu GJ, Werry EL, Bennett MR. Secretion of ATP from Schwann cells in response to uridine triphosphate. *Eur J Neurosci* 2005;21:151–160
41. Monif M, Reid CA, Powell KL, Smart ML, Williams DA. The P2X7 receptor drives microglial activation and proliferation: a trophic role for P2X7R pore. *J Neurosci* 2009;29:3781–3791
42. Ferrari D, Villalba M, Chiozzi P, Falzoni S, Ricciardi-Castagnoli P, Di Virgilio F. Mouse microglial cells express a plasma membrane pore gated by extracellular ATP. *J Immunol* 1996;156:1531–1539
43. Ferrari D, Chiozzi P, Falzoni S, et al. Extracellular ATP triggers IL-1 beta release by activating the purinergic P2Z receptor of human macrophages. *J Immunol* 1997;159:1451–1458
44. Hide I, Tanaka M, Inoue A, et al. Extracellular ATP triggers tumor necrosis factor-alpha release from rat microglia. *J Neurochem* 2000;75:965–972
45. Suzuki T, Hide I, Ido K, Kohsaka S, Inoue K, Nakata Y. Production and release of neuroprotective tumor necrosis factor by P2X7 receptor-activated microglia. *J Neurosci* 2004;24:1–7
46. Pellegatti P, Falzoni S, Pinton P, Rizzuto R, Di Virgilio F. A novel recombinant plasma membrane-targeted luciferase reveals a new pathway for ATP secretion. *Mol Biol Cell* 2005;16:3659–3665
47. Humphreys BD, Dubyak GR. Modulation of P2X7 nucleotide receptor expression by pro- and anti-inflammatory stimuli in THP-1 monocytes. *J Leukoc Biol* 1998;64:265–273
48. Xu H, Zhang X, Mannon RB, Kirk AD. Platelet-derived or soluble CD154 induces vascularized allograft rejection independent of cell-bound CD154. *J Clin Invest* 2006;116:769–774
49. Yngen M, Ostenson CG, Hu H, Li N, Hjemdahl P, Wallén NH. Enhanced P-selectin expression and increased soluble CD40 ligand in patients with type 1 diabetes mellitus and microangiopathy: evidence for platelet hyperactivity and chronic inflammation. *Diabetologia* 2004;47:537–540
50. Boeri D, Maiello M, Lorenzi M. Increased prevalence of microthromboses in retinal capillaries of diabetic individuals. *Diabetes* 2001;50:1432–1439