

RESEARCH ARTICLE

Deletion of pH Regulator *pac-3* Affects Cellulase and Xylanase Activity during Sugarcane Bagasse Degradation by *Neurospora crassa*

Amanda Cristina Campos Antoniêto¹, Wellington Ramos Pedersoli², Lílian dos Santos Castro¹, Rodrigo da Silva Santos², Aline Helena da Silva Cruz², Karoline Maria Vieira Nogueira¹, Rafael Silva-Rocha³, Antonio Rossi², Roberto Nascimento Silva¹*

1 Department of Biochemistry and Immunology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil, **2** Department of Genetics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil, **3** Systems and Synthetic Biology Laboratory, Department of Cell and Molecular Biology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, Brazil

* rsilva@fmrp.usp.br



OPEN ACCESS

Citation: Campos Antoniêto AC, Ramos Pedersoli W, dos Santos Castro L, da Silva Santos R, Cruz AHdS, Nogueira KMV, et al. (2017) Deletion of pH Regulator *pac-3* Affects Cellulase and Xylanase Activity during Sugarcane Bagasse Degradation by *Neurospora crassa*. PLoS ONE 12(1): e0169796. doi:10.1371/journal.pone.0169796

Editor: Katherine A. Borkovich, University of California Riverside, UNITED STATES

Received: June 20, 2016

Accepted: December 21, 2016

Published: January 20, 2017

Copyright: © 2017 Campos Antoniêto et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was supported by grants from the Brazilian Funding Agencies: São Paulo Research Foundation (FAPESP), (Grant No 2014/09493-2, ACCA) (Grant No 2014/03847-7; AR, RSS, AHSC), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), (Grant No 304222/2013-5), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior

Abstract

Microorganisms play a vital role in bioethanol production whose usage as fuel energy is increasing worldwide. The filamentous fungus *Neurospora crassa* synthesizes and secretes the major enzymes involved in plant cell wall deconstruction. The production of cellulases and hemicellulases is known to be affected by the environmental pH; however, the regulatory mechanisms of this process are still poorly understood. In this study, we investigated the role of the pH regulator PAC-3 in *N. crassa* during their growth on sugarcane bagasse at different pH conditions. Our data indicate that secretion of cellulolytic enzymes is reduced in the mutant $\Delta pac-3$ at alkaline pH, whereas xylanases are positively regulated by PAC-3 in acidic (pH 5.0), neutral (pH 7.0), and alkaline (pH 10.0) medium. Gene expression profiles, evaluated by real-time qPCR, revealed that genes encoding cellulases and hemicellulases are also subject to PAC-3 control. Moreover, deletion of *pac-3* affects the expression of transcription factor-encoding genes. Together, the results suggest that the regulation of holocellulase genes by PAC-3 can occur as directly as in indirect manner. Our study helps improve the understanding of holocellulolytic performance in response to PAC-3 and should thereby contribute to the better use of *N. crassa* in the biotechnology industry.

Introduction

The filamentous fungus *Neurospora crassa* is an appealing microorganism for use by the bioethanol industry, because it degrades biomass releasing fermentable sugars [1]. Many isolates of this species were collected from sugarcane plantations [2–4] exhibiting the synthesis and secretion of holocellulolytic enzymes involved in plant cell wall degradation [5].

(CAPES), Fundação de Apoio ao Ensino, Pesquisa e Assistência (FAEPA).

Competing Interests: The authors have declared that no competing interests exist.

The *N. crassa* genome contains 171 genes encoding glycosylhydrolases, about 15.5% less than the cellulolytic model fungus *Trichoderma reesei*. However, the *N. crassa* genome possesses 35 genes encoding cellulolytic enzymes, almost twice that of *T. reesei* [6, 7]. Three types of enzymes are common to all cellulose-degrading fungi, including *N. crassa*: endoglucanases, exoglucanases, and β -glucosidases, which act synergistically to degrade the cellulolytic complex into glucose monomers [5].

The whole process of lignocellulosic biomass deconstruction is finely regulated at the transcriptional level. In *N. crassa*, the transcription factor XLR-1 (xylan degradation regulator 1) is necessary for growth on hemicellulose, whereas its orthologs, XlnR from *Aspergillus* sp. and XYR1 from *Trichoderma* sp., are the primary transcriptional regulators of genes encoding both cellulases and hemicellulases [8, 9]. Two other transcription factors, CLR-1 and CLR-2, also have an important role for the growth and induction of cellulases and some hemicellulases on cellulose; however, they are not required for growth on xylan [9, 10]. It is also known that the zinc-finger transcription factor CRE-1 is responsible for carbon catabolite repression (CCR) in *N. crassa* and that deletion of this regulator results in increased expression of cellulolytic genes and enhanced enzymatic activity during growth on cellulose [11].

In some species like *T. reesei*, *Aspergillus fumigatus*, and *N. crassa* [12–15], cellulase production is affected not only by available carbon source but also by the pH of the culture medium, indicating that pH is an important factor in the effective expression of these enzymes [12, 16]. The transcription factor responsible for the pH-signaling pathway is the protein PacC. In *Aspergillus nidulans*, this regulator mediates cell adaptation to pH by both activation of genes that are preferentially expressed in alkaline medium and repression of specific acid-associated genes [17]. At alkaline pH, PAC-3 is activated by PalB-catalyzed proteolysis, and active PAC-3 binds to the DNA of its target genes through the core consensus 5'-BGCCVAGV-3' in *N. crassa* [18].

Although a link has been demonstrated between PacC and lignocellulolytic enzyme production in species such as *A. nidulans*, *T. reesei*, and *Humicola grisea* var. *thermoideae* [12, 13, 19, 20], little is known about the role of PAC-3 in the expression of cellulases or hemicellulases in *N. crassa*. Here, we investigated the role of the transcription factor PAC-3 in the activity of holocellulolytic enzymes as well as in the regulation of cellulase and hemicellulase genes in *N. crassa*. We show that in alkaline environments, *pac-3* deletion results in decreased activity of endoglucanase and xylanase enzymes, as well as differential regulation of cellulase and hemicellulase gene expression. This study improves our understanding of the regulation of lignocellulosic biomass degradation in response to ambient pH in *N. crassa* for further application of metabolically engineered strains in the cellulosic production of ethanol.

Materials and Methods

Strains and growth conditions

Wild-type strain, St.L.74-OR23-1VA (FGSC No. 2489) and Δ *mus-52* (FGSC N° 9568) (control) of *Neurospora crassa* were purchased from the Fungal Genetics Stock Center (FGSC; University of Missouri, MO, USA, <http://www.fgsc.net>) [21]. A *N. crassa* Δ *pac-3* knockout strain (*pac-3*^{KO}) was generated following the knockout procedures described previously [22], based on the strain FGSC N° 9568 (mat a, *mus-52::hyg*) [23]. Strains were maintained on agar-based Vogel's minimal (VM) medium at pH 5.8 [24], containing 2% sucrose at 30°C. For gene expression assays, conidia from each strain (approximately 10⁷ cells per mL) were germinated in 100 mL of liquid VM medium, supplemented with 1% (w/v) sugarcane bagasse (i.e., the carbon source) at 30°C in an orbital shaker (200 rpm) for eight days at different pH conditions (pH 3, 5, 7, 8, and 10). After this culture period, the resulting mycelia were collected by filtration, frozen, and stored at -80°C until RNA isolation. Supernatants were used for secretome

and enzymatic activity analysis. Fungal growth was indirectly measured by total protein quantification using the Quick Start Bradford protein assay kit (Bio-Rad) with bovine serum albumin (BSA) as a standard. All experiments were performed in three biological replicates.

RNA extraction and quantitative real-time analysis (RT-qPCR)

Total RNA was isolated from mycelia with the TRIzol RNA kit (Invitrogen Life Technologies, CA, USA), per the manufacturer’s instructions. RNA concentration was determined by spectrophotometric OD 260/280, and RNA integrity was verified by gel electrophoresis in 1% agarose. For RT-qPCR analysis, 1 µg of RNA was treated with DNase I (Thermo Fisher, MA, USA) and reverse-transcribed to cDNA with the First Strand cDNA kit Maxima Synthesis, per manufacturer’s instructions. The cDNA was diluted 1/50-fold and used for RT-qPCR analysis in the Bio-Rad CFX96™ System, using SsoFast™ EvaGreen® Supermix (Bio-Rad, CA, USA) for signal detection. An actin-encoding gene (NCU04173) was used as an endogenous control. The following amplification reaction was used: 95°C for 10 minutes followed by 39 cycles of 95°C for 10 seconds and 60°C for 30 seconds, followed by a dissociation curve of 60°C to 95°C with an increment of 0.5°C for 10 seconds. Gene expression values were calculated according to the 2^{-ΔΔCT} method [25] and fold change expressed as Log₂ using the 74A strain grown on sugarcane bagasse as the reference sample. Primers utilized in the RT-qPCR experiments are described in Table 1. Data analysis was performed using GraphPad Prism v 5.0 software. All samples were analyzed in triplicate.

Enzymatic activities

Total cellulase activity (FPase) was determined by an enzymatic reaction employing Whatman filter paper no. 1, 30 µL of 100 mM citrate-phosphate buffer pH 5.0, and 30 µL of sample. The reaction was incubated at 50°C for 30 minutes. Next, 60 µL of dinitrosalicylic acid (DNS) was added to the reaction, which was then heated at 95°C for 5 minutes. Carboxymethylcellulase (CMCase) activity was determined following the protocol previously described [26] with some modifications. The reaction consisted of 30 µL of carboxymethylcellulose (CMC) prepared in 1% sodium acetate buffer (50 mM and pH 5.0) and 30 µL of sample. The reaction was

Table 1. qRT-PCR primers used in this study.

GENES	Sequence 5'→3' (Forward)	Sequence 5'→3' (Reverse)
<i>cbh-1</i>	ACCCCTCCACCAACG	GTGAACGCTGTAGAGACTTTGT
<i>gh7-1</i>	GCCCTACTCCAGCATTCAT	GTGAAGCGGTGGCG
<i>gh3-2</i>	GGTGGAGAGGATGGATGAT	CCATCTGTTTCGTTTCGTCC
<i>gh11-1</i>	GGCTCTTACTCTGTGAAC TGG	GTAGCCGTTGCCCGAG
<i>gh10-4</i>	GGGTGGGAGGAATGAAGA	ATCCTCGCCGCACACA
<i>gh43-5</i>	GAATGTTTATCAGGGCACG	GCTCATACCCAGTCCCG
<i>act</i>	ATGTCGACGTCAGGAAGGATCT	GAGCAGTGATCTCCTTCTGCAT
<i>clr-1</i>	GTTGTATTGTGCTTCTCGTCTTG	AACGCTGTTGGAGGTCATC
<i>clr-2</i>	ACGCAGTGTGGTTTCTCTAC	TCCAATACCGCCTCAATCTG
<i>cre-1</i>	GAGGATCACGATGACCACTATG	GAGTAGCAAGTGGAGTGTGATC
<i>xlr-1</i>	AGAGTACCACAGCAACCAAG	TTCGCTCATATCGCTCATGG

cbh-1 = Cellobiohydrolase 1 (exoglucanase) NCU07340; *gh7-1* = Endoglucanase 1 NCU05057; *gh3-2* = β-glucosidase 1 NCU08054; *gh11-1* = Endo-1,4-β-xylanase 1 NCU02855; *gh10-4* = Endo-1,4-β-xylanase 2 NCU07130; *gh43-5* = β-xylosidase NCU09652. *clr-1* = Cellulose degradation regulator-1 NCU07705; *clr-2* = Cellulose degradation regulator-2 NCU08042; *cre-1* = Carbon catabolite regulation NCU08807; *xlr-1* = Xylan degradation regulator-1 NCU06971; *act* = actin NCU04173

doi:10.1371/journal.pone.0169796.t001

incubated at 50°C for 30 minutes, followed by the addition of 60 μ L of DNS, and an additional heating step at 95°C for 5 minutes. Xylanase activity was determined by reacting 50 μ L of xylan substrate (xylan from beechwood [Sigma, MO, USA] at 50 mg/mL + 100 mM sodium acetate at pH 5.0) with 25 μ L of sample at 50°C for 30 minutes. After incubation, 75 μ L of DNS was added, and the reaction was heated at 95°C for 5 minutes. All enzymatic activities were performed in a 96-well microplate, and absorbance was read at 540 nm using the xMark™ Microplate Spectrophotometer (Bio-Rad, CA, USA). One enzyme unit was defined as the amount of enzyme capable of liberating 1 μ mol of reducing sugar per minute [27].

Western blotting

The proteins secreted by *N. crassa* were transferred to nitrocellulose (GE Healthcare, Buckinghamshire, UK). Rabbit secondary antibodies and anti-cellulase-*Trichoderma viride* antibodies (LS-C147688, LifeSpan Biosciences, WA, USA) were used as previously described [28]. Antibodies were visualized by enhanced chemiluminescence (Santa Cruz Biotechnology, TX, USA).

In silico analysis of putative PAC-3 binding sites in *N. crassa* promoters

The complete genome sequence of *N. crassa* OR74A [29] was downloaded from <https://www.broadinstitute.org/fungal-genome-initiative/neurospora-crassa-genome-project> and was used to obtain the 5' -upstream regions (1 kb) for each gene. The occurrence of the PAC-3 motifs 5' -BGCCVAGV-3' [30] were determined in the genes analyzed in this study, as well as for the 9730 genes in *N. crassa* genome, using an *ad hoc* Perl script. The motif frequency was calculated as the number of motif occurrences per kb.

Statistical analysis

All tests of statistical significance were performed using one-way ANOVA (nonparametric) followed by the Bonferroni test (comparing all pairs of columns), available in the Prism software v. 5.0. Significance was indicated in the form of probability values (*P < 0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001).

Results

pac-3 deletion affects cellulase and xylanase activities

Since the growth rate of the $\Delta pac-3$ mutant on solid medium is affected by pH [31], we investigated the final growth (total protein content) of 74A, $\Delta pac-3$, and $\Delta mus-52$ in the cultured medium at pH 3.0, 5.0, 7.0, 8.0, and 10.0. The $\Delta pac-3$ mutant grew better at acidic (3.0 and 5.0) rather than neutral (7) or alkaline (8.0 and 10.0) pH when compared with 74A and $\Delta mus-52$ strains (Fig 1A). We conclude that the different growth rates observed in the $\Delta pac-3$ mutant can affect enzyme production; thus, all enzymatic activities were normalized by total protein content and expressed as specific activity. We also observed changes in the medium pH after the *N. crassa* strains 74A, $\Delta pac-3$, and $\Delta mus-52$ were cultivated for eight days on sugarcane bagasse. However, only the $\Delta pac-3$ strain showed a significant difference at both pH 3 and 8, probably because of its poor growth under these culture conditions (Fig 1B).

To determine whether PAC-3 affects the secretion of holocellulolytic enzymes by the fungus, we analyzed the profile of total cellulolytic (FPase), endoglucanase (CMCase), and xylanolytic activities in strains 74A, $\Delta pac-3$, and $\Delta mus-52$ (Fig 2).

With respect to total cellulolytic activity, we observed that the $\Delta pac-3$ strain showed nearly a 4-fold decrease in activity at an acidic pH of 3.0 and a 2.5-fold decrease at pH 5.0 as compared to that in the 74A strain (Fig 2A). However, after raising the pH to 7.0, 8.0, or 10.0, we




Fig 1. Total protein content (A) and pH changes (B) after growth of *N. crassa* for eight days on sugarcane bagasse. Strains 74A, $\Delta mus-52$, and $\Delta pac-3$, were cultivated on sugarcane bagasse at pH 3.0, 5.0, 7.0, 8.0, and 10.0 (initial pH). The final growth and pH was measured on the eighth day. The error bar indicates the standard deviation. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

doi:10.1371/journal.pone.0169796.g001

observed an increase in specific FPase activity in the $\Delta pac-3$ strain (approximately 1-fold, 3-fold, and 4.5-fold, respectively) as compared to that in the parental strains, suggesting that the PAC-3 protein acts in neutral to alkaline pH by modulating the enzymatic activity and, consequently, the total specific cellulolytic activity (Fig 2A). As with FPase activity, the same




Fig 2. PAC-3 influences holocellulase activity in *N. crassa*. Total cellulolytic (FAPase) (A), endoglucanase (CMCase) (B), and xylanolytic activities (C) were assayed in the supernatant of the strains 74A, $\Delta mus-52$, and $\Delta pac-3$ after their cultivation for eight days on sugarcane bagasse. Values show the mean of three replicates. The error bar indicates the standard deviation. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

doi:10.1371/journal.pone.0169796.g002

pattern was observed for the specific activity of endoglucanase (CMCase). At acidic pH environments, endoglucanase showed about 3-fold (pH 3.0) and 2.5-fold (pH 5.0) less activity than the parental strains (Fig 2B). At neutral pH, CMCase activity was approximately 2-fold higher than that of the parental strains. On the other hand, at alkaline pH, the endoglucanase activity only showed a significant difference at pH 10 (approximately 3-fold higher than the parental strains) (Fig 2B).

When the xylanase activity profile was evaluated, a significant decrease was observed in the mutant strain $\Delta pac-3$ as compared to the 74A and $\Delta mus-52$ strains at pH 3.0 (6.5-fold), pH 5.0 (5.5-fold), pH 7.0 (2.7-fold), and pH 10.0 (2-fold) (Fig 2C). Except at pH 8.0, the protein PAC-3 appears to upregulate the expression of xylanase activity regardless of the initial culture pH value (Fig 2C).

Since cellobiohydrolase I (CBH-1) is the most highly produced extracellular protein during *N. crassa* growth on cellulose [32], we performed western blot analysis to evaluate the influence of PAC-3 on cellulase secretion. Western blot analysis showed a high level of cellulase secretion in a $pac-3^+ mus52^+$ background (strain 74A) irrespective of the initial pH but preferentially at neutral to alkaline pH (Fig 3). In a $pac-3^+ mus52^-$ background (strain $\Delta mus-52$), cellulase secretion preferentially occurred at neutral to alkaline pH. Unlike to observed for both strains, a decreased cellulase secretion was observed in the $\Delta pac-3$ strain at alkaline pH (pH 10.0). Thus,




Fig 3. Western blot analysis shows a reduced amount of cellulase (CBH-1) at both neutral and alkaline pH. The analysis was performed using supernatants collected from the *N. crassa* 74A, $\Delta mus-52$, and $\Delta pac-3$ strains after growth for eight days on sugarcane bagasse at pH 3.0, 5.0, 7.0, 8.0, and 10.0. Anti-cellulase antibody from *Trichoderma viride* were used to detect CBH-1.

doi:10.1371/journal.pone.0169796.g003

regardless of the initial culture pH, the protein PAC-3 affects the secretion of CBHI (Fig 3) as well as cellulase activity (Fig 2A).

PAC-3 affects holocellulolytic genes in different ways

To evaluate the influence of the transcription factor PAC-3 on the expression of holocellulolytic genes, we used RT-qPCR to determine the expression profiles of the cellulases-encoding genes *cbh1*, *gh7-1* and *gh3-2* and the hemicellulases-encoding genes *gh11-1*, *gh10-4* and *gh43-5* in the *N. crassa* strains 74A, $\Delta pac-3$, and $\Delta mus-52$ (Fig 4).

The behavior of holocellulolytic genes was quite varied in response to deletion of the *pac-3* gene. Regarding cellulases, we observed that the absence of *pac-3* decreased the expression of *cbh1* in acidic (4-fold at pH 3.0 and 5.6-fold at pH 5.0), neutral (4-fold), and alkaline (7.5-fold at pH 8.0) media (Fig 4A). These results indicate that PAC-3 can serve as a positive regulator of *cbh1* in acidic, neutral, and alkaline media. Expression of the endoglucanase gene (*gh7-1*) in $\Delta pac-3$ revealed a reduction at pH 3.0 (0.5-fold), pH 5.0 (2.7-fold), and pH 8.0 (1.6-fold) and an upregulation at pH 7.0 (1.6-fold) and pH 10.0 (approximately 6.5-fold) (Fig 4B). In contrast, the expression of β -glucosidase 1 (*gh3-2*) increased as compared to the parental strains at all tested pH conditions—except at pH 5.0—reaching a maximum expression at pH 10.0 (approximately 7.5-fold increase) (Fig 4C). This result suggests that PAC-3 can act as a repressor of β -glucosidase during the growth of *N. crassa* on sugarcane bagasse, principally in alkaline medium.

Concerning the behavior of hemicellulases during the growth of *N. crassa* on sugarcane bagasse, we observed that expression of the *gh11-1* gene decreased at all tested pH conditions (reaching an almost 5-fold reduction at pH 7.0), except at pH 10 (approximately 2-fold higher) (Fig 4D). In contrast, expression of the *gh10-4* gene was null in the $\Delta pac-3$ strain in acidic and




Fig 4. Gene expression profiles of holocellulolytic enzymes in *N. crassa*. The strains 74A, $\Delta mus-52$, and $\Delta pac-3$ grown on sugarcane bagasse at different pH (3.0, 5.0, 7.0, 8.0, and 10.0) were used for RT-qPCR experiments. (A) *cbh-1* = Cellobiohydrolase 1 (exoglucanase) NCU07340; (B) *gh7-1* = Endoglucanase 1 NCU05057; (C) *gh3-2* = β -glucosidase 1 NCU08054; (D) *gh11-1* = Endo-1,4- β -xylanase 1 NCU02855; (E) *gh10-4* = Endo-1,4- β -xylanase 2 NCU07130; (F) *gh43-5* = β -xylosidase NCU09652. Values show the mean of three replicates. The error bar indicates the standard deviation. *P < 0.05, **P < 0.01, ***P < 0.001.

doi:10.1371/journal.pone.0169796.g004

alkaline cultures (pH 3.0, pH 5.0, pH 8.0, and pH 10.0) as compared to that in 74A. However, at pH 7.0, the expression was dramatically reduced as compared to the parental strain (almost 4-fold) (Fig 4E). Thus, we believe that PAC-3 may be important in inducing the expression of the *ebx2* gene at neutral pH. Similar to the *gh11-1* gene, *pac-3* deletion caused a reduction of *gh43-5* expression in acidic and alkaline media (reaching 2.5-fold reduction at pH 5.0) (Fig 4F), suggesting that PAC-3 upregulates the expression of these genes.

Our data reveal that PAC-3 affects the expression of holocellulolytic genes and can serve as a positive or negative regulator, depending on the initial pH of the cultures. Therefore, this transcription factor may be involved in fungal adaptation to different pH conditions, implicating a critical role for PAC-3 in the regulation of lignocellulosic biomass deconstruction in *N. crassa*.

In order to understand how PAC-3 regulates cellulase and xylanase gene expression, we performed a genome-wide analysis of PAC-3 binding motifs. The results are presented in Fig 5. As can be observed, all studied genes showed potential motifs for PAC-3 binding [18]. The cellulase genes *cbh-1* and *gh7-1* contained three potential sites, while the *gh3-2* glucosidase-encoding gene contained only two sites. For xylanases-encoding genes, the gene *gh11-1*




Fig 5. Schematic representation of putative PAC-3 binding sites (5' -BGCCVAGV-3'). The analysis was performed using the region 1.0-kbp upstream of cellulolytic and xylanolytic genes. The position of the motifs is relative to the translation initiation codon (ATG). B = C or G or T; V = A or C or G, according to International Union of Pure and Applied Chemistry (IUPAC) norms.

doi:10.1371/journal.pone.0169796.g005

showed four potential PAC-3 binding sites, whereas *gh10-4* and *gh43-5* only contained one site. This data suggests that the transcription factor PAC-3 can directly regulate the expression of cellulases and xylanases.

The deletion of *pac-3* affects the expression of transcription factor-encoding genes

Although PAC-3 appears to directly regulate cellulase and xylanase genes, we decided to investigate if PAC-3 also influences the expression of the major transcription factors that are known to regulate holo-cellulases genes [9, 11]: *xlr-1*, *cre-1*, *clr-1*, and *clr-2*. First, we performed genome-wide analysis of the PAC-3 binding motifs in the promoters (1 kb upstream the start codon) of genes encoding these transcription factors. Only the gene *cre-1* did not contain a potential binding site for PAC-3, whereas *xlr-1* contained four sites, followed by two sites for *clr-2*, and one site for *clr-1* (data not shown).

The expression of transcription factor-encoding genes evaluated by RT-qPCR showed that, indeed, PAC-3 can regulate the expression of these factors, depending on pH conditions. Expression of *xlr-1* decreased in all tested pH conditions, showing a 0.5-fold reduction at pH 8.0 when compared to that in the parental 74A strain (Fig 6A). Although, *cre-1* does not contain a potential binding site for PAC-3, the expression of this gene was significantly decreased in acidic (pH 3.0) and alkaline (pH 8.0 and 10.) media, suggesting an indirect regulation of carbon catabolite repression mediated by CRE-1 by PAC-3 (Fig 6B). The expression of *clr-1* and *clr-2* were decreased in acidic (pH 3.0 and 5.0) and alkaline (pH 8.0) conditions. Notably, at pH 7.0, both transcription factors, *clr-1* and *clr-2*, were upregulated when compared to the strain 74A (Fig 6C and 6D). Also, we noted that the deletion of the gene *mus52* affected the expression of both *cre-1* and *clr-2* (Fig 6B and 6D). These results indicate that cellulase- and xylanase-encoding genes can be regulated by PAC-3 in both a direct and indirect manner.

Discussion

Different pH conditions lead microorganisms to adapt physiologically to ensure their survival and reproduction. For this purpose, organisms have developed signal transduction pathways, allowing the secretion of enzymes with optimal activity at specific pH values. The regulation of gene expression and secretion of extracellular hydrolytic enzymes is part of this process and is crucial to improving lignocellulosic biomass degradation [33].

Our data show that the transcription factor PAC-3, a well-defined pH regulator in several species of filamentous fungi [12, 13, 19, 20], is also involved in the regulation of holo-cellulolytic enzyme production in *Neurospora crassa*. A study by Mello-de-Souza et al. [19] involving the thermophilic fungus *H. grisea* var. *thermoidea* showed that the transcription factor PacC is responsible for the regulation of those genes encoding cellulases and xylanases, which are strongly induced in sugarcane bagasse cultures at alkaline pH. This regulation can occur in a direct or an indirect manner. Another transcriptome study with the phytopathogen fungus *Colletotrichum gloeosporioides* showed that *pacC* deletion affects the expression of several genes, including cell wall-degrading enzymes [34]. Considering cellulase secretion, we observed that PAC-3 does not interfere with the expression of endoglucanase (CMCase activity) in acidic media, similar to that found by He and collaborators [12] in *T. reesei* during growth on cellulose. In this same study, they observed that total cellulolytic activity was higher in the mutant strain $\Delta Trpac1$ at neutral pH, as we showed for *N. crassa*, suggesting that *T. reesei* and *N. crassa* may share a similarity in PAC-3. However, the role of PAC1 in regulating cellulase-encoding genes in *T. reesei* is still unclear, since cellulase was not affected by *pac1* deletion in another study [20].




Fig 6. Effect of *pac-3* deletion on transcriptional factors gene expression. The strains 74A, $\Delta mus-52$, and $\Delta pac-3$ grown on sugarcane bagasse at different pH (3.0, 5.0, 7.0, 8.0, and 10.0) were used for RT-qPCR experiments. (A) *xlr-1* = Xylan degradation regulator-1 NCU06971; (B) *cre-1* = Carbon catabolite regulator NCU08807; (C) *clr-1* = Cellulose degradation regulator-1 NCU07705 and (D) *clr-2* = Cellulose degradation regulator-2 NCU08042. Values show the mean of three replicates. The error bar indicates the standard deviation. *P < 0.05, **P < 0.01, ***P < 0.001.

doi:10.1371/journal.pone.0169796.g006

PAC-3 both positively and negatively regulates the expression of holocellulolytic genes during the cultivation of *N. crassa* on sugarcane bagasse. A study of *T. reesei* using genome-wide transcriptional analysis showed that, although *pac1* deletion affected several genes, few cellulases and hemicellulases were PAC1-regulated at pH 6.0, whereas β -glucosidase (Tr_47268) and α -1,6-mannanase (Tr_122495) were both downregulated in the $\Delta pac1$ mutant [20]. Analyzing the expression of a β -glucosidase (*bgl1*), we noted that PAC-3 served as a repressor in both neutral and alkaline pH, different from that observed for the candidate β -glucosidase Tr_47268 from *T. reesei*. However, it is important to highlight that there are at least 11 β -glucosidases annotated in the *T. reesei* genome that could potentially be targeted for regulation. In contrast, our results agree with a previous report where the expression of *bgl1* increased dramatically in the mutant $\Delta Trpac1$ at neutral pH [12].

In *A. nidulans*, the induction or repression of cellulase genes is delayed in absence of the regulator PacC. RNA-sequencing analysis showed that genes encoding β -endoglucanases, cellobiohydrolases, and β -endoxyanases are among the major CAZy genes downregulated in $\Delta pacC$ mutant during growth on cellobiose at alkaline pH [13]. Our results also showed that in *N. crassa* there is a reduction in the expression of the genes *cbh-1*, *gh11-1*, and *gh10-4* in $\Delta pac-3$ compared to that of the parental strain at all analyzed pH conditions, except for *gh11-1* at pH 10.0. Also in *A. nidulans*, Perez-Gonzalez et al. [35] showed that during the growth on D-xylose, expression of β -xylosidase (*xlnD*) is partially reduced in the mutant strain *pacC*^{C14}, suggesting that this transcription factor serves to positively regulate *xlnD*. Our results suggest a similar regulation in *N. crassa*, considering that the expression of the β -xylosidase gene *gh43-5* was also downregulated in the *N. crassa* $\Delta pac-3$ mutant.

A study involving the phytopathogen fungus *Alternaria alternata* showed that environmental pH has a large influence on the expression of the endoglucanase *AaK1* gene [36]. Examining the effect of PAC-3 on the expression of the *eg1* gene in *N. crassa*, we observed that the $\Delta pac-3$ mutant showed reduced expression of *eg1* at pH 8.0, whereas it increased considerably at pH 10.0. Thus, the effect of PAC-3 on the regulation of endoglucanase in *N. crassa* remains unclear.

Overall, the transcription factor PAC-3 is involved in secretion and gene expression regulation of cell wall-degrading enzymes. Notably, genes with similar functions can be positively or negatively regulated by PAC-3 depending on pH conditions, as found in *A. nidulans* [13]. This observation may be related to the dual regulatory role of PacC, previously described by McCabe et al. [37], in which two xylanases from *A. nidulans* have opposite expression patterns depending on environmental pH. This interesting finding should promote more detailed studies in understanding how pH is sensed by *N. crassa* and how PAC-3 is involved in synergism of holocellulase action.

In Summary, our results suggest that PAC-3 acts inducing total cellulolytic activity and endoglucanase activity at acidic pH environments whereas acting as a repressor of total cellulolytic and endoglucanase activity at neutral and alkaline pH. On the other hand, for xylanase activity, PAC-3 acts as a positive regulator in all pH environments (except at pH 8.0). It is well known that holocellulolytic enzymes are both directly and indirectly regulated [13, 19], by PacC. Our study showed that all studied genes can be directly targeted by PAC-3, probably at transcriptional level. On the other hand, PAC-3 can also regulate the major transcription factors that regulate the expression of holocellulase-encoding genes. Given the regulons of these transcription factors [9, 11] and our results, we propose a simple model of holocellulase regulation by PAC-3 during sugarcane bagasse degradation (Fig 7).

Changes in the media pH activate the transcription factor PAC-3, which can then directly activate cellulase- and xylanases-encoding genes. In addition, PAC-3 can also directly activate *xlr-1*, which activates β -glucosidase (*gh3-1*) as well as endo-xylanase (*gh11-1*) and β -xylosidase (*gh43-5*). In this regulatory network, the transcription factor CLR-1 seems to have an important role. Directly, CLR-1 can activate only cellulases (*cbh-1* and *gh7-1*); however CLR-1 can also activate *xyr-1* and *clr-2* [9], which in turn will activate holocellulases-encoding genes. Finally, CLR-2 can activate both cellulases (*cbh-1* and *gh7-1*) and endo-xylanases (*gh11-1*) (Fig 7). CRE-1 mediates carbon catabolite repression, and PAC-3 can be involved in *cre-1* expression in an indirect manner. However, despite these evidences of regulation, how the overlapping regulation between PAC-3 and other transcription factors, especially CLR-1, is governed remains unclear. Indirect regulation or overlapping regulation with transcriptional factors seems to be related to the availability of low molecular weight inducers in *A. nidulans* [13]. However, more detailed studies will be necessary to elucidate the mechanism involved.




Fig 7. The model proposed for the regulation of holocellulolytic-encoding genes in *N. crassa* by PAC-3. Changes in the pH environment lead to activation of PAC-3, which directly or indirectly regulates transcription of cellulases and xylanases as well as the transcription factors XLR-1, CLR-1, and CLR-2 genes. The transcription factor CLR-1 regulate transcription of both *xlr-1* and *clr-2* genes. These transcription factors can also regulate cellulases and xylanases. XLR-1 regulate transcription of *gh3-1*, *gh11-1*, and *gh43-5* genes (purple arrows); CLR-1 regulate transcription of *cbh-1* and *gh7-1* genes (green arrows), and CLR-2 regulate transcription of *cbh-1*, *gh7-1*, and *gh11-1* genes (red arrows). (?) Missing component in regulation of *cre-1* by PAC-3.

doi:10.1371/journal.pone.0169796.g007

Conclusion

Our data shows that the transcription factor PAC-3 from *N. crassa* influences the expression profile and secretion of cell wall-degrading enzymes during growth in sugarcane bagasse. At the different pH levels examined, we show that cellulase secretion is reduced in the absence of *pac-3* at an alkaline pH, and xylanolytic activity is positively regulated by PAC-3 at pH 5.0, 7.0, and 10.0. Regarding holocellulolytic genes, we observed differential regulation depending on the pH of the culture medium, suggesting that *N. crassa* can adapt to different environmental conditions. PAC-3 can also activate the major transcription factors that regulate holocellulase gene expression, suggesting that regulation occurs in both a direct and indirect manner. Ambient pH is a key factor in the industrial production of biomass-degrading enzymes. Our study contributes to a better understanding of the mechanisms involved in holocellulase production in response to pH, thereby improving its use in several industrial processes.

Acknowledgments

This work was supported by grants from the following Brazilian Funding Agencies: São Paulo Research Foundation (FAPESP; Grant No 2014/09493-2, ACCA; Grant No 2014/03847-7, AR, RSS, and AHSC), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; Grant No 304222/2013-5), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Fundação de Apoio ao Ensino, Pesquisa e Assistência (FAEPA). We thank Professor Maria Celia Bertolini (Department of Biochemistry and Chemical Technology, Institute of Chemistry, Universidade Estadual Paulista, UNESP, Araraquara, São Paulo, Brazil) for providing the *N. crassa pac-3* knockout mutant (*pac-3^{KO}*).

Author Contributions

Conceptualization: RNS AR.

Data curation: ACCA WRP KMVN LSC RSS AHSC.

Funding acquisition: AR.

Methodology: RSR.

Project administration: RNS.

Writing – original draft: RNS AR ACCA.

Writing – review & editing: AR RNS.

References

1. Segato F, Damásio ARL, de Lucas RC, Squina FM, Prade RA. Genomics Review of Holocellulose Deconstruction by Aspergilli. *Microbiology and Molecular Biology Reviews*. 2014; 78(4):588–613. doi: [10.1128/MMBR.00019-14](https://doi.org/10.1128/MMBR.00019-14) PMID: [25428936](https://pubmed.ncbi.nlm.nih.gov/25428936/)
2. Tian C, Beeson WT, Iavarone AT, Sun J, Marletta MA, Cate JH, et al. Systems analysis of plant cell wall degradation by the model filamentous fungus *Neurospora crassa*. *Proc Natl Acad Sci U S A*. 2009; 106(52):22157–62. Epub 2009/12/19. doi: [10.1073/pnas.0906810106](https://doi.org/10.1073/pnas.0906810106) PMID: [20018766](https://pubmed.ncbi.nlm.nih.gov/20018766/)
3. Perkins DD, Turner BC, Barry EG. Strains of *Neurospora* Collected from Nature. *Evolution*. 1976; 30(2):281.
4. Pandit A, Maheshwari R. Life-history of *Neurospora intermedia* in a sugar cane field. *Journal of Biosciences*. 1996; 21(1):57–79.
5. Dogaris I, Mamma D, Kekos D. Biotechnological production of ethanol from renewable resources by *Neurospora crassa*: an alternative to conventional yeast fermentations? *Appl Microbiol Biotechnol*. 2013; 97(4):1457–73. doi: [10.1007/s00253-012-4655-2](https://doi.org/10.1007/s00253-012-4655-2) PMID: [23318834](https://pubmed.ncbi.nlm.nih.gov/23318834/)
6. Kubicek CP. Systems biological approaches towards understanding cellulase production by *Trichoderma reesei*. *J Biotechnol*. 2013; 163(2):133–42. doi: [10.1016/j.jbiotec.2012.05.020](https://doi.org/10.1016/j.jbiotec.2012.05.020) PMID: [22750088](https://pubmed.ncbi.nlm.nih.gov/22750088/)
7. Gutiérrez-Rojas I, Moreno-Sarmiento N, Montoya D. Mecanismos y regulación de la hidrólisis enzimática de celulosa en hongos filamentosos: casos clásicos y nuevos modelos. *Rev Iberoam Micol*. 2015; 32(1):1–12.
8. Sun JP, Tian CG, Diamond S, Glass NL. Deciphering Transcriptional Regulatory Mechanisms Associated with Hemicellulose Degradation in *Neurospora crassa*. *Eukaryot Cell*. 2012; 11(4):482–93. doi: [10.1128/EC.05327-11](https://doi.org/10.1128/EC.05327-11) PMID: [22345350](https://pubmed.ncbi.nlm.nih.gov/22345350/)
9. Craig JP, Coradetti ST, Starr TL, Glass NL. Direct Target Network of the *Neurospora crassa* Plant Cell Wall Deconstruction Regulators CLR-1, CLR-2, and XLR-1. *mBio*. 2015; 6(5):e01452–15.
10. Coradetti ST, Craig JP, Xiong Y, Shock T, Tian C, Glass NL. Conserved and essential transcription factors for cellulase gene expression in ascomycete fungi. *Proceedings of the National Academy of Sciences*. 2012; 109(19):7397–402.
11. Sun JP, Glass NL. Identification of the CRE-1 Cellulolytic Regulon in *Neurospora crassa*. *Plos One*. 2011; 6(9).
12. He R, Ma L, Li C, Jia W, Li D, Zhang D, et al. Trpac1, a pH response transcription regulator, is involved in cellulase gene expression in *Trichoderma reesei*. *Enzyme Microb Technol*. 2014; 67:17–26. Epub 2014/12/03. doi: [10.1016/j.enzmictec.2014.08.013](https://doi.org/10.1016/j.enzmictec.2014.08.013) PMID: [25442944](https://pubmed.ncbi.nlm.nih.gov/25442944/)
13. Kunitake E, Hagiwara D, Miyamoto K, Kanamaru K, Kimura M, Kobayashi T. Regulation of genes encoding cellulolytic enzymes by Pal-PacC signaling in *Aspergillus nidulans*. *Appl Microbiol Biotechnol*. 2016; 100(8):3621–35. doi: [10.1007/s00253-016-7409-8](https://doi.org/10.1007/s00253-016-7409-8) PMID: [26946171](https://pubmed.ncbi.nlm.nih.gov/26946171/)
14. Eberhart BM, Beck RS, Goolsby KM. Cellulase of *Neurospora crassa*. *J Bacteriol*. 1977; 130(1):181–6. Epub 1977/04/01. PMID: [15975](https://pubmed.ncbi.nlm.nih.gov/15975/)
15. Stewart JC, Parry JB. Factors influencing the production of cellulase by *Aspergillus fumigatus* (Fresenius). *J Gen Microbiol*. 1981; 125(1):33–9. Epub 1981/07/01. doi: [10.1099/00221287-125-1-33](https://doi.org/10.1099/00221287-125-1-33) PMID: [7038036](https://pubmed.ncbi.nlm.nih.gov/7038036/)
16. Ferreira SMP, Duarte AP, Queiroz JA, Domingues FC. Influence of buffer systems on *Trichoderma reesei* Rut C-30 morphology and cellulase production. *Electron J Biotechnol*. 2009; 12(3).
17. Rossi A, Cruz AHS, Santos RS, Silva PM, Silva EM, Mendes NS, et al. Ambient pH Sensing in Filamentous Fungi: Pitfalls in Elucidating Regulatory Hierarchical Signaling Networks. *Iubmb Life*. 2013; 65(11):930–5. doi: [10.1002/iub.1217](https://doi.org/10.1002/iub.1217) PMID: [24265200](https://pubmed.ncbi.nlm.nih.gov/24265200/)
18. Virgilio S, Cupertino FB, Bernardes NE, Freitas FZ, Takeda AAS, Fontes MRD, et al. Molecular Components of the *Neurospora crassa* pH Signaling Pathway and Their Regulation by pH and the PAC-3 Transcription Factor. *PloS one*. 2016; 11(8).

19. Mello-de-Sousa TM, Silva-Pereira I, Poças-Fonseca MJ. Carbon source and pH-dependent transcriptional regulation of cellulase genes of *Humicola grisea* var. *thermoidea* grown on sugarcane bagasse. *Enzyme Microb Technol.* 2011; 48(1):19–26. doi: [10.1016/j.enzmictec.2010.08.007](https://doi.org/10.1016/j.enzmictec.2010.08.007) PMID: [22112766](https://pubmed.ncbi.nlm.nih.gov/22112766/)
20. Hakkinen M, Sivasiddharthan D, Aro N, Saloheimo M, Pakula TM. The effects of extracellular pH and of the transcriptional regulator PAC1 on the transcriptome of *Trichoderma reesei*. *Microb Cell Fact.* 2015; 14.
21. McCluskey K. *The Fungal Genetics Stock Center: From Molds to Molecules.* 2003; 52:245–62.
22. Colot HV, Park G, Turner GE, Ringelberg C, Crew CM, Litvinkova L, et al. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proceedings of the National Academy of Sciences.* 2006; 103(27):10352–7.
23. Cupertino FB, Freitas FZ, de Paula RM, Bertolini MC. Ambient pH Controls Glycogen Levels by Regulating Glycogen Synthase Gene Expression in *Neurospora crassa*. *New Insights into the pH Signaling Pathway.* *PLoS One.* 2012; 7(8):e44258. doi: [10.1371/journal.pone.0044258](https://doi.org/10.1371/journal.pone.0044258) PMID: [22952943](https://pubmed.ncbi.nlm.nih.gov/22952943/)
24. Vogel HJ. A convenient growth medium for *Neurospora crassa* (medium N). *Microbiol Genet Bull.* 1956; 13:42–3.
25. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta$ CT Method. *Methods.* 2001; 25(4):402–8. doi: [10.1006/meth.2001.1262](https://doi.org/10.1006/meth.2001.1262) PMID: [11846609](https://pubmed.ncbi.nlm.nih.gov/11846609/)
26. Xiao Z, Storms R, Tsang A. Microplate-based carboxymethylcellulose assay for endoglucanase activity. *Analytical Biochemistry.* 2005; 342(1):176–8. doi: [10.1016/j.ab.2005.01.052](https://doi.org/10.1016/j.ab.2005.01.052) PMID: [15958198](https://pubmed.ncbi.nlm.nih.gov/15958198/)
27. Castro LdS, Antoniêto ACC, Pedersoli WR, Silva-Rocha R, Persinoti GF, Silva RN. Expression pattern of cellulolytic and xylanolytic genes regulated by transcriptional factors XYR1 and CRE1 are affected by carbon source in *Trichoderma reesei*. *Gene Expression Patterns.* 2014; 14(2):88–95. doi: [10.1016/j.gep.2014.01.003](https://doi.org/10.1016/j.gep.2014.01.003) PMID: [24480777](https://pubmed.ncbi.nlm.nih.gov/24480777/)
28. Nogueira KMV, Costa MdN, de Paula RG, Mendonça-Natividade FC, Ricci-Azevedo R, Silva RN. Evidence of cAMP involvement in cellobiohydrolase expression and secretion by *Trichoderma reesei* in presence of the inducer sophorose. *BMC Microbiol.* 2015; 15(1).
29. Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, et al. The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature.* 2003; 422(6934):859–68. doi: [10.1038/nature01554](https://doi.org/10.1038/nature01554) PMID: [12712197](https://pubmed.ncbi.nlm.nih.gov/12712197/)
30. Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, et al. Determination and inference of eukaryotic transcription factor sequence specificity. *Cell.* 2014; 158(6):1431–43. doi: [10.1016/j.cell.2014.08.009](https://doi.org/10.1016/j.cell.2014.08.009) PMID: [25215497](https://pubmed.ncbi.nlm.nih.gov/25215497/)
31. Cupertino FB, Freitas FZ, de Paula RM, Bertolini MC. Ambient pH Controls Glycogen Levels by Regulating Glycogen Synthase Gene Expression in *Neurospora crassa*. *New Insights into the pH Signaling Pathway.* *PloS one.* 2012; 7(8).
32. Tian CG, Beeson WT, Iavarone AT, Sun JP, Marletta MA, Cate JHD, et al. Systems analysis of plant cell wall degradation by the model filamentous fungus *Neurospora crassa*. *Proceedings of the National Academy of Sciences of the United States of America.* 2009; 106(52):22157–62. doi: [10.1073/pnas.0906810106](https://doi.org/10.1073/pnas.0906810106) PMID: [20018766](https://pubmed.ncbi.nlm.nih.gov/20018766/)
33. Peñalva MA, Tilburn J, Bignell E, Arst HN. Ambient pH gene regulation in fungi: making connections. *Trends in Microbiology.* 2008; 16(6):291–300. doi: [10.1016/j.tim.2008.03.006](https://doi.org/10.1016/j.tim.2008.03.006) PMID: [18457952](https://pubmed.ncbi.nlm.nih.gov/18457952/)
34. Alkan N, Meng X, Friedlander G, Reuveni E, Sukno S, Sherman A, et al. Global Aspects of pacC Regulation of Pathogenicity Genes in *Colletotrichum gloeosporioides* Revealed by Transcriptome Analysis. *Molecular Plant-Microbe Interactions.* 2013; 26(11):1345–58. doi: [10.1094/MPMI-03-13-0080-R](https://doi.org/10.1094/MPMI-03-13-0080-R) PMID: [23902260](https://pubmed.ncbi.nlm.nih.gov/23902260/)
35. Perez-Gonzalez JA, van Peij NN, Bezoen A, MacCabe AP, Ramon D, de Graaff LH. Molecular cloning and transcriptional regulation of the *Aspergillus nidulans* xlnD gene encoding a beta-xylosidase. *Applied and Environmental Microbiology.* 1998; 64(4):1412–9. Epub 1998/04/18. PMID: [9546179](https://pubmed.ncbi.nlm.nih.gov/9546179/)
36. Eshel D, Miyara I, Ailing T, Dinoor A, Prusky D. pH regulates endoglucanase expression and virulence of *Alternaria alternata* in persimmon fruit. *Mol Plant Microbe Interact.* 2002; 15(8):774–9. Epub 2002/08/17. doi: [10.1094/MPMI.2002.15.8.774](https://doi.org/10.1094/MPMI.2002.15.8.774) PMID: [12182334](https://pubmed.ncbi.nlm.nih.gov/12182334/)
37. MacCabe AP, Orejas M, Perez-Gonzalez JA, Ramon D. Opposite patterns of expression of two *Aspergillus nidulans* xylanase genes with respect to ambient pH. *J Bacteriol.* 1998; 180(5):1331–3. Epub 1998/03/12. PMID: [9495775](https://pubmed.ncbi.nlm.nih.gov/9495775/)