

## Dynamics of Multiple *lin* Gene Expression in *Sphingomonas paucimobilis* B90A in Response to Different Hexachlorocyclohexane Isomers

Mrutyunjay Suar,<sup>1</sup> Jan Roelof van der Meer,<sup>2,3\*</sup> Kirsten Lawlor,<sup>2</sup>  
Christof Holliger,<sup>4</sup> and Rup Lal<sup>1</sup>

Department of Zoology, University of Delhi, Delhi, India,<sup>1</sup> and Process of Environmental Microbiology and Molecular Ecotoxicology, Swiss Federal Institute for Environmental Science and Technology (EAWAG), Duebendorf,<sup>2</sup> and Laboratory for Environmental Biotechnology, ENAC-ISTE, EPFL,<sup>4</sup> and Department of Fundamental Microbiology, University of Lausanne,<sup>3</sup> Lausanne, Switzerland

Received 10 March 2004/Accepted 5 July 2004

*Sphingomonas paucimobilis* B90A is able to degrade the  $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ -isomers of hexachlorocyclohexane (HCH). It contains the genes *linA*, *linB*, *linC*, *linD*, *linE*, and *linR*, which have been implicated in HCH degradation. In this study, dynamic expression of the *lin* genes was measured in chemostat-grown *S. paucimobilis* B90A by RNA dot blot hybridization and real-time reverse transcriptase PCR upon exposure to a pulse of different HCH isomers. Irrespective of the addition of HCH, *linA*, *linB*, and *linC* were all expressed constitutively. In contrast, *linD* and *linE* were induced with  $\alpha$ -HCH (2 mg/liter) and  $\gamma$ -HCH (7 mg/liter). A sharp increase in mRNA levels for *linD* and *linE* was observed from 10 to 45 min after the addition of  $\alpha$ - or  $\gamma$ -HCH. Induction of *linD* and *linE* was not detectable upon the addition of 0.7 mg of  $\gamma$ -HCH per liter, although the compound was degraded by the cells. The addition of  $\beta$ -HCH (5 mg/liter) or  $\delta$ -HCH (20 mg/liter) did not lead to *linE* and *linD* induction, despite the fact that 50% of the compounds were degraded. This suggests that degradation of  $\beta$ - and  $\delta$ -HCH proceeds by a different pathway than that of  $\alpha$ - and  $\gamma$ -HCH.

Hexachlorocyclohexane (HCH) has been extensively used for the control of insect pests on agriculturally important crops, seeds, and vegetables, in forestry, and in vector control (34). Mainly two forms of HCH, lindane ( $\gamma$ -HCH) and a technical mixture of all isomers, have been applied. Technical grade HCH (33) largely consists of  $\alpha$ -HCH (60 to 70%), with  $\beta$ -HCH (5 to 12%),  $\gamma$ -HCH (10 to 15%), and  $\delta$ -HCH (6 to 10%) (16). As a result of the extensive use of lindane dust and technical HCH over the years and despite the recent ban on the use of HCH, several countries are currently faced with two very serious problems: (i) soil contamination with small amounts of HCH and (ii) highly contaminated sites where lindane was produced and purified or disposed of (5, 25).

Although HCH is persistent and difficult to biodegrade, a few microorganisms have been isolated which can degrade one or more HCH isomers under aerobic conditions (27, 29, 31). Most of the strains, such as *Sphingomonas paucimobilis* UT26, degrade  $\alpha$ - and  $\gamma$ -HCH but do not degrade  $\beta$ -HCH. Only one strain has been described, *S. paucimobilis* B90A (9), the parent strain of strain B90 (14, 15), that is able to degrade the four  $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ -HCH isomers, although with different rates and not to completion for the  $\beta$ - and  $\delta$ -HCH isomers. The degradation pathway for  $\gamma$ -HCH is well established from work on *S. paucimobilis* strain UT26 (20–24). Degradation of  $\gamma$ -HCH is mediated by the products of the so-called *lin* genes (23). It is assumed that  $\alpha$ -HCH is degraded through the same pathway,

but this has not been proven for  $\beta$ - and  $\delta$ -HCH. *S. paucimobilis* strains UT26 and B90A have very similar *lin* gene sequences and organizations (9, 15). In contrast to typical degradation pathways in, for example, *Pseudomonas* or *Ralstonia*, where very long polycistronic operons are common (10, 17, 36), the genes for lindane degradation in *S. paucimobilis* are not organized within one or a few operons (Fig. 1). At least five different transcriptional units encode the lindane degradation pathway in strain UT26, e.g., *linXA*, *linB*, *linC*, and *linDE* and *linR* (21, 22). The last three genes form part of a small inducible regulon with *linR* as its main transcriptional activator (21). As far as is known, one of the main differences between strains UT26 and B90A is that strain B90A contains two copies of the *linA* gene and three copies of a *linX*-like gene (9). Since the two *linA* gene copies in B90A are not part of the same organizational unit, this increases the number of possible transcriptional units for the lindane degradation pathway in strain B90A to six (Fig. 1A). This makes it an interesting system to determine if and how *S. paucimobilis* B90A is capable of coordinating *lin* gene expression in response to HCH isomers.

Here we have studied the expression dynamics of the *lin* genes in *S. paucimobilis* B90A. Our first interest was to determine if *lin* gene expression from the different operons in strain B90A is regulated in a coordinated way in response to  $\gamma$ -HCH. Second, we wanted to know whether low  $\gamma$ -HCH concentrations would still lead to detectable *lin* gene expression. As a third aspect of this work, we studied *lin* gene expression in response to the other HCH isomers ( $\alpha$ -,  $\beta$ -, and  $\delta$ -HCH). Gene expression studies were all carried out on chemostat-grown cultures of *S. paucimobilis* B90A cells, which could be sampled in a rapid time sequence after the addition of the HCH isomers

\* Corresponding author. Mailing address: University of Lausanne, Department of Fundamental Microbiology, Bâtiment de Biologie, CH-1015 Lausanne, Switzerland. Phone: 41 21 692 5630. Fax: 41 21 692 5605. E-mail: JanRoelof.VanDerMeer@unil.ch.




FIG. 1. Organization of the *lin* genes in *S. paucimobilis* strain B90A. (A) Organization of the relevant regions analyzed by DNA sequencing with the *lin* genes indicated (5). The only known regulatable promoter among the *lin* genes is in front of *linE* and is activated by LinR (21). The function of the *linX* gene product is not known, but based on sequence homology, it was predicted to be a dehydrogenase perhaps similar in function to LinC (24). (B) Simplified version of the degradation pathway for  $\gamma$ -HCH ([http://umbbd.ahc.umn.edu/ghch/ghch\\_image\\_map.html](http://umbbd.ahc.umn.edu/ghch/ghch_image_map.html)). Reactions catalyzed by Lin enzymes are indicated as such. When more than one arrow is drawn, reactions are supposed to have intermediate steps involving unstable intermediates. The part shown within brackets remains speculative and is based on analogies to maleylacetate degradation. 2,5-DCHQ, 2,5-dichloroquinone; MAR, maleylacetate reductase.

for *lin* mRNA synthesis and HCH concentration without disturbing the culture growth, as described previously (12, 19). Amounts of *lin* mRNAs were determined by quantitative dot blot hybridization, and *linD* and *linB* levels were determined by quantitative real-time reverse transcriptase PCR (qRT-PCR).

#### MATERIALS AND METHODS

**Bacterial strains and culture medium.** *S. paucimobilis* B90A (9) was grown at 28°C in mineral salt medium containing the following (per liter): 0.5 g of  $(\text{NH}_4)_2\text{HPO}_4$ , 0.2 g of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.1 g of  $\text{K}_2\text{HPO}_4$ , 0.01 g of  $\text{Ca}(\text{NO}_3)_2$ , and 0.01 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  (pH 7.0), supplemented with 1 g of glucose and 1 g of peptone. *Escherichia coli* DH5 $\alpha$  was used as general host for plasmid cloning. *E. coli* strains were grown routinely at 37°C on Luria broth Luria agar (28), supplemented with 100  $\mu\text{g}$  of ampicillin per ml when necessary.

**Chemostat cultivation of *S. paucimobilis* B90A.** In order to study *lin* gene expression, *S. paucimobilis* B90A was grown in continuous culture in a 0.5-liter reactor under carbon-limited conditions. The reactor had a working volume of 200 ml at a dilution rate of 0.05  $\text{h}^{-1}$  and was maintained at 30°C with a dissolved oxygen concentration at 90% of saturation. Growth medium consisted of mineral salt medium containing 0.1% glucose. *S. paucimobilis* B90A was grown for at least 7 volume changes before being exposed to HCH isomers. The turbidity of the culture at 600 nm under steady-state conditions was 1.2. HCH was added to the cultures as one single pulse of 500  $\mu\text{l}$  of a solution of the respective HCH isomer in dimethyl sulfoxide, which was introduced directly into the chemostat vessel while continuing the supply of normal growth medium. The final concen-

trations of the HCH isomers were 2 mg/liter for  $\alpha$ -HCH, 5 mg/liter for  $\beta$ -HCH, and 20 mg/liter for  $\delta$ -HCH; these concentrations approach the aqueous solubilities of the isomers as reported in the literature (35). In induction experiments with  $\gamma$ -HCH, two final concentrations of 7 (the maximal aqueous solubility) and 0.7 mg per liter were established. Experiments with  $\gamma$ -HCH were performed twice independently, and induction experiments with  $\alpha$ -,  $\beta$ -, and  $\delta$ -HCH were performed only once.

**Total RNA extraction.** Total RNA from chemostat-grown *S. paucimobilis* B90A was isolated 50 and 100 min before and 5, 10, 15, 20, 30, 45, 60, 120, 180, and 240 min after the addition of HCH to the culture. Samples (3 ml) were taken directly from the chemostat, immediately pelleted by centrifugation ( $10,000 \times g$ , 30 s) at 4°C, and resuspended in 50  $\mu\text{l}$  of RNAlater (Ambion, Inc., Austin, Tex.). When all the samples were collected, the cells in RNAlater were again pelleted by centrifugation, resuspended in buffer, and extracted by acid phenol at 60°C as described by Aiba et al. (1). Total nucleic acids were further purified by phenol-chloroform extraction. After that, the RNA was precipitated with ethanol, treated with DNase I to remove traces of contaminating DNA (RNase free; Roche Biochemicals, Rotkreuz, Switzerland), and once more precipitated with ethanol. Aqueous RNA concentrations were measured from their UV absorption at 260 nm in a regular UV-visible light spectrophotometer (Kontron Instruments AG, Zürich, Switzerland).

**In vitro synthesis of sense and antisense *lin* probes.** The *linA1*, *linB*, *linC*, *linX*, *linD*, *linE*, and *linR* open reading frames (ORFs) were cloned from plasmids pLINA1, pLINB, pLINC, pLINX, pLIND, pLINE, and pLINR (9, 15) in pGEM-7Zf(+) (Promega Corp., Madison, Wis.) which were maintained in *E. coli* DH5 $\alpha$  (28). All the clones were verified by restriction analysis, and the orientation of *lin*

TABLE 1. PCR primers used for real-time RT-PCR analysis

Primer sequence (5' to 3')	Designation	GenBank accession no.
GTC CGA ATC GCC CAT GCC GA	RT <i>linB</i>	AY150581
GTC GAA CCC TTC GGA ATC TT	RT <i>linD</i>	AY150583
TGG CGA GAA GAA ATT CAT TGA GA	<i>linBf</i>	AY150581
TCG CCG GTC CCT TCA TC	<i>linBr</i>	AY150581
GAA CTG TTC CAC TTC GTG TTC TCA	<i>linDf</i>	AY150583
GGT CAC GCC CTT CTC CAT TA	<i>linDr</i>	AY150583

ORFs in pGEM7Zf(+) was analyzed by DNA sequencing. One microgram of each pGEM-derived plasmid was linearized by restriction digestion to obtain an antisense transcript from the respective *lin* ORF. Antisense-labeled RNAs were then synthesized by *in vitro* transcription with biotin-16-UTP and either T7 or SP6 RNA polymerase (Roche Biochemicals). After synthesis, the template DNAs were degraded by incubation with RNase-free DNase I (Roche Biochemicals), and the mRNAs were aliquoted, precipitated in absolute ethanol, and stored at  $-20^{\circ}\text{C}$ . This procedure was followed for probes of all *lin* genes, except *linA2*. This gene is too similar to *linA1* and, therefore, cannot be distinguished by hybridization from *linA1* (15).

**Dot blot hybridization of *lin* mRNAs.** Total RNA from all the extractions was diluted in diethylpyrocarbonate-treated double-distilled water to final concentrations of 4 and 0.4  $\mu\text{g}$  in 25  $\mu\text{l}$  and blotted onto positively charged nylon membranes (QIAGEN, Basel, Switzerland) in a 96-well dot blot manifold (Gibco Life Technologies, Gaithersburg, Md.). Dilutions of denatured total genomic DNA from *S. paucimobilis* B90A and of pGEM-T-easy plasmid DNA containing the respective *lin* inserts were included on the same blot. The membrane was prehybridized for 1 h at  $65^{\circ}\text{C}$  in a solution of 7% sodium dodecyl sulfate, 10 g of bovine serum albumin (Fraction V; Sigma Chemical Co., St. Louis, Mo.) per liter, 0.5 M sodium phosphate buffer (pH 7.2), and 1 mM EDTA (pH 8.0) and then hybridized for 15 h at  $65^{\circ}\text{C}$  in the same solution with one aliquot of *in vitro* synthesized biotin-labeled single-stranded antisense RNA probes (approximately 100 ng) as described previously (4). Sense RNAs from the same plasmids were used in separate hybridizations as negative controls. After hybridization, the membranes were washed twice at  $65^{\circ}\text{C}$  in 0.1% sodium dodecyl sulfate and  $0.2\times$  SSC ( $1\times$  SSC is 0.15 M NaCl plus 0.015 M sodium citrate) for 30 min and then incubated for detection as described for the Southern-Light chemiluminescence system (Tropix, Bedford, Mass.). Membranes were exposed at room temperature to Hyperfilm (Amersham Life Sciences, Little Chalfont, Buckinghamshire, United Kingdom). Exposed films were scanned and transferred to 8-bit TIFF files, and the densities of the dots on the exposures were quantified with Meta-view version 4.5 (Universal Imaging Corporation, West Chester, Pa.). In all cases, the sense RNA hybridizations did not show any signals above the background.

Signal intensities of hybridization spots were measured across the exact spot size (i.e., total gray value divided by spot area) but were subsequently standardized for one spot size throughout the membrane. The chromosomal and plasmid DNA standards were used to prepare a standard curve for the amount of DNA versus average gray value per pixel, according to a similar method described by Leveau et al. (19). Signal intensities of RNA samples were converted from the standard curve to equivalent DNA amounts and divided by the amount of RNA spotted. When possible, values derived from different RNA dilutions were averaged, and standard deviations were calculated.

**Analytical method.** The disappearance of individual HCH isomers during the induction of chemostat-grown cells of *S. paucimobilis* B90A was measured by a gas-liquid chromatograph (GC-5890; Perkin Elmer) equipped with an electron capture detector and DB5 capillary column (15). For gas chromatography analysis, chemostat samples (200  $\mu\text{l}$ ) were extracted with 500  $\mu\text{l}$  of hexane, of which 1  $\mu\text{l}$  was injected into the gas chromatograph as described previously (15).

**Real-time PCR analysis.** Based on results from dot blot hybridization, only *linB* (an example of a constitutively expressed gene) and *linD* mRNA (an inducible gene) were chosen for quantification by qRT-PCR analysis. Total RNA was taken from samples of the cultures induced with 7 mg of  $\gamma$ -HCH per liter, which were also used for dot blot studies. All purified RNA samples were used in two separate reactions, one to synthesize cDNA and the other to test for DNA contamination. The reverse transcription step was performed with TaqMan reverse transcription reagents (Applied Biosystems) with (for the cDNA synthesis reaction) and without (for the control reaction) reverse transcriptase with the help of gene-specific primers (Table 1). Mixtures for cDNA synthesis and the

control reactions were heated in a standard thermocycler (Applied Biosystems) for 10 min at  $25^{\circ}\text{C}$  and then for 30 min at  $48^{\circ}\text{C}$  (transcription step), followed by 5 min at  $95^{\circ}\text{C}$  (inactivation). Following cDNA synthesis, real-time PCR was performed (ABI Prism 7000; Applied Biosystems) for primer optimization for the *linB* and *linD* genes with gene-specific primers (Table 1) in new tubes to which cDNA and control preparation were added, as described above. Once the correct primer concentration was identified for each of the genes, real-time PCR was performed with an initial hold temperature of  $95^{\circ}\text{C}$  for 10 min, followed by 40 amplification cycles of 15 s at  $95^{\circ}\text{C}$  and 1 min at  $60^{\circ}\text{C}$ .

## RESULTS AND DISCUSSION

**Constitutive expression of the *linA*, *linB*, *linC*, and *linX* genes.** In order to find out whether the different *lin* operons in strain B90A were coordinately regulated in response to  $\gamma$ -HCH induction, chemostat-grown cells of *S. paucimobilis* B90A were treated with a single dose of  $\gamma$ -HCH at a concentration of 7 mg/liter. This concentration (25  $\mu\text{M}$ ) allows its maximum solubility in water but is still relatively low in comparison to the inducer concentrations usually applied in gene expression studies, which are normally tested in concentrations above 100  $\mu\text{M}$  (8). The formation of individual *lin* gene transcripts before and up to 4 h after induction was analyzed by quantitative dot blotting with antisense labeled *lin* probes. Quantitative dot blotting is a simple but relatively accurate technique to monitor gene induction (12, 19). In contrast to the *tfd* or *hbp* regulons which were previously studied by this method (12, 19), most of the *lin* genes of *S. paucimobilis* B90A were expressed to the same level, irrespective of the addition of  $\gamma$ -HCH (Fig. 2A). Hence, we conclude that *linA* (here taken as the combination of both *linA1* and *linA2*), *linB*, *linC*, and *linX* are constitutively expressed and are not under the control of lindane-inducible promoters. This situation is similar to the expression of the *lin* genes in strain UT26, although the *lin* genetic organizations in strains UT26 and B90A are not identical (9). Approximations based on standard DNA curves showed that the amounts of *linA*, *linB*, and *linC* transcripts were in the range of  $0.5 \times 10^8$  to  $2.0 \times 10^8$  mean DNA equivalent copy numbers per microgram of RNA (Fig. 2A). The range measured for *linX* was slightly lower ( $1 \times 10^7$  to  $5 \times 10^7$ ) (data not shown). Despite standard deviations of, in some cases, 20% for individual spot measurements (calculated from hybridization intensities of different RNA dilutions on the same blot), independently repeated induction experiments resulted in the same relative transcript levels.

**Inducible expression of *linD* and *linE*.** In contrast to the expression of *linA*, *linB*, *linC*, and *linX*, that of *linD* and *linE* appeared to be inducible (Fig. 2B and C). The amount of transcripts while the cells were growing on glucose was only around or below  $10^7$  DNA equivalent copy numbers per  $\mu\text{g}$  of RNA (Fig. 2B and C), but a sharp increase of *linD* and *linE* transcripts became visible 10 and 5 min, respectively, after the addition of  $\gamma$ -HCH. The faster appearance of the *linE* transcript compared to *linD* mRNA might be due to the fact that it is the first gene of that polycistronic mRNA and the time required for RNA polymerase to reach *linD* is longer, which is similar to previous observations of other systems (9, 19, 21). The relative induction level for *linD* and *linE* with 7 mg of  $\gamma$ -HCH per liter compared to expression on glucose alone was about 10-fold higher (although the uninduced expression level could not be measured accurately due to the low transcript




FIG. 2. Amounts of *lin*-specific mRNAs before and after induction with  $\gamma$ -HCH (7 mg/liter). At time zero a single HCH dose was added to the chemostat culture of *S. paucimobilis* B90A. Panels show mRNAs probed with different antisense mRNAs, as indicated in the right top of each panel. The corresponding autoradiogram used for scanning is included within the graphs. The mRNA amounts are represented as mean DNA equivalent copy number (by comparing to a DNA standard, as explained in Materials and Methods) per microgram of RNA loaded in each spot. A different symbol is used for data from each of three independent induction experiments.




FIG. 3. Real-time PCR analysis of *linB* and *linD* gene expression in *S. paucimobilis* B90A. The mRNAs were purified from the induction experiment with  $\gamma$ -HCH (7 mg/liter) as described in the legend of Fig. 2. The time sequence is that described in the legend of Fig. 2. The y axis shows the threshold cycle number during which exponential amplification starts.

abundance). To more accurately measure induction of the *linD* transcript, we used qRT-PCR, which resulted in a value of the threshold cycle number which was six lower at the peak of mRNA abundance (Fig. 3). Theoretically, this would correspond to a 64-fold increase in induction (i.e.,  $2^6$ ). Both the dot blot hybridizations and qRT-PCR showed that abundance of the *linB* mRNA remained the same throughout the experiment. The 64-fold induction increase measured with qRT-PCR is similar to the magnitude of induction observed for the *tfdA*, *tfdC<sub>1</sub>*, and *tfdCD* mRNAs in *Ralstonia eutropha* after the addition of 10  $\mu$ M 2,4-dichlorophenoxyacetate (2,4-D) (19). Probably, since the uninduced level cannot be measured very accurately by dot blotting, the ratios for the *linE* and *linD* transcripts were underestimated by that technique. A slight difference in the transcript levels of *linE* and *linD* was calculated even though both mRNAs are part of the same original transcript, which might be due to different mRNA stabilities, as observed before for the *tfd* system in *R. eutropha* (19).

The increased level of *linD* and *linE* mRNA was only temporary and disappeared after 1 to 2 h, paralleling the degradation of  $\gamma$ -HCH (Fig. 4). This pulse-like transcript appearance is probably a reflection of the time course of the intracellular concentration of the inducing compound, which—by analogy to that shown for strain UT26 (21)—might be 2,5-dichlorohydroquinone, a metabolite formed during  $\gamma$ -HCH degradation. One can envision that upon first entry of  $\gamma$ -HCH in the cells, the metabolite is produced and accumulates due to constitutive LinA, LinB, and LinC activity (Fig. 1). Once formed, 2,5-dichlorohydroquinone will induce LinR-mediated expression from the *linE* promoter, which in turn will lead to formation of the dichlorohydroquinone metabolizing enzymes and breakdown of the inducer. If we assume that the rate of transcription from the *linE* promoter is directly related to the intracellular concentration of 2,5-dichlorohydroquinone, the net effect during  $\gamma$ -HCH degradation will be an initial sharp increase of *linE* and *linD* mRNAs, followed by a decrease. A comparison, for example, of the amount of the *linA*




FIG. 4. Disappearance of HCH isomers after pulsing to a continuous culture of *S. paucimobilis* B90A (depicted as the percentage of HCH remaining compared to the initial concentration). The washout curve depicts the theoretical dilution of HCH in the chemostat due to medium replenishment. Initial established concentrations were as follows:  $\alpha$ -HCH, 2 mg/liter;  $\beta$ -HCH, 5 mg/liter;  $\gamma$ -HCH, 7 and 0.7 mg/liter; and  $\delta$ -HCH, 20 mg/liter.

transcript and that of *linD* makes clear that the level of constitutive expression is similar to or only slightly less than the maximum expression level of the inducible *lin* genes reached with 7 mg of  $\gamma$ -HCH per liter. This might mean that metabolites from HCH are produced faster than they can be degraded by the inducible pathways. No significant fluctuations were observed for the *linR* transcript, but the general abundance of the *linR* mRNA was much lower than for the other *lin* genes (data not shown).

**Lin gene expression after exposure of strain B90A to other HCH isomers.** When the cells were exposed to a pulse of  $\alpha$ -HCH (2 mg/liter), a similar response was seen as for  $\gamma$ -HCH (Fig. 5). The transcript levels of all *lin* genes previously observed to be constitutively expressed were again similar irrespective of  $\alpha$ -HCH application to the chemostat (data not shown) and the same as after exposure to  $\gamma$ -HCH (i.e., around  $2 \times 10^8$  mean DNA equivalent copies per  $\mu\text{g}$  of RNA). Induction of *linE* and *linD* also took place with  $\alpha$ -HCH and with an induction factor of about 10-fold, although the measured *linD* transcript was slightly less abundant than with  $\gamma$ -HCH (maximum  $10^8$  mean DNA equivalent copies per  $\mu\text{g}$  of RNA). In-




FIG. 5. Induction of *linE*- and *linD*-specific mRNAs after addition of different HCHs. At time zero a single dose of HCH was added to the chemostat culture of *S. paucimobilis* B90A. (A and B)  $\alpha$ -HCH (filled symbols) and  $\delta$ -HCH (open symbols) at concentrations of 2 and 20 mg per liter, respectively. (C and D)  $\gamma$ -HCH (0.7 mg/liter) and  $\beta$ -HCH (5 mg/liter). The corresponding autoradiogram used for scanning is included within the graphs. The mRNA amounts are represented as the mean DNA equivalent copy number (by comparing to a DNA standard, as explained in Materials and Methods) per microgram of RNA loaded in each spot. Induction experiments with 0.7 mg of  $\gamma$ -HCH per liter were carried out twice; other experiments were carried out once.

duction and subsequent disappearance of the *linD* and *linE* transcripts followed the measured  $\alpha$ -HCH concentration in the chemostat (Fig. 4). This strongly suggests that  $\alpha$ -HCH is metabolized by *S. paucimobilis* B90A cells via a pathway similar to that of  $\gamma$ -HCH, giving rise to similar metabolites, which can induce the *linE* promoter (21). In contrast, no induction of *linD* and *linE* was observed in RNA isolated from cells exposed to  $\beta$ - and  $\delta$ -HCH isomers (Fig. 5), although their concentrations were of the same order as those of  $\alpha$ - and  $\gamma$ -HCH and both compounds were at least partially degraded by the cells (Fig. 4). This was surprising but indicates that  $\beta$ - and  $\delta$ -HCH are either partially degraded by LinA, LinB, and LinC activity without formation of the metabolite necessary to achieve *linED* induction or are degraded via a different metabolic pathway not involving the *lin* gene products. The disappearance of  $\beta$ - and  $\delta$ -HCH in the chemostat was rather different. Whereas  $\beta$ -HCH did not seem to be touched by the cells for about 1 h but then began disappearing at a higher rate (Fig. 4),  $\delta$ -HCH was only partially degraded for about 1 h, after which degradation stopped completely. This could be an indication for induction of a different degradation pathway in the case of  $\beta$ -HCH. The apparent complete stop of further  $\delta$ -HCH degradation after 1 h might have been due to the accumulation of a toxic intermediate. After exposure to both  $\beta$ - and  $\delta$ -HCH, the cells accumulated a metabolite in the culture medium, which was detectable by a gas chromatograph equipped with an electron capture detector but is not further identified here (data not shown). It could be that the higher concentration of  $\delta$ -HCH (20 mg/liter) resulted in higher concentrations of a  $\delta$ -HCH intermediate, which was toxic for the cells, similar to observations of 2,4-dichlorophenol in 2,4-D degradation (19). In other ongoing studies in our laboratories, we have observed that *S. paucimobilis* cultures exposed to  $\beta$ - and  $\delta$ -HCH release only half of all chlorine atoms of HCH as chloride (C. Holliger, unpublished data), which supports the idea that the limited degradation of  $\beta$ - and  $\delta$ -HCH in chemostat cultures of strain B90A did not lead to formation of the (di)chlorohydroquinones necessary for induction of the *linE* promoter.

**Induction of *linD* and *linE* in response to a low  $\gamma$ -HCH concentration.** Finally, it was determined whether *linD* and *linE* would still be induced at a 10-fold-lower  $\gamma$ -HCH concentration (0.7 mg/liter), which is indicative of what may happen at the low micromolar concentrations more often found in the environment. Although *linD* and *linE* genes were induced by  $\gamma$ -HCH at a concentration of 7 mg/liter (25  $\mu$ M), no measurable change in mRNA was observed upon exposing the cells to a concentration of 0.7 mg/liter (2.5  $\mu$ M). This was reproducibly observed (Fig. 5C and D), but since quantitation of individual spots resulted in standard deviations of up to 20%, very low mRNA bursts may have remained undetectable. Interestingly,  $\gamma$ -HCH was still degraded by the cells (Fig. 4), although we cannot conclude whether full mineralization took place on the basis of the gas chromatograph-electron capture detector measurements.

There is good evidence from other experimental systems to believe that such a low concentration (2.5  $\mu$ M) might be at or below a threshold for induction of the *linD* and *linE* genes and similarly for other catabolic genes in general. We conclude this from experimental studies on catabolic promoter response with reporter gene fusions, which independently discovered

the lowest detectable induction to take place in a concentration range of 1 to 10  $\mu$ M. For example, by using promoter fusions to the *luxAB* bacterial luciferase genes, the lowest response from the *tfdD<sub>1</sub>* promoter in *R. eutropha* JMP134 to 2,4-D was found to occur at 2  $\mu$ M (11). Similarly, the lowest inducer concentration for the *hbpC* promoter in *Pseudomonas azealica* was 9  $\mu$ M 2-hydroxybiphenyl (13), for the Po promoter in *Pseudomonas putida* was 3.2  $\mu$ M phenol (30), and for the *todX* promoter of *P. putida* F1 was 1  $\mu$ M toluene (2). It might be that a threshold for specific gene activation occurs at low substrate concentrations (around 1  $\mu$ M), because when most of the substrate is channeled through the metabolic pathway, only a small fraction of inducer is left to be detected by the regulatory protein. For example, a previous study calculated a diffusion rate of uncharged 2,4-D through the cytoplasmic membrane of *R. eutropha* at a 1  $\mu$ M outside concentration in the order of 100 molecules per s per cell (18), which, in fact, was mostly driven by the rate of 2,4-D metabolism.

**Orchestrating expression of the lindane degradation pathway.** In conclusion, the results of this work have shown that expression of the *lin* genes behaves rather differently than what is usually observed for catabolic pathways of, e.g., *Beta*- and *Gammaproteobacteria*. Probably due to the absence of regulatable promoters and a scattering of the different genes involved in the pathway, most of the *lin* genes are expressed to a constitutive level (i.e., *linA1* and/or *linA2*, *linB*, *linC*, and *linX*). Only the *linE* and *linD* genes, as observed before for *S. paucimobilis* UT26 (21), were inducibly expressed. The scattering of pathway genes is more commonly found in sphingomonads, for example, for the pentachlorophenol degradation pathway in *Sphingobium chlorophenolicum* ATCC 39723 (7), for the genes involved in the dioxin degradation pathway in *Sphingomonas* sp. strain RW1 (3), or for the catabolic plasmid of *Sphingomonas aromaticivorans* (26), and may reflect a recent acquisition of gene sequences (in which case regulatable expression may evolve in the end) or simply be a more common property for this class of bacteria. Constitutively expressing part of the *lin* genes does not provide an apparent physiological disadvantage for the cells, except perhaps at the highest  $\delta$ -HCH concentration, where a toxicity effect cannot be excluded. The present genetic organization is sufficiently advantageous, and *Sphingomonas* strains with related *lin* genes and expression characteristics have been isolated from all over the world (e.g., from Japan [29], India [27], and France [31]). On the contrary, it might be that by providing constitutive expression, the cells escape at least part of the phenomena of apparent noninduction at low substrate concentrations. An absence of measurable gene induction does not necessarily mean that complete degradation is brought to a halt. *S. paucimobilis* B90A cells could rely on small amounts of LinD and LinE enzymes that are present because of low constitutive gene expression. However, it is currently not known whether *S. paucimobilis* cells gain any energy, carbon, or reducing equivalents by cometabolizing  $\gamma$ - or  $\alpha$ -HCH partially, without inducing the lower pathway, or whether they will still completely metabolize HCH at concentrations below 2.5  $\mu$ M in the environment. Mineralization studies with radiolabeled  $\gamma$ -HCH and with  $\gamma$ -HCH concentrations in the microgram-per-liter range are on their way. *Pseudomonas* sp. strain B13 metabolized 3-chlorobenzoate even in the nanomolar range but with completely different

uptake and metabolic rates prevailing at the nanomolar range than at the micromolar range (32). Although this is a very relevant environmental question, it remains unclear which biological phenomena are taking place at very low pollutant concentrations (micromolar and lower) (6).

#### ACKNOWLEDGMENTS

The authors thank Christoph Werlen and Alexandra Bähler, EAWAG, Dübendorf, Switzerland, for helping with chemostat cultivation.

Part of this work was supported by grants under the Indo-Swiss Collaboration in Biotechnology Program.

#### REFERENCES

- Aiba, H., S. Adhya, and B. de Crombrughe. 1981. Evidence for two functional *gal* promoters in intact *Escherichia coli* cells. *J. Biol. Chem.* **256**:11905–11910.
- Applegate, B., C. Kelly, L. Lackey, J. McPherson, S. Kehrmeyer, F.-M. Menn, P. Bienkowski, and G. Saylor. 1997. *Pseudomonas putida* B2: a *tod-lux* bioluminescent reporter for toluene and trichloroethylene co-metabolism. *J. Ind. Microbiol.* **18**:4–9.
- Armengaud, J., B. Happe, and K. N. Timmis. 1998. Genetic analysis of dioxin dioxygenase of *Sphingomonas* sp. strain RW1: catabolic genes dispersed on the genome. *J. Bacteriol.* **180**:3954–3966.
- Baumann, B., M. Snozzi, A. J. B. Zehnder, and J. R. van der Meer. 1996. Dynamics of denitrification activity of *Paracoccus denitrificans* in continuous culture during aerobic-anaerobic changes. *J. Bacteriol.* **178**:4367–4374.
- Braga, A. M., T. Krauss, C. R. Reis dos Santos, and P. Mesquita de Souza. 2002. PCDD/F-contamination in a hexachlorocyclohexane waste site in Rio de Janeiro, Brazil. *Chemosphere* **46**:1329–1333.
- Button, D. K. 1998. Nutrient uptake by microorganisms according to kinetic parameters from theory as related to cytoarchitecture. *Microbiol. Mol. Biol. Rev.* **62**:636–645.
- Cai, M., and L. Xun. 2002. Organization and regulation of pentachlorophenol-degrading genes in *Sphingobium chlorophenolicum* ATCC 39723. *J. Bacteriol.* **184**:4672–4680.
- Dauert, S., G. Barrett, J. S. Feliciano, R. S. Shetty, S. Shrestha, and W. Smith-Spencer. 2000. Genetically engineered whole-cell sensing systems: coupling biological recognition with reporter genes. *Chem. Rev.* **100**:2705–2738.
- Dogra, C., V. Raina, R. Pal, M. Suar, S. Lal, K.-H. Gartemann, C. Holliger, J. R. van der Meer, and R. Lal. 2004. Organization of *lin* genes and IS6100 among different strains of hexachlorocyclohexane-degrading *Sphingomonas paucimobilis*: evidence for horizontal gene transfer. *J. Bacteriol.* **186**:2225–2235.
- Franklin, F. C. H., M. Bagdasarian, M. M. Bagdasarian, and K. N. Timmis. 1981. Molecular and functional analysis of the TOL plasmid pWW0 from *Pseudomonas putida* and cloning of genes for the entire regulated aromatic ring-cleavage pathway. *Proc. Natl. Acad. Sci. USA* **78**:7458–7462.
- Hay, A. G., J. F. Rice, B. M. Applegate, N. G. Bright, and G. S. Saylor. 2000. A bioluminescent whole-cell reporter for detection of 2,4-dichlorophenoxyacetic acid and 2,4-dichlorophenol in soil. *Appl. Environ. Microbiol.* **66**:4589–4594.
- Jaspers, M. C., A. Schmid, M. H. Sturme, D. A. Goslings, H. P. Kohler, and J. R. van der Meer. 2001. Transcriptional organization and dynamic expression of the *hbpCAD* genes, which encode the first three enzymes for 2-hydroxybiphenyl degradation in *Pseudomonas azelaica* HBP1. *J. Bacteriol.* **183**:270–279.
- Jaspers, M. C., W. A. Suske, A. Schmid, D. A. Goslings, H. P. Kohler, and J. R. van der Meer. 2000. HbpR, a new member of the XylR/DmpR subclass within the NtrC family of bacterial transcriptional activators, regulates expression of 2-hydroxybiphenyl metabolism in *Pseudomonas azelaica* HBP1. *J. Bacteriol.* **182**:405–417.
- Johri, A. K., M. Dua, D. Tuteja, R. Saxena, D. M. Saxena, and R. Lal. 1998. Degradation of  $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ -hexachlorocyclohexane by *Sphingomonas paucimobilis*. *Biotechnol. Lett.* **20**:885–887.
- Kumari, R., S. Subudhi, M. Suar, G. Dhingra, V. Raina, C. Dogra, S. Lal, C. Holliger, J. R. van der Meer, and R. Lal. 2002. Cloning and characterization of *lin* genes responsible for the degradation of hexachlorocyclohexane isomers in *Sphingomonas paucimobilis* strain B90. *Appl. Environ. Microbiol.* **68**:6021–6028.
- Kutz, F. W., P. H. Wood, and D. P. Bottimore. 1991. Organochlorine pesticides and polychlorinated biphenyls in human adipose tissue. *Rev. Environ. Contam. Toxicol.* **120**:1–82.
- Laemmli, C. M., J. H. J. Leveau, A. J. B. Zehnder, and J. R. van der Meer. 2000. Characterization of a second *tfd* gene cluster for chlorophenol and chlorocatechol metabolism on plasmid pJP4 in *Ralstonia eutropha* JMP134(pJP4). *J. Bacteriol.* **182**:4165–4172.
- Leveau, J. H., A. J. B. Zehnder, and J. R. van der Meer. 1998. The *tfdK* gene product facilitates uptake of 2,4-dichlorophenoxyacetate by *Ralstonia eutropha* JMP134(pJP4). *J. Bacteriol.* **180**:2237–2243.
- Leveau, J. H. J., F. König, H.-P. Fuchslin, C. Werlen, and J. R. van der Meer. 1999. Dynamics of multigene expression during catabolic adaptation of *Ralstonia eutropha* JMP134 (pJP4) to the herbicide 2,4-dichlorophenoxyacetate. *Mol. Microbiol.* **33**:396–406.
- Miyauchi, K., Y. Adachi, Y. Nagata, and M. Takagi. 1999. Cloning and sequencing of a novel *meta*-cleavage dioxygenase gene whose product is involved in degradation of  $\gamma$ -hexachlorocyclohexane in *Sphingomonas paucimobilis*. *J. Bacteriol.* **181**:6712–6719.
- Miyauchi, K., H. S. Lee, M. Fukuda, M. Takagi, and Y. Nagata. 2002. Cloning and characterization of *linR*, involved in regulation of the downstream pathway for  $\gamma$ -hexachlorocyclohexane degradation in *Sphingomonas paucimobilis* UT26. *Appl. Environ. Microbiol.* **68**:1803–1807.
- Nagata, Y., K. Miyauchi, and M. Takagi. 1999. Complete analysis of genes and enzymes for  $\gamma$ -hexachlorocyclohexane degradation in *Sphingomonas paucimobilis* UT26. *J. Ind. Microbiol. Biotechnol.* **23**:380–390.
- Nagata, Y., T. Nariya, R. Ohtomo, M. Fukuda, K. Yano, and M. Takagi. 1993. Cloning and sequencing of a dehalogenase gene encoding an enzyme with hydrolase activity involved in the degradation of  $\gamma$ -hexachlorocyclohexane in *Pseudomonas paucimobilis*. *J. Bacteriol.* **175**:6403–6410.
- Nagata, Y., R. Ohtomo, K. Miyauchi, M. Fukuda, K. Yano, and M. Takagi. 1994. Cloning and sequencing of a 2,5-dichloro-2,5-cyclohexadiene-1,4-diol dehydrogenase gene involved in the degradation of  $\gamma$ -hexachlorocyclohexane in *Pseudomonas paucimobilis*. *J. Bacteriol.* **176**:3117–3125.
- Oliveira, R. M., O. M. Brilhante, J. C. Moreira, and A. C. Miranda. 1995. Hexachlorocyclohexane contamination in urban areas of the south eastern region of Brazil. *Rev. Saude Publica* **29**:228–233. (In Portuguese.)
- Romine, M. F., L. C. Stillwell, K. K. Wong, S. J. Thurston, E. C. Sisk, C. Sensen, T. Gaasterland, J. K. Fredrickson, and J. D. Saffer. 1999. Complete sequence of a 184-kilobase catabolic plasmid from *Sphingomonas aromati-covorans* F199. *J. Bacteriol.* **181**:1585–1602.
- Sahu, S. K., K. K. Patnaik, M. Sharmila, and N. Sethunathan. 1990. Degradation of  $\alpha$ -,  $\beta$ -, and  $\gamma$ -hexachlorocyclohexane by a soil bacterium under aerobic conditions. *Appl. Environ. Microbiol.* **56**:3620–3622.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Senoo, K., and H. Wada. 1989. Isolation and identification of an aerobic  $\gamma$ -HCH decomposing bacterium from soil. *Soil Plant Nutr.* **35**:79–87.
- Shingler, V., and T. Moore. 1994. Sensing of aromatic compounds by the DmpR transcriptional activator of phenol-catabolizing *Pseudomonas* sp. strain CF600. *J. Bacteriol.* **176**:1555–1560.
- Thomas, J. C., F. Berger, M. Jacquier, D. Bernikson, F. Baud-Grasset, N. Truffault, P. Normand, T. M. Vogel, and P. Simonet. 1996. Isolation and characterization of a novel  $\gamma$ -hexachlorocyclohexane-degrading bacterium. *J. Bacteriol.* **178**:6049–6055.
- Tros, M. E., G. Schraa, and A. J. B. Zehnder. 1996. Transformation of low concentrations of 3-chlorobenzoate by *Pseudomonas* sp. strain B13: kinetics and residual concentrations. *Appl. Environ. Microbiol.* **62**:437–442.
- Walker, K., D. A. Vallerio, and R. G. Lewis. 1999. Factors influencing the distribution of lindane and other hexachlorocyclohexanes in the environment. *Environ. Sci. Technol.* **33**:4373–4378.
- Ware, G. W. 1989. *The pesticide book*, 3rd ed. Thomas Publications, Fresno, Calif.
- Windholz, M., S. Budavari, L. Y. Stroumstos, and M. N. Fertig (ed.). 1976. *The Merck index*, 9th ed. Merck & Co., Inc., Rahway, N.J.
- Zylstra, G. J., W. R. McCombie, D. T. Gibson, and B. A. Finette. 1988. Toluene degradation by *Pseudomonas putida* F1: genetic organization of the *tod* operon. *Appl. Environ. Microbiol.* **54**:1498–1503.