

## Detection and Selection of Microsatellites in the Genome of *Paracoccidioides brasiliensis* as Molecular Markers for Clinical and Epidemiological Studies

Érika Nascimento,<sup>1</sup> Roberto Martinez,<sup>2</sup> André Rodrigues Lopes,<sup>1</sup>  
Luciano Angelo de Souza Bernardes,<sup>1</sup> Carolina Pomponio Barco,<sup>3</sup>  
Maria Helena S. Goldman,<sup>4</sup> John W. Taylor,<sup>5</sup> Juan G. McEwen,<sup>6</sup>  
Marina Pasetto Nobrega,<sup>3</sup> Francisco G. Nobrega,<sup>3</sup>  
and Gustavo H. Goldman<sup>1\*</sup>

*Faculdade de Ciências Farmacêuticas de Ribeirão Preto,<sup>1</sup> Faculdade de Medicina de Ribeirão Preto,<sup>2</sup> and Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto,<sup>4</sup> Universidade de São Paulo, São Paulo, and Universidade do Vale do Paraíba, UNIVAP, Vale do Paraíba,<sup>3</sup> Brazil; Department of Plant and Microbial Biology, University of California, Berkeley, California<sup>5</sup>; and Corporacion para Investigaciones Biológicas and Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia<sup>6</sup>*

Received 5 April 2004/Returned for modification 13 June 2004/Accepted 5 July 2004

***Paracoccidioides brasiliensis*, a thermodimorphic fungus, is the causative agent of the prevalent systemic mycosis in Latin America, paracoccidioidomycosis (PCM). Here, we describe the microsatellite patterns observed in a collection of *P. brasiliensis* random sequence tags. We identified 1,117 microsatellite patterns in about 3.8 Mb of unique sequences (0.47% of the total DNA used in the analysis). The majority of these microsatellites (87.5%) are found in noncoding sequences. We used two polymorphic microsatellites located on noncoding and coding sequences, as well as two microsatellites located on introns, as molecular markers to discriminate *P. brasiliensis* isolates, to look for relationships between the genetic background of the strains and the types of human disease they cause. We did not observe any correlation between the clinical form of human PCM and four simple sequence repeat patterns analyzed.**

*Paracoccidioides brasiliensis*, a thermodimorphic fungus, is the causative agent of the prevalent systemic mycosis in Latin America, paracoccidioidomycosis (PCM). Epidemiological data indicate a broad geographic distribution of *P. brasiliensis* in Central and South America, from Mexico to Argentina (31). The pathogen apparently has its natural habitat in soil or in plants in areas where PCM is endemic, and rural workers appear to become infected by inhaling dust containing the infecting propagules (32). It is estimated that as many as 10 million individuals could be infected with *P. brasiliensis*, acquired by inhalation of airborne microconidia, which reach the pulmonary alveolar epithelium and transform into the parasitic yeast form (27). The human form of (PCM) caused by this fungus is characterized by a range of clinical manifestations from benign or asymptomatic forms to severe and disseminated disease that is often fatal. The development of PCM depends on interactions between fungus and host components. Many authors have tried to correlate certain characteristics of *P. brasiliensis* isolates with virulence without success (22, 37, 42, 44). In *P. brasiliensis*, restriction fragment length polymorphism and random amplified polymorphic DNA (RAPD) markers have been used in attempts to establish epidemiological and phylogenetic relationships between isolates that have

different degrees of virulence and that come from distinct geographical regions (3, 23, 25, 26). Although, it has been proposed that *P. brasiliensis* isolates differ in their ability to cause human disease (36), the issue is far from settled (25). Recently, Hebel-Barbosa et al. (14) completed the first genetic analysis of *P. brasiliensis* isolates from 10 armadillos and confirmed their similarity with 19 clinical isolates by DNA sequencing. These authors showed by sequence comparison of the internal transcribed spacer 1 and internal transcribed spacer 2 regions that eight isolates differed by one or three sites among the five polymorphic sites found, suggesting the existence of two genetic groups.

The main antigenic component described in *P. brasiliensis* is gp43, an exocellular glycoprotein containing a single oligosaccharide chain; it elicits a strong humoral response and can be detected in PCM patient serum (for a review, see reference 41). gp43 is a potential virulence factor because it binds murine laminin, resulting in increased pathogenicity of yeast cells (43). Morais et al. (24) reported *P. brasiliensis* gp43 gene polymorphism in a variety of *P. brasiliensis* isolates from patients suffering from chronic and acute PCM. These authors observed that the *P. brasiliensis* gp43 gene sequences of three isolates from patients with pulmonary or chronic PCM were phylogenetically distant from the sequences of other isolates. These results suggest a possible correlation between *P. brasiliensis* gp43 gene polymorphism and the degree of pathogenicity of these strains in the animal model.

Microsatellites or simple sequence repeats (SSRs) are tandemly repeated tracts of DNA composed of 1- to 6-bp-long

\* Corresponding author. Mailing address: Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Av. do Café S/N, CEP 14040-903, Ribeirão Preto, São Paulo, Brazil. Phone: 0055-016-6024280/81. Fax: 0055-016-6331092. E-mail: address: ggoldman@usp.br.

TABLE 1. *P. brasiliensis* isolates analyzed in this study from patients located in São Paulo and south of the Minas Gerais State, Brazil

Clinical isolate	Isolation date (mo/day/yr)	Classification	Material
Pb18			
Pb51	8/27/99	Chronic	Skin biopsy
Pb52	8/30/99	Acute/HIV	Skin biopsy
Pb61	9/24/00	Acute/HIV	Skin biopsy
Pb66	3/15/01	Chronic	Skin biopsy
Pb67	4/18/01	Acute/HIV	Mouth lesion
Pb71	8/08/01	Acute	Abscess
Pb72	7/04/01	Chronic	Skin biopsy
Pb74	7/30/01	Chronic	Bronchoalveolar lavage
Pb76	1/11/02	Acute/HIV	Duodenal biopsy
Pb78	1/15/02	Acute/HIV	Sputum
Pb79	2/12/02	Chronic	Sputum
Pb80	9/27/01	Acute	Skin biopsy
Pb82	11/29/01	Chronic	Lymph node
Pb85	9/25/01	Acute	Abscess
Pb89	3/28/02	Acute	Blood
Pb91	9/24/02	Chronic	Sputum
Pb92	11/21/02	Acute	Blood
Pb93	12/02/02	Chronic	Peritoneal fluid
PbIbiá	6/97		Soil from Ibiá region, Minas Gerais State
PbT1	6/99		Armadillo from Ibiá region, Minas Gerais State
PbT2	6/99		Armadillo from Ibiá region, Minas Gerais State
PbT3	6/99		Armadillo from Ibiá region, Minas Gerais State

units. They are omnipresent in prokaryotes and eukaryotes, even in the smallest bacterial genomes, and are found anywhere in the genome in both protein-encoding and noncoding regions (40). SSRs are considered to be evolutionarily neutral DNA markers (20). Length polymorphism arises from variations in the number of repeated units, probably due to DNA polymerase slippage during the replication of SSRs (19). They have been used for both population genetics and typing studies because they have several advantages as markers, such as that they are highly polymorphic, multiallelic, highly reproducible, and detectable by PCR (29).

Recently, we established a collection of about 3.8 Mb of

unique random sequence tags (RSTs) (M. P. Nobrega et al., unpublished data) and decided to investigate the microsatellite occurrence in this set. Here, we describe the microsatellite patterns observed and use some of them as molecular markers to discriminate *P. brasiliensis* isolates in a search for correlations between the genetic background of the strains and the types of human disease they cause.

## MATERIALS AND METHODS

**Fungal strains and DNA preparation.** We analyzed 23 isolates (4 environmental isolates, 18 clinical isolates, and the Pb18 isolate) (Pb18 was kindly provided by Z. P. Camargo, Universidade Federal de São Paulo, Brazil) that are listed in Table 1. Yeast cells were grown to the logarithmic phase in 125-ml Erlenmeyer flasks containing 25 ml of Fava-Neto's medium as previously described (35) at 37°C with constant shaking. DNA was prepared according to a glass bead protocol (33).

**PCRs.** Using Primer Express design software, version 1.0 (Applied Biosystems), we designed PCR primers for amplifying each DNA fragment that contains microsatellites. The 40- $\mu$ l amplification mixture included 1 $\times$  *Taq* DNA platinum buffer (Invitrogen), 0.5  $\mu$ M of each primer (Table 2), a 0.2 mM deoxynucleotide triphosphate mixture, 2.5 U of *Taq* DNA platinum polymerase (Invitrogen), and 100 ng of genomic DNA. PCR amplification was carried out with a PTC100 96-well thermal cycler (MJ Research) at 95°C for 1 min; for 38 cycles at 95°C for 1 min, 39 to 54.4°C (depending on the fragment) for 1 min, and 72°C for 1 min; and followed by an extension step at 72°C for 10 min. After the reaction, the PCR products were purified with a QIAGEN PCR cleanup kit, following the manufacturer's instructions. Sequencing reactions were prepared with the BigDye Terminator Cycle Sequencing kit (Applied Biosystems), with the primers listed in Table 2. The nucleotide sequences in both strands were determined by primer elongation with an ABI3100 automated DNA sequencer (Applied Biosystems).

**Data handling and analysis.** A pipeline was built to analyze and assemble the *P. brasiliensis* RST sequences. Sequences were automatically edited for each RST with the programs Phred-Phrap (4, 5), Consed (12), and Crossmatch from Phrap (13). The sequences were cleaned from the pUC18 vector sequences with Crossmatch (13); RSTs with a quality value of at least 20 were considered for further analysis. Edited sequences were clustered with the Phrap program (5). To identify if the microsatellite was at either a coding or noncoding region, the *P. brasiliensis* clusters containing microsatellites were compared with the BLASTX and BLASTN algorithms (1) with the the National Center for Biotechnology Information (NCBI) nonredundant database (<http://ncbi.nlm.nih.gov/>), several fungal genome databases ([www.broad.mit.edu](http://www.broad.mit.edu)), and the *P. brasiliensis* expressed sequence tag project databases (<http://143.107.203.68/est/default.html>). When *E* values greater than  $10^{-5}$  were obtained, they were considered not statistically significant (no significant match).

We have extracted all the microsatellites from our RST databank (Table 3) by using the methodology described by Jurka and Pethiyagoda (16). Essentially, we extracted only simple repeats composed of tandemly repeated basic units 1 to 6 nucleotides (nt) long. Most simple repeats and their complementary counter-

TABLE 2. List of primers used in this work

Primer	Sequence	Amplified SSR (bp)	NCBI accession no.
MS5	5'-TGCCCGAAGCAGCCCCCGGG-3'	ATTT (236)	BQ503230
MS6	5'-GAGAAAGTGAGTTGGTTTACG-3'		
MS11	5'-TTTGCTACACTTCCCTCTCCC-3'	AT (410)	CL524685
MS12	5'-CTTCCCCATTCTGATTCTCG-3'		
MG13	5'-CACGTGTCAAGTCATAATAAATAG-3'	AT, ATTT (251)	CL524686
MG14	5'-AATCTGCTGCCAATAGTCAT-3'		
MG15	5'-GCACAGACGCAAAATATGC-3'	TAAA (152)	CL524687
MG16	5'-GGTGGAAAAAGATATGCGAA-3'		
MG21	5'-ACAATCAAACCGGGAG-3'	AAAAGG (214)	CL524688
MG22	5'-GGAGGGATAGGAACGAATT-3'		
MGI125	5'-TCAAACCTGACAACCTCAGCC-3'	TCA (203)	CL524689
MGI126	5'-TAAGAAGATGGATGAGGCC-3'		
MGI127	5'-AAAAATAACTACGCAGACGC-3'	CCCA (251)	CL524690
MGI128	5'-TTAAGCTGGGCTTTGGGTAC-3'		

TABLE 3. Summary of the data for 67 microsatellite patterns with two occurrences and with two or more sizes

Msat <sup>a</sup>	Maximum <sup>b</sup>	Mean <sup>c</sup>	SD	Frequency <sup>d</sup>	Repeats <sup>e</sup>	Total length <sup>f</sup>	Abundance (%) <sup>g</sup>
A	39	17.1	6.0	149	23	2,552	14.1
C	20	14.2	1.9	34	7	483	2.6
AC	74	25.0	14.1	28	14	702	3.8
AG	68	20.2	11.2	32	12	648	3.5
AT	40	17.8	6.1	110	12	1,958	10.8
ACG	24	17	4.4	6	5	102	0.5
CAA	39	17.3	6.6	23	6	399	2.2
CCA	21	14.8	2.4	17	4	252	1.3
CTA	36	24	12	2	2	48	0.2
GAA	54	20	12.3	23	8	462	2.5
GCA	18	13.3	2.2	24	3	321	1.7
GCC	15	13.5	1.5	4	2	54	0.3
GGA	18	13.6	1.9	11	3	150	0.8
TAA	108	31.2	26.5	30	14	936	5.1
TCA	51	18.8	9.0	22	6	414	2.2
AACC	32	14.4	5.9	10	3	144	0.8
AACT	20	12.7	1.9	21	3	268	1.4
ATGC	20	13.3	2.9	6	2	80	0.4
CAAA	24	21.3	3.7	3	2	64	0.3
GAAA	44	14.3	6.4	24	3	344	1.9
GACA	40	24	11.7	3	3	72	0.4
GCCA	24	15	5.1	4	2	60	0.3
GGGA	20	14	3.4	4	2	56	0.3
TAAA	36	16.3	6.7	24	6	392	2.1
TACA	20	13.6	2.6	10	3	136	0.7
TCAA	16	12.8	1.6	5	2	64	0.3
TCCA	20	15	3.3	4	3	60	0.3
TGAA	20	14.6	3.7	3	2	44	0.2
ACAGC	20	17.5	2.5	2	2	35	0.1
ATGCC	30	22.5	7.5	2	2	45	0.2
CAAAA	30	22.5	7.5	2	2	45	0.2
GAAAA	40	19.8	6.4	25	5	495	2.7
TAAAA	35	18	5.0	15	3	270	1.4
TTAAA	20	17.5	2.5	2	2	35	0.1
AAAAGG	42	15.3	9.4	9	2	138	0.7
AAAATC	24	15	5.1	4	2	60	0.3
AAAATT	18	14	2.8	3	2	42	0.2
AAAGAG	30	14.5	6.2	7	2	102	0.5
AAAGGG	18	15	3	2	2	30	0.1
AAATAG	18	15	3	2	2	30	0.1
AAATAT	18	13.5	2.5	4	2	54	0.3
AAATGG	18	13	2.2	6	2	78	0.4
AACAGC	18	13.7	2.7	7	2	96	0.5
AACCAC	30	19.5	7.7	4	3	78	0.4
AACCGC	24	15	5.1	4	2	60	0.3
AACCTC	30	24	6	2	2	48	0.2
AACGAC	18	14	2.8	3	2	42	0.2
AAGATG	18	14	2.8	3	2	42	0.2
AATAAC	18	13.5	2.5	4	2	54	0.3
AATATT	18	14	2.8	3	2	42	0.2
AATCTC	18	13.2	2.4	5	2	66	0.3
AATTAT	24	16	5.6	3	2	48	0.2
ACACCC	18	14	2.8	3	2	42	0.2
ACACTC	24	18	6	2	2	36	0.2
ACCGGG	18	15	3	2	2	30	0.1
ACGAGG	18	15	3	2	2	30	0.1
ACTGCT	30	20	7.4	3	3	60	0.3
ATATCC	24	20	2.8	3	2	60	0.3
ATATGC	18	15	3	2	2	30	0.1
ATCCAC	18	12.8	2.0	7	2	90	0.5
ATCCGC	24	15	5.1	4	2	60	0.3
ATCGCC	18	15	3	2	2	30	0.1
ATCTCC	18	14	2.8	3	2	42	0.2
CCCCAA	36	20.4	8.9	5	4	102	0.5
GAAAAA	24	12.6	2.4	27	3	342	1.8
TAAAAA	30	14.8	5.3	25	4	372	2.0
TATACA	18	14.4	2.9	5	2	72	0.4

<sup>a</sup> Of a total of 501 patterns, 58 have two occurrences and a single size, 101 have a single occurrence, and 275 patterns were not detected. Msat, type of basic pattern in the simple repeat.

<sup>b</sup> Maximum length of segments.

<sup>c</sup> Average length of segments.

<sup>d</sup> The number of times the pattern appears independently of the length of the size of the segment.

<sup>e</sup> Number of different length ranges ( $N_e$ ) found among repeats longer than 12 nt. For example, CAAA and its complement are represented by two fragments ( $N_e = 2$ ): one is 12 nt long and the other is 24 nt long. (AC)<sub>n</sub> is present in 14 different length fragments ( $N_e = 14$ ). Only patterns with  $N \geq 2$  are shown.

<sup>f</sup> Total length of repeat units over 12 nt long.

<sup>g</sup> Relative abundance of repeats over 12 nt long.

TABLE 4. Main features of seven clusters chosen for microsatellite characterization

Microsatellite primer pair <sup>a</sup>	Microsatellite pattern (bp) <sup>b</sup>	No. of repeats at microsatellite locus <sup>c</sup>	Microsatellite features <sup>d</sup>
MS5XMS6	TAAA (236)	4–11	Coding region; NP_587920.1, probable 6-phosphogluconolactonase <i>Schizosaccharomyces pombe</i> ; 3e-64/52/66
MS11XMS12	AT (410)	10–14	Noncoding region
MG13XMG14	AT, ATTT, AT (251)	9–19, 1–6, 2–20	Located in a Noncoding region
MG15XMG16	TAAA (152)	3–6	Noncoding region
MG21XMG22	AAAAGG (214)	1–8	coding region; EAA60777 hypothetical protein AN4735.2, <i>Aspergillus nidulans</i> ; 5e-14/70/88
MGI25XMG126	TCA (203)	4–5	Intron region; EAA61572.1 hypothetical protein AN7784.2, <i>Aspergillus nidulans</i> ; 5e-19/50/62
MGI27XMG128	CCCA (251)	3	Intron region; EAA28067.1 hypothetical protein, <i>Neurospora crassa</i> ; 5e-36/61/66

<sup>a</sup> Each primer pair is shown with an X linking the two primers (e.g., MS5 and MS6).

<sup>b</sup> Size of DNA fragment amplified by PCR in strain Pb18.

<sup>c</sup> Range in the number of simple repeats in the clinical and environmental isolates listed in Table 1.

<sup>d</sup> Features are listed as follows: microsatellite location; NCBI accession number, homologue; *E*-value/% identity/% similarity.

parts can be represented by several different basic unit patterns. For example, the pattern (GCC)<sub>n</sub>, listed by its unit name GCC in Table 3, also represents (CCG)<sub>n</sub>, (CGC)<sub>n</sub>, (GGC)<sub>n</sub>, (GCG)<sub>n</sub>, and (CGG)<sub>n</sub>. Each simple sequence was counted on one strand only and consequently the length is given by the number of nucleotides. Furthermore, whenever tandemly repeated patterns with different unit sizes were identical, they were listed under the smallest unit size. For example, patterns like (ACACAC)<sub>n</sub> or (ACAC)<sub>n</sub> were included into the category (AC)<sub>n</sub>. As a result, the total number of theoretically possible, nonoverlapping patterns was reduced to 501 (2 monomeric, 4 dimeric, 10 trimeric, 33 tetrameric, 102 pentameric, and 350 hexameric patterns).

Phylogenetic analysis was carried out with the MEGA2 (Molecular Evolutionary Genetics Analysis) software, version 2.1 (<http://www.megasoftware.net>; 18). The SSR sequences were aligned and the dendrogram was determined by using the ClustalX and the neighbor-joining method, respectively (30, 39). A bootstrap analysis (6) was performed (for 1,025 repeats) to evaluate the topology of the phylogenetic tree.

## RESULTS

**Identification of microsatellites.** We used the methodology described by Jurka and Pethiyagoda (16) to identify microsatellite patterns in a collection of 6,689 clusters of RSTs that correspond to 3.8 Mb (Nobrega et al., unpublished; <http://143.107.203.68/rstpb/frame2.htm>). We selected repeat segments over 12 nt long and represented by more than a length size in the *P. brasiliensis* Pb18 isolate RST database. The SSR patterns range from 1 to 6 nt. The SSRs represent about 0.47% of the total DNA in the analysis. Of 501 possible types of simple repeats, only 125 met these criteria. Table 3 shows 67 of these microsatellite patterns that displayed two occurrences and two or more different sizes.

According to Jurka and Pethiyagoda (16), a possible indicator of sequence variability in microsatellites is the maximal observed lengths per given microsatellite pattern (Table 3, columns “Frequency” and “Repeats”). These numbers must also be used separately for each group of patterns (monomeric, dimeric, etc.) to reduce the overall impact of sample sizes on their absolute values. In Table 3, the column “Abundance” is equivalent to the column “Total length,” as the former lists the proportions of each individual pattern that are >12 nt long relative to the sum of all pattern lengths of ≥12 nt listed in the column “Total length.” Overall, the most abundant are hexanucleotide repeats representing 29.4% of the total simple repeats, followed by dinucleotide (18.3%), trinucleotide (17.3%), mononucleotide (16.8%), tetranucleotide (11.8%),

and pentanucleotide (6.4%) repeats. Noticeably, pentanucleotide repeats are underrepresented in our database when compared to the others. As can be seen from Table 3 (columns “Frequency” and “Repeats”), TAA repeats are probably more polymorphic than GCC repeats because they have 108 nt, as opposed to 15 nt (the longest observable length), and 14, as opposed to 2, types of length size. The same can be observed with TAAA when compared to TCAA, where 36 nt as opposed to 16 nt is the longest observable length, and there are six, as opposed to two, types of length size.

Comparison of the genomic sequence that comprises a specific microsatellite plus 200 bp upstream and downstream from the microsatellite, against a collection of 4,692 *P. brasiliensis* expressed sequence tags (11), the NCBI databank (<http://ncbi.nlm.nih.gov/>), and fungal genome databases ([www.broad.mit.edu](http://www.broad.mit.edu)) allowed us to summarize the genomic distribution of the *P. brasiliensis* microsatellites (Table 4). There are 887 clusters with microsatellites and 1,117 clusters with more than one SSR, because some sequences carry more than one microsatellite pattern (Table 4). Most of these microsatellite patterns (87.5%) are found in noncoding sequences, 10.9% are found in coding sequences, 1.3% are found in intron sequences, and 0.3% are located in transposons (Table 4).

**Characterization of microsatellite loci.** Based mainly on the criterion of sequence variability proposed above, we chose seven clusters that contained microsatellites to analyze their degrees of polymorphism. Two of the corresponding microsatellite patterns were located on coding sequences (TAAA and AAAAGG), three were on noncoding sequences (AT, AT/ATTT/AT, and TAAA), and two were in intron sequences (TCA and CCCA) (Table 4). We designed oligonucleotide primers (Table 2) based on the sequences that limit these microsatellite patterns and PCR amplified the DNA fragments. Size distribution and sequence analysis of several clinical and environmental isolates (Table 1) revealed that clusters MG13XMG14 (consisting of MG13 and MG14) and MG21XMG22 have the most polymorphic microsatellite patterns (Fig. 1 and Table 5). The MG13XMG14 cluster has three different microsatellite patterns (AT, ATTT, and another AT) that can range from 6 to 19, 1 to 6, and 2 to 20 repeats, respectively (Table 5). The MG21XMG22 cluster has an




FIG. 1. Agarose gel (2.0%) showing the PCR amplification products of the corresponding DNA fragments from microsatellite primer pair MG13XMG14. The molecular weights indicated at the left refer to the PCR product amplified from strain Pb18 (A) and primer pair MG21XMG22 (B). The numbers and letters refer to the different isolates described in Table 1. I, PbIbiá; T1, PbT1; and T2, PbT2.

AAAAGG simple repeat that can range from 1 to 8 repeats (Tables 4 and 5).

**Use of microsatellites as markers to evaluate correlation between genetic background of strains and virulence degree.** We have used these two most polymorphic clusters (MG13XMG14 and MG21XMG22) as well as the DNA sequences of the two clusters that have microsatellite patterns in the introns (MGI25XMG126 and MGI27XMG128) to evaluate relatedness among clinical and environmental isolates of *P. brasiliensis*. The clinical isolates were different from each other in the type of human disease they caused: 8 caused chronic disease, and 10 caused acute disease (Table 1). In addition, four iso-

lates were environmental isolates: three were isolated from armadillos, and one was isolated from soil (Table 1). Two phylogenetic trees were constructed by the neighbor-joining method based on the number of repeats obtained by PCR amplification of the microsatellites MG13XMG14 and MG21XMG22 (Fig. 2). All isolates had different multilocus genotypes, but there was no clustering of isolates associated with chronic or acute disease or with the environment. The number of repeats was randomly distributed among clinical versus non-clinical isolates (Table 5). The reliability of the phylogenetic trees inferred was verified by the bootstrap method (Fig. 2). In the first tree (MG13XMG14) (Fig. 2A), two main groups were identified. (i) In the first group, three environmental isolates (PbIbiá, PbT1, and PbT3) clustered in two different subgroups (A and B). (ii) In the second group, there are seven subgroups (C to I) which have not shown clustering between isolates that cause chronic or acute disease. Furthermore, the fourth environmental isolate in the second group clustered with isolates that cause chronic or acute disease (subgroup H). In the second tree (MG21XMG22) (Fig. 2B), two main groups were also observed. (i) In the first group, one environmental isolate, an isolate causing chronic disease, and an isolate causing acute disease (PbIbiá, Pb51, and Pb85) clustered into two different subgroups (A and B). (ii) In the second group, there are seven subgroups (C to G) which have not shown clustering between isolates that cause chronic or acute disease. Comparable results were observed for the microsatellite clusters MGI25XMG126 and MGI27XMG128 (data not shown). Taken together, these results suggest there is no clear correlation between the genetic background of the isolates, as measured here, and the types of human disease they cause.



TABLE 5. Repeats in microsatellites MG13XMG14 and MG21XMG22 of *P. brasiliensis*

Isolate	No. of repeats in:			
	MG13XMG14			MG21XMG22 <sup>a</sup>
	AT	ATTT	AT	AAAAGG
Pb18	10	6	14	7
Pb51	15	2	16	2
Pb52	14	4	16	7
Pb61	18	3	20	8
Pb66	10	6	14	ND
Pb67	18	3	16	7
Pb71	15	4	16	7
Pb72	15	4	13	7
Pb74	11	6	14	7
Pb76	13	4	16	7
Pb78	16	4	16	7
Pb79	17	3	16	ND
Pb80	11	1	2	5
Pb82	16	4	17	ND
Pb85	17	3	16	1
Pb89	15	4	16	ND
Pb91	19	2	17	3
Pb92	18	3	16	7
Pb93	19	3	17	2
PbIbiá	17	4	16	3
PbT1	15	4	14	6
PbT2	19	1	2	7
PbT3	9	1	2	7

<sup>a</sup> ND, not determined.

**DISCUSSION**

The main objective of this study was to develop microsatellite or SSR markers for the pathogenic dimorphic fungus *P. brasiliensis*. To identify SSR markers, we digitally screened an RST collection composed of 3.8 Mb of unique sequences, which should represent about 15 to 20% of the *P. brasiliensis* genome. SSR markers have been successfully applied for typ-



ing, mapping, and population studies of several fungal species, such as *Saccharomyces cerevisiae*, *Candida* spp., *Aspergillus fumigatus*, *Magnaporthe grisea*, *Coccidioides* spp., and *Histoplasma capsulatum* (2, 7–10, 15, 17, 21, 28, 34, 38). SSRs constitute a rather large fraction of noncoding DNA and are relatively rare in protein-encoding regions (20). Accordingly, most of the observed *P. brasiliensis* SSRs are located in such noncoding regions. The majority of SSRs found in many species are dinucleotides; in primates, mononucleotides [mainly poly(A-T) tracts] are the most copious classes of SSRs (20). We have observed that the most abundant SSRs in *P. brasiliensis* are hexanucleotides (about 29%). More fungal genomes are being sequenced (see the White Paper Initiative at [www.wi.mit.edu](http://www.wi.mit.edu)), which will provide more information about the type, length, and frequency of SSRs in fungi.

*P. brasiliensis* usually reaches the human host through the respiratory route by inhalation of airborne mycelial propagules that convert to the tissue yeast form, initiating infection; with time, infection may give rise to clinical PCM, a disease that may adopt different clinical forms (31). Once established, the disease may be acute or chronic, depending on the severity and localization of lesions. Variations in the intensity, extension, dissemination, and characteristics of the lesions in PCM will occur in a given patient depending on fungal virulence, fluctuations of the host defense mechanisms, and environmental factors (23, 32). In the present work, we evaluate SSR profiles as genetically associated elements with the potential to discriminate *P. brasiliensis* isolates according to their degree of virulence as determined from the corresponding clinical forms of patients with PCM. In typing with SSRs, an entire stretch of sequence is surveyed for genetic variation through length polymorphisms. Recently, Calcagno et al. (3) using RAPD analysis demonstrated that genetic differentiation could be associated with geographical region but not with different clinical manifestations of human PCM. Motta et al. (25) found no correlation between RAPD patterns and the type of pathology as observed with experimental infection in mice or in the clinical form of human PCM. In contrast with these authors, Molinari-Madlum et al. (23) have shown that RAPD patterns correlated with the virulence degree of *P. brasiliensis* isolates. Nevertheless, our results are similar to those of Motta et al. (25) and Calcagno et al. (3); we observed no correlation between the clinical form of human PCM and four SSR patterns. Interestingly, the *P. brasiliensis* isolates derived from AIDS patients (in this work, these isolates are classified as isolates that cause acute disease) have also not clustered, suggesting the possibility that any *P. brasiliensis* isolate could behave as an opportunistic pathogen in immunocompromised HIV patients.

The *P. brasiliensis* SSRs now available will provide new opportunities for epidemiological and phylogenetic studies of this organism.

#### ACKNOWLEDGMENTS

We thank the following agencies for their financial support: Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), both of Brazil (to M.H.S.G., M.P.N., F.G.N., and G.H.G.), and NIH Fogarty International Center grant R03TW001308 (to J.W.T. and J.G.M.).

We also thank Leila M. Toffoli for expert technical help and Diógenes Custódio Duarte Ribeiro for help with the bioinformatics.

#### REFERENCES

- Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Bart-Delabesse, E., J. F. Humbert, E. Delabesse, and S. Bretagne. 1998. Microsatellite markers for typing *Aspergillus fumigatus* isolates. *J. Clin. Microbiol.* **36**:2413–2418.
- Calcagno, A. M., G. Niño-Vega, F. San-Blas, and G. San-Blas. 1998. Geographic discrimination of *Paracoccidioides brasiliensis* strains by randomly amplified polymorphic DNA analysis. *J. Clin. Microbiol.* **36**:1733–1736.
- Ewing, B., L. Hillier, M. C. Wendl, and P. Green. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* **8**:175–185.
- Ewing, B., and P. Green. 1998. Base calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res.* **8**:186–194.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**:783–791.
- Fisher, M. C., G. Koenig, T. J. White, and J. W. Taylor. 2000. A test for concordance between the multilocus genealogies of genes and microsatellites in the pathogenic fungus *Coccidioides immitis*. *Mol. Biol. Evol.* **17**:1164–1174.
- Fisher, M. C., G. L. Koenig, T. J. White, G. San-Blas, R. Negroni, I. G. Alvarez, B. Wanke, and J. W. Taylor. 2001. Biogeographic range expansion into South America by *Coccidioides immitis* mirrors New World patterns of human migration. *Proc. Natl. Acad. Sci. USA* **98**:4558–4562.
- Fisher, M. C., G. L. Koenig, T. J. White, and J. W. Taylor. 2002. Molecular and phenotypic description of *Coccidioides posadasii* sp. nov., previously recognized as the non-California population of *Coccidioides immitis*. *Mycologia* **94**:73–84.
- Fisher, M. C., B. Rannala, V. Chaturvedi, and J. W. Taylor. 2002. Disease surveillance in recombining pathogens: multilocus genotypes identify sources of *Coccidioides* infections. *Proc. Natl. Acad. Sci. USA* **99**:9067–9071.
- Goldman, G. H., E. d. R. Marques, D. C. Ribeiro, L. A. de Souza Bernardes, A. C. Quiapin, P. M. Vitorelli, M. Savoldi, C. P. Semighini, R. C. de Oliveira, L. R. Nunes, L. R. Travassos, R. Puccia, W. L. Batista, L. E. Ferreira, J. C. Moreira, A. P. Bogossian, F. Tekaia, M. P. Nobrega, F. G. Nobrega, and M. H. S. Goldman. 2003. Expressed sequence tag analysis of the human pathogen *Paracoccidioides brasiliensis* yeast phase: identification of putative homologues of *Candida albicans* virulence and pathogenicity. *Eukaryot. Cell* **2**:34–48.
- Gordon, D., C. Abajian, and P. Green. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* **8**:195–202.
- Green, P. 1996. Phred, Phrap, Consed. <http://bozeman.mbt.washington.edu/phredphrapconsed.html>.
- Hebeler-Barbosa, F., F. V. Morais, M. R. Montenegro, E. E. Kuramae, B. Montes, J. G. McEwen, E. Bagagli, and R. Puccia. 2003. Comparison of the sequences of the internal transcribed spacer regions and *PbGP43* genes of *Paracoccidioides brasiliensis* from patients and armadillos (*Dasypus novemcinctus*). *J. Clin. Microbiol.* **41**:5735–5737.
- Hennequin, C., A. Thierry, G. F. Richard, G. Lecointre, H. V. Nguyen, C. Gaillardin, and B. Dujon. 2001. Microsatellite typing as a new tool for identification of *Saccharomyces cerevisiae* strains. *J. Clin. Microbiol.* **39**:551–559.
- Jurka, J., and C. Pethiyagoda. 1995. Simple repetitive DNA sequences from primates: compilation and analysis. *J. Mol. Evol.* **40**:120–126.
- Kaye, C., J. Milazzo, S. Rozenfeld, M.-H. Lebrun, and D. Tarreau. 2003. The development of simple sequence repeat markers for *Magnaporthe grisea* and their integration into an established genetic linkage map. *Fungal Genet. Biol.* **40**:207–214.

FIG. 2. Phylogenetic trees based on the MG13XMG14 (A) and MG21XMG22 (B) microsatellite sequences show the relationships of the 23 *P. brasiliensis* isolates. These unrooted trees were constructed by the neighbor-joining method. Topology was also evaluated by bootstrap analysis (MEGA2 program; 1,025 repeats). The numerical values in the trees represent bootstrap results. The distance between two strains is the sum of the branch lengths between them. The identified subgroups are indicated by the letters at the right.

18. Kumar, S., K. Tamura, I. B. Jakobsen, and M. Nei. 2001. MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics* **17**:1244–1245.
19. Levinson, G., and G. A. Gutman. 1987. Slipped-strand mispairing: a major mechanism for DNA sequence evolution. *Mol. Biol. Evol.* **4**:203–221.
20. Li, Y.-C., A. B. Korol, T. Fahima, A. Beiles, and E. Nevo. 2002. Microsatellites: genomic distribution, putative functions and mutational mechanism: a review. *Mol. Ecol.* **11**:2453–2465.
21. Lunel, F. V., L. Licciardello, S. Stefani, H. A. Verbrugh, W. J. Melchers, J. F. Meis, S. Scherer, and A. van Belkum. 1998. Lack of consistent short sequence repeat polymorphisms in genetically homologous colonizing and invasive *Candida albicans* strains. *J. Bacteriol.* **180**:3771–3778.
22. Manocha, M. S., G. San-Blas, and S. Centeno. 1980. Lipid composition of *Paracoccidioides brasiliensis*: possible correlation with virulence of different strains. *J. Gen. Microbiol.* **117**:147–154.
23. Molinari-Madlum, E. E. W. I., M. S. S. Felipe, and C. M. A. Soares. 2000. Virulence of *Paracoccidioides brasiliensis* isolates can be correlated to groups by random amplified polymorphic DNA analysis. *Med. Mycol.* **37**:269–276.
24. Moraes, F. V., T. F. Barros, M. K. Fukada, P. S. Cisalpino, and R. Puccia. 2000. Polymorphism in the gene coding for the immunodominant antigen gp43 from the pathogenic fungus *Paracoccidioides brasiliensis*. *J. Clin. Microbiol.* **38**:3960–3966.
25. Motta, T. R., C. A. Moreira-Filho, R. P. Mendes, L. R. Souza, M. F. Sugizak, S. Baueb, V. L. G. Calich, and C. A. C. Vaz. 2002. Evaluation of DNA polymorphisms amplified by arbitrary primers (RAPD) as genetically associated elements to differentiate virulent and non-virulent *Paracoccidioides brasiliensis* isolates. *FEMS Immunol. Med. Microbiol.* **33**:151–157.
26. Niño-Vega, G. A., A. M. Calcagno, G. San-Blas, F. San-Blas, G. W. Gooday, and N. A. R. Gow. 2000. RFLP analysis reveals marked geographical isolation between strains of *Paracoccidioides brasiliensis*. *Med. Mycol.* **38**:437–441.
27. Restrepo, A., J. G. McEwen, and E. Castañeda. 2001. The habitat of *Paracoccidioides brasiliensis*: how far from solving the riddle? *Med. Mycol.* **39**:233–241.
28. Rosehart, K., M. H. Richards, and M. J. Bidochka. 2002. Microsatellite analysis of environmental and clinical isolates of the opportunist fungal pathogen *Aspergillus fumigatus*. *J. Med. Microbiol.* **51**:1128–1134.
29. Saghai Maroof, M. A., R. M. Biyashev, G. Yang, Q. Zhang, and R. W. Allard. 1994. Extraordinarily polymorphic microsatellite DNA in barley: species diversity, chromosomal locations, and population dynamics. *Proc. Natl. Acad. Sci. USA* **91**:5466–5470.
30. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**:406–425.
31. San-Blas, G., and G. Niño-Vega. 2001. *Paracoccidioides brasiliensis*: virulence and host response, p. 205–226. In R. L. Cihlar and R. A. Calderone (ed.), *Fungal pathogenesis: principles and clinical applications*. Marcel Dekker, Inc., New York, N.Y.
32. San-Blas, G., G. Niño-Vega, and T. Iturriaga. 2002. *Paracoccidioides brasiliensis* and paracoccidioidomycosis: molecular approaches to morphogenesis, diagnosis, epidemiology, taxonomy and genetics. *Med. Mycol.* **40**:225–242.
33. Semighini, C. P., Z. P. De Camargo, R. Puccia, M. H. Goldman, and G. H. Goldman. 2002. Molecular identification of *Paracoccidioides brasiliensis* by 5' nuclease assay. *Diagn. Microbiol. Infect. Dis.* **44**:383–386.
34. Shemer, R., Z. Weissman, N. Hashman, and D. Kornitzer. 2001. A highly polymorphic degenerate microsatellite for molecular strain typing of *Candida krusei*. *Microbiology* **147**:2021–2028.
35. Silva, S. P., M. S. S., M. Pereira, M. O. Azevedo, and C. M. A. Soares. 1999. Phase transition and stage-specific protein synthesis in the dimorphic fungus *Paracoccidioides brasiliensis*. *Exp. Mycol.* **18**:294–299.
36. Soares, C. M. A., E. E. W. I. Molinari Madlum, S. P. da Silva, M. Pereira, and M. S. S. Felipe. 1995. Characterization of *Paracoccidioides brasiliensis* isolates by random amplified polymorphic DNA analysis. *J. Clin. Microbiol.* **33**:505–507.
37. Svidzinsky, T. I. E., and Z. P. Camargo. 1995. Isoenzyme profile of *Paracoccidioides brasiliensis*. *J. Med. Vet. Mycol.* **33**:281–285.
38. Taylor, J. W., D. M. Geiser, A. Burt, and V. Koufonapou. 1999. The evolutionary biology and population genetics underlying fungal strain typing. *Clin. Microbiol. Rev.* **12**:126–146.
39. Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **24**:4876–4882.
40. Tóth, G., Z. Gáspári, and J. Jurka. 2000. Microsatellites in different eukaryotic genomes: survey and analysis. *Genome Res.* **10**:967–981.
41. Travassos, L. R., C. P. Taborda, L. K. Iwai, E. C. Cunha-Neto, and R. Puccia. 2004. The gp43 from *Paracoccidioides brasiliensis*: a major diagnostic antigen and vaccine candidate, p. 279–296. In J. E. Dömer and G. S. Kobayashi (ed.), *The Mycota XII: human fungal pathogens*. Springer-Verlag, Berlin, Germany.
42. Vaz, C. A. C., D. W. Mackenzie, V. M. Hearn, Z. P. Camargo, L. M. Singer-Vermes, E. Burger, and V. L. G. Calich. 1994. Gelatinase activity of exoantigens from virulent and non-virulent isolates of *Paracoccidioides brasiliensis*. *J. Med. Vet. Mycol.* **32**:65–69.
43. Vicentini, A. P., J. L. Gesztes, M. Franco, W. Souza, J. Moraes, L. R. Travassos, and J. D. Lopes. 1994. Binding of *Paracoccidioides brasiliensis* to laminin through surface glycoprotein gp43 leads to enhancement of fungal pathogenesis. *Infect. Immun.* **62**:4–7.
44. Zaccharias, D., A. Ueda, M. Moschardi-Bacchi, M. Franco, and G. San-Blas. 1986. A comparative histopathological, immunological and biochemical study of experimental intravenous paracoccidioidomycosis induced in mice by three *Paracoccidioides brasiliensis* isolates. *J. Med. Vet. Mycol.* **24**:445–454.