

# Aquaporin-3 mediates hydrogen peroxide-dependent responses to environmental stress in colonic epithelia

Jay R. Thiagarajah<sup>a,1</sup>, Jeffrey Chang<sup>a</sup>, Jeremy A. Goettel<sup>a</sup>, Alan S. Verkman<sup>b,c</sup>, and Wayne I. Lencer<sup>a</sup>

<sup>a</sup>Department of Gastroenterology, Nutrition, and Hepatology, Boston Children's Hospital, Harvard Medical School, Boston, MA 02115; <sup>b</sup>Department of Medicine, University of California, San Francisco, CA 94143; and <sup>c</sup>Department of Physiology, University of California, San Francisco, CA 94143

Edited by Lora V. Hooper, The University of Texas Southwestern Medical Center, Dallas, TX, and approved December 1, 2016 (received for review August 3, 2016)

The colonic epithelium provides an essential barrier against the environment that is critical for protecting the body and controlling inflammation. In response to injury or gut microbes, colonic epithelial cells produce extracellular hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), which acts as a potent signaling molecule affecting barrier function and host defense. In humans, impaired regulation of H<sub>2</sub>O<sub>2</sub> in the intestine has been associated with early-onset inflammatory bowel disease and colon cancer. Here, we show that signal transduction by H<sub>2</sub>O<sub>2</sub> depends on entry into the cell by transit through aquaporin-3 (AQP3), a plasma membrane H<sub>2</sub>O<sub>2</sub>-conducting channel. In response to injury, AQP3-depleted colonic epithelial cells showed defective lamellipodia, focal adhesions, and repair after wounding, along with impaired H<sub>2</sub>O<sub>2</sub> responses after exposure to the intestinal pathogen *Citrobacter rodentium*. Correspondingly, AQP3<sup>-/-</sup> mice showed impaired healing of superficial wounds in the colon and impaired mucosal innate immune responses against *C. rodentium* infection, manifested by reduced crypt hyperplasia, reduced epithelial expression of IL-6 and TNF- $\alpha$ , and impaired bacterial clearance. These results elucidate the signaling mechanism of extracellular H<sub>2</sub>O<sub>2</sub> in the colonic epithelium and implicate AQP3 in innate immunity at mucosal surfaces.

aquaporin-3 | hydrogen peroxide | colon | epithelium | microbe

In the mammalian colon, a single-cell-thick continuous layer of epithelial cells (ECs) forms the physical barrier that constitutes the interface between the body and the gut lumen, a compartment heavily laden with environmental microbes and other foreign and potentially toxic substances. Major gastrointestinal diseases, including infections and inflammatory bowel disease (IBD), are associated with dysfunction of the epithelial barrier (1). In both health and disease, mechanisms are in place to restore barrier function after injury and to defend against microbial colonization and invasion.

One of the pathways underlying barrier restitution and innate host defense against gut microbes depends on the generation of extracellular hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by the epithelial monolayer. Exactly how extracellular H<sub>2</sub>O<sub>2</sub> signals are transduced to cellular responses in the host epithelium remains incompletely characterized, especially at mucosal surfaces, where commensal microbes can produce H<sub>2</sub>O<sub>2</sub> in the absence of epithelial injury or pathogenic infection.

H<sub>2</sub>O<sub>2</sub> is a stable reactive oxygen species (ROS) molecule that acts as an extracellular and intracellular signal that mediates pleiotropic effects in tissues, including recruitment of immune cells to damaged areas and cell migration (2–4). In the intestine, the generation of extracellular H<sub>2</sub>O<sub>2</sub> at the mucosal or serosal surface is carried out by NADPH oxidase (NOX) enzymes, such as NOX1 and dual oxidase 2 (DUOX2) (5, 6). NOX1 is highly expressed in the mammalian colon, and the NOX enzymes are important in the defense against microbial pathogens and for regenerative homeostasis (7, 8). In addition, NOX enzymes are thought to play a role in the pathogenesis of IBD, with inactivating mutations recently described in children with early-onset

IBD and strong DUOX2 up-regulation found in inflamed pediatric ileal biopsy specimens (9, 10).

Aquaporin-3 (AQP3), a member of the aquaporin water channel family, is a water-, glycerol-, and H<sub>2</sub>O<sub>2</sub>-transporting protein implicated in various cellular functions (11, 12). AQP3-mediated H<sub>2</sub>O<sub>2</sub> transport has been reported to modulate epidermal growth factor signaling in ECs (13) and TNF- $\alpha$  signaling in skin keratinocytes (14, 15). In addition, AQP3-mediated H<sub>2</sub>O<sub>2</sub> transport has been shown to be important for T-cell and breast cancer cell migration (16, 17), although how AQP3 contributes to these changes in cellular function remains undefined. In the colon, AQP3 is expressed in ECs, with changes in expression found in response to inflammation, and our previous studies showed that mice lacking AQP3 experienced impaired recovery after chemical-induced colitis (18, 19).

In the present study, we tested whether AQP3 mediates the entry of extracellular H<sub>2</sub>O<sub>2</sub> into ECs lining the colon, and, if this explains the H<sub>2</sub>O<sub>2</sub>-dependent mucosal responses to epithelial injury and microbial infection.

## Results

**AQP3 Renders the Colonic Epithelium Responsive to Extracellular H<sub>2</sub>O<sub>2</sub>.** H<sub>2</sub>O<sub>2</sub> transport was measured by live cell imaging using a ratiometric H<sub>2</sub>O<sub>2</sub>-sensitive fluorescent protein (HyPer) stably transfected in Caco-2 colon ECs depleted of AQP3 (shRNA; AQP3-KD) or in control Caco-2 cells expressing AQP3 (scrambled shRNA; AQP3-SC). HyPer fluorescence ratios reflect changes in cytosolic H<sub>2</sub>O<sub>2</sub> concentration and represent an indirect measure of extracellular H<sub>2</sub>O<sub>2</sub> entry into cells. AQP3 is localized to the plasma membrane in Caco-2 cells, and analysis of AQP3 protein expression demonstrated efficient knockdown (Fig. S14). When exposed to medium containing H<sub>2</sub>O<sub>2</sub>, cytosolic increases in H<sub>2</sub>O<sub>2</sub> were significantly faster in cells expressing AQP3 compared with cells depleted of AQP3 or in which AQP3 transport was acutely

## Significance

The extracellular production of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by membrane or secreted oxidases has been linked to epithelial wound repair, defense against infection, and inflammation. Here, we elucidate the pathway that explains how extracellular H<sub>2</sub>O<sub>2</sub> transduces a signal into the cell to induce these critical cell functions. Our study shows the central component of this pathway in the mammalian colon is the H<sub>2</sub>O<sub>2</sub>-conducting aquaglyceroporin AQP3 and implicates this channel in innate immune responses at mucosal surfaces.


Author contributions: J.R.T. and W.I.L. designed research; J.R.T., J.C., and J.A.G. performed research; A.S.V. contributed new reagents/analytic tools; J.R.T. and J.C. analyzed data; and J.R.T., A.S.V., and W.I.L. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: jay.thiagarajah@childrens.harvard.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1612921114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1612921114/-DCSupplemental).



**Fig. 1.** AQP3 transports extracellular  $\text{H}_2\text{O}_2$  into colonic epithelia. (A) Caco-2 cells expressing  $\text{H}_2\text{O}_2$ -sensitive YFP (HyPer) show increased intracellular  $\text{H}_2\text{O}_2$  (ratio fluorescence) on exposure to  $100 \mu\text{M}$   $\text{H}_2\text{O}_2$ , which is inhibited by  $\text{AgNO}_3$  ( $5 \mu\text{M}$ ), supporting AQP3-facilitated  $\text{H}_2\text{O}_2$  transport. (B) Representative traces of increases in the HyPer ratio after the addition of  $100 \mu\text{M}$   $\text{H}_2\text{O}_2$  in Caco-2 cells transfected with control shRNA (AQP-SC) or after treatment with  $\text{AgNO}_3$  ( $5 \mu\text{M}$ ) or in AQP3-KD cells (Upper), and mean initial rates of increase in cellular  $\text{H}_2\text{O}_2$  (Lower);  $n = 6$ . (C) Representative traces of increases in CM- $\text{H}_2$ -DCFDA fluorescence after addition of  $100 \mu\text{M}$   $\text{H}_2\text{O}_2$  in AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> intestinal crypts (Upper) and mean initial rates of increase in cellular  $\text{H}_2\text{O}_2$  (Lower);  $n = 4$ . (D) Representative images showing increased  $\text{H}_2\text{O}_2$  entry in intestinal crypts from AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice measured using CM- $\text{H}_2$ -DCFDA after the addition of extracellular  $100 \mu\text{M}$   $\text{H}_2\text{O}_2$  pulse.

inhibited by  $\text{AgNO}_3$  (Fig. 1 A and B). Similarly, cells lacking endogenous AQP3 (FRT cells) showed greatly increased rates of  $\text{H}_2\text{O}_2$  elevation after transfection with AQP3 (Fig. S1B).

Because antioxidant cytosolic proteins, such as glutathione peroxidases (Gpx) and peroxiredoxins (Prx), rapidly remove free cellular  $\text{H}_2\text{O}_2$ , the rate-limiting step for the cytosolic increases in  $\text{H}_2\text{O}_2$  in our model systems is determined primarily by  $\text{H}_2\text{O}_2$  membrane permeability. Thus, increased  $\text{H}_2\text{O}_2$  membrane permeability in the presence of AQP3 should cause increases in cytosolic  $\text{H}_2\text{O}_2$  at much lower concentrations of extracellular  $\text{H}_2\text{O}_2$ . Dose-response studies have shown this prediction to be correct. AQP3-containing Caco-2 cells responded to extracellular  $\text{H}_2\text{O}_2$  at 100-fold lower concentrations compared with Caco-2 cells depleted of AQP3 (Fig. S1C). Importantly, uptake was observed in the concentration range critical for cellular  $\text{H}_2\text{O}_2$  signaling ( $1$ – $100 \mu\text{M}$ ) (3, 13, 21), and the expression of antioxidant proteins was not different between AQP3-KD and AQP3-SC cells (Fig. S1 D and E). At much higher and toxic concentrations of  $\text{H}_2\text{O}_2$  ( $>1 \text{ mM}$ ), cellular entry was not rate-limited by AQP3, but was governed by the intrinsic membrane permeability to  $\text{H}_2\text{O}_2$ . Thus, the presence or absence of membrane AQP3 allows colonic ECs to discriminate extracellular  $\text{H}_2\text{O}_2$  signals.

To test this idea in primary ECs in situ, we isolated colonic crypts from AQP3<sup>-/-</sup> and AQP3<sup>+/+</sup> mice. ECs in WT (AQP3<sup>+/+</sup>)

mice express AQP3 on both apical and basolateral membranes (Fig. S1F). Colonic crypts from AQP3<sup>-/-</sup> mice and exposed to exogenous  $\text{H}_2\text{O}_2$  showed a much slower increase in cytoplasmic  $\text{H}_2\text{O}_2$  compared with crypts from AQP3<sup>+/+</sup> mice, consistent with AQP3-facilitated  $\text{H}_2\text{O}_2$  entry in AQP3<sup>+/+</sup> mice (Fig. 1 C and D).


**Epithelial Repair and Wound Healing Depends on  $\text{H}_2\text{O}_2$  Transport via AQP3.** To test whether AQP3-dependent  $\text{H}_2\text{O}_2$  transport underlies signal transduction in the intestine, we investigated cell migration and barrier repair in response to injury, an important EC function modulated by extracellular  $\text{H}_2\text{O}_2$ . Studies on monolayers of WT Caco-2 cells after an initial scratch wounding showed that the addition of  $\text{H}_2\text{O}_2$  increased cell migration and wound closure (Fig. 2 A and B), consistent with previous reports (20). Wound closure was slowed in AQP3-KD cells, as well as in control AQP3-SC cells after inhibition of AQP3 channel function by  $\text{AgNO}_3$  or Auphen (21, 22). Wound closure was similarly slowed in control (AQP3-SC) Caco-2 cells after the addition of diphenyleneiodonium (DPI), which inhibits the NOX enzymes normally involved in production of extracellular  $\text{H}_2\text{O}_2$  after cell injury, and also by catalase, which enzymatically depletes extracellular  $\text{H}_2\text{O}_2$  (Fig. 2C). For each of these inhibitors, the  $\text{H}_2\text{O}_2$  specificity of their effect on wound healing was confirmed by showing their lack of effect on AQP3-dependent water transport (Fig. S1C). Finally, imaging of intracellular  $\text{H}_2\text{O}_2$  at the wound edge of scratch-injured Caco-2 monolayers showed reduced  $\text{H}_2\text{O}_2$  accumulation in AQP3-KD cells, consistent with the interpretation that wound healing is facilitated by  $\text{H}_2\text{O}_2$  transport via AQP3 (Fig. 2 D and E and Fig. S2A).

Closure of epithelial wounds after injury begins with migration of the wound-associated ECs from either side of the wound (23), which requires the production of numerous actin-driven lamellopodial projections. If AQP3 underlies wound signal transduction by  $\text{H}_2\text{O}_2$ , then the induction of lamellopodia also should be AQP3-dependent. To test this possibility, we imaged lamellopodia by actin staining in wound-associated Caco-2 cells depleted or not in AQP3 and in corresponding cells expressing eGFP-actin (Life-Act). Depletion of AQP3 or inhibition of  $\text{H}_2\text{O}_2$  signaling by incubation with catalase or DPI decreased the numbers and sizes of actin-containing lamellopodia (Fig. 3A and Fig. S2 B and C). For lamellopodial assembly and cell migration, wound-associated ECs require active turnover of focal adhesion complexes controlled by phosphorylation of key focal adhesion proteins, such as paxillin and focal adhesion kinase. We tested for this by examining the induction of focal adhesion complexes. Both proteins were found to be depleted in wound-associated ECs lacking AQP3 (AQP3-KD cells) (Fig. 3B and Fig. S2B). In addition, the ratio of phosphorylated to total paxillin was reduced in AQP3-KD cells, suggesting reduced active focal adhesions at the wound edge (Fig. 3B). The assembly of lamellopodia at the wound edge was accompanied by the expression and colocalization of AQP3 and NOX1 at these sites in AQP3-expressing cells (Fig. 3C), suggesting active regulation of cell migration by spatially coordinated  $\text{H}_2\text{O}_2$  production (NOX1) and membrane  $\text{H}_2\text{O}_2$  permeability (AQP3).

To test this pathway in vivo, we created endoscopic wounds in the colons of AQP3<sup>-/-</sup> and AQP3<sup>+/+</sup> mice and assessed wound healing by endoscopic visualization as described previously (24). AQP3<sup>-/-</sup> mice had distinctly impaired superficial wound healing, as assessed by analysis of the healed wound area at 6 d after injury (Fig. 3 D and E and Fig. S3A). ECs at the edge of the induced wounds showed a small increase in phosphorylated paxillin on day 1 postinjury in AQP3<sup>+/+</sup> mice compared with AQP3<sup>-/-</sup> mice (Fig. S3 B and C). Thus, AQP3-facilitated  $\text{H}_2\text{O}_2$  entry underlies the native NOX/ $\text{H}_2\text{O}_2$ -dependent epithelial response to injury and barrier repair.

**Host Defense Against *Citrobacter rodentium* Infection Depends on  $\text{H}_2\text{O}_2$  Transport via AQP3.** Because colonic ECs form the primary barrier against microbes and microbial products, we next tested





**Fig. 4.** fMLF and *C. rodentium* induce epithelial H<sub>2</sub>O<sub>2</sub> entry via AQP3. (A) Representative traces of HyPer ratio change in Caco-2 colonic ECs after the addition of fMLF (1 μM) (Upper), and summary of the mean initial rate of H<sub>2</sub>O<sub>2</sub> increase (Lower). *n* = 5. (B) Images of mouse distal colonoids from AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice after microinjection into the colonoid lumen of media (LB) or *C. rodentium* (CR, 1 × 10<sup>7</sup> cfu) showing CM-H<sub>2</sub>-DCFDA fluorescence and brightfield. (C) Summary of mean CM-H<sub>2</sub>-DCFDA fluorescence in AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> colonoids microinjected with LB, CR, or DH5α. *n* = 3. (D) Summary of accumulation of colonic epithelial ROS species in colon from AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice measured by ROS<sub>550</sub> fluorescence at 3 d post-infection with CR. *n* = 3.

leads to activation of NOX1 and generation of extracellular H<sub>2</sub>O<sub>2</sub>, which mediates the epithelial response to wound healing. In alignment with this and our earlier results (Figs. 1–3), the addition of fMLF to colon ECs produced a time-dependent increase in cellular H<sub>2</sub>O<sub>2</sub> that was attenuated in AQP3-KD monolayers and blocked in WT Caco-2 cells by depletion of extracellular H<sub>2</sub>O<sub>2</sub> using catalase and inhibition of NOX1 using DPI (Fig. 4A).

To test whether a colonic pathogen can induce such a response, and whether this action is AQP3-dependent, colonoids generated from AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice were exposed to the noninvasive, attaching, and effacing pathogen *C. rodentium*, the murine equivalent of enteropathogenic *Escherichia coli* in humans. *C. rodentium* or nonpathogenic *E. coli* (DH5α) were microinjected into the lumen of the colonoids (Fig. 4B and Fig. S4A). Luminal incubation with *C. rodentium*, but not with growth medium alone or *E. coli* DH5α, induced a robust increase in cellular H<sub>2</sub>O<sub>2</sub> that was significantly attenuated in AQP3<sup>-/-</sup> colonoids (Fig. 4B and C and Fig. S4B). Similarly, colonic tissue from AQP3<sup>-/-</sup> mice infected with *C. rodentium* showed reduced accumulation of ROS in surface colonic ECs compared with infected AQP3<sup>+/+</sup> mice (Fig. 4D).

In vivo, colonic infection by *C. rodentium* induces a robust early mucosal inflammatory response required for subsequent bacterial clearance and recovery. Induction of this response depends on direct interaction of the bacteria with the epithelium


and ROS signaling (25, 26). To test whether AQP3 mediates this recovery, we studied *C. rodentium* infection in AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice. AQP3<sup>+/+</sup> mice infected with *C. rodentium* showed an early loss in body weight at 3–6 d postinfection with subsequent crypt hyperplasia, indicative of the early inflammatory response. Body weight then recovered, as described previously (27) and in contrast to AQP3<sup>-/-</sup> mice (Fig. 5A and B). AQP3<sup>-/-</sup> mice showed no initial loss of weight or crypt hyperplasia at early stages of infection, and no weight gain later (days 18–25), consistent with the lack of an initial inflammatory response and persistent *C. rodentium* infection. This was further evidenced by the failure of AQP3<sup>-/-</sup> mice to show increased expression of the inflammatory cytokines IL-6 and TNF-α at early stages of infection, as well as the failure to clear the bacteria from the colon at later stages (Fig. 5C and D). At later stages of infection (day 25), AQP3<sup>+/+</sup> mice exhibited reduced crypt hyperplasia in conjunction with resolution of the infection. In contrast, AQP3<sup>-/-</sup> mouse crypts showed a delayed increase in crypt length, possibly a result of persistent bacterial colonization (Fig. S5B). Importantly, in both AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice, Nox1 expression was up-regulated in surface ECs after infection (Fig. 5E and Fig. S5C). Furthermore, AQP3<sup>+/+</sup> mice given *N*-acetyl cysteine (NAC) in drinking water, a well-tolerated antioxidant that removes free H<sub>2</sub>O<sub>2</sub>, phenocopied AQP3<sup>-/-</sup> mice at early stages of *C. rodentium* infection (Fig. 5F and G) and showed impaired bacterial clearance and cytokine expression at later stages of infection (Fig. S5A, B, and D). Thus, AQP3 is required for H<sub>2</sub>O<sub>2</sub> signaling in the colon and for the epithelial response to infection by *C. rodentium*.

## Discussion

The innate immune function of ECs lining the colon is thought to be critical for the maintenance of epithelial transport and barrier functions in the face of a complex and potentially hostile luminal environment (1). Epithelial generation of H<sub>2</sub>O<sub>2</sub> has been shown to be an important signal for wound repair and recruitment of inflammatory cells after injury. NOX and DUOX complexes at the intestinal epithelial plasma membrane generate H<sub>2</sub>O<sub>2</sub> or superoxide (O<sup>-</sup>), which is rapidly converted to H<sub>2</sub>O<sub>2</sub> by extracellular superoxide dismutase. In the distal colon, this appears to occur primarily via NOX1 based on its high regional expression (6). Studies in *Drosophila*, *Caenorhabditis elegans*, and zebrafish intestine have shown that epithelial production of H<sub>2</sub>O<sub>2</sub> is part of the host response to gut infection and is involved in microbial clearance and responses to gut symbionts and pathogens (28–30). In mammals, a similar role has been postulated based on studies of pathogen infection in stomach and airway ECs (31, 32).

In the present study, we have shown that AQP3 mediates H<sub>2</sub>O<sub>2</sub> signal transduction in colonic epithelia. The increased membrane permeability to H<sub>2</sub>O<sub>2</sub> in AQP3-containing colonic ECs allows them to respond to external H<sub>2</sub>O<sub>2</sub> at concentrations (1–100 μM) relevant for cellular signaling processes, such as migration and pathogen recognition. This was demonstrated both in vitro and in vivo by studying wound repair after injury and host defense against *C. rodentium* infection.

In the case of wound repair, various signaling proteins may be involved in translating the rapid H<sub>2</sub>O<sub>2</sub> uptake facilitated by AQP3 into the signals required for the observed epithelial cytoskeletal and cell migratory responses. Previous studies have identified various candidate ROS-sensitive intracellular proteins, including the src family kinases and several tyrosine phosphatases, including SHP2 and PTP-PEST (4, 15, 26). These proteins contain free cysteines that can alter function and are affected by changes in cellular H<sub>2</sub>O<sub>2</sub> levels. Given that the specific proteins involved are often cell type-specific, further studies are needed in primary EC systems to fully understand the signaling pathways and effectors of AQP3-facilitated H<sub>2</sub>O<sub>2</sub> cellular entry.



**Fig. 5.** AQP3-dependent mucosal response to *C. rodentium* infection in vivo. (A) Time course of body weight change after *C. rodentium* (CR) infection in AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice, and mean bodyweight change at days 3 and 25 postinfection. *n* = 6. (B) Crypt length (μm) in AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice at 12 d postinfection and representative H&E images of crypt hyperplasia in AQP3<sup>+/+</sup>, but not AQP3<sup>-/-</sup>, distal colon at 12 d postinfection. *n* = 6. (C) CR bacterial load in stool at 5 and 25 d postinfection and cecum tissue at 25 d postinfection. *n* = 6. (D) Relative expression of IL-6 and TNF-α mRNA in isolated ECs at 5 d postinfection. *n* = 6. (E) Images of NOX1 (red) expression in distal colon at 5 d postinfection. Arrows indicate increased expression in surface ECs after infection. (F) Time course of mean body weight change after CR infection in AQP3<sup>+/+</sup> mice, AQP3<sup>+/+</sup> mice administered NAC, and AQP3<sup>-/-</sup> mice. *n* = 5. (G) Relative expression of IL-6 and TNF-α mRNA in isolated ECs from AQP3<sup>+/+</sup> mice, AQP3<sup>+/+</sup> mice administered NAC, and AQP3<sup>-/-</sup> mice at 5 d postinfection. *n* = 5. (H) Relative expression of IL-6 and TNF-α mRNA at 25 d postinfection. *n* = 4.

Recent studies also have pointed to an important role for ROS signaling in relation to pathogen infection and microbial recognition, as well as in the mucosal inflammation seen in chronic inflammatory bowel diseases, which is associated with alterations in gut microbes (9, 28, 31, 33, 34). In the case of *C. rodentium* infection, ROS signaling has been shown to be required for a robust TH<sub>17</sub> response and bacterial clearance (26). The results reported here suggest that this response is likely dependent on AQP3-mediated H<sub>2</sub>O<sub>2</sub> signal transduction. In the skin epidermis, another bacterially colonized mucosal surface, AQP3-dependent H<sub>2</sub>O<sub>2</sub> responses have been shown to play a role in the pathogenesis of psoriasis and in T-cell migration (14, 15). Further studies will help characterize how colonic epithelial cytokine secretion is regulated by AQP3-dependent H<sub>2</sub>O<sub>2</sub> signals, and how this may modulate other epithelial innate immune pathways, such as Toll-like receptor signaling and activation of the Nlrp6 inflammasome (35, 36).

A recent study identified several mutations in epithelial NOXs in children with very-early-onset IBD. In these patients, defective production of H<sub>2</sub>O<sub>2</sub> by ECs was linked to impaired initial host resistance to pathogenic infections (9), as we found here for AQP3<sup>-/-</sup> mice. Although AQP3 has not been identified as a monogenic cause of IBD, patients with ulcerative colitis have significantly lower expression of epithelial AQP3 during both inflammation and clinical remission (17). Conversely, AQP3 expression is highly increased in many colon carcinomas and is correlated with metastatic spread (37). As shown here, AQP3-dependent H<sub>2</sub>O<sub>2</sub> cellular entry increases colonic epithelial migration, and it is plausible that AQP3-

facilitated H<sub>2</sub>O<sub>2</sub> transport may play a role in the development of colonic cancers and metastases.

Given that AQP3 is expressed in mucosal surface cells lining the colon, airway, genitourinary tract, and skin, we speculate that AQP3-facilitated H<sub>2</sub>O<sub>2</sub> entry represents a general mechanism for mediating EC responses to the external environment. Environmental changes in commensal microbial composition or pathogenic infection generate a variety of signals that mucosal ECs are poised to recognize, including changes in redox status and the production of stable ROS species, such as H<sub>2</sub>O<sub>2</sub>, in the extracellular milieu. The mechanisms by which the host ECs discriminate between pathogenic “danger” H<sub>2</sub>O<sub>2</sub> signals and homeostatic commensal H<sub>2</sub>O<sub>2</sub> signals remains unknown, however. Accumulation of H<sub>2</sub>O<sub>2</sub> at colonic EC surfaces can occur in a cell-autonomous manner through activation of NOX1 (5, 6), in a paracrine manner after release from gut commensal bacteria (34), or as a product of secreted enzymes, such as xanthine oxidase (38). In this context, and perhaps dependent on the source of H<sub>2</sub>O<sub>2</sub> (cell-autonomous or paracrine) and/or its localization (luminal or subepithelial), AQP3-facilitated H<sub>2</sub>O<sub>2</sub> transport in the epithelium may be intrinsic to host responses to the environment, including the acute and chronic adaptive response to gut commensal or pathogenic microbes.

### Materials and Methods

The materials and experimental procedures used in this study are described in detail in *SI Materials and Methods*. Primer sequences for qRT-PCR are shown in *Table S1*.

**Mice.** AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice in a CD1 genetic background were generated as described previously. Animals were maintained in a specific pathogen-free animal facility at Boston Children's Hospital. Experiments were conducted after approval from Animal Care Systems and in accordance with regulations of the Institutional Animal Care and Use Committee.

**H<sub>2</sub>O<sub>2</sub> Fluorescence Ratio Imaging.** Stably transfected cells containing the H<sub>2</sub>O<sub>2</sub>-sensitive probe HyPer (39) on glass coverslips were placed in a micro-perfusion chamber (Warner Instruments) with HBSS with glucose at 37 °C and imaged.

**In Vivo Colonoscopy.** AQP3<sup>+/+</sup> and AQP3<sup>-/-</sup> mice ( $n = 3-5$ ) were anesthetized by i.p. injection of a ketamine (100 mg/kg)/xylazine (10 mg/kg) solution. The distal colon was imaged using a high-resolution colonoscopy system (Karl Storz). To create superficial wounds, endoscopic biopsies (three to six per colon) using biopsy forceps were made in the distal colon and followed at 1 and 6 d postwounding by repeat endoscopy and image capture.

**Colonoid Microinjection.** Epithelial colonoids were generated and plated in Matrigel on glass-bottom Petri dishes. At 4–5 d after plating, colonoids were microinjected with a pulled glass pipette containing injection solutions (*C. rodentium*  $1 \times 10^7$  cfu, LB, *E. coli* DH5 $\alpha$ ) attached to a Nanoject (World Precision Instruments). Colonoids (25–40 per condition) were injected with 2 nL of solution. Some colonoids were injected with FluoSpheres (Thermo Fisher Scientific) to mark injections (Fig. S4B). Injected colonoids were incubated at 37 °C for 6 h.

**Statistics.** Quantitative data are expressed as mean  $\pm$  SEM for each treatment group. Statistical comparisons were performed using the two-tailed Student's *t* test or ANOVA with Tukey's multiple-comparison post hoc test. *P* values < 0.05 were considered significant (\**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001).

**ACKNOWLEDGMENTS.** We thank Dr. Dan Chinnapen for synthesis of the aquaporin inhibitor Auphen. This work was supported by National Institutes of Health Grants R01 DK048106 (to W.I.L.), R01 DK035124 (to A.S.V.), P30 DK034854 (to the Harvard Digestive Diseases Center), and Pediatric Scientist Development Program K12-HD000850 (to J.R.T.).

- Peterson LW, Artis D (2014) Intestinal epithelial cells: Regulators of barrier function and immune homeostasis. *Nat Rev Immunol* 14(3):141–153.
- Niethammer P, Grabher C, Look AT, Mitchison TJ (2009) A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459(7249):996–999.
- van der Vliet A, Janssen-Heininger YMW (2014) Hydrogen peroxide as a damage signal in tissue injury and inflammation: Murderer, mediator, or messenger? *J Cell Biochem* 115(3):427–435.
- Yoo SK, Freisinger CM, LeBert DC, Huttenlocher A (2012) Early redox, Src family kinase, and calcium signaling integrate wound responses and tissue regeneration in zebrafish. *J Cell Biol* 199(2):225–234.
- Rada B, Leto TL (2008) Oxidative innate immune defenses by Nox/Duox family NADPH oxidases. *Contrib Microbiol* 15:164–187.
- Szanto I, et al. (2005) Expression of NOX1, a superoxide-generating NADPH oxidase, in colon cancer and inflammatory bowel disease. *J Pathol* 207(2):164–176.
- Corcionivoschi N, et al. (2012) Mucosal reactive oxygen species decrease virulence by disrupting *Campylobacter jejuni* phosphotyrosine signaling. *Cell Host Microbe* 12(1):47–59.
- Jones RM, et al. (2013) Symbiotic lactobacilli stimulate gut epithelial proliferation via Nox-mediated generation of reactive oxygen species. *EMBO J* 32(23):3017–3028.
- Hayes P, et al. (2015) Defects in NADPH oxidase genes NOX1 and DUOX2 in very early onset inflammatory bowel disease. *Cell Mol Gastroenterol Hepatol* 1(5):489–502.
- Haberman Y, et al. (2014) Pediatric Crohn disease patients exhibit specific ileal transcriptome and microbiome signature. *J Clin Invest* 124(8):3617–3633.
- Verkman AS (2011) Aquaporins at a glance. *J Cell Sci* 124(Pt 13):2107–2112.
- Miller EW, Dickinson BC, Chang CJ (2010) Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proc Natl Acad Sci USA* 107(36):15681–15686.
- Hara-Chikuma M, Watanabe S, Satooka H (2016) Involvement of aquaporin-3 in epidermal growth factor receptor signaling via hydrogen peroxide transport in cancer cells. *Biochem Biophys Res Commun* 471(4):603–609.
- Hara-Chikuma M, et al. (2015) Aquaporin-3-mediated hydrogen peroxide transport is required for NF- $\kappa$ B signaling in keratinocytes and development of psoriasis. *Nat Commun* 6:7454.
- Hara-Chikuma M, et al. (2012) Chemokine-dependent T cell migration requires aquaporin-3-mediated hydrogen peroxide uptake. *J Exp Med* 209(10):1743–1752.
- Satooka H, Hara-Chikuma M (2016) Aquaporin-3 controls breast cancer cell migration by regulating hydrogen peroxide transport and its downstream cell signaling. *Mol Cell Biol* 36(7):1206–1218.
- Ricanek P, et al. (2015) Reduced expression of aquaporins in human intestinal mucosa in early stage inflammatory bowel disease. *Clin Exp Gastroenterol* 8:49–67.
- Thiagarajah JR, Zhao D, Verkman AS (2007) Impaired enterocyte proliferation in aquaporin-3 deficiency in mouse models of colitis. *Gut* 56(11):1529–1535.
- Knock GA, Ward JPT (2011) Redox regulation of protein kinases as a modulator of vascular function. *Antioxid Redox Signal* 15(6):1531–1547.
- Basuroy S, Dunagan M, Sheth P, Seth A, Rao RK (2010) Hydrogen peroxide activates focal adhesion kinase and c-Src by a phosphatidylinositol 3 kinase-dependent mechanism and promotes cell migration in Caco-2 cell monolayers. *Am J Physiol Gastrointest Liver Physiol* 299(1):G186–G195.
- Niemietz CM, Tyerman SD (2002) New potent inhibitors of aquaporins: Silver and gold compounds inhibit aquaporins of plant and human origin. *FEBS Lett* 531(3):443–447.
- Martins AP, et al. (2012) Targeting aquaporin function: Potent inhibition of aquaporin-3 by a gold-based compound. *PLoS One* 7(5):e37435.
- Cordeiro JV, Jacinto A (2013) The role of transcription-independent damage signals in the initiation of epithelial wound healing. *Nat Rev Mol Cell Biol* 14(4):249–262.
- Leoni G, et al. (2013) Annexin A1, formyl peptide receptor, and NOX1 orchestrate epithelial repair. *J Clin Invest* 123(1):443–454.
- Collins JW, et al. (2014) *Citrobacter rodentium*: Infection, inflammation and the microbiota. *Nat Rev Microbiol* 12(9):612–623.
- Atarashi K, et al. (2015) Th17 cell induction by adhesion of microbes to intestinal epithelial cells. *Cell* 163(2):367–380.
- Brown JB, et al. (2011) Epithelial phosphatidylinositol-3-kinase signaling is required for  $\beta$ -catenin activation and host defense against *Citrobacter rodentium* infection. *Infect Immun* 79(5):1863–1872.
- Ha E-M, Oh C-T, Bae YS, Lee W-J (2005) A direct role for dual oxidase in *Drosophila* gut immunity. *Science* 310(5749):847–850.
- Flores MV, et al. (2010) Dual oxidase in the intestinal epithelium of zebrafish larvae has anti-bacterial properties. *Biochem Biophys Res Commun* 400(1):164–168.
- Hoeven RV, McCallum KC, Cruz MR, Garsin DA (2011) Ce-Duox1/BLI-3 generated reactive oxygen species trigger protective SKN-1 activity via p38 MAPK signaling during infection in *C. elegans*. *PLoS Pathog* 7(12):e1002453.
- Grasberger H, El-Zaatari M, Dang DT, Merchant JL (2013) Dual oxidases control release of hydrogen peroxide by the gastric epithelium to prevent *Helicobacter felis* infection and inflammation in mice. *Gastroenterology* 145(5):1045–1054.
- Geiszt M, Witta J, Baffi J, Lekstrom K, Leto TL (2003) Dual oxidases represent novel hydrogen peroxide sources supporting mucosal surface host defense. *FASEB J* 17(11):1502–1504.
- Jostins L, et al.; International IBD Genetics Consortium (IBDGC) (2012) Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 491(7422):119–124.
- Voltan S, et al. (2008) *Lactobacillus crispatus* M247-derived H<sub>2</sub>O<sub>2</sub> acts as a signal-transducing molecule activating peroxisome proliferator activated receptor-gamma in the intestinal mucosa. *Gastroenterology* 135(4):1216–1227.
- Joo J-H, et al. (2012) Dual oxidase 2 is essential for the Toll-like receptor 5-mediated inflammatory response in airway mucosa. *Antioxid Redox Signal* 16(1):57–70.
- Kuwano Y, et al. (2008) Tumor necrosis factor alpha activates transcription of the NADPH oxidase organizer 1 (NOXO1) gene and up-regulates superoxide production in colon epithelial cells. *Free Radic Biol Med* 45(12):1642–1652.
- Kang BW, et al. (2015) Expression of aquaporin-1, aquaporin-3, and aquaporin-5 correlates with nodal metastasis in colon cancer. *Oncology* 88(6):369–376.
- Crane JK, Naehre TM, Broome JE, Boedeker EC (2013) Role of host xanthine oxidase in infection due to enteropathogenic and Shiga-toxicogenic *Escherichia coli*. *Infect Immun* 81(4):1129–1139.
- Belousov VV, et al. (2006) Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods* 3(4):281–286.
- Block B, Bailar JC (1951) The reaction of gold(III) with some bidentate coordinating groups. *J Am Chem Soc* 73(3):4722–4725.