



OPEN ACCESS

Epigenetic treatment of pancreatic cancer: is there a therapeutic perspective on the horizon?

Elisabeth Hessmann,¹ Steven A Johnsen,² Jens T Siveke,^{3,4} Volker Ellenrieder¹

¹Department of Gastroenterology and Gastrointestinal Oncology, University Medical Center Goettingen, Goettingen, Germany

²Department of General, Visceral and Pediatric Surgery, University Medical Center Goettingen, Goettingen, Germany

³Division of Solid Tumor Translational Oncology, German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ), Heidelberg, Germany

⁴West German Cancer Center, University Hospital Essen, Essen, Germany

Correspondence to

Professor Volker Ellenrieder, Department of Gastroenterology and Gastrointestinal Oncology, University Medical Center Goettingen, Robert-Koch-Strasse 40, Goettingen 37075, Germany; volker.ellenrieder@med.uni-goettingen.de

Received 30 June 2016

Accepted 30 August 2016

Published Online First

3 November 2016

ABSTRACT

Pancreatic ductal adenocarcinoma (PDAC) constitutes one of the most aggressive malignancies with a 5-year survival rate of <7%. Due to growing incidence, late diagnosis and insufficient treatment options, PDAC is predicted to soon become one of the leading causes of cancer-related death. Although intensified cytostatic combinations, particularly gemcitabine plus nab-paclitaxel and the folinic acid, fluorouracil, irinotecan, oxaliplatin (FOLFIRINOX) protocol, provide some improvement in efficacy and survival compared with gemcitabine alone, a breakthrough in the treatment of metastatic pancreatic cancer remains out of sight. Nevertheless, recent translational research activities propose that either modulation of the immune response or pharmacological targeting of epigenetic modifications alone, or in combination with chemotherapy, might open highly powerful therapeutic avenues in GI cancer entities, including pancreatic cancer. Deregulation of key epigenetic factors and chromatin-modifying proteins, particularly those responsible for the addition, removal or recognition of post-translational histone modifications, are frequently found in human pancreatic cancer and hence constitute particularly exciting treatment opportunities. This review summarises both current clinical trial activities and discovery programmes initiated throughout the biopharma landscape, and critically discusses the chances, hurdles and limitations of epigenetic-based therapy in future PDAC treatment.

INTRODUCTION

Pancreatic ductal adenocarcinoma (PDAC) is a dismal disease with a 5-year survival rate of <7%.¹ Despite tremendous efforts in PDAC research and therapeutic development, no significant improvement in survival has been achieved within the last decades. Research efforts in PDAC have traditionally focused on genetic alterations underlying malignant transformation, carcinogenesis and tumour progression. Early studies on these genetic abnormalities defined common mechanisms of oncogenesis in PDAC, such as activating mutations of *Kras* or inactivation of the tumour suppressor genes *TP53*, *DPC4* or *CDKN2A*.² In addition to these well-characterised ‘driver mutations’, a plethora of diverse genetic events occurs in each pancreatic tumour, characterising PDAC as one of the most heterogeneous malignant diseases.^{1–3} These studies led to the testing of targeted therapies, which have been evaluated in preclinical and clinical settings as monotherapies or in combination with standard chemotherapy.^{4–5} While small subgroups of patients with PDAC show responsiveness towards selected therapeutic regimes, the majority

of patients with PDAC is refractory to these treatments a priori or rapidly acquires therapeutic resistance.

Drug resistance in PDAC is mediated by pronounced plasticity enabling PDAC cells to switch between phenotypic states and to select for cellular clones that eventually evade therapy.⁶ Importantly, apart from the restrictive contribution of genetic alterations, the acquisition of a drug-tolerant phenotype is frequently largely reversible. Indeed, the dynamic character of cell plasticity and drug resistance suggests the involvement of epigenetic regulation in controlling phenotypic heterogeneity in PDAC.⁷ The term ‘epigenetics’ describes processes that comprise non-genetic, heritable information through alterations that do not change the DNA sequence⁸ and refers to mechanisms involving DNA-methylation, post-transcriptional control of gene expression via non-coding RNAs as well as modifications and remodelling of chromatin. In addition to its functions to package DNA into a cell, its crucial involvement in mitosis and prevention of DNA damage, chromatin significantly impacts on the transcriptional activity of a gene. The nucleosome constitutes the functional subunit of chromatin and consists of approximately 147 bp of DNA wrapped around an octameric structure composed of two histones each of H2A, H2B, H3 and H4.⁹ Regulation of chromatin conformation is facilitated through DNA methylation, chromatin remodelling or histone modifications, such as acetylation, methylation, ubiquitination and phosphorylation.⁸ These post-translational modifications alter chromatin architecture by non-covalent interactions within and between nucleosomes, resulting in changes of nucleosome structure and accessibility of the transcription machinery.⁸ The major effectors of post-translational histone modifications are chromatin-modifying enzymes that either add (‘writers’) or remove (‘erasers’) modifications or factors that recognise specific histone modifications or combinations of modifications (‘readers’) (figure 1). Defects in these chromatin modulating proteins can have profound effects on vital cellular processes and hence contribute to development and progression of numerous diseases including cancer.¹⁰ Consequently, several recent preclinical studies investigated the impact of epigenetic alterations in diverse cancer models and emphasised the significance of epigenetics in malignant transformation and tumour progression in haematological malignancies and solid tumours including PDAC.^{1–11} Importantly, in contrast to genetic defects, epigenetic alterations are reversible and therefore represent bona fide targets for novel cancer therapies. Not surprisingly, first clinical trials



CrossMark

To cite: Hessmann E, Johnsen SA, Siveke JT, et al. *Gut* 2017;**66**:168–179.




Figure 1 Schematic illustration of chromatin-associated regulation of gene transcription. The chromatin conformation significantly determines the accessibility of the transcription machinery to the DNA. The switch between condensed and transcriptional inactive heterochromatin (left panel) and open, accessible euchromatin (right panel) is controlled by chromatin regulators that establish ('writers'), maintain ('readers') or remove ('erasers') post-translational modifications on the lysine residues of histone tails. Exemplary chromatin regulators are illustrated to show their impact on chromatin conformation and gene transcription. All inhibitors depicted in the blue boxes are evaluated in clinical trials in pancreatic ductal adenocarcinoma (PDAC). ac, acetylation; BET, family of bromodomain and external terminal proteins; HDAC, histone deacetylase; H3, histone 3; KDM6, lysine demethylase 6; K27, lysine 27; me3, 3 methyl groups; PRC1/2, polycomb repressor complex 1/2; SWI/SNF, SWI/Itch/sucrose non-fermentable chromatin complex.

aiming at altered epigenetic signalling in PDAC are underway. Furthermore, epigenetic alterations are increasingly being recognised for their predictive value in molecular tumour stratification in many cancer entities and thus extend the scientific and translational relevance of epigenetics in this particular tumour entity.

In this review, we summarise recent advances in our understanding of altered chromatin regulation in PDAC development and progression, and critically discuss ongoing initiatives to tackle epigenetic dysregulation in PDAC disease. Moreover, we highlight the potential of epigenetic approaches in therapeutic PDAC stratification strategies with a particular focus on chromatin-associated mechanisms (box 1).

TRANSLATIONAL APPROACHES TO TACKLE EPIGENETIC DYSREGULATION IN PANCREATIC CANCER

Histone acetylation

Acetylation and deacetylation of lysine residues within histone tails represents a crucial mechanism within a plethora of epigenetic regulatory mechanisms controlling gene expression.¹⁰ While histone acetylation mediated by histone acetyltransferases (HATs), for example, CREB-binding protein (CBP), p300 and pCAF is associated with accessible chromatin and transcriptional activation, deacetylation by different groups of histone deacetylases (HDACs) is responsible for repression of gene transcription (figure 1). Histone acetylation and deacetylation levels are tightly controlled by antagonistic activities of HATs and HDACs and hence, unbalanced enzyme activation in one way or the other can foster malignant transformation and tumour progression.¹²

HATs control gene expression by catalysing acetylation of histones and non-histone proteins. Specifically, acetylation

neutralises the positive charge on the amino group of specific lysine residues, thus weakening the DNA-chromatin complex and creating an open chromatin configuration.¹³ The p300 protein belongs to the best-studied HATs and represents a ubiquitously expressed global transcriptional coactivator with critical involvement in a wide variety of cellular mechanisms.¹⁴ Compared with HDACs, the contribution of HATs to pancreatic cancer formation and progression is multifaceted and highly dependent on the cellular context and the selection of regulated target genes.¹² While certain studies designate a tumour-promoting function of p300 in PDAC, for example, by transcriptional activation of the c-Myc promoter,¹⁵ other reports describing p300 as a metastasis repressive protein¹⁶ as well as the frequent occurrence of loss of function mutations in the *EP300* gene in PDAC cell lines¹⁴ argue for tumour-suppressive HATs functions in PDAC. Due to the contradictory preclinical findings as well as the unavailability of highly specific HAT inhibitors, little progress has been made in evaluating the potential utility of HAT inhibition for the treatment of patients with PDAC.

The natural turmeric-derived polyphenol compound curcumin represents a potent inhibitor of p300 HAT activity.¹⁴ A number of preclinical data have demonstrated antitumourigenic effects of curcumin in PDAC using in vitro and in vivo systems.^{17–19} These findings combined with a minimal toxicity profile led to the initiation of a few clinical trials to investigate the safety and efficacy of curcumin in PDAC therapy. The first-in-patient study performed with the natural compound tested the efficiency and feasibility of curcumin application in combination with gemcitabine in chemotherapy-naïve patients with advanced PDAC (NCT00192842) (compare tables 2 and 3).²⁰ In contrast to the following trials, the daily oral dose of 8 g caused severe and

Table 1 Exemplary HDAC targets in cancer

Oncogenic function	HDAC target gene
Cell cycle regulation	p21 Ku70
Inhibition of epithelial differentiation	GATA4 GATA6 DAB2 Muc2
Inhibition of apoptosis	p53 Bcl-xl
EMT and metastasis	CDH1 CXCR4 miR-203

EMT, epithelial-mesenchymal transition.

intractable abdominal pain, indicating an increased GI toxicity of the drug when applied together with gemcitabine.²¹ Dhillon *et al* conducted a subsequent monotherapy trial with curcumin in patients with PDAC (NCT00094445). In a phase II setting, pre-treated or untreated patients received 8 g curcumin daily, which was well tolerated and, despite its limited bioavailability, showed biological activity in some patients with PDAC with stable disease and a brief, but remarkable response (73% reduction of liver metastasis size) as the best outcome.^{21–23} Another group performed two clinical trials with curcumin and a nanoparticle-based curcumin (Theracurmin). In a phase I/II study, patients who became resistant to gemcitabine-based chemotherapy were treated with a combined curcumin/gemcitabine regime.²³ No cumulative toxicity from curcumin was observed, but unfortunately, no patient experienced a complete or partial response.²³ The described improvement of quality-of-life scores following Theracurmin administration needs to be confirmed in a randomised placebo-controlled trial.²¹

Overall, the clinical trials evaluating curcumin in PDAC therapy reflect the conflicting results obtained in preclinical studies. Although high p300 specificity of curcumin has been reported *in vitro*,¹⁴ it remains unclear whether the activity of the drug is limited to acetyltransferase inhibition. Careful preclinical investigations are required to understand the involvement of p300 in tumour-promoting functions in PDAC. If further evidence were to support a definitive function of HATs as oncogenic drivers in subgroups of PDAC, more specific and potent inhibitors would need to be developed to increase the chances of antitumourigenic activity of HAT blockade.

HDAC proteins counteract HAT activity and have been extensively described as significant drivers of transformation and tumour progression in multiple tissues, including the pancreas. Hyperactivity of HDAC proteins can foster proliferation and impair cell death regulation in pancreatic cancer^{24–26} and HDAC-mediated transcriptional control is associated with the regulation of epithelial-mesenchymal transition (EMT) programmes in PDAC cells (table 1),^{27–30} thereby contributing to PDAC invasion and metastasis. Based on their homology with yeast deacetylases, their subcellular localisation and diverse functions, HDAC proteins are grouped into three different classes.³¹ As a result of the frequently observed dysregulation of HDAC family members in cancer,^{32–33} and prompted by the finding that HDACs control various key oncogenic features of cancer cells,³⁴ inhibition of HDAC activity has been evaluated as a therapeutic strategy to restore the balance of histone acetylation and to subsequently interfere with HDAC target gene expression. Several natural as well as synthetic compounds that inhibit

HDAC activity are now available.^{35–36} HDAC inhibitors (HDACi) have been identified which either target all HDAC family members (pan-HDACi) or selectively interfere with subgroups of HDAC isoforms, thus facilitating a more specific manipulation of particular oncogenic HDAC functions (figure 2).^{12–31} Several HDACi have been investigated in clinical trials, where they have shown beneficial effects in selected haematological malignancies and solid tumour entities.³⁴ Three HDACi, vorinostat (Zolinza), romidepsin (Istodax) and panobinostat (Farydak) have reached approval by the US Food and Drug Administration (FDA) or by the European Medicines Agency for the treatment of cutaneous T-cell lymphoma (vorinostat and romidepsin), peripheral T-cell lymphoma (romidepsin) and multiple myeloma (vorinostat and panobinostat).³⁷

Based on the oncogenic activity of HDACs in PDAC, several clinical trials were initiated to evaluate the safety and efficacy of HDACi in patients with PDAC (tables 2 and 3). While HDACi monotherapy showed clinical activity in haematological malignancies, results have largely been disappointing in most solid tumours including PDAC.³⁴ Consequently, recent clinical trials investigating HDACi in PDAC focus on combined approaches of HDACi with small-molecule inhibitors or chemotherapeutic agents. In fact, three clinical trials are currently evaluating HDAC inhibition in combination with gemcitabine (NCT00379639, NCT00372437, NCT00004861). While monotherapy with mocetinostat (MGCD0103) stabilised tumour disease at best in some cases,³⁸ it was significantly more active in combination with gemcitabine, as described in a recent phase I/II study on advanced solid tumours (NCT00372437). Among 14 evaluable phase I patients, 2 out of 5 patients with PDAC showed partial response with tumour shrinking and 1 patient had stable disease (American Society of Clinical Oncology Meeting 2008, abstract 4625). In contrast to these results that suggest synergism of the gemcitabine/HDACi combination in patients with PDAC,³⁹ the majority of studies revealed restricted effects in a limited number of patients with distinct histological PDAC subgroups^{40–41} or even characterised combination with HDACi to be inferior to gemcitabine monotherapy in this tumour entity (NCT00004861).⁴²

As an alternative approach to chemotherapeutic agents, several trials have been conducted to evaluate the combination of HDACi and targeted therapies in PDAC treatment. Unfortunately, many HDACi combinatory treatment regimens with strong and promising mechanistic and functional synergism in preclinical PDAC models failed in first-in-patient studies, thus reflecting the difficulties of translating these findings into the clinical setting (eg, NCT00667082, NCT01056601).⁴³ Nevertheless, a plethora of preclinical data on synthetic lethal interactions exist in the context of HDACi that can be used to optimise the potential of HDAC inhibition in PDAC treatment. For instance, a combinatory treatment of HDACi with LY294002-mediated phosphoinositide 3-kinase (PI3-kinase) inhibition has recently been reported to induce apoptosis in renal cancer cells⁴⁴ and in a xenograft model of endometrial cancer.⁴⁵ These findings might be of particular interest for PDAC therapy, as dysregulation of PI3-kinase activity occurs in subgroups of PDAC⁴⁶ and might therefore represent an interesting target for synthetic-lethal approaches in the context of HDAC inhibition. Moreover, one of the neoadjuvant clinical trials investigating the efficiency of epigenetic drugs as a therapeutic option in PDAC treatment combines vorinostat-mediated HDACi with standard therapy (gemcitabine plus nab-paclitaxel or gemcitabine plus radiation) and sorafenib in patients with stage I–III PDAC (NCT02349867). The combination of HDACi

Table 2 Inhibitors of epigenetic regulators validated in clinical trials in pancreatic cancer (terminated, completed trials and active, but not recruiting clinical trials)

Compound	Epigenetic target	Combination	Treatment	NCT number	Outcome measures	Results in PDAC	Enrolled tumour entities	No. of enrolled patients
Phase I and combined phase I/II								
Vorinostat	HDAC class I/II	Marizomib	Palliative	00667082	MTD, pharmacokinetics, pharmacodynamics, toxicity and antitumour activity of the combination therapy	No antitumour activity	PDAC, NSCLC Multiple myeloma Melanoma Lymphoma	22
Vorinostat	HDAC class I/II	Radiation capecitabine	Palliative +adjuvant	00983268	MTD, tumour response, toxicity	No results yet	Periapillary adenocarcinoma PDAC	21
Vorinostat	HDAC class I/II	Radiation 5-FU	Palliative	00948688	MTD, progression-free survival 7 months after registration	No results yet	PDAC	50*
Vorinostat	HDAC class I/II	Radiation	Palliative	00831493	MTD, MOS, correlation of serum cytokine levels with symptoms and outcome	No results yet	PDAC	3
Valproate	HDAC class I	Epirubicin 5-FU cyclophosphamide	Palliative	00246103	Safety, tolerability, MTD, pharmacokinetic profile, VPA effect on histone acetylation in blood cells and tumour biopsies	Low potency, but preferable toxicity profile, might be beneficial for patients with HDAC2 overexpression	Advanced neoplasms	44
Entinostat	HDAC class I	13- <i>cis</i> retinoic acid	Palliative	00098891	MTD, dose-limiting toxicity, pharmacokinetics, tumour response	Stable disease in one patient with chemotherapy-resistant PDAC	Metastatic or advanced solid tumours or lymphomas	24
Entinostat	HDAC class I		Palliative	00020579	MTD and dose-limiting toxicity, pharmacokinetics, acetylation in blood cells, tumour response	No results yet	Not specified cancer	75*
Mocetinostat	HDAC classes I +IV	Gemcitabine	Palliative	00372437	Phase I: MTD, response rate, determination of recommended phase II dose Phase II: Overall response rate in combination with gemcitabine in gemcitabine-naïve patients in stage III and IV PDAC	Partial response and stabilised disease	Solid tumours where gemcitabine is considered as standards of care, phase II limited to PDAC	47
CHR-3996	HDAC class I		Palliative	00697879	Safety, tolerability, dose-limiting toxicity, MTD, pharmacokinetics, antitumour activity	Reduction of liver metastases in one patient with acinar pancreatic cancer	Solid tumours	40
Romidepsin	HDAC class I	Gemcitabine	Palliative	00379639	Dose-limiting toxicity, number of patients with adverse events, best overall response	4/9 patients showed stable disease; additive haematological side effects	Solid tumours	36
OTX015	Pan-BET		Palliative	02259114	Number of dose-limiting toxicities at cycle 1	No results yet	NUT midline carcinoma Breast cancer NSCLC, castrate-resistant prostate cancer, PDAC	98*
Bay1238097	Pan-BET		Palliative	02369029	Incidence of dose-limiting toxicity, tumour response	No results yet	Neoplasms	8
BI-2536	BRD4	Gemcitabine	Palliative	02215044	Occurrence of dose-limiting toxicity, survival, tumour response, CA19-9 levels, maximal concentration of the analytes in plasma	No results yet	Pancreatic neoplasms	12
Phase II non-randomised								
Panobinostat	HDAC class I/II	Bortezomib	Palliative	01056601	PFS, number of participants with tumour response, duration of response, characterisation of quality and quantity toxicity	No treatment response and severe treatment-related toxicity, study was closed ahead of schedule	PDAC	7

Continued

Table 2 Continued

Compound	Epigenetic target	Combination	Treatment	NCT number	Outcome measures	Results in PDAC	Enrolled tumour entities	No. of enrolled patients
Curcumin	p300, CBP	Gemcitabine	Palliative	00192842	Time to tumour progression, response rate, survival, clinical benefit, toxicity	In chemotherapy-naïve patients with advanced PDAC. Increased GI toxicity	PDAC	17
Curcumin	p300, CBP		Palliative	00094445	6 months survival, response rate, assessment of biological activity in blood mononuclear cells via signalling and apoptotic pathways, pharmacokinetics	No severe toxicity, curcumin showed biological activity with stable disease (1/22) and brief response (1/22) with 73% reduction of liver metastasis size	PDAC	50
BI-2536	BRD4		Palliative	00710710	Objective response, PFS, dose-limiting toxicity, overall survival, clinical benefit response	First-line therapy in patients with unresectable tumours, stable disease in 24.4% of patients; the second stage of the study which was supposed to test BI-2536 as a second-line therapy was not initiated	Pancreatic neoplasms	89
Phase II randomised CI-994	HDAC class I	Gemcitabine	Palliative	00004861	Efficiency and safety of CI-994 in combination with gemcitabine	Inferior compared with gemcitabine monotherapy with decreased quality of life	PDAC	†

* Estimated enrolment.

† Enrolment not specified.

BET, bromodomain and extraterminal; FU, fluorouracil; MOS, medium overall survival; MTD, maximum tolerated dose; NSCLC, non-small cell lung cancer; PDAC, pancreatic ductal adenocarcinoma; PFS, progression-free survival; VPA, valproate.

with sorafenib-mediated tyrosine-proteinase inhibition has been reported to show additive effects in preclinical hepatocellular carcinoma (HCC) models.⁴⁷ Until today, the mechanism of the reported synergistic or additive effects of combined HDAC inhibition and tyrosine protein kinase inhibition remains elusive. Nevertheless, the fact that erlotinib-resistant glioblastoma cells could be resensitised for tyrosine kinase inhibition upon blockade of HDAC activity⁴⁸ and preclinical data on a vorinostat-driven overcoming of erlotinib resistance in PDAC cells,⁴⁹ suggests that HDACi might represent a promising treatment option in combination with erlotinib-mediated inhibition of epidermal growth factor receptor (EGFR) signalling in patients with PDAC. However, clinical trials investigating combined EGFR inhibition/HDAC inhibition are currently limited to glioblastoma multiforme (NCT01110876), lung cancer (eg, NCT00251589) and head and neck tumours (NCT00738751).

Overall, preclinical and clinical data on HDACi as a therapeutic strategy in PDAC treatment are disappointing. To improve the potential of HDACi in PDAC treatment, preclinical studies and clinical trials need to (1) systematically dissect the impact of the different HDAC proteins on PDAC progression in order to clarify which HDAC classes represent preferable targets for HDACi and (2) should concentrate on the identification of predictive markers that allow therapeutic stratification of patients with PDAC that might benefit from HDAC inhibition. One such surrogate is the multiubiquitin chain receptor protein RAD23B, which has been recently characterised as a determinant for HDACi-induced apoptosis and is highly expressed in cutaneous T-cell lymphoma, a tumour which responds favourably towards HDACi-based therapy.⁵⁰ These results suggest that therapy response prediction to HDACi is feasible, thus potentially opening new avenues to better translate HDAC inhibition strategies into the clinic. Moreover, future translational studies on HDACi need to extend the application of HDACi in combination with established and novel drugs in order to increase the potential of HDAC blockade in PDAC treatment. As indicated in the following sections of this review, the antitumourigenic potential of HDACi is highly dependent on the molecular context of the tumour and can be significantly enhanced when combined with the right therapeutic agents (box 2).

BET proteins

While acetylation levels are regulated by HATs ('writers') and HDACs ('erasers'), acetylation marks are recognised by bromodomains, which can be found in chromatin-associated and transcription-associated proteins that drive the formation of protein complexes that mediate active transcription.¹³ The bromodomain and extraterminal (BET) domain family of proteins (BRD2, BRD3, BRD4 and BRDT) constitutes the probably best characterised group of chromatin 'reader' proteins in cancer.⁵¹ By binding to acetylated chromatin via their tandem-bromodomains, BET proteins regulate the transcription of specific subsets of genes, including those that promote cell-cycle progression and the evasion of apoptosis.⁵² Furthermore, BET proteins function as critical mediators of transcriptional elongation by promoting the recruitment and activation of the positive transcription elongation factor-b complex (P-TEFb).⁵³ Based on the importance of BET proteins in controlling important numerous cancer-relevant genes such as c-Myc, Bcl2, FosL1⁵⁴ and others, as well as the activity of the EMT-related transcription factor Twist1,⁵⁵ several potent and selective inhibitors of BET proteins (BETi) have been developed.¹³ The apparent selectivity of BETi for tumour cells appears to originate from a particular dependence of many tumour-relevant genes (notably

Table 3 Inhibitors of epigenetic regulators validated in clinical trials in pancreatic cancer (recruiting trials)

Compound	Epigenetic target	Combination	Treatment	NCT number	Outcome measures	Enrolled tumour entities	No. of enrolled patients
Phase I and combined phase I/II							
Vorinostat	HDAC I/II	Sorafenib Gemcitabine n-Paclitaxel Radiation	Neoadjuvant	02349867	Recommended phase II doses and schedule	PDAC	35*
OBP-801	HDAC (pan)		Palliative	02414516	MTD, pharmacokinetics, objective response, durability of objective response	Advanced solid tumours	36*
Romidepsin	HDAC class I		Palliative	01638533	MTD; dose-limiting toxicity, pharmacokinetics, antitumour activity, Child-Pugh classification	Lymphoma, CLL, solid tumours	132*
TEN-010	Pan-BET		Palliative	01987362	MTD, toxicity, pharmacokinetics, efficiency	Solid tumours	66*
Tazemetostat	EZH2		Palliative	01897571	MTD, objective response rate, effect of high-fat-meal on EPZ6438 bioavailability, OS, PFS, duration to response, effect of exposure to midazolam	Phase I: B-cell lymphoma, advanced solid tumours; phase II: diffuse large B-cell lymphoma, follicular lymphoma	225*
GSK2816126	EZH2		Palliative	02082977	Number of subjects with adverse events, withdrawal caused by adverse events, dose-limiting toxicity, change of clinical, cardiac and laboratory parameters, objective response rate, pharmacokinetics, compared with baseline	Not specified cancer	169*
Phase II non-randomised							
Tazemetostat	EZH2		Palliative	02601950	Objective response, PFS, pharmacokinetics, response duration, pharmacodynamics†	Malignant rhabdoid tumours, rhabdoid tumours of the kidney, atypical teratoid rhabdoid tumours, selected tumours with rhabdoid features, synovial sarcoma, INI1-negative tumours malignant rhabdoid tumour of ovary, renal medullary carcinoma, epithelioid sarcoma	150*

*Estimated enrolment.

†Fifty-five per cent disease control rate in solid tumours, responses have been restricted to SMARCB1-mutated and SMARCA4-mutated tumours. No data reported for PDAC so far. BET, bromodomain and extraterminal; CLL, chronic lymphocytic leukemia; MTD, maximum tolerated dose; OS, overall survival; PDAC, pancreatic ductal adenocarcinoma; PFS, progression-free survival.

Box 1 Inventory of epigenetic therapy in pancreatic cancer treatment

- ▶ A plethora of phase I/II clinical trials evaluate epigenetic therapies in pancreatic ductal adenocarcinoma (PDAC)
- ▶ Targeting chromatin dysregulation in PDAC can overcome cell plasticity and primary or secondary therapeutic resistance
- ▶ A hitherto underestimated vulnerability towards inhibitors of chromatin regulatory proteins exists in PDAC
- ▶ Monotherapies using epigenetic drugs have been shown to be not beneficial, but the pharmaceutical targeting of epigenetic modifications can significantly alter the susceptibility of PDAC towards standard chemotherapy
- ▶ Combination of drugs targeting diverse chromatin regulators (eg, combined inhibitors of BET proteins/histone deacetylase inhibitors) to date represents the most promising concept for epigenetic therapies in PDAC

c-Myc) on the recruitment of BRD4 to larger, composite distal regulatory regions frequently referred to as 'super enhancers'. This requirement is gene-specific and context-specific, where even for c-Myc different tumour types are dependent upon different BRD4-dependent enhancer regions.^{56 57} One of the

founding BETi molecules was JQ1, which was initially described to be a potent suppressor of NUT midline carcinoma,⁵³ B-cell lineage malignancies⁵⁸ and other tumour entities.⁵⁹ In a subcutaneous xenograft model of PDAC, Garcia *et al*⁶⁰ observed a remarkable decrease of tumour size upon JQ1 administration, indicating antitumourigenic activity in this tumour entity. Moreover, administration of JQ1 blocked acinar-to-ductal metaplasia in the pancreas, a key event in PDAC initiation, decreased formation of PanIN lesions as well as PDAC cell proliferation⁶¹ and attenuated the progenitor phenotype of dedifferentiated PDAC in favour of enhanced cellular maturity.⁶² A more recent study confirmed a beneficial effect of BETi in mouse PDAC xenograft studies and suggested a specific importance of BET proteins in controlling the activity of Gli transcription factors downstream of oncogenic Hedgehog signalling.⁶³ These data strongly support a role of the BET proteins as important drivers of both PDAC development and progression and clinical trials using different BETi agents have been initiated with great hope and expectation (NCT01987362, NCT02259114, NCT02369029). However, first results of clinical studies testing monotherapeutic applications of BETi in PDAC are discouraging: a randomised phase II trial in patients with PDAC with unresectable tumours using BI-2536, an inhibitor of the Polo-like kinase that has been shown to block BRD4 activity in vitro (NCT00710710), yielded poor response rates and the




Figure 2 Histone deacetylase (HDAC) classes and their inhibitors. According to their structural and functional qualities, HDAC proteins are grouped into three classes. Class III deacetylases (silent mating type information regulation two (SIRT) proteins) are not depicted here. Asterisks indicate inhibitors that are investigated as therapeutic options in clinical trials.

second stage of the study was not even initiated.¹³ Since it is unclear whether the dose of BI-2536 used in this study is sufficient to adequately inhibit BET protein activity *in vivo*, this result does not exclude the biological activity of BETi in PDAC. However, a recent preclinical approach in a genetically engineered mouse model of PDAC combining JQ1-mediated BET inhibition with gemcitabine also did not show a benefit in survival, when compared with single applications of the inhibitors,⁶¹ suggesting that neither BETi monotherapies nor BETi/chemotherapy combinations may represent beneficial therapeutic strategies in PDAC treatment.

In contrast to chemotherapy-based combinatory anticancer therapies, BET inhibitors applied together with non-chemotherapeutic agents have recently attracted much attention in various preclinical tumour models. Surprisingly, despite apparent opposing mechanistic effects, BET inhibitors seem to synergise with HDACi in defined tumour entities. For instance, Myc-induced murine lymphoma mouse models are highly responsive to BET inhibition, while BETi application sensitised Myc-overexpressing lymphoma cells to HDAC inhibition.⁵² Similar results have been reported in acute myelogenous leukaemia (AML), where panobinostat-mediated HDAC inhibition synergised with JQ1 to induce apoptosis in human AML cells.⁶⁴ Most importantly, this mechanism of synergism has recently been confirmed in PDAC, where therapeutic co-application of JQ1 and the FDA-approved HDACi vorinostat potently suppressed tumour growth in advanced PDAC.⁶¹ Strikingly, PDAC bearing *Kras*^{G12D};p53^{ko} mice, an established genetically

Box 2 Essential aspects that need to be considered to increase the potential of epigenetics-based therapeutic approaches in PDAC

- ▶ Molecular stratification is inevitable to perform therapeutic response prediction in PDAC and thus to alleviate therapy decision making in PDAC
- ▶ An intensified investigation of synthetic lethal interactions is essential to understand how epigenetic alterations influence each other and how they interfere with standard therapy
- ▶ Alterations in the epigenetic landscape can be used to define certain PDAC subtypes with distinct response towards standard therapy
- ▶ The elucidation of context-specific dependencies that facilitate maximal exhaustion of the therapeutic and predictive potential of epigenetic-based mechanisms in PDAC will significantly determine the success of the epigenetic translational PDAC research

engineered mouse model of PDAC,⁶⁵ displayed a significantly reduced tumour volume upon combined JQ1 and vorinostat treatment. Importantly, mice that received the combination of both epigenetic drugs did not show any signs of tumour relapse, such as regularly seen with other therapies and died as a consequence of neurological symptoms rather than tumour burden.⁶¹ This strongly indicates that BETi/vorinostat cotreatment may be sufficient to overcome therapeutic resistance in PDAC. A similar effect could be observed in a second transgenic PDAC model that is driven by deletion of the *CDKN2A* gene locus in combination with oncogenic *Kras* activation, a genetic event that regularly occurs in PDAC¹ and in experimental lung cancer models, arguing that the antitumorigenic activity of BETi/HDACi treatment is an attractive strategy in otherwise highly resistant Ras-driven cancers.⁶¹

Several mechanisms have been proposed to explain the antitumour activity of BETi in general and of combined BETi/HDACi efficiency in particular. While several reports suggest transcriptional downregulation of oncogenic c-Myc upon BET inhibition as the crucial mechanism of antitumour activity (reviewed in),⁶⁶ other studies strongly support the notion that BETi activity is frequently independent of effects on c-Myc expression.^{52–57} Recently, JQ1 has been demonstrated to suppress tumour cell growth specifically in colon cancers that are characterised by a CpG island methylator phenotype (CIMP), one of the main subtypes of colorectal cancer.⁵⁷ Genome-wide analyses led to identification of a specific BRD4-bound ‘super enhancer’ in CIMP colon cancers, which serves to promote the expression of the long non-coding RNA colon cancer-associated transcript 1 (CCAT1) and can be utilized as a marker for sensitivity to BETi. These data characterise CCAT1 as a potential clinical marker that predicts BETi responsiveness and might have a strong impact on the selection of patients who could benefit from BETi.⁵⁷ Recent studies have also uncovered resistance mechanisms against BETi,⁶⁷ demonstrating that their effective clinical application requires additional information about their precise, context-specific molecular mechanisms and biomarkers both predictive for and indicative of their biological activity.

The unexpected synergism of HDACi and BETi in cancer therapy prompted several groups to dissect the mechanisms that underlie this phenomenon. Mazur *et al*⁶¹ proposed that de-repression of p57 upon combined JQ1 and vorinostat

treatment is responsible for cell death induction upon treatment. Further mechanistic evidence comprises synergistic attenuation of c-Myc and BCL-2 expression.⁶⁴ In their lymphoma model, Bhadury *et al* identified gene sets that are similarly induced upon isolated BET or HDAC inhibition. To explain this phenomenon, the authors used a model which has been proposed for BETi in HIV, inflammation and arteriosclerosis. Specifically, they propose that P-TEFb, which is inactivated by a complex containing HEXIM1 and the 7SK snRNP in untreated cells, is transiently released from this complex following HDAC or BET inhibition, thus enabling P-TEFb recruitment to and subsequent transcriptional induction of a defined set of alternative target genes.⁵² A more recent study demonstrated that HDACi treatment blocks transcriptional elongation and results in a re-distribution of BRD4 across the genome.⁶⁸ Based on these findings, it is conceivable that BETi synergise with HDACi by targeting non-redundant regulatory mechanisms controlling transcriptional elongation of a specific subset of genes relevant for cancer progression. However, further studies will be necessary to definitively address the complex mechanisms by which BET and HDAC inhibition synergise in cancer treatment.

A BETi-similar effect on the transcriptional activity in cancer cells might be achieved by the cyclin-dependent kinase (CDK) inhibitor Dinaciclib, which also displays high potency against CDK9. BET bromodomains recruit CDK9, the catalytic subunit of P-TEFb, which catalyses the phosphorylation of serine 2 in the heptapeptide repeat sequence within the C-terminal domain of RNA polymerase II, a post-translational modification associated with transcriptional elongation.⁶⁶ Importantly, Dinaciclib has been reported to abrogate PDAC growth and progression *in vitro* and *in vivo*.^{69 70} and is currently being evaluated for its therapeutic potential in patients with PDAC in combination with the poly [ADP-ribose] polymerase (PARP) inhibitor veliparib (NCT01434316) and MK2206-mediated Akt inhibition (NCT01783171). Interestingly, CDK9 and BET inhibitors were also shown to function synergistically.⁷¹ Thus, it will be necessary to analyse overlapping and diverse functions of CDK9 inhibition and BET inhibition in cancer to determine in which context the combination of different therapeutic strategies may optimally work to have synergistic or additive effects in PDAC treatment.

Altogether, combined epigenetic targeting appears to be beneficial in certain tumour entities. Prospective clinical trials need to consider the promising effects of combined HDAC and BET targeting in preclinical PDAC models and have to validate whether this therapeutic strategy can be translated into the clinic.

Polycomb proteins

Monomethylation, dimethylation and trimethylation of histones elicit distinct changes in chromatin conformation and transcriptional activity dependent on the particular location and context of the modified lysine residues.¹⁰ For example, trimethylation of histone 3 lysine 4 by Trithorax group proteins is strongly associated with active gene expression, while the trimethylation of lysine 27 (H3K27me3) induces gene silencing.⁷² The histone mark H3K27me3 is catalysed and maintained by the polycomb repressor complex-2 (PRC2).¹⁰ The PRC2 constitutes an approximately 600 kDa complex with four subunits: Suz12, embryonic ectoderm development (EED), RaAp46/48 and the catalytic component enhancer of zeste homologue-2 (EZH2), which mediates H3K27 methylation.^{73–75} Physiologically, the polycomb proteins (PcG) function as crucial epigenetic repressors of differentiation and are responsible for the maintenance

of stem cell capacities.⁷⁶ Importantly, PcG proteins in general and EZH2 in particular are frequently overexpressed in cancer.⁷⁷ High expression levels of the histone methyltransferase regularly correlate with tumour stage and poor prognosis, while depletion of EZH2 can restrict proliferation and tumour progression.⁷⁸ The role of PRC2 proteins as epigenetic regulators of tumour progression has only recently become a focus in PDAC.⁷⁹ Ougolkov *et al*⁸⁰ showed nuclear overexpression of EZH2 in PDAC cell lines and in 68% of investigated human PDAC samples, while the nuclear accumulation of the enzyme was more prevalent in dedifferentiated PDAC (91%). Accordingly, an inverse expression profile of EZH2 and the epithelial marker protein E-cadherin⁸¹ and a robust EZH2-dependent promotion of cancer stem cell self-renewal⁷⁶ have been demonstrated in PDAC, characterising EZH2 as a strong promoter of cancer cell plasticity in the pancreas. Consequently, EZH2 inactivation sensitised PDAC cells to chemotherapy in preclinical PDAC models.^{76 80} Reports demonstrating a significant longer survival of gemcitabine-treated patients with PDAC with low pancreatic EZH2 levels⁸¹ emphasise the oncogenic capacities of the histone methyltransferase.

Its involvement in a wide range of oncogenic functions in several cancer entities has prompted the development of specific inhibitors of EZH2 activity. With tazemetostat (EPZ-6438), CPI-1205 and GSK2816126 three different EZH2 inhibitors are being assessed, for their tolerability and efficiency in clinical trials for haematological and solid malignancies. The potent inhibitor of EZH2 methyltransferase activity tazemetostat⁸² is currently being evaluated in a multicentre, open-label phase I/II study (NCT01897571). Patients with haematological and solid tumours including PDAC are enrolled in the phase I arm, while the phase II portion is restricted to lymphoma patients. Preliminary results for a tazemetostat trial investigating dose escalation and expansion of the drug in B-cell lymphoma and refractory or relapsed solid tumour patients (NCT02601950) have been presented at the European Society for Medical Oncology (ESMO)'s European Cancer Congress in Vienna, Austria in September 2015. Response assessment was conducted after 8 weeks of application. Importantly, tazemetostat monotherapy induced a 55% disease control rate in solid tumours, including rhabdoid tumours and epitheloid sarcoma. It is worth mentioning that the subgroup of responding tumours exhibited inactivating mutations of either SMARCB1 (also known as INI1 or SNF5) or SMARCA4 (also designated as BRG1). Both proteins belong to the SWItch/sucrose non-fermentable chromatin complex (SWI/SNF) chromatin remodelling complex and have been characterised for their tumour-suppressive potential. Intriguingly, mutations of the SWI/SNF complex can be found in approximately 20% of all human cancers.⁸³ The dependency of antitumour activity of EZH2 inhibition on SWI/SNF mutations has recently been described in a preclinical approach. In a detailed mechanistic study, Kim *et al* demonstrate that genetic depletion of EZH2 sufficiently blocked proliferation of a wide panel of SWI/SNF mutation containing cancer cells, while loss of EZH2 had no impact on cells harbouring wild-type expression of SWI/SNF proteins. Surprisingly, the dependency of SWI/SNF-mutated cells on PRC2 expression did not necessarily always require EZH2 enzymatic activity, but was rather based on the stabilisation of the PRC2 complex in the context of Ras or Raf mutation.⁷⁸ These data suggest the existence of methyltransferase-independent oncogenic EZH2 functions and indicate that EZH2 inhibitors may not fully repress the oncogenic activity of the enzyme under some contexts, unless they are able to intervene with EZH2 expression or its interaction

with the PRC2 complex. Whether this observation holds true for pancreatic cancer is currently unclear. Although patients with PDAC are eligible for the EZH2 inhibitor trials, no data are currently available on the enrolment of patients with PDAC into the trials or the efficiency of EZH2 blockade in PDAC. Nevertheless, since recent whole genome sequencing approaches reported frequent occurrence of SWI/SNF mutations (of complex members such as ARID1A, ARID1B, SMARCA2 and SMARCA4) in PDAC,¹⁻³ beneficial effects of EZH2 inhibition in this subgroup of patients with PDAC are worth further investigation.

Synthetic lethal interactions between EZH2 inhibition and the disturbance of other epigenetic mechanisms or signalling pathways are not limited to the SWI/SNF complex, but have been described in other contexts as well. For example, EZH2 inhibition sensitised colorectal cancer cells to EGFR inhibition by the induction of autophagy⁸⁴ and blockade of EZH2 activity increases the susceptibility of small cell lung cancer with EGFR gain-of-function mutations towards topoisomerase II inhibition.⁸⁵ Moreover, an interesting connection between PRC2 integrity and rat sarcoma (RAS) activity has been recently revealed in malignant peripheral nerve sheath tumours. Herein, deficiency of the non-catalytic PRC2 subunit Suz12 or EED has been shown to amplify RAS-driven transcription.⁸⁶ As attenuation of the H3K27me3 mark upon Suz12 or EED ablation resulted in an increase of H3K27 acetylation and subsequent recruitment of bromodomain proteins, the authors hypothesised that Suz12-deficient tumours might respond to pharmacological bromodomain blockade. Indeed, JQ1 treatment counteracted activation of RAS signalling upon Suz12 deficiency by suppressing RAS signature genes.⁸⁶ These preclinical studies indicate that the functional consequences of PRC2 inactivation are diverse across tumour entities and underscore that methyltransferase-independent functions of EZH2 should be considered to exhaust the therapeutic potential of PRC2 disruption in PDAC treatment. Moreover, the experience and knowledge gained from preclinical and clinical studies in the context of EZH2 highlight the utmost importance of molecular stratification approaches as a central component of epigenetically based cancer therapy.

CONCLUSION AND PERSPECTIVE

Epigenetic mechanisms in general, and chromatin alterations in particular, are involved in all aspects of cancer development and progression. A number of elegantly conducted translational

studies have provided significant evidence that dysregulation of chromatin organisation strongly contributes to the aggressive behaviour of PDAC, for instance, by promoting cellular plasticity and therapeutic resistance. Consequently, several clinical trials have been initiated to determine the safety and efficiency of epigenetic drugs in patients with PDAC. The small number of predominantly non-randomised studies on epigenetic targeting in this tumour entity and restrictive enrolment of patients with PDAC in some of the studies both limit the informative value of these trials. Nevertheless, the following conclusions can be drawn: (1) similar to chemotherapy, monotherapeutic targeting of epigenetic alterations in PDAC has thus far not proven to be beneficial; (2) efficiency of epigenetic drugs could be reported in selected patients with PDAC, especially following targeting of HDAC or BET proteins; (3) the mechanistic basis of the tumour cell selectivity of some epigenetic inhibitors remains largely unclear.

As demonstrated in diverse preclinical studies, a striking vulnerability of PDAC cells exists towards epigenetic-based therapy in this disease. However, targeting chromatin alterations in cancer to date is limited to a subset of histone writers, readers and erasers, while the potential targetability of other epigenetic regulators in the therapeutic landscape of cancer treatment is unknown. Whether this is a consequence of less academic and pharmaceutical research interest or a result of reduced vulnerability compared with the established targets of epigenetic drugs is unclear. For instance, it is conceivable that inhibition of selective histone demethylase activities may be effective in the treatment of tumours displaying mutations in histone methyltransferases such as *MLL2*, *MLL3* and *SETD2*,¹⁻³ while tumours with mutations in *KDM6A* might be expected to respond well to EZH2 inhibitors. It remains a tremendous hurdle to identify the most beneficial therapeutic targets of epigenetic cancer therapy and to detect those subpopulations of patients that will profit from targeting of a particular epigenetic alteration in PDAC. While initial indications suggest that *SMARCB1* and/or *SMARCA4* mutations may be predicative of tumour responsiveness to EZH2 inhibitors in rhabdoid tumours, it remains to be determined whether pancreatic tumours displaying mutations in *KDM6A*, *ARID1A*, *ARID2* or other BAF complex components also depend upon EZH2 activity. Significant preclinical and clinical efforts are required to develop a paradigm of molecular stratification to address this challenge. First advances in the prediction of responses towards epigenetic therapies⁵⁷ indicate that this is a difficult, but feasible task. Recently developed high-throughput

Table 4 Side effects of exemplary clinical trials testing epigenetic inhibitors in cancer treatment

Drug	Tumour entity	Side effects	References
AN-9 (butyric acid prodrug)	Melanoma, lung cancer, leukaemia	Moderate vomiting, fever, nausea and fatigue, mild elevation of hepatic transaminase, hyperglycaemia, transient visual toxicity, combination with docetaxel resulted in severe toxicity	28-87
Valproic acid	Leukaemia, myelodysplasia, cervical cancer	Neurological toxicity, bone pain, delayed haematological recovery	88-89
Vorinostat (plus paclitaxel plus carboplatin)	NSCLC	Febrile neutropenia, asthenia, diarrhoea, thrombocytopenia compared with the placebo group (paclitaxel+carboplatin+placebo)	NCT00473889
Panobinostat	Hodgkin's lymphoma, multiple myeloma	Thrombocytopenia, vomiting, diarrhoea, headache and abdominal pain more frequent compared with the placebo group, no significant increase of side effects when combined with bortezomib	NCT01034163, NCT01023308
CI-994 (plus gemcitabine)	Pancreatic cancer	Increased neutropaenia and thrombocytopenia compared with placebo group	42
BI-2536	Pancreatic cancer	Neutropenia, thrombocytopenia, leucopenia, fatigue	90

NSCLC, non-small cell lung cancer.

techniques can be used to identify molecular signatures that allow therapeutic response prediction to certain epigenetic drugs and alleviate therapy decision making in PDAC. This strategy is extremely important to enrol patients into the right clinical trials and to learn about the determinants of epigenetic drug responses. Intriguingly, considering molecular stratification approaches in PDAC, chromatin deregulation contributes to the development of epigenetic targeting strategies, and also influences therapy decision making in standard care, as alterations in the epigenetic landscape might characterise certain subtypes of PDAC, which are susceptible to defined (non-epigenetic) therapies. Therefore, epigenetic dysregulation in PDAC can function as both a therapeutic target and therapy-response predictor.

As a result of its tremendous cellular plasticity, monotherapeutic strategies usually fail in PDAC. Therefore, the identification of the right drug combinations should significantly impact the therapeutic response of patients with PDAC. Combinations of epigenetic drugs with standard chemotherapy or targeted therapies or even combinations of different epigenetic drugs with one another are conceivable approaches that need to be validated in preclinical and clinical settings. Due to the dynamic character of epigenetic modifications, their pharmaceutical targeting can significantly alter the susceptibility of the tumour towards standard chemotherapy, when applied in the right combination and sequence.

Intriguingly, a rising number of preclinical and clinical studies in the past years emphasise the relevance and chances of therapeutic strategies that combine epigenetic inhibitors with each other to combat cancer. The increasing understanding of the mechanistic and functional consequences of targeting central chromatin regulatory proteins in PDAC cells and the subsequent implications of these therapeutic strategies on the epigenetic balance of a tumour reveals a hitherto underestimated vulnerability of these cancer cells towards inhibitors of other chromatin-modifying or remodelling proteins with similar or inverse functions. In this context, the identification and characterisation of synthetic lethal interactions represents a crucial component to understanding how epigenetic mechanisms interact with each other and how they function within the genetic background of a tumour cell. Unfortunately, in contrast to the good tolerability of epigenetic monotherapies, combinatorial applications of epigenetic drugs with other antitumourigenic agents frequently show additive and dose-limiting toxicities (table 4).²⁸ Considering both the documented chances of epigenetic combination therapies in PDAC treatment as well as their toxicity-related limitations, extensions of synthetic lethality studies and their translation to clinical settings might significantly expand the therapeutic potential and the feasibility of epigenetic therapeutic approaches in PDAC treatment. Furthermore, the utilisation of epigenetic drugs at or near the maximum tolerated dose may not be the best approach. Given their mechanisms of action, which function primarily through gene expression changes rather than cytotoxic effects, the definition of an 'effective dose' based on biomarkers may significantly increase the potential utility of combinatorial approaches and reduce toxicity.

Without a doubt, the explosive scientific and pharmaceutical interest in epigenetic mechanisms in PDAC biology and therapy represents a significant advance in the field and contributes to our understanding of key mechanisms of PDAC biology. Finally, the success of the epigenetic era in translational PDAC research will be determined by the elucidation and creation of molecular and context-specific dependencies that guarantee maximal exhaustion of the therapeutic and predictive potential of epigenetic-based mechanisms in pancreatic cancer.

Contributors EH, SJ, JS and VE wrote the manuscript and designed the images for this review article.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Waddell N, Pajic M, Patch AM, *et al.* Whole genomes redefine the mutational landscape of pancreatic cancer. *Nature* 2015;518:495–501.
- Jones S, Zhang X, Parsons DW, *et al.* Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science* 2008;321:1801–6.
- Bailey P, Chang DK, Nones K, *et al.* Genomic analyses identify molecular subtypes of pancreatic cancer. *Nature* 2016;531:47–52.
- Weinberg BA, Yabar CS, Brody JR, *et al.* Current standards and novel treatment options for metastatic pancreatic adenocarcinoma. *Oncology* 2015;29:809–20, 886.
- Ciliberto D, Staropoli N, Chiellino S, *et al.* Systematic review and meta-analysis on targeted therapy in advanced pancreatic cancer. *Pancreatology* 2016;16:249–58.
- Singh SK, Chen N, Hessmann E, *et al.* Antithetical NFATc1-Sox2 and p53-miR200 signaling networks govern pancreatic cancer cell plasticity. *EMBO J* 2015;34:517–30.
- Sharma SV, Lee DY, Li B, *et al.* A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 2010;141:69–80.
- Roy DM, Walsh LA, Chan TA. Driver mutations of cancer epigenomes. *Protein Cell* 2014;5:265–96.
- Margueron R, Reinberg D. The polycomb complex PRC2 and its mark in life. *Nature* 2011;469:343–9.
- Lomber G, Mathison AJ, Grzenda A, *et al.* The sunset of somatic genetics and the dawn of epigenetics: a new frontier in pancreatic cancer research. *Curr Opin Gastroenterol* 2009;24:597–602.
- Moffitt RA, Marayati R, Flate EL, *et al.* Virtual microdissection identifies distinct tumor- and stroma-specific subtypes of pancreatic ductal adenocarcinoma. *Nat Genet* 2015;47:1168–78.
- Schneider G, Krämer OH, Schmid RM, *et al.* Acetylation as a transcriptional control mechanism-HDACs and HATs in pancreatic ductal adenocarcinoma. *J Gastrointest Cancer* 2011;42:85–92.
- Smith SG, Zhou M. The bromodomain: a new target in emerging epigenetic medicine. *ACS Chem Biol* 2016;11:598–608.
- Balasubramanyam K, Varier RA, Altaf M, *et al.* Curcumin, a novel p300/CREB-binding protein-specific inhibitor of acetyltransferase, represses the acetylation of histone/nonhistone proteins and histone acetyltransferase-dependent chromatin transcription. *J Biol Chem* 2004;279:51163–71.
- Koenig A, Linhart T, Schlegelmann K, *et al.* NFAT-induced histone acetylation relay switch promotes c-Myc-dependent growth in pancreatic cancer cells. *Gastroenterology* 2009;138:1189–99.e1–2.
- Mees ST, Mardin WA, Wendel C, *et al.* EP300—a miRNA-regulated metastasis suppressor gene in ductal adenocarcinomas of the pancreas. *Int J Cancer* 2009;126:114–24.
- Li M, Zhang Z, Hill DL, *et al.* Curcumin, a dietary component, has anticancer, chemosensitization, and radiosensitization effects by down-regulating the MDM2 oncogene through the PI3K/mTOR/ETS2 pathway. *Cancer Res* 2007;67:1988–96.
- Sahu RP, Batra S, Srivastava SK. Activation of ATM/Chk1 by curcumin causes cell cycle arrest and apoptosis in human pancreatic cancer cells. *Br J Cancer* 2009;100:1425–33.
- Kunnumakkara AB, Guha S, Krishnan S, *et al.* Curcumin potentiates antitumor activity of gemcitabine in an orthotopic model of pancreatic cancer through suppression of proliferation, angiogenesis, and inhibition of nuclear factor-kappaB-regulated gene products. *Cancer Res* 2007;67:3853–61.
- Epelbaum R, Schaffer M, Vizek B, *et al.* Curcumin and gemcitabine in patients with advanced pancreatic cancer. *Nutr Cancer* 2010;62:1137–41.
- Kanai M. Therapeutic applications of curcumin for patients with pancreatic cancer. *World J Gastroenterol* 2014;20:9384–91.
- Dhillon N, Aggarwal BB, Newman RA, *et al.* Phase II trial of curcumin in patients with advanced pancreatic cancer. *Clin Cancer Res* 2008;14:4491–9.
- Kanai M, Yoshimura K, Asada M, *et al.* A phase I/II study of gemcitabine-based chemotherapy plus curcumin for patients with gemcitabine-resistant pancreatic cancer. *Cancer Chemother Pharmacol* 2010;68:157–64.
- Sato N, Ohta T, Kitagawa H, *et al.* FR901228, a novel histone deacetylase inhibitor, induces cell cycle arrest and subsequent apoptosis in refractory human pancreatic cancer cells. *Int J Oncol* 2004;24:679–85.
- Haefner M, Blüthner T, Niederhagen M, *et al.* Experimental treatment of pancreatic cancer with two novel histone deacetylase inhibitors. *World J Gastroenterol* 2008;14:3681–92.

- 26 Kumagai T, Wakimoto N, Yin D, *et al.* Histone deacetylase inhibitor, suberoylanilide hydroxamic acid (vorinostat, SAHA) profoundly inhibits the growth of human pancreatic cancer cells. *Int J Cancer* 2007;121:656–65.
- 27 Glozak MA, Seto E. Histone deacetylases and cancer. *Oncogene* 2007;26:5420–32.
- 28 Minucci S, Pelicci PG. Histone deacetylase inhibitors and the promise of epigenetic (and more) treatments for cancer. *Nat Rev Cancer* 2006;6:38–51.
- 29 Meidhof S, Brabletz S, Lehmann W, *et al.* ZEB1-associated drug resistance in cancer cells is reversed by the class I HDAC inhibitor mocetinostat. *EMBO Mol Med* 2015;7:831–47.
- 30 von Burstin J, Eser S, Paul MC, *et al.* E-cadherin regulates metastasis of pancreatic cancer in vivo and is suppressed by a SNAIL/HDAC1/HDAC2 repressor complex. *Gastroenterology* 2009;137:361–71, 371.e1–5.
- 31 Schölz C, Weinert BT, Wagner SA, *et al.* Acetylation site specificities of lysine deacetylase inhibitors in human cells. *Nat Biotechnol* 2015;33:415–23.
- 32 Witkiewicz AK, McMillan EA, Balaji U, *et al.* Whole-exome sequencing of pancreatic cancer defines genetic diversity and therapeutic targets. *Nat Commun* 2015;6:6744.
- 33 Nakagawa M, Oda Y, Eguchi T, *et al.* Expression profile of class I histone deacetylases in human cancer tissues. *Oncol Rep* 2007;18:769–74.
- 34 Damaskos C, Karatzas T, Nikolidakis L, *et al.* Histone deacetylase (HDAC) inhibitors: current evidence for therapeutic activities in pancreatic cancer. *Anticancer Res* 2015;35:3129–35.
- 35 Bolden JE, Peart MJ, Johnstone RW. Anticancer activities of histone deacetylase inhibitors. *Nat Rev Drug Discov* 2006;5:769–84.
- 36 Balasubramanian S, Verner E, Buggy JJ. Isoform-specific histone deacetylase inhibitors: the next step? *Cancer Lett* 2009;280:211–21.
- 37 Walkinshaw DR, Yang XJ. Histone deacetylase inhibitors as novel anticancer therapeutics. *Curr Oncol* 2008;15:237–43.
- 38 Siu LL, Pili R, Duran I, *et al.* Phase I study of MGCD0103 given as a three-times-per-week oral dose in patients with advanced solid tumors. *J Clin Oncol* 2008;26:1940–7.
- 39 Tan J, Cang S, Ma Y, *et al.* Novel histone deacetylase inhibitors in clinical trials as anti-cancer agents. *J Hematol Oncol* 2010;3:5.
- 40 Banerji U, van Doorn L, Papadatos-Pastos D, *et al.* A phase I pharmacokinetic and pharmacodynamic study of CHR-3996, an oral class I selective histone deacetylase inhibitor in refractory solid tumors. *Clin Cancer Res* 2012;18:2687–94.
- 41 Münster P, Marchion D, Bicaku E, *et al.* Phase I trial of histone deacetylase inhibition by valproic acid followed by the topoisomerase II inhibitor epirubicin in advanced solid tumors: a clinical and translational study. *J Clin Oncol* 2007;25:1979–85.
- 42 Richards DA, Boehm KA, Waterhouse DM, *et al.* Gemcitabine plus CI-994 offers no advantage over gemcitabine alone in the treatment of patients with advanced pancreatic cancer: results of a phase II randomized, double-blind, placebo-controlled, multicenter study. *Ann Oncol* 2006;17:1096–102.
- 43 Nawrocki ST, Carew JS, Pino MS, *et al.* Aggresome disruption: a novel strategy to enhance bortezomib-induced apoptosis in pancreatic cancer cells. *Cancer Res* 2006;66:3773–81.
- 44 Yamada T, Horinaka M, Shinnoh M, *et al.* A novel HDAC inhibitor OBP-801 and a PI3K inhibitor LY294002 synergistically induce apoptosis via the suppression of survivin and XIAP in renal cell carcinoma. *Int J Oncol* 2013;43:1080–6.
- 45 Yoshioka T, Yagosawa S, Yamada T, *et al.* Combination of a novel HDAC inhibitor OBP-801/YM753 and a PI3K inhibitor LY294002 synergistically induces apoptosis in human endometrial carcinoma cells due to increase of Bim with accumulation of ROS. *Gynecol Oncol* 2013;129:425–32.
- 46 Weiss GA, Rossi MR, Khushalani NI, *et al.* Evaluation of phosphatidylinositol-3-kinase catalytic subunit (PIK3CA) and epidermal growth factor receptor (EGFR) gene mutations in pancreaticobiliary adenocarcinoma. *J Gastrointest Oncol* 2013;4:20–9.
- 47 Lachenmayer A, Toffanin S, Cabellos L, *et al.* Combination therapy for hepatocellular carcinoma: additive preclinical efficacy of the HDAC inhibitor panobinostat with sorafenib. *J Hepatol* 2012;56:1343–50.
- 48 Liffers K, Kolbe K, Westphal M, *et al.* Histone deacetylase inhibitors resensitize EGFR/EGFRVIII-overexpressing, erlotinib-resistant glioblastoma cells to tyrosine kinase inhibition. *Target Oncol* 2016;11:29–40.
- 49 Park SJ, Kim SM, Moon JH, *et al.* SAHA, an HDAC inhibitor, overcomes erlotinib resistance in human pancreatic cancer cells by modulating E-cadherin. *Tumour Biol* 2016;37:4323–30.
- 50 Stimson L, La Thangue NB. Biomarkers for predicting clinical responses to HDAC inhibitors. *Cancer Lett* 2009;280:177–83.
- 51 Belkina AC, Denis GV. BET domain co-regulators in obesity, inflammation and cancer. *Nat Rev Cancer* 2012;12:465–77.
- 52 Bhadury J, Nilsson LM, Muralidharan SV, *et al.* BET and HDAC inhibitors induce similar genes and biological effects and synergize to kill in Myc-induced murine lymphoma. *Proc Natl Acad Sci USA* 2014;111:E2721–30.
- 53 Filippakopoulos P, Qi J, Picaud S, *et al.* Selective inhibition of BET bromodomains. *Nature* 2010;468:1067–73.
- 54 Shi J, Vakoc CR. The mechanisms behind the therapeutic activity of BET bromodomain inhibition. *Mol Cell* 2014;54:728–36.
- 55 Shi J, Wang Y, Zeng L, *et al.* Disrupting the interaction of BRD4 with diacetylated Twist suppresses tumorigenesis in basal-like breast cancer. *Cancer Cell* 2014;25:210–25.
- 56 Chapuy B, McKeown MR, Lin CY, *et al.* Discovery and characterization of super-enhancer-associated dependencies in diffuse large B cell lymphoma. *Cancer Cell* 2013;24:777–90.
- 57 McClelland ML, Mesh K, Lorenzana E, *et al.* CCAT1 is an enhancer-templated RNA that predicts BET sensitivity in colorectal cancer. *J Clin Invest* 2016;126:639–52.
- 58 Delmore JE, Issa GC, Lemieux ME, *et al.* BET bromodomain inhibition as a therapeutic strategy to target c-Myc. *Cell* 2011;146:904–17.
- 59 Jung M, Gelato KA, Fernández-Montalván A, *et al.* Targeting BET bromodomains for cancer treatment. *Epigenomics* 2015;7:487–501.
- 60 Garcia PL, Miller AL, Kreitzburg KM, *et al.* The BET bromodomain inhibitor JQ1 suppresses growth of pancreatic ductal adenocarcinoma in patient-derived xenograft models. *Oncogene* 2016;35:833–45.
- 61 Mazur PK, Herner A, Mello SS, *et al.* Combined inhibition of BET family proteins and histone deacetylases as a potential epigenetics-based therapy for pancreatic ductal adenocarcinoma. *Nat Med* 2015;21:1163–71.
- 62 Roy N, Malik S, Villanueva KE, *et al.* Brg1 promotes both tumor-suppressive and oncogenic activities at distinct stages of pancreatic cancer formation. *Genes Dev* 2015;29:658–71.
- 63 Huang Y, Nahar S, Nakagawa A, *et al.* Regulation of GLI underlies a role for BET bromodomains in pancreatic cancer growth and the tumor microenvironment. *Clin Cancer Res* 2016;22:4259–70.
- 64 Fiskus W, Sharma S, Qi J, *et al.* Highly active combination of BRD4 antagonist and histone deacetylase inhibitor against human acute myelogenous leukemia cells. *Mol Cancer Ther* 2014;13:1142–54.
- 65 Hingorani SR, Petricoin EF, Maitra A, *et al.* Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* 2004;4:437–50.
- 66 Hessmann E, Schneider G, Ellenrieder V, *et al.* MYC in pancreatic cancer: novel mechanistic insights and their translation into therapeutic strategies. *Oncogene* 2016;35:1609–18.
- 67 Kurimchak AM, Shelton C, Duncan KE, *et al.* Resistance to BET bromodomain inhibitors is mediated by kinome reprogramming in ovarian cancer. *Cell Rep* 2016;16:1273–86.
- 68 Greer CB, Tanaka Y, Kim YJ, *et al.* Histone deacetylases positively regulate transcription through the elongation machinery. *Cell Rep* 2015;13:1444–55.
- 69 Allaway RJ, Fischer DA, de Abreu FB, *et al.* Genomic characterization of patient-derived xenograft models established from fine needle aspirate biopsies of a primary pancreatic ductal adenocarcinoma and from patient-matched metastatic sites. *Oncotarget* 2016;7:17087–102.
- 70 Feldmann G, Mishra A, Bisht S, *et al.* Cyclin-dependent kinase inhibitor Dinaciclib (SCH727965) inhibits pancreatic cancer growth and progression in murine xenograft models. *Cancer Biol Ther* 2011;12:598–609.
- 71 Lu H, Xue Y, Xue Y, *et al.* Compensatory induction of MYC expression by sustained CDK9 inhibition via a BRD4-dependent mechanism. *Life* 2015;4:e06535.
- 72 Grzenda A, Ordog T, Urrutia R. Polycomb and the emerging epigenetics of pancreatic cancer. *J Gastrointest Cancer* 2011;42:100–11.
- 73 Margueron R, Justin N, Ohno K, *et al.* Role of the polycomb protein EED in the propagation of repressive histone marks. *Nature* 2009;461:762–7.
- 74 Pasini D, Bracken AP, Jensen MR, *et al.* Suz12 is essential for mouse development and for EZH2 histone methyltransferase activity. 2004;23:4061–71.
- 75 Cao R, Zhang Y. SUZ12 is required for both the histone methyltransferase activity and the silencing function of the EED-EZH2 complex. *Mol Cell* 2004;15:57–67.
- 76 Avan A, Crea F, Paolicchi E, *et al.* Molecular mechanisms involved in the synergistic interaction of the EZH2 inhibitor 3-deazaneplanocin A with gemcitabine in pancreatic cancer cells. *Mol Cancer Ther* 2012;11:1735–46.
- 77 Völkel P, Dupret B, Le Bourhis X, *et al.* Diverse involvement of EZH2 in cancer epigenetics. *Am J Transl Res* 2015;7:175–93.
- 78 Kim KH, Kim W, Howard TP, *et al.* SWI/SNF-mutant cancers depend on catalytic and non-catalytic activity of EZH2. *Nat Med* 2015;21:1491–6.
- 79 Neureiter D, Jäger T, Ocker M, *et al.* Epigenetics and pancreatic cancer: pathophysiology and novel treatment aspects. *World J Gastroenterol* 2014;20:7830–48.
- 80 Ougolkov AV, Bilim VN, Billadeau DD. Regulation of pancreatic tumor cell proliferation and chemoresistance by the histone methyltransferase enhancer of zeste homologue 2. *Clin Cancer Res* 2008;14:6790–6.
- 81 Toll AD, Dasgupta A, Potoczek M, *et al.* Implications of enhancer of zeste homologue 2 expression in pancreatic ductal adenocarcinoma. *Hum Pathol* 2010;41:1205–9.
- 82 Knutson SK, Warholik NM, Johnston LD, *et al.* Synergistic anti-tumor activity of EZH2 inhibitors and glucocorticoid receptor agonists in models of germinal center non-Hodgkin lymphomas. *PLoS ONE* 2014;9:e111840.
- 83 Zinzalla G. A new way forward in cancer drug discovery: inhibiting the SWI/SNF chromatin remodelling complex. *Chembiochem* 2016;17:677–82.
- 84 Katona BW, Liu Y, Ma A, *et al.* EZH2 inhibition enhances the efficacy of an EGFR inhibitor in suppressing colon cancer cells. *Cancer Biol Ther* 2014;15:1677–87.
- 85 Fillmore CM, Xu C, Desai PT, *et al.* EZH2 inhibition sensitizes BRG1 and EGFR mutant lung tumours to Topoll inhibitors. *Nature* 2015;520:239–42.

- 86 De Raedt T, Beert E, Pasmant E, *et al.* PRC2 loss amplifies Ras-driven transcription and confers sensitivity to BRD4-based therapies. *Nature* 2014;514:247–51.
- 87 Patnaik A, Rowinsky EK, Villalona MA, *et al.* A phase I study of pivaloyloxymethyl butyrate, a prodrug of the differentiating agent butyric acid, in patients with advanced solid malignancies. *Clin Cancer Res* 2002;8:2142–8.
- 88 Tassara M, Döhner K, Brossart P, *et al.* Valproic acid in combination with all-trans retinoic acid and intensive therapy for acute myeloid leukemia in older patients. *Blood* 2014;123:4027–36.
- 89 Pilatino C, Cilloni D, Messa E, *et al.* Increase in platelet count in older, poor-risk patients with acute myeloid leukemia or myelodysplastic syndrome treated with valproic acid and all-trans retinoic acid. *Cancer* 2005;104:101–9.
- 90 Mross K, Dittrich C, Aulitzky WE, *et al.* A randomised phase II trial of the Polo-like kinase inhibitor BI 2536 in chemo-naïve patients with unresectable exocrine adenocarcinoma of the pancreas—a study within the Central European Society Anticancer Drug Research (CESAR) collaborative network. *Br J Cancer* 2012; 107:280–6.