

# Leveraging Health-Related Quality of Life in Population Health Management: The Case for Healthy Days

S. Lane Slabaugh, PharmD, MBA,<sup>1</sup> Mona Shah, MPH, PhD,<sup>2</sup> Matthew Zack, MD, MPH,<sup>3</sup>  
Laura Happe, PharmD, MPH,<sup>1</sup> Tristan Cordier, MPH,<sup>1</sup> Eric Havens, MA,<sup>1</sup>  
Evan Davidson, MD,<sup>1</sup> Michael Miao, MD,<sup>1</sup> Todd Prewitt, MD,<sup>1</sup> and Haomiao Jia, PhD<sup>4,5</sup>

## Abstract

Measuring population health with morbidity and mortality data, often collected at the site of care, fails to capture the individual's perspective on health and well-being. Because health happens outside the walls of medical facilities, a holistic and singular measure of health that can easily be captured for an entire population could aid in understanding the well-being of communities. This paper postulates that Healthy Days, a health-related quality of life measure developed and validated by the Centers for Disease Control and Prevention, is an ideal survey instrument to advance population health. A systematic literature review was conducted and revealed a strong evidence base using Healthy Days with significant correlations to chronic disease conditions. Building on the literature base and experience, methods for analyzing Healthy Days data are discussed, including stratified sampling techniques, statistical measures to account for variance, and modeling techniques for skewed distributions. Using such analytic techniques, Healthy Days has been used extensively in national health surveillance. As the health care system faces increasing costs and constrained resources, the Healthy Days survey instrument can be used to inform public policies and allocate health service resources. Because Healthy Days captures broad dimensions of health from the individual's perspective, it is a simple way to holistically measure the health and well-being of a population and its trend over time. Expanded use of Healthy Days can aid population health managers and contribute to the understanding of the broader determinants of the nation's and individual community's health and aid in evaluating progress toward health goals.

## Introduction

SEVERAL DIFFERENT GROUPS ARE ACCOUNTABLE for the health and medical care of communities. Some, such as the Indian Health Service, have wide responsibility to provide all health programs for a distinct population. Most, however, manage only certain aspects of health programs or work within less defined communities. For example, state or local health departments greatly influence public health initiatives but generally are not responsible for directly providing health care. Health care providers, on the other hand,

provide medical care to their patients but usually cannot affect social determinants of health in their communities. Given this fragmented system of responsibility for overall health in the United States, it can be difficult for any of these groups that are accountable for community medical care to get a complete look at the health of their community.<sup>1,2</sup>

Traditional measures of health, such as morbidity and mortality, are important health indicators but fall short of incorporating the voice of individuals. Health-related quality of life (HRQOL)—a multidimensional concept encompassing physical, mental, emotional, and social functioning—is

---

<sup>1</sup>Humana Inc., Louisville, Kentucky.

<sup>2</sup>Robert Wood Johnson Foundation, Princeton, New Jersey.

<sup>3</sup>Centers for Disease Control and Prevention, Atlanta, Georgia.

<sup>4</sup>Mailman School of Public Health, Columbia University, New York, New York.

<sup>5</sup>School of Nursing, Columbia University, New York, New York.

widely accepted as a valid measure of the burden of disease as experienced directly by individuals, providing a more holistic view of overall health.<sup>3-5</sup> Furthermore, HRQOL is highly correlated with traditional measures of health, including morbidity, mortality, and health care costs.<sup>6-10</sup>

Given its multidimensional nature, HRQOL is an important measure of population health.<sup>3</sup> When applied to a population, HRQOL assessment provides the global picture that is missing from the somewhat limited perspective of various agencies and professionals. HRQOL is one of the 4 Foundation Health Measures used by Healthy People 2020 to assess the health of communities.<sup>11</sup> HRQOL can be used to measure health disparities, track the influence of social determinants on overall health, and shed light on the ultimate impact of the health care system. However, it can be challenging to select the right HRQOL instrument. Although several available instruments have demonstrated validity and reliability, they vary in scope, intended purpose, applicability, and general ease of use. To measure HRQOL, the ideal instrument for those who manage population health would be holistic and easy to administer, and would measure the individual's perspective while being understandable to health care providers and the general public alike.

### Healthy Days

During the late 1980s, the Centers for Disease Control and Prevention (CDC) sought to develop a survey instrument to capture self-perceived HRQOL in a short questionnaire that was easy for people to understand and answer. The resulting tool—the CDC-HRQOL-4, or “Healthy Days”—consists of 4 questions that ask people about how they perceive their own recent health (Table 1). Answers to the second question about recent physical health and the third question about recent mental health provide a summary index of *unhealthy days* (UHD) for an individual. When a respondent replies with zero days to questions 2 and 3, the respondent is not asked the final question on activity limitation, which is imputed as zero.

The CDC developed this questionnaire as an alternative to other, longer HRQOL measures for large population-based surveys; it is easily understood by the public and policy makers. Its value lies in its simplicity. It is also unique in that it measures physical and mental health as separate domains. Starting in 1993, Healthy Days was incorporated into the state-based Behavioral Risk Factor Surveillance System (BRFSS), a continuous, state-based, random telephone survey of community-dwelling US adults aged 18 and older. The inclusion of Healthy Days in the BRFSS survey has become one of the greatest strengths of the instrument, as this survey provides health-related information for a multi-year sample across the US population. A response rate of >98% to the Healthy Days questions demonstrates that these

questions are easy for people to answer.<sup>12</sup> No other measures of HRQOL, such as the Medical Outcomes Study Short-forms (SF-12 and SF-36) or the EuroQoL Group's EQ-5D, are available in a similarly large annual sample as in the BRFSS, which tracks trends over time and in different geographic areas. Another national survey, the National Health and Nutrition Examination Survey (NHANES), also included the Healthy Days questions from 2000 through 2011, and the Medicare Health Outcomes Survey has included them since 2003.

In 2000 the CDC National Center for Chronic Disease Prevention and Health Promotion's Division of Adult and Community Health published a report, *Measuring Healthy Days: Population Assessment of Health-Related Quality of Life*.<sup>11</sup> This report describes the aims of the Healthy Days survey instrument and the research that has demonstrated its validity and reliability. This research shows that the questions have high construct validity (do a good job of measuring the intended concepts) and concurrent validity (its results align with other validated instruments that measure the same concepts). The validity of Healthy Days has been confirmed across elderly, adolescent, and disabled population segments, as well as in specific clinical populations such as those with spinal cord injuries.<sup>9,13-17</sup> Furthermore, the Healthy Days questions are a valid and reliable measure of population health in distinct geographic areas including Canada and Puerto Rico, as well as at the state and local level within the United States.<sup>18-20</sup> Healthy Days was recently suggested as a potential measure of population health within the Triple Aim (population health, patient experience, and per capita cost).<sup>21</sup>

Given its demonstrated validity and reliability, its unique simplicity, and its ability to reflect people's perceptions of their own health, Healthy Days can be an effective metric of overall health for those who manage the health of population groups, whether an organization's responsibilities involve community interventions, disease management, or delivery of clinical care. This paper will review the existing published research using Healthy Days as a measure of HRQOL, describe practical considerations in collection and analysis of the data, and show how the survey instrument can be used to inform public policy.

### Use of Healthy Days in the Literature

To quantify the scope and current use of the Healthy Days survey instrument, a systematic literature review was conducted using the PubMed and Google Scholar databases and the following search terms *Healthy Days*, *Unhealthy Days*, *HRQOL-4*, and the 4 questions. The search was limited to English, and no constraints were placed on publication date. Two independent reviewers screened the publications retrieved and excluded publications of study populations younger than 18 years of age or in other than North American

TABLE 1. CDC-HRQOL-4 SURVEY INSTRUMENT—HEALTHY DAYS

1	Would you say that in general your health is excellent, very good, good, fair, or poor?
2	Now thinking about your physical health, which includes physical illness and injury, for how many days during the past 30 days was your physical health not good?
3	Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?
4	During the past 30 days, for about how many days did poor physical or mental health keep you from doing your usual activities, such as self-care, work, or recreation?

CDC-HRQOL = Centers for Disease Control and Prevention Health-Related Quality of Life.



TABLE 2. STUDIES REPORTING MEAN UNHEALTHY DAYS BY DISEASE STATE  
(ALPHABETICALLY BY DISEASE STATE)

Study	Study Overview	Data Source	Disease State	Physical Unhealthy Days	Mental Unhealthy Days	Total Unhealthy Days
Austin S, 2012	Retrospective, cross-sectional study of respondents age >45 years with self-reported arthritis	BRFSS, 2007	Arthritis	Arthritis: 7.7 days Arthritis, physically active: 5.3 days Arthritis, not physically active: 9.3 days	Arthritis: 4.4 days Arthritis, physically active: 3.3 days Arthritis, not physically active: 5.1 days	—
Mili F, 2003	Retrospective study of 54,154 respondents	BRFSS, 1996–1999	Arthritis	—	—	Arthritis: 7.5 days* No arthritis: 3.7 days
Abeil JE, 2005	Retrospective study of 212,000 respondents aged >18 years	BRFSS, 2001	Arthritis	Arthritis: 6.7 days No arthritis: 1.8 days	Arthritis: 4.9 days No arthritis: 2.7 days	—
Shih M, 2008	Retrospective study of 8648 households in California	Los Angeles County Healthy Survey, 2005	Arthritis	Arthritis: 6.4 days No arthritis: 3.3 days	Arthritis: 4.0 days No arthritis: 3.3 days	—
Quick V, 2015	Retrospective, case-control study of 2625 respondents aged 18–26 years	Survey	Bowl disease	Bowel condition: 5.7 days No bowel condition: 3.0 days	Bowel condition: 8.5 days No bowel condition: 6.0 days	—
Richardson LC, 2011	Retrospective study of 12,191 respondents living in North Carolina	BRFSS, 2001–2002	Cancer	Cancer survivors: 6.4 days No history of cancer: 3.4 days	Cancer survivors: 3.7 days No history of cancer: 2.6 days	—
Brown DW, 2004	Retrospective study of 37,054 respondents aged >50 years	BRFSS, 2001	Diabetes	Diabetes: 8.4 days No diabetes: 4.0 days	Diabetes: 3.3 days No diabetes: 2.4 days	Diabetes: 10.0 days No diabetes: 5.6 days
Cook EL, 2008	Retrospective study of 4833 respondents aged >18 years	NHANES, 2003–2004	Diabetes	—	—	Diabetes: 12.3 days No diabetes: not reported
Huang DL, 2013	Retrospective, cross-sectional study of 379,021 respondents aged >65 years	BRFSS, 2006, 2008, 2010	Diabetes	Diabetes 8.4: days No diabetes: 4.7 days	Diabetes: 3.0 days No diabetes: 2.0 days	Diabetes: 9.5 days No diabetes: 5.8 days

(continued)

TABLE 2. (CONTINUED)

<i>Study</i>	<i>Study Overview</i>	<i>Data Source</i>	<i>Disease State</i>	<i>Physical Unhealthy Days</i>	<i>Mental Unhealthy Days</i>	<i>Total Unhealthy Days</i>
Smith DW, 2004	Retrospective study of 84,308 participants	BRFSS, 1996–2000	Diabetes	Diabetes: 8.5 days No diabetes: 4.1 days	Diabetes: 4.2 days No diabetes: 2.6 days	—
Shih M, 2008	Retrospective study of 8648 households in California	Los Angeles County Healthy Survey, 2005	Diabetes	Diabetes: 6.6 days No diabetes: 3.6 days	Diabetes: 4.0 days No diabetes: 3.4 days	—
Wubben DP, 2005	Retrospective study of 1035 participants with self-reported diabetes	BRFSS, 1998–2001	Diabetes	Diabetes: 9.3 days No diabetes: not reported	Diabetes 4.5 days No diabetes: not reported	—
Quinlivan EB, 2015	Retrospective, cross-sectional study of 921 HIV+ women	Women of Color Initiative	HIV	Urban: 5.2 days Rural: 4.4 days	Urban: 9.1 days Rural: 9.1 days	—
Cook EL, 2008	Retrospective study of 4833 respondents aged >18 years	NHANES, 2003–2004	Heart Disease	—	—	Hypertension: 12.2 days No hypertension: not reported
Shih M, 2008	Retrospective study of 8648 households in California	Los Angeles County Healthy Survey, 2005	Heart Disease	Heart disease: 6.3 days No heart disease: 3.7 days	Heart disease: 4.1 days No heart disease: 3.4 days	—
Shih M, 2008	Retrospective study of 8648 households in California	Los Angeles County Healthy Survey, 2005	Mental Health	Serious psychologic disorder: 8.4 days No serious psychologic disorder: 3.6 days Depression: 5.9 days No depression: 3.6 days	Serious psychologic disorder: 14.9 No serious psychologic disorder: 2.9 days Depression: 8.2 days No depression: 2.8 days	—
Cook EL, 2008	Retrospective study of 4833 respondents aged >18 years	NHANES, 2003–2004	Mental Disorder	—	—	Mental health disorder: 17.6 days No mental health disorder: not reported

(continued)

TABLE 2. (CONTINUED)

Study	Study Overview	Data Source	Disease State	Physical Unhealthy Days	Mental Unhealthy Days	Total Unhealthy Days
Hassan MK, 2003	Retrospective, cross-sectional study of 182,372 respondents	BRFSS, 2000	Obesity	Severely obese: 6.5 days Obese: 4.3 days Overweight 3.3 days	Severely obese: 5.2 days Obese: 3.6 days Overweight: 2.9 days	—
Ford ES, 2001	Retrospective study of 109,076 respondents	BRFSS, 1996	Obesity	Non-overweight: 3.0 days BMI Ranges <18.5: 4.0 days 18.5 to <25: 2.5 days 25-29.9: 2.7 day 30 to <35: 3.8 day 35 to <40: 5.3 day ≥40: 6.5 days	Non-overweight: 3.2 days BMI Ranges <18.5: 4.0 days 18.5 to <25: 2.8 days 25 to <30: 2.5 days 30 to <35: 3.3 35 to <40: 4.4 days ≥40: 4.9 days	BMI Ranges <18.5: 6.0 days 18.5 to <25: 4.9 days 25 to <30: 5.1 30 to <35: 5.9 35 to <40: 7.4 ≥40: 8.3 days Back and neck pain: 12.2 days
Cook EL, 2008	Retrospective study of 4833 respondents aged >18 years	NHANES, 2003–2004	Pain	—	—	No back and neck pain: not reported Asthma: 10.1 days No asthma: not reported COPD: 13.7 days No COPD: 6.1 days
Ford ES, 2004	Retrospective study of 12,111 participants with asthma	BRFSS, 2000	Pulmonary Disease	Asthma: 6.5 days No asthma: not reported	Asthma: 5.3 days No asthma: not reported	—
Brown DW, 2010	Retrospective, cross-sectional study of 13,887 respondents aged >18 years living in North Carolina	BRFSS, 2007	Pulmonary Disease	—	—	—
Ford ES, 2003	Retrospective study of 163,773 respondents aged >18 years	BRFSS, 2000	Pulmonary Disease	Current asthma: 6.5 days Former asthma: 3.8 days No asthma: 2.9 days Chronic respiratory conditions: 6.0 days No chronic respiratory conditions: 3.8 days	Current asthma: 5.2 days Former asthma: 3.8 days No asthma: 3.0 days Chronic respiratory conditions: 4.5 days No chronic respiratory conditions: 3.4 days	Current asthma: 10.0 days Former asthma: 6.8 days No asthma: 5.4 days
Shih M, 2008	Retrospective study of 8648 households in California	Los Angeles County Healthy Survey, 2005	Pulmonary disease	Current asthma: 5.6 days No current asthma: 3.7 days	Current asthma: 3.6 days No current asthma: 3.4 days	—

(continued)

TABLE 2. (CONTINUED)

Study	Study Overview	Data Source	Disease State	Physical Unhealthy Days	Mental Unhealthy Days	Total Unhealthy Days
Morgen K, 2007	Retrospective study of 303 participants from 3 therapeutic community substance abuse programs in New York, NY	Custom Survey and BRFSS, 2002–2006	Substance Abuse	Substance use: 6.34 days No substance use: 3.5 days	Substance use: 9.31 days No substance use: 3.3 days	—

\*Subset of population aged >65 years  
BMI=body mass index; BRFSS=Behavioral Risk Factor Surveillance System; COPD=chronic obstructive pulmonary disorder; HIV=human immunodeficiency disorder; NHANES=National Health and Nutrition Examination Survey.

## References

- Abell JE, Hootman JM, Zack MM, Moriarty D, Helmick CG. Physical activity and health related quality of life among people with arthritis. *J Epidemiol Community Health*. 2005;59:380–385.
- Austin S, Qu H, Shewchuk RM. Association between adherence to physical activity guidelines and health-related quality of life among individuals with physician-diagnosed arthritis. *Qual Life Res*. 2012;21:1347–1357.
- Brown DW, Balluz LS, Giles WH, et al. Diabetes mellitus and health-related quality of life among older adults. Findings from the behavioral risk factor surveillance system (BRFSS). *Diabetes Res Clin Pract*. 2004;65:105–115.
- Brown DW, Pleasants R, Ohar JA, et al. Health-related quality of life and chronic obstructive pulmonary disease in North Carolina. *N Am J Med Sci*. 2010;2:60–65.
- Cook EL, Harman JS. A comparison of health-related quality of life for individuals with mental health disorders and common chronic medical conditions. *Public Health Rep*. 2008;123:45–51.
- Ford ES, Moriarty DG, Zach MM, Mokdad AH. Self-reported body mass index and health-related quality of life: findings from the Behavioral Risk Factor Surveillance System. *Obes Res*. 2001;9:21–31.
- Ford ES, Mannino DM, Homa DM, et al. Self-reported asthma and health-related quality of life: findings from the behavioral risk factor surveillance system. *Chest*. 2003;123:119–127.
- Ford ES, Mannino DM, Redd SC, Moriarty DG, Mokdad AH. Determinants of quality of life among people with asthma: findings from the Behavioral Risk Factor Surveillance System. *J Asthma*. 2004;41:327–336.
- Hassan MK, Joshi AV, Madhavan SS, Amonkar MM. Obesity and health-related quality of life: a cross-sectional analysis of the US population. *Int J Obes Relat Metab Disord*. 2003;27:1227–1232.
- Huang DL, Chan KC, Young BA. Poor oral health and quality of life in older U.S. adults with diabetes mellitus. *J Am Geriatr Soc*. 2013;61:1782–1788.
- Mili F, Helmick CG, Moriarty DG. Health related quality of life among adults reporting arthritis: analysis of data from the Behavioral Risk Factor Surveillance System, US, 1996–99. *J Rheumatol*. 2003;30:160–166.
- Morgen K, Astone-Twerell J, Hermiche T, Gunnesson L, Santangelo K. Health-related quality of life among substance abusers in residential drug abuse treatment. *Appl Res Qual Life*. 2007;2:239–246.
- Quick V, McWilliams R, Byrd-Bredbenner C. A case-control study of current psychological well-being and weight-teasing history in young adults with and without bowel conditions. *J Hum Nutr Diet*. 2015;28(1):28–36.
- Quinlivan EB, Fletcher J, Eastwood EA, Blank AE, Verdecias N, Roytburd K. Health status of HIV-infected women entering care: baseline medical findings from the women of color initiative. *AIDS Patient Care STDS*. 2015;29 suppl 1:S11–S19.
- Richardson LC, Townsend JS, Fairley TL, et al. Use of 2001–2002 Behavioral Risk Factor Surveillance System data to characterize cancer survivors in North Carolina. *N C Med J*. 2011;72:20–27.
- Shih M, Simon PA. Health-related quality of life among adults with serious psychological distress and chronic medical conditions. *Qual Life Res*. 2008;17:521–528.
- Smith DW. The population perspective on quality of life among Americans with diabetes. *Qual Life Res*. 2004;13:1391–400.
- Wubben DP, Porterfield D. Health-related quality of life among North Carolina adults with diabetes mellitus. *N C Med J*. 2005;66:179–185.

Adjustments for differential sampling probabilities driven by stratified random sampling and adjustment for nonresponse must be considered for sample surveys so that aggregate results can be considered representative of the underlying population. Adjustments for oversampling typically present little issue when the base population is known. In this case, each participant's response can be weighted according to the relative size, in the underlying population, of the stratum (subgroup) to which that participant belongs. However, adjustments for nonresponse can present a larger challenge. If survey nonresponse within each stratum is associated with particular participant characteristics, then further adjustment, or reweighting, is necessary after adjustment for oversampling so that the reweighted survey results are fully applicable to the underlying population. Iterative proportional fitting is the technique currently used in the BRFSS survey to adjust for nonresponse. Numerous additional methods can be considered to adjust for nonresponse including poststratification, such as weighting using logistic regression and replicate weighting. Ultimately the decision is driven by available data for the survey sample and the target population, in addition to the volume of the observed differences. Data necessary for weighting may have to be drawn from outside sources. For example, organizations conducting surveys in Medicare populations can obtain data such as race/ethnicity or income from the Centers for Medicare & Medicaid Services (CMS).

Once results have been weighted, researchers need to consider several factors in analyzing Healthy Days data. Mean UHD values and percentages for response categories for the target population can be estimated along with 95% confidence intervals accounting for the survey design using variance estimation techniques now available in most standard statistical software packages.

As an alternative to a mean value for each Healthy Days measure, many researchers construct "total" UHD as a sum of mentally UHD and physically UHD. Although the calculated sum of UHD can exceed 30 days, most investigators cap this sum at 30 days for purposes of interpretability. However, capping in this way can be misleading. Capping precludes identification of those with the poorest health (ie, those with the equivalent of more than 30 overall UHD). As an example, the response of 15 physically UHD and 16 mentally UHD implies an overlap of between 1 day (minimal overlap) and 15 days (maximal overlap). In other words, the respondent experienced anywhere from 1 to 15 days that were *both* physically and mentally unhealthy and 1 day that was only mentally unhealthy. A cap of 30 would not reflect the fact that at least 1 day was characterized by both physical and mental poor health.

For more complex analysis of the factors associated with UHD, different forms of multivariate regression allow researchers to simultaneously take into account numerous factors and to estimate the degree of impact for each factor. Generalized linear modeling (GLM) using ordinary least squares regression has been used in several published studies to examine mean UHD. However, each Healthy Days question has a highly skewed distribution, with zero being the most common response in most populations. Because of the nature of the question requiring recall, many respondents report Healthy Days in intervals of 5. The resulting distribution has peaks and therefore may violate the assumption of normal residuals in the GLM. Converting UHD data into a binary (frequently <14 UHD or  $\geq 14$  UHD), ordinal, or categorical variable to perform

logistic, multinomial, or ordinal regression can be a helpful way to relax assumptions regarding the distribution of UHD in continuous form. Many popular software packages today include a survey GLM and logistic regression procedures, making these an attractive option.

Less well-known modeling techniques also can be considered. One option is the zero-inflated negative binomial (ZINB) regression model, which was developed for count variables with excessive zero-value observations. Although one published study has shown the skewed nature of Healthy Days to fit well to the ZINB,<sup>25</sup> none of the popular software packages (at the time of writing) have this capability built in for survey data. The ZINB model also is complicated to communicate, and makes several underlying assumptions that do not apply to UHD; for example, the data range from zero to infinity. Additionally, ZINB does not present a uniform solution, as some subpopulations, particularly those with disability or chronic disease, will not present with the excess zeroes seen in healthier populations. Tobit, Spline and censored least absolute deviations models make no assumptions regarding the underlying distribution and both are commonly used to analyze HRQOL scores, particularly in studies with censored/truncated data or data with strong ceiling/floor effects.<sup>26-28</sup> The type of censoring addressed by these models occurs with use of a scale in which all values above or below a threshold value are counted as equal to the threshold. Ceiling/floor effects occur when the bulk of values are at either end of the scale, which would be the case in very healthy or very unhealthy populations. Although these models have not been used in the published literature to analyze Healthy Days, their use is plausible as the scores can be considered censored at zero and 30 days and responses are often skewed toward zero. However, few software packages have these methods available.

An alternative modeling approach is use of regression methodologies that do not make assumptions regarding the underlying distribution. The bootstrap method provides results that are at once robust and easy to explain.<sup>29</sup> Bootstrapping is a nonparametric approach that uses simulated resampling of the study sample and recalculation of the estimate many times to replace the traditional distributional assumptions when calculating a confidence interval or other measure of the variance in estimate. Complications related to the survey design can be accounted for using estimates from survey regression procedures as the basis of the simulation. One shortcoming that may be encountered in more complex regression models such as bootstrapping is their computationally intensive nature, which can slow the modeling process if multiple simulations are required.

If variance estimation is less important and the analyst is primarily concerned with understanding what factors are associated with increased or decreased UHD, decision tree regression provides an additional alternative for analysis without assumptions regarding underlying distribution. In decision tree regression, a large number of factors with potential association with UHD can be considered simultaneously and ranked according to their prevalence in the population and the extent to which they differentiate between individuals with more or fewer UHD. For trended analysis, structured time-series autoregressive integrated moving average modeling has been used successfully to highlight the seasonal effect associated with Healthy Days data while accounting for worsening physical and mental health over time.<sup>30</sup>



As with all instruments, the Healthy Days survey instrument has limitations that should be noted. First, the instrument was validated for each question individually. Thus, although the sum of mentally UHD and physically UHD is commonly used to summarize overall health in a single value, this application has not yet been fully validated. Furthermore, unlike other preference-based HRQOL measures such as EQ-5D and SF-6D, total UHD cannot be used directly to calculate quality-adjusted life-years (QALYs). However, regression analysis using physically and mentally UHD as explanatory variables can be used to predict measures like the EQ-5D, which can in turn be converted to QALYs for cost-utility studies.<sup>31,26</sup> Another limitation is that Healthy Days has not been validated in languages other than English. Syntax and diction varies between languages. Non-English speakers can potentially respond differently to a translated HRQOL-4, especially the first question assessing general health. Lastly, wording changes and question order could affect responses; thus, enforcing the standardization of HRQOL-4 is important.

### Using Healthy Days to Inform Public Policy

The Institute of Medicine's 2012 *Living Well with Chronic Illness* report concluded that it is critical to implement new public policies to promote living well with chronic illnesses.<sup>32</sup> Accordingly, the Healthy Days measures have broad applications for federal, state, and local governments to better understand the needs of their communities and to identify vulnerable subpopulations. Several groups are using Healthy Days to inform such policy efforts. For more than a decade, CMS has included the Healthy Days questions in the Medicare Health Outcomes Survey, a longitudinal patient-reported outcomes tool used to measure the quality of care provided by Medicare Advantage organizations.<sup>33</sup> The National Women's Law Center also uses this measure in its report card on women's health to advocate on behalf of women's health and corresponding policy blueprints.<sup>34</sup> Similarly, the University of Wisconsin's Mobilizing Action Toward Community Health (MATCH) project uses the Healthy Days measures in its county-level health rankings and uses the information to implement programs and policies addressing community health needs.<sup>35</sup>

Now more than ever, the health care system is focused on finding ways to improve well-being from the limited resources available. In January of 2015, US Secretary of Health and Human Services Sylvia M. Burwell announced a goal to have 50% of all Medicare payments tied to quality or value through alternative payment models by the end of 2018.<sup>36</sup> As the health care systems shifts to a value-based purchasing model, individuals' perceptions of their health are a critical element of defining value. Patient-reported assessments of their health such as Healthy Days should play an important role in determining where value is being delivered and should be reflected in reimbursement models. Accurate, reliable, and consistent measurement of quality of life using a validated measure like Healthy Days will help to identify promising payment solutions going forward.

Given that quality of life is closely associated to measurable health outcomes, policy makers may find Healthy Days a powerful tool for predicting the needs and use of health care in a community. Expanded utilization of Healthy Days measures as an indicator of the current perceived health of communities could help health planners and leg-

islators allocate resources for health services and assess impact of population-based policies or interventions.

### Opportunity for Broader Use of Healthy Days

The Healthy Days survey instrument offers several strengths for measuring population health:

- has demonstrated validity and reliability,
- incorporates the individual's current perspective of health and well-being,
- is easy to administer and respond to, allowing for large population-based surveys,
- produces interpretable and meaningful results that are easily understood by policy makers and the public,
- is correlated to health outcomes, and
- can be tracked over time and compared to national and state benchmarks.

Given these merits, several organizations with important roles in managing and improving population health have adopted Healthy Days as a measurement instrument. Such entities are actively using the data collected to advocate on behalf of the health care needs of their communities. Despite these adopters, great opportunity remains across a wide range of settings for those who manage population health to leverage the Healthy Days tool to track and improve health within their settings. In this time of exponential growth in biometric data collection and the digitization of health records, the addition of data related to the individual's perception of health and well-being fills an important gap in what can be gathered from electronic health records and billing data. Such a holistic view can more accurately isolate the most pressing health care needs of a population, measure improvements resulting from interventions that may otherwise take years to realize the outcomes benefits, and allocate the provision of health care services toward those that improve not only outcomes, but also quality of life.

### Author Disclosure Statement

The authors declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Drs. Davidson, Happe, Miao, Prewitt, and Slabaugh, Mr. Cordier and Mr. Havens are employees of Humana Inc. Drs. Happe and Slabaugh own Humana stock. No external financial support was provided to Humana for this research. Dr. Jia received compensation from Humana for his contributions. Drs. Shah and Zack report no conflict of interest.

The investigators retained full independence in the conduct of this research. The findings and conclusions in this report are those of the author and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

### References

1. Remington PL, Booske BC. Measuring the health of communities—how and why? *J Public Health Manag Pract.* 2011;17:397–400.
2. Berwick DM, Nolan TW, Whittington J. The triple aim: care, health, and cost. *Health Aff (Millwood).* 2008;27:759–769.

3. Healthy People 2020. Healthy People 2020 Foundation Health Measure Report: Health-Related Quality of Life and Well-Being. November 2010. <https://www.healthypeople.gov/sites/default/files/HRQoLWBFullReport.pdf> Accessed November 16, 2015.
4. Moriarty DG, Zack MM, Kobau R. The Centers for Disease Control and Prevention's Healthy Days Measures—population tracking of perceived physical and mental health over time. *Health Qual Life Outcomes*. 2003;1:37.
5. Zahran HS, Kobau R, Moriarty DG, Zack MM, Holt J, Donehoo R. Health-related quality of life surveillance—United States, 1993–2002. *MMWR Surveill Summ*. 2005;54(4):1–35.
6. DeSalvo KB, Fan VS, McDonnell MB, Fihn SD. Predicting mortality and healthcare utilization with a single question. *Health Serv Res*. 2005;40:1234–1246.
7. Latham K, Peek CW. Self-rated health and morbidity onset among late midlife U.S. adults. *J Gerontol B Psychol Sci Soc Sci*. 2013;68(1):107–116.
8. Schron E, Friedmann E, Thomas SA. Does health-related quality of life predict hospitalization or mortality in patients with atrial fibrillation? *J Cardiovasc Electrophysiol*. 2014;25(1):23–28.
9. Dominick KL, Ahern FM, Gold CH, Heller DA. Relationship of health-related quality of life to health care utilization and mortality among older adults. *Aging Clin Exp Res*. 2002;14:499–508.
10. Idler EL, Benyamini Y. Self-rated health and mortality: a review of twenty-seven community studies. *J Health Soc Behav*. 1997;38(1):21–37.
11. HealthyPeople.gov. Foundation Health Measures. 2015. <http://www.healthypeople.gov/2020/about/Foundation-Health-Measures>. Accessed December 14, 2015.
12. Centers for Disease Control and Prevention. Measuring Healthy Days: Population Assessment of Health-related Quality of Life. November 2000. <http://www.cdc.gov/hrqol/pdfs/mhd.pdf>. Accessed November 16, 2015.
13. Zullig KJ, Valois RF, Huebner ES, Drane JW. Evaluating the performance of the Centers for Disease Control and Prevention core health-related quality of life scale with adolescents. *Public Health Rep*. 2004;119:577–584.
14. Andresen EM, Fitch CA, McLendon PM, Meyers AR. Reliability and validity of disability questions for US Census 2000. *Am J Public Health*. 2000;90:1297–1299.
15. Newschaffer CJ. Validation of Behavioral Risk Factor Surveillance System (BRFSS) HRQOL Measures in a Statewide Sample. Atlanta, GA: Centers for Disease Control and Prevention; 1998.
16. Jia H, Lubetkin EI, Moriarty DG, Zack MM. A comparison of Healthy Days and EuroQol EQ-5D measures in two US adult samples. *App Res Qual Life*. 2007;2:209–221.
17. Andresen EM, Fouts BS, Romeis JC, Brownson CA. Performance of health related quality-of-life instruments in a spinal cord injured population. *Arch Phys Med Rehabil*. 1999;80:877–884.
18. Ounpuu S, Chambers LW, Chan D, Yusuf S. Validity of the US Behavioral Risk Factor Surveillance System's health related quality of life survey tool in a group of older Canadians. *Chronic Dis Can*. 2001;22:93–101.
19. Health-related quality of life—Puerto Rico, 1996–2000. *MMWR Morb Mortal Wkly Rep*. 2002;51(8):166–168.
20. Jia H, Muennig P, Lubetkin EI, Gold MR. Predicting geographical variations in behavioural risk factors: an analysis of physical and mental healthy days. *J Epidemiol Community Health*. 2004;58:150–155.
21. Stiefel M, Nolan K. Measuring the triple aim: a call for action. *Popul Health Manag*. 2013;16:219–220.
22. Borawski E, Bowlin S, Wu G, Jia H, Chen H. The use of the Behavioral Risk Factor Surveillance System (BRFSS) in estimating disability at the state and substate levels. Poster presented at: American Public Health Association annual meeting; Indianapolis, IN; 1997:350.
23. Self-reported frequent mental distress among adults—United States, 1993–1996. *MMWR Morb Mortal Wkly Rep*. 1998;47(16):326–331.
24. Segev Z, Arif AA, Rohrer JE. Activity limitations and healthcare access as correlates of frequent mental distress in adults 65 years and older: a Behavioral Risk Factor Surveillance study—2008. *J Prim Care Community Health*. 2012;3(1):17–22.
25. Zhou H, Siegel PZ, Barile J, et al. Models for count data with an application to Healthy Days measures: are you driving in screws with a hammer? *Prev Chronic Dis*. 2014;11:E50.
26. Jia H, Zack MM, Moriarty DG, Fryback DG. Predicting the EuroQol Group's EQ-5D index from CDC's "Healthy Days" in a US sample. *Med Decis Making*. 2011;31:174–185.
27. Sullivan PW, Ghushchyan V. Preference-based EQ-5D index scores for chronic conditions in the United States. *Med Decis Making*. 2006;26:410–420.
28. Li L, Fu AZ. Some methodological issues with the analysis of preference-based EQ-5D index scores. *Health Serv Outcomes Res Method*. 2009;9:162–176.
29. Freedman D.A. Bootstrapping regression models. *Ann Statist*. 1981;9:1218–1228.
30. Jia H, Lubetkin EI. Time trends and seasonal patterns of health-related quality of life among U.S. adults. *Public Health Rep*. 2009;124:692–701.
31. Jia H, Lubetkin EI. Estimating EuroQol EQ-5D scores from population Healthy Days data. *Med Decis Making*. 2008;28:491–499.
32. The National Academies of Sciences, Engineering, and Medicine. Report Brief. Living Well with Chronic Illness: A Call for Public Action. January 31, 2012. <http://iom.nationalacademies.org/Reports/2012/Living-Well-with-Chronic-Illness/Report-Brief.aspx>. Accessed November 16, 2015.
33. Centers for Medicare & Medicaid Services. Medicare Health Outcomes Survey. <http://www.hosonline.org/> Accessed November 16, 2015.
34. National Women's Law Center. Making the Grade on Women's Health: A National and State by State Report Card. <http://hrc.nwlc.org/>. Accessed November 16, 2015.
35. University of Wisconsin Population Health Institute, School of Medicine and Public Health. Mobilizing Action Toward Community Health (MATCH). <http://uwphi.pophealth.wisc.edu/programs/match/>. Accessed November 16, 2015.
36. US Department of Health and Human Services. Better, Smarter, Healthier: In historic announcement, HHS sets clear goals and timeline for shifting Medicare reimbursements from volume to value. January 26, 2016. <http://www.hhs.gov/news/press/2015pres/01/20150126a.html>. Accessed November 16, 2015.

Address correspondence to:  
 Dr. Laura Happe  
 Humana, Inc.  
 500 W. Main  
 Louisville, KY 40202

E-mail: lhappe@humana.com