

RESEARCH

Open Access



Molecular characterization of infectious pancreatic necrosis virus strains isolated from the three types of salmonids farmed in Chile

René A. Manríquez^{1,2}, Tamara Vera^{1,2}, Melina V. Villalba¹, Alejandra Mancilla^{1,2}, Vikram N. Vakharia³, Alejandro J. Yañez^{1,2} and Juan G. Cárcamo^{1,2*} 

Abstract

Background: The infectious pancreatic necrosis virus (IPNV) causes significant economic losses in Chilean salmon farming. For effective sanitary management, the IPNV strains present in Chile need to be fully studied, characterized, and constantly updated at the molecular level.

Methods: In this study, 36 Chilean IPNV isolates collected over 6 years (2006–2011) from *Salmo salar*, *Oncorhynchus mykiss*, and *Oncorhynchus kisutch* were genotypically characterized. Salmonid samples were obtained from freshwater, estuary, and seawater sources from central, southern, and the extreme-south of Chile (35° to 53°S).

Results: Sequence analysis of the VP2 gene classified 10 IPNV isolates as genogroup 1 and 26 as genogroup 5. Analyses indicated a preferential, but not obligate, relationship between genogroup 5 isolates and *S. salar* infection. Fifteen genogroup 5 and nine genogroup 1 isolates presented VP2 gene residues associated with high virulence (i.e. Thr, Ala, and Thr at positions 217, 221, and 247, respectively). Four genogroup 5 isolates presented an oddly long VP5 deduced amino acid sequence (29.6 kDa). Analysis of the VP2 amino acid motifs associated with clinical and subclinical infections identified the clinical fingerprint in only genogroup 5 isolates; in contrast, the genogroup 1 isolates presented sequences predominantly associated with the subclinical fingerprint. Predictive analysis of VP5 showed an absence of transmembrane domains and plasma membrane tropism signals. WebLogo analysis of the VP5 BH domains revealed high identities with the marine birnavirus Y-6 and Japanese IPNV strain E1-S. Sequence analysis for putative 25 kDa proteins, coded by the ORF between VP2 and VP4, exhibited three putative nuclear localization sequences and signals of mitochondrial tropism in two isolates.

Conclusions: This study provides important advances in updating the characterizations of IPNV strains present in Chile. The results from this study will help in identifying epidemiological links and generating specific biotechnological tools for controlling IPNV outbreaks in Chilean salmon farming.

Keywords: IPNV, Birnavirus, Genotyping, Phylogenetic characterization, VP2, VP5

* Correspondence: gcarcamo@uach.cl

¹Instituto de Bioquímica y Microbiología, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile

²Centro FONDAP, Interdisciplinary Center for Aquaculture Research (INCAR), Valdivia, Chile

Full list of author information is available at the end of the article



Background

Infectious pancreatic necrosis virus (IPNV) is the causative agent of a contagious fish disease by the same name and is responsible for economic losses in the aquaculture industry worldwide. In 2015, this viral agent represented about 30% of disease diagnoses reported from Chilean salmon farming centers and caused about 8% of pathogen-associated mortalities [1]. Infections with IPNV mainly occur in the freshwater or fry first-feeding stages, but fingerlings of most salmonids can also become infected [2]. The transfer of young salmon to saltwater is a particularly stressful stage, and high mortalities associated with IPNV outbreaks are commonly reported. Notably, infected fish that overcome IPN remain asymptomatic carriers, with some authors reporting that the virus continues replicating in hematopoietic tissues [3–5]. Carrier fish could possibly be the most important mechanism through which this virus is maintained and remains contagious.


IPNV is the prototype virus of the *Birnaviridae* family; the virion consists of two genome segments of double-stranded RNA (termed A and B) that are encased in a non-enveloped icosahedral capsid [6]. Segment B of the IPNV genome is monocistronic, contains approximately 2783 nucleotides, and encodes for the RNA-dependent RNA polymerase VP1. In turn, segment A contains approximately 3097 nucleotides, is bicistronic and has an open reading frame (ORF) that encodes a 106-kDa polyprotein (NH₂-pVP2-VP4-VP3-COOH) (Fig. 1). This polyprotein is co-translationally cleaved by viral protease 4 (termed VP4 or NS) [7] to generate the viral capsid proteins pVP2 (comprised by amino acids 1 to 512), VP3 (756 to 1012), and protease VP4 (513 to 755) [8, 9]. Subsequently, pVP2 is further processed at the C terminus to obtain the mature VP2 protein [10, 11]. In Sp IPNV serotypes, an ORF coding for a putative 25 kDa protein exists between the VP2 and VP4 coding regions that contains a nuclear localization signal (NLS) [12].

The VP2 gene encodes the principal capsid protein of IPNV, which is responsible for virulence and contains all

neutralizing epitopes. The key residues involved in virulence are Thr, Ala, Thr/Ala, and Tyr/His at positions 217, 221, 247, and 500, respectively [12–14].

Also in relation to segment A, some IPNV isolates present a smaller ORF that partially overlaps and precedes the major ORF. This smaller ORF encodes a single peptide termed VP5 (Fig. 1), which ranges in size from 15 kDa to 3.3 kDa [15–17]. VP5 might also participate in the virulence mechanism, but studies with VP5-deficient mutants gave a similar mortality that VP5-producing strains [18]. Likewise, some IPNV isolates lack the reading-frame for VP5 [19, 20], while some highly virulent IPNV strains have truncated forms of this peptide [17]. While these results seem to exclude VP5 as a factor linked to virulence, discrepancies between studies remain. Therefore, additional research is required to fully elucidate the roles of VP2 and VP5 in the virulence of IPNV infection. The VP5 sequence does contain the Bcl-2 homology (BH) domains BH1, BH2, BH3, and BH4 typical of class I and II Bcl-2 family proteins [21, 22]. Accurate understandings on the role of these sequences in the cellular infection mechanism remain lacking, as is also the case for the functional effects of structural motif variations in VP5.

Serological studies have identified two aquatic birnavirus serogroups, A and B [23]. IPNV belongs to serogroup A, which contains nine serotypes (A1 to A9). However, perhaps the most significant contribution towards the classification of IPNV was proposed by Blake et al. [24] through phylogenetic analysis of deduced amino acid sequences. This analysis revealed six genogroups that generally correlate with the geographical origin of the virus and previous serological classifications. Subsequent phylogenetic analysis conducted by Nishizawa et al. [25] revealed the existence of a new genogroup, G.7, for Japanese *aquabirnavirus* isolates. Therefore, according to the classification proposed by Blake et al. [24], genogroup 1 is composed of USA isolates (A1) and Jasper strains from Canada; genogroup 2 is composed of isolates from Europe (A3) and Asia; genogroup 3 is composed of two isolates from



Canada (C1) (A6) and the European Tellina strain (A5); genogroup 4 is composed of isolates Canada 2 (A7) and Canada 3 (A8) isolates; genogroup 5 is composed of five European isolates and one isolate from Asia (A2); genogroup 6 is composed by only the Hecht isolate (A4); and genogroup 7 is composed of isolates mainly from Japan and Korea [25, 26].

Several studies link Chilean IPNV isolates with sero- or genotypes from North America and Europe. Early reports found that Chilean IPNV isolates from rainbow trout were similar to the North American serotype VR-299, but this generated debate regarding the possible presence of serotype Sp, which is of European origin [27, 28]. According to the classification of Blake et al. [24], serotypes VR-299 and Sp belong to genogroups 1 and 5, respectively. Later, Mutoloki and Evensen [29] grouped some Chilean isolates from Atlantic salmon with Norwegian and Irish isolates, a classification consistent with the Sp serotype and genogroup 5. Likewise, researchers from Valparaiso, Chile [30, 31] phylogenetically analyzed five and 18 IPNV samples, respectively, from different locations from the south-Austral region of the country, and, as in previous studies, most of the sequenced isolates were of European origin (genogroup 5), whereas the remaining were of North American origin (genogroup 1). Furthermore, real-time RT-PCR assays based on Univeral ProbeLibrary probes successfully identifies and discriminates between Chilean isolates from both IPNV genogroups [32], and the present study more deeply assessed several of these same isolates.

Chile currently has approximately 241 freshwater-phase salmonid farms and 485 seawater salmonid farming concessions, resulting in 726 salmonid farming centers nationwide. Considering the wide prevalence of this industry, there is a persisting need to widely extend the number of analyzed IPNV isolates. Additionally, novel studies should be more extensive and include many more sampling points, a vast amount of different isolates, various salmonid sources, and diverse water types, in addition to ideally collecting samples over an extended period of time and over a wide territory. Studies should also try to combine analysis techniques to expand on the available knowledge regarding molecular underpinnings, genome characterization, VP5 characterization, VP2 virulence markers, sequence and motif conservations, and phylogeny. Such studies would be highly relevant in fully elucidating the roles of VP2 and VP5 in the virulence mechanisms associated with IPNV infection.

To advance the available knowledge on IPNV, this study carried out genotypic and phylogenetic characterizations for 36 Chilean IPNV isolates collected over a 6 year period (2006–2011). Additional analyses of VP2 and comparative genotyping of VP5, including the putative 25 kDa protein, were also performed for all isolates.

Methods

Virus isolation and cell cultures

For growth and titration of IPNV from sampled tissues, the RTG2 cell line, from rainbow trout gonads (ATCC CCL-55) and CHSE-214, from Chinook salmon (*Oncorhynchus tshawytscha*) embryo cells (European Collection of Cell Cultures, ECACC 91041114) were used. Cells were propagated in Leibovitz's L-15 culture medium (Life Technologies) supplemented with 10% fetal bovine serum. The incubation temperature used for growth was 17 °C. Confluent cell layers were sub-cultured twice a week.

The virus isolates were kindly provided by the company Biovac S.A., and the Institute of Pathology, Faculty of Veterinary Sciences, Universidad Austral de Chile. According to the information provided by them, these samples were obtained from *S. salar*, *O. mykiss*, and *O. kisutch*, in different locations along the southern Chilean, from 2006 to 2011 (Table 1). The virus isolates were inoculated on monolayers of RTG2 cell cultures. Once an extensive cell cytopathic effect occurred (3–7 days post-infection), the cell culture medium was harvested and stored in aliquots until use. Concomitantly, the viral cultures were titrated by an end-point dilution assay, using CHSE-214 cells.

Viral RNA extraction and reverse-transcription (RT)

Viral RNA was isolated with the E.Z.N.A. Total RNA Kit (Omega Bio-tek) according to the manufacturer's instructions. For cDNA synthesis, 10 µl of the RNA suspension were subjected to reverse-transcription using the M-MLV Reverse Transcriptase Kit (Life Technologies) and random primers according to the manufacturer's instructions. The cDNA was stored at –20 °C until use.

RT-qPCR used primers PrD1 5'-AAAGCCATAG CCGCCCATGAAC-3' and PrD2 5'-TCTCATCAG CTGGCCAGGTAC-3' to detect IPNV-positive samples [33] using a StepOnePlus™ System (Thermo Fisher, USA). Reactions (10 µl) included 1 µl of template cDNA, 0.5 µM primers and components provided by Fast SYBR Green Master Mix (Thermo Fisher, USA). Cycle conditions were for 20 s at 95 °C, followed by 40 cycles including denaturation for 3 s at 95 °C and annealing for 10 s at 60 °C. Melting curve analysis of amplification products was performed at the end of PCR to confirm that only one PCR product was amplified and detected. For partial sequencing of the VP2 gene in the different isolates, PCR was applied using KOD Hot start DNA Polimerasa (CALBIOCHEM/USA) and the primers PrA1 5'-TGA GATCCATTATGCTT CCCGA-3' and PrA2 5'-GACAG GATCATCTTGGCATAGT-3'. Cycle conditions were for 5 min at 95 °C, followed by 40 cycles including denaturation for 30 s at 95 °C, annealing for 30 s at 60 °C and final extension for 30 s at 72 °C as described by Blake et al. [33].

Table 1 The Chilean infectious pancreatic necrosis virus isolates were obtained from different geographical regions and salmonid species. Additionally, the tissue, stage, weight, water source, and year of the isolate are reported

Isolate code	Tissue used for isolation	Geographical area ^a	Species	Stage	Water source	Year	Accession number
1096-BC1	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>O. kisutch</i>	Juvenile	NA	2009	KX523824
1096-BC2	Mix (kidney, liver, spleen)	Comau Fjord (X Region)	<i>S. salar</i>	Juvenile	Freshwater	2010	KX523802
1096-BC3	Mix (whole fish)	Maule R. (VII Region)	<i>S. salar</i>	Alevin	Freshwater	2009	KX523827
1096-BC4	Mix (kidney, liver, spleen)	Comau Fjord (X Region)	<i>S. salar</i>	Alevin	Freshwater	2010	KX523803
1096-BC5	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>O. kisutch</i>	Juvenile	NA	2009	KX523816
1096-BC6	Mix (kidney, liver, spleen)	Los Angeles (VIII Region)	<i>O. mykiss</i>	Alevin	Freshwater	2009	KX523804
1096-BC7	Mix (kidney, liver, spleen)	Comau Fjord (X Region)	<i>O. mykiss</i>	Juvenile	Freshwater	2010	KX523818
1096-BC8	Mix (kidney, liver, spleen)	Rupanco Lake (X region)	<i>O. mykiss</i>	Juvenile	Freshwater	2010	KX523828
1096-BC9	Mix (kidney, liver, spleen)	Reloncaví Cove (X Region)	<i>S. salar</i>	Adult	Freshwater	2010	KX523829
1096-BC10	Mix (kidney, liver, spleen)	Rupanco Lake (X region)	<i>O. mykiss</i>	Juvenile	Freshwater	2010	KX523801
1096-BC11	Mix (kidney, liver, spleen)	Cisnes (XI Region)	<i>S. salar</i>	NA	Seawater	2010	KX523819
1096-BC12	Mix (kidney, liver, spleen)	Calbuco (X Region)	<i>S. salar</i>	Alevine	Freshwater	2010	KX523800
1096-BC13	Mix (kidney, liver, spleen)	Reloncaví Estuary (X Region)	<i>S. salar</i>	Adult	Estuary	2010	KX523833
1096-BC14	Mix (kidney, liver, spleen)	Comau Fjord (X Region)	<i>S. salar</i>	Alevine	Freshwater	2010	KX523821
1096-BC15	Mix (kidney, liver, spleen)	Reloncaví Estuary (X Region)	<i>S. salar</i>	Adult	Estuary	2010	KX523830
1096-BC16	Mix (kidney, liver, spleen)	Cisnes (XI Region)	<i>S. salar</i>	Smolt	Seawater	2010	KX523805
1096-BC17	Mix (kidney, liver, spleen)	Nehuentue (VIII Region)	<i>S. salar</i>	Alevin	Freshwater	2011	KX523822
1096-BC18	Mix (kidney, liver, spleen)	Calbuco (X Region)	<i>S. salar</i>	Alevin	Freshwater	2010	KX523820
1096-V1	Mix (whole fish)	Araucanía R. (IX Region)	<i>O. mykiss</i>	Alevin	Freshwater	2006	KX523809
1096-V2	Mix (kidney/spleen)	Magallanes R. (XII Region)	<i>S. salar</i>	Pre-smolt	Freshwater	2009	KX523810
1096-V3	Mix (kidney/spleen)	Magallanes R. (XII Region)	<i>S. salar</i>	Pre-smolt	Freshwater	2009	KX523811
1096-V4	Mix (kidney/spleen)	Los Ríos R. (XIV Region)	<i>O. kisutch</i>	Pre-smolt	Estuary	2008	KX523799
1096-V5	Mix (kidney/spleen)	Los Ríos R. (XIV Region)	<i>S. salar</i>	Smolt	Estuary	2008	KX523807
1096-V6	Mix (kidney/spleen)	Los Lagos R. (X Region)	<i>O. mykiss</i>	Smolt	Freshwater	2006	KX523808
1096-V7	Mix (whole fish)	Los Lagos R. (X Region)	<i>O. mykiss</i>	Alevin	Freshwater	2006	KX523825
1096-V8	Mix (kidney/spleen)	Los Ríos R. (XIV Region)	<i>O. mykiss</i>	Pre-smolt	Estuary	2007	KX523806
1096-V9	Mix (kidney/spleen)	Los Lagos R. (X Region)	<i>S. salar</i>	Pre-smolt	Estuary	2006	KX523826
1096-V10	Mix (kidney/spleen)	Los Lagos R. (X Region)	<i>S. salar</i>	Alevin	Freshwater	2006	KX523823
1096-BC20	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>S. salar</i>	NA	NA	2009	KX523817
1096-BC21	Mix (kidney, liver, spleen)	Araucanía R. (IX Region)	<i>O. kisutch</i>	Alevin	Freshwater	2009	KX523813
1096-BC22	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>S. salar</i>	Adult	Seawater	2009	KX523812
1096-BC23	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>O. kisutch</i>	Juvenile	Freshwater	2009	KX523815
1096-BC24	Mix (kidney, liver, spleen)	Aysén R. (XI Region)	<i>O. kisutch</i>	Alevin	Freshwater	2009	KX523814
1096-BC25	Mix (kidney, liver, caeca)	Los Ríos R. (XIV Region)	<i>S. salar</i>	Alevin	Freshwater	2010	KX523832
1096-BC26	Mix (kidney, liver, caeca)	Araucanía R. (IX Region)	<i>O. kisutch</i>	Alevin	Freshwater	2009	KX523831
1096-BC27	NA	NA	NA	NA	NA	2010	KX523834

NA: Not available. ^aChile is geopolitically divided from north to south into 15 distinct regions equally recognized by Roman numerals or written names. The regions included in this study are as follows: VII = Maule Region; VIII = Biobío Region; IX = Araucanía Region; X = Los Lagos Region; XI = Aysén Region; XII = Magallanes Region; and XIV = Los Ríos Region

Nucleotide sequencing and analysis of VP2

PCR products of 1180 base pairs were recovered from agarose gel, purified using the E.Z.N.A. Extraction Kit (Omega Biotek), and cloned into the pCR2.1 vector using a TOPO TA Cloning Kit for Sequencing (Invitrogen by Life Technologies). The pCR2.1 plasmid, with the 36 different

inserts, were sent to Macrogen Inc. (Korea) for sequencing using the M13 universal primers (M13F: 5'-GTAAAAC GACGGCCAGT-3' and M13R: 5'-AACAGCTATGAC CATG-3'). The obtained nucleotide sequences were assembled using DNA Baser v4.12, and the deduced amino acid sequences were obtained using the online ExPASy tool

(<http://web.expasy.org/translate/>). The obtained VP2 sequences and Genbank strains representative of all seven IPNV genogroups and of the previously reported Chilean IPNV isolates [24, 29, 30] were included in comparative and phylogenetic analyses.

The deduced amino acid sequences were aligned using ClustalW in the Bioedit software and analyzed in relation to previously described amino acid virulence markers for IPNV [12–14]. Pairwise distances were calculated for the percentage of nucleotide and amino acid identities of total sequences. Phylogenetic and molecular evolutionary analyses were conducted using the MEGA v5 software [34]. Prior to the construction of the phylogenetic tree, the best model for DNA/Protein to VP2 sequences was computed. Then, phylogenetic analysis was performed using the deduced amino acid sequences, with evolutionary history inferred by using the maximum likelihood method based on the JTT matrix-based model [35]. One thousand bootstrap replicates were performed, and a condensed phylogenetic tree was constructed for the genogroups comparison with previously described reference strains (Table 2).

Nucleotide sequencing and analysis of VP5

The previously synthesized viral cDNA was additionally used to amplify the region encoding VP5. The primers used in PCR analysis (VP5F: 5'- TCCGTCGATGGC GAAAGCCC-3' and VP5R: 5'- CTCCACCTCAGACA GACTGCC -3') generated a product of 460 base pairs. This product was recovered from agarose gel, purified using the E.Z.N.A. Extraction Kit (Omega Biotek), and cloned into the pCR2.1 vector using a TOPO TA Cloning Kit for Sequencing (Invitrogen Life Technologies). The pCR2.1 vectors with the different inserts were sent to Macrogen Inc. (Korea) for sequencing, using a similar analysis to the one performed for VP2. To obtain the complete deduced amino acid sequences of VP5 isoforms longer than 269 amino acid residues (around 29.6 kDa), the nucleotide sequences obtained for VP5 were

overlapped to the VP2 nucleotide sequences previously obtained. To assess if the VP5 sequences of the collected Chilean isolates contained transmembrane domains, public transmembrane domain predictors (TMpred and TOPCONS) were run with VP5 amino acid sequences. Furthermore, to determine the presence of a cationic cluster for plasmatic membrane tropism, the VP5 sequences were aligned and analyzed for the presence of polycationic regions using the ClustalW tool and the VP5 sequence of the infectious bursal disease virus [36].

WebLogo analyses

To study the variability of the deduced amino acid sequences of four BH domains in VP5, the VP5 sequences of Chilean IPNV isolates were aligned using the MUSCLE software [37]. The obtained results were run in the online WebLogo tool (<http://weblogo.berkeley.edu/>), where each residue of the BH domains is represented as a stack of one letter for each amino acid and where size indicates sequence conservation at each position. The sequences for each VP5 BH domain of Chilean IPNV isolates were compared with conserved BH1, BH2, BH3, and BH4 sequences described for these domains, which included the BH domains of IPNV E1-S (Ab strain), MABV Y-6, and other Bcl-2-related proteins sharing homology in one or more of these domains, in accordance with previously described methodologies [38, 39].

Analysis of the ORF for putative 25 kDa protein

Nine of the 36 Chilean IPNV isolates (1096-BC1, BC2, BC4, BC6, BC10, BC12, BC16, and 1096-V4 and -V8) were subjected to full sequencing of segment A. This provided information on the possible presence of the ORF for the putative 25 kDa protein between the VP2 and VP4 coding regions. The nucleotide sequence of the putative 25 kDa protein described for isolate Sp103 (Accession number AY354519.1) was used in multiple alignment analyses with the online ClustalW tool. Then, the deduced amino acid sequence of each ORF was

Table 2 Reference aquabirnavirus strains used

Strain	Geographical origin	Host	Genogroup	Serotype	GenBank accession N°
WB	Maine, USA	Trout	1	A1	AF342727
Sp	Denmark	Trout	5	A2	AF342728
Ab	Denmark	Trout	2	A3	AF342729
He	Germany	Pike	6	A4	AF342730
Te	England	Tellina	3	A5	AF342731
C1	Canada	Trout	3	A6	AF342732
C2	Canada	Trout	4	A7	AF342733
C3	Canada	Arctic char	4	A8	AF342734
Jasper	Canada	Trout	1	A9	AF342735
YTAV	Japan	Yellowtail	7		AY283781

identified using the online ExPASy tool [40]. To determine possible NLS, the online CNLS Mapper (http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi) and NLStradamus [41] tools were used. Moreover, to identify possible mitochondrial location signal sequences, the online predictor TargetP 1.1 was used (<http://www.cbs.dtu.dk/services/TargetP/>).


Results


A virus collection with 36 Chilean IPNV isolates was constituted. These isolates were sampled from the three most important farmed salmonids in Chile: Atlantic salmon (*S. salar*), rainbow trout (*O. mykiss*), and coho salmon (*O. kisutch*). These fish were sampled from waters with different salinities (freshwater, estuaries, and seawater) and over a wide expanse of territory extending along central to southern Chile (35° to 53° S; Fig. 2).

Using molecular phylogenetic analyses, 10 of the 36 Chilean IPNV isolates were classified within genogroup 1 (around 28% of all isolates; Fig. 3), a group that also includes the North American isolates Jasper and West

Buxton (Table 2). The remaining 26 IPNV isolates were classified within genogroup 5 (72%), close to the European isolate Sp. The 10 Chilean isolates belonging to genogroup 1 had nucleotide identity percentages ranging from 76.9 to 99.4% (Table 3), whereas, those within genogroup 5 had nucleotide identities ranging from 87.7 to 100% (Table 4), with isolates BC-11 and BC-13 being identical. Nucleotide sequence comparisons between the VP2 of Chilean isolates and their respective reference strains found that the highest nucleotide variability between the 10 isolates of genogroup 1 and the West Buxton strain is 19.7% (Table 3). This variability was around 11.0% for the 26 isolates of genogroup 5 as compared to the Sp strain (Table 4). The percentages of identity for deduced amino acid sequences between the isolates of genogroup 1 were higher than nucleotide identities (90.61 to 99.45%; Table 3), and this situation was even more marked for isolates in genogroup 5, with several isolates showing 100% amino acid identity (Table 4).

Despite only having precise mortality (%) information for two isolates, the VP2 amino acid residues with





described virulence involvement in the Sp serotype, were compared between the deduced amino acid sequences of genogroup 5 isolates. This analysis revealed that 15 presented the Thr, Ala, and Thr residues at positions 217, 221, and 247, respectively (Table 5). Sequencing beyond amino acid position 500 was only accomplished in eight of these isolates, with seven presenting Tyr in this position and one presenting His. Similarly, comparative analysis of the 10 genogroup 1 isolates revealed nine to have

Ala, Thr, and Ala at positions 217, 221, and 247, respectively. Only one isolate was sequenced beyond position 500, showing Arg in this position (Table 5).

Comparative analyses of the VP5 deduced amino acid sequences showed that six had a truncated VP5 with 28 amino acid residues (around 3.1 kDa), five of which belonged to genogroup 1 (1096-BC1, BC3, BC8, BC26, and BC27) and one to genogroup 5 (BC24; Figs. 4 and 5). Moreover, 26 isolates showed a VP5 with 133 amino acid

Table 3 Pairwise comparison of percent identity between the nucleotide and deduced amino acid sequences of VP2 from IPNV isolates belonging to genogroup 1. The West Buxton strain (GenBank Accession Number AF342727) was included as a reference. The highest and lowest identity percentages are highlighted in boxes

		Percent identity nucleotide sequence of VP2										
		Wb	1096-V1	1096-V7	1096-V9	1096-BC1	1096-BC3	1096-BC8	1096-BC9	1096-BC15	1096BC26	1096-BC27
Percent identity deduced amino acid sequence of VP2	Wb		80.25	94.3	94.38	98.36	97.63	93.90	98.14	98.64	98.20	98.03
	1096-V1	97.79		83.9	83.98	79.66	78.9	76.93	79.41	79.83	79.56	79.79
	1096-V7	95.86	96.13		99.29	94.33	94.75	90.42	94.75	95.25	93.87	93.72
	1096-V9	95.86	96.13	98.90		94.41	94.75	90.51	94.83	95.34	93.87	93.72
	1096-BC1	97.51	98.07	97.79	97.79		98.73	95.00	99.24	99.75	99.28	99.12
	1096-BC3	95.86	96.41	97.24	97.24	98.34		94.49	98.47	98.98	98.68	98.59
	1096-BC8	90.61	91.44	90.61	90.61	92.82	91.16		94.66	95.17	96.12	96.4
	1096-BC9	96.13	96.69	96.41	96.41	98.62	96.96	91.71		99.49	98.85	98.77
	1096-BC15	98.34	97.79	97.51	97.51	99.17	97.51	92.27	97.79		99.38	99.3
	1096BC26	97.51	98.34	97.24	97.24	99.45	97.79	92.54	98.07	99.17		98.8
	1096-BC27	96.69	97.79	96.96	96.96	99.17	97.51	93.65	97.79	98.34	98.90	

residues (around 14.6 kDa). Surprisingly, four isolates (1096-BC20, BC21, BC22, and BC23) had a VP5 isoform longer than 269 amino acid residues (around 29.6 kDa) (Fig. 5).

WebLogo analysis of the BH domains of VP5 showed that the BH4 domain had a highly conserved sequence, with slight variations present only in position 4 of this

sequence (Fig. 6). Specifically, Gln was in this position in 86% of cases (Fig. 6a; larger letters in position 4), whereas Lys was in the same position in 14% of cases (Fig. 6a; the small letter k that appears below the Q in position 4). When comparing this sequence with BH4 domains from viruses and other Bcl-2-related proteins (Table 6), a very high identity to IPNV E1-S and even

Table 4 Pairwise comparison of percent identity between the nucleotide and deduced amino acid sequences of VP2 from IPNV isolates belonging to genogroup 5. The Sp strain was included as a reference (GenBank Accession Number AF342728). The highest and lowest identity percents are highlighted in boxes, as are isolates with full identity (100%)

		Percent identity nucleotide sequence of VP2																											
		1096-BC5	1096-V5	Sp	1096-BC24	1096-V6	1096-BC7	1096-BC17	1096-BC25	1096-BC18	1096-BC22	1096-BC20	1096-BC21	1096-BC23	1096-V10	1096-BC10	1096-V8	1096-BC4	1096-V3	1096-V2	1096-BC14	1096-BC2	1096-BC11	1096-BC16	1096-V4	1096-BC6	1096-BC12	1096-BC13	
Percent identity deduced amino acid sequence of VP2	1096-BC5		91.78	93.64	91.86	94.17	95.08	95.25	95.43	94.83	95.68	95.68	95.42	95.76	95.68	94.83	92.8	95.51	95.59	95.51	96.02	95.93	95.95	95.85	90.08	90.76	92.63	95.97	
	1096-v5	87.85		96.23	91.33	97.85	94.49	93.89	95.72	94.72	95.08	95.54	95.08	95.17	95.54	91.04	90.57	90.71	96.09	95.89	95.17	96.23	96.4	96.3	87.68	89.23	88.42	96.33	
	Sp	90.33	94.20		93.06	98.19	96.27	95.79	96.75	96.53	97.03	97.36	96.95	97.12	97.36	91.67	91.91	92.15	97.58	97.37	97.2	98.14	97.6	98.07	89.6	91.91	89.02	97.61	
	1096-BC24	90.06	93.37	96.13		93.38	94.66	93.61	94.53	95.35	95.68	94.87	95.42	95.76	95.02	93.72	91.57	93.99	94.96	94.72	95.34	95.13	94.53	95.05	94.25	89.68	91.41	95.06	
	1096-V6	91.16	95.58	98.07	97.24		96.64	96.22	97.25	96.9	97.44	97.76	97.7	97.53	97.76	97.08	95.1	97.51	97.59	97.33	97.53	97.76	98.02	97.85	91.75	92.78	94.5	98.02	
	1096-BC7	91.16	93.92	96.69	96.96	97.51		99.66	98.95	98.31	98.47	98.31	98.22	98.56	98.47	97.12	95.34	97.88	98.22	98.14	98.47	98.56	98.5	98.31	92.37	93.22	94.66	98.42	
	1096-BC17	91.44	94.20	96.69	97.24	97.79	99.72		98.37	97.64	98.64	97.87	98.39	98.73	98.03	96.53	95.03	97.31	97.69	97.52	98.64	98.02	98.04	97.77	91.72	93.06	94.22	97.95	
	1096-BC25	91.71	94.48	96.69	97.24	97.79	98.90	99.17		98.97	98.59	98.8	98.68	98.68	98.97	97.86	95.98	98.29	98.63	98.46	98.59	98.97	98.89	98.72	92.73	93.84	95.38	98.89	
	1096-BC18	91.71	94.48	97.24	97.79	98.34	99.17	99.45	99.45		98.31	98.35	98.05	98.39	98.5	97.03	95.11	97.63	98.27	98.1	98.31	98.6	98.29	98.35	93.14	93.14	94.63	98.47	
	1096-BC22	91.16	94.48	97.24	97.79	98.34	98.07	98.34	98.34	98.90		99.49	99.41	99.75	99.49	97.63	95.68	98.39	98.9	98.73	99.15	99.07	99.12	98.98	92.97	93.81	95.25	99.12	
	1096-BC20	91.71	95.03	97.79	98.34	98.90	98.62	98.90	98.90	99.45	99.45		99.24	99.58	99.68	98.02	96.21	98.82	99.26	99.01	99.15	99.42	99.51	99.34	92.59	94.32	95.71	99.49	
	1096-BC21	91.71	95.03	97.79	98.34	98.90	98.62	98.90	98.90	99.45	99.45	100.0		99.49	99.24	97.37	95.42	98.14	98.64	98.47	98.9	98.81	99.21	98.73	92.71	93.56	95	99.21	
	1096-BC23	91.71	95.03	97.79	98.34	98.90	98.62	98.90	98.90	99.45	99.45	100.0	100.0		99.58	97.71	95.76	98.47	98.98	98.81	99.24	99.15	99.21	99.07	93.05	93.9	95.34	99.21	
	1096-V10	91.44	94.75	97.51	98.07	98.62	98.34	98.62	98.62	99.17	99.17	99.72	99.72		99.72		98.02	96.21	98.82	99.26	99.01	99.15	99.42	99.51	99.34	92.73	94.47	95.71	99.49
	1096-BC10	91.16	93.37	95.86	95.86	96.69	96.69	96.96	96.96	96.96	97.51	97.51	97.51	97.51		97.24		95.63	93.59	93.87	93.67	97.97	92.98	98.97	92.98	96.1	94.34	92.74	98.81
	1096-V8	90.88	93.37	96.13	96.13	96.96	96.96	97.24	96.96	97.24	97.24	97.79	97.79	97.79	97.79	97.51	97.51		93.74	93.2	93.0	96.1	93.08	96.98	93.12	94.41	97.51	92.37	96.76
	1096-BC4	91.44	94.20	96.41	96.41	97.24	97.24	97.51	97.24	97.51	97.51	98.07	98.07	98.07	98.07	97.79	98.34	97.51		93.4	93.2	98.64	93.52	99.43	93.52	91.31	93.57	90.06	99.23
	1096-V3	91.16	94.75	96.96	96.96	97.79	97.79	98.07	97.79	98.07	98.07	98.62	98.62	98.62	98.34	97.24	97.51	97.79			99.39	99.07	99.6	99.57	99.53	90.1	91.38	91.04	99.57
	1096-V2	91.44	93.92	96.69	96.69	97.51	98.07	98.34	98.07	98.34	98.07	98.34	98.34	98.34	98.07	96.96	97.24	97.51	98.07			98.98	99.46	99.32	99.33	89.9	91.18	90.84	99.32
	1096-BC14	91.99	94.48	97.51	97.51	98.34	98.34	98.62	98.62	99.17	98.62	99.17	99.17	99.17	98.90	97.51	97.79	97.79	98.34	98.62			99.58	99.3	99.15	93.05	93.81	95.59	99.3
1096-BC2	92.27	94.75	97.51	97.51	98.34	98.90	99.17	98.90	99.17	98.62	99.17	99.17	99.17	98.90	97.79	98.07	98.34	98.90	99.17	99.45			99.74	99.39	90.95	93.08	90.17	99.74	
1096-BC11	91.99	95.03	97.79	97.79	98.62	98.62	98.90	98.62	98.90	98.90	99.45	99.45	99.45	99.17	98.07	98.34	98.62	99.17	98.90	99.17	99.72			99.83	93.31	94.69	96.49	100.0	
1096-BC16	91.71	94.75	97.51	97.51	98.34	98.34	98.62	98.34	98.62	98.62	99.17	99.17	99.17	98.90	97.79	98.07	98.34	98.90	98.62	98.90	99.45	99.72			90.95	93.08	90.17	99.83	
1096-V4	88.67	90.88	93.09	94.75	93.92	93.92	94.20	93.92	94.20	94.20	94.75	94.75	94.75	94.75	94.48	95.86	95.30	96.13	94.48	94.20	94.75	95.03	95.30	95.03	95.03	94.18	92.54	93.52	
1096-BC6	89.78	92.27	94.48	95.03	95.58	95.86	96.13	95.86	96.13	96.13	96.69	96.69	96.69	96.69	96.41	95.86	97.24	95.86	95.86	95.58	96.13	96.41	96.69	96.41	95.58		92.74	94.37	
1096-BC12	90.06	91.71	94.20	94.20	95.03	95.03	95.30	95.03	95.30	95.30	95.86	95.86	95.86	95.86	95.58	97.24	96.41	96.69	95.58	95.30	95.86	96.13	96.41	96.13	96.96	97.79		96.33	
1096-BC13	91.99	95.03	97.79	97.79	98.62	98.62	98.9	98.62	98.9	98.9	99.45	99.45	99.45	99.17	98.07	98.34	98.62	99.17	98.9	99.17	99.72	100.0			99.72	95.3	96.69	96.41	

Table 5 Amino acid variations in the VP2 region of the of Chilean IPNV isolates, the positions showed were proposed as linked to virulence [13, 14]. ND: not determined

Amino acid position					Amino acid position				
Isolate	217	221	247	500	Isolate	217	221	247	500
Genogroup 5					Genogroup 5				
1096-BC2	T	A	T	Y	1096-V2	T	A	T	ND
1096-BC4	T	A	T	Y	1096-V3	T	A	T	ND
1096-BC5	S	S	T	ND	1096-V4	T	A	T	Y
1096-BC6	T	A	T	Y	1096-V5	P	S	A	ND
1096-BC7	T	A	T	ND	1096-V6	T	A	T	ND
1096-BC10	T	A	T	Y	1096-V8	T	A	T	Y
1096-BC11	T	A	T	ND	1096-V10	T	T	T	ND
1096-BC12	T	A	T	Y					
1096-BC13	T	A	T	ND	Genogroup 1				
1096-BC14	T	T	T	ND	1096-V1	T	A	T	ND
1096-BC16	T	A	T	H	1096-V7	A	T	A	ND
1096-BC17	T	A	T	ND	1096-V9	A	T	A	ND
1096-BC18	T	T	T	ND	1096-BC1	A	T	A	R
1096-BC20	T	T	T	ND	1096-BC3	A	T	A	ND
1096-BC21	T	T	T	ND	1096-BC8	A	T	A	ND
1096-BC22	T	T	T	ND	1096-BC9	A	T	A	ND
1096-BC23	T	T	T	ND	1096-BC15	A	T	A	ND
1096-BC24	T	T	T	ND	1096-BC26	A	T	A	ND
1096-BC25	T	S	T	ND	1096-BC27	A	T	A	ND


higher identity to MABV Y-6 were observed (Fig. 6b). Isolate analysis for BH3 was similar to that described above for BH4, with a high identity value to the BH3 domain of MABV Y-6. In slight contrast, the sequence Logo for the BH1 domain in Chilean IPNV isolates (Fig. 6c) was more similar to IPNV E1-S than MABV Y-6. Finally, the situation for the BH2 domain was very different (Fig. 6d). The principal sequence Logo obtained for the Chilean VP5 sequences did not present similarity with any of the BH2 domains used as canonical BH domain sequences (Table 6).

For the nine isolates in which segment A was fully sequenced, analysis showed that only two (1096-BC2 and BC16) evidenced the existence of an ORF at position 1326 to 1982 from the VP2 start, coding for a complete additional 25 kDa protein, as previously described by Shivappa et al. [12]. The remaining isolates presented a truncated sequence at position 38 of the putative 25 kDa protein (data not shown). According to alignment analyses, and using the online NLS Mapper and NLStradamus tools, four possible NLS sequences were recognized in the putative 25 kDa protein of the Chilean isolates (Fig. 7), which were also present in the nonstructural proteins of the *Drosophila X* virus and IPNV strain Sp103 [12].

Discussion


In 2015, Chile produced 834,000 t of salmonids. This production was the result of 726 freshwater/seawater culturing centers distributed along a distance covering roughly 2000 Km. Due to the economic importance of this industry, and considering the underreported presence of IPNV, a national characterization of the different IPNV strains present in Chile is of considerable importance. In this study, 36 Chilean IPNV isolates were characterized following sample collection over a 5-year period. Samples were collected from Atlantic salmon, rainbow trout, and coho salmon, the three most important salmonid exports for Chile.

In accordance with previous reports, only two types of Chilean-isolated IPNV strains were found. Specifically, the obtained isolates were classified as either genogroup 1, which is highly similar to the American reference strain West Buxton, or genogroup 5, which is highly similar to the European reference strain Sp. Most of the genogroup 1 isolates originated from the Los Lagos Region in southern Chile, but some were also of isolated from sites in central and the extreme-south of Chile. Genogroup 5 isolates were present in all of the analyzed regions (Table 1).



Amino acid sequence analysis showed two independent groups with 100% identities within genogroup 5: BC20 (*S. salar*), BC21 (*O. kisutch*), and BC23 (*O. kisutch*); and BC11 (*S. salar*) and BC13 (*S. salar*). Only BC11 and BC13 maintained 100% nucleotide identity, whereas BC20,

BC21, and BC23 showed identities ranging from 99.24 to 99.58%. Interestingly, the 1096-BC11 and BC13 isolates were obtained from different geographical zones: the Aysén (72.7° S; 45.4° W) and Los Lagos (72.3° S; 41.5° W) Regions, respectively. Nevertheless, since data on egg







Fig. 6 Sequence Logo diagram showing amino acid sequences of Bcl-2 Homology (BH) domains in VP5 of Chilean IPNV isolates. In A, B, C and D, consensus sequences of the BH4, BH3, BH1 and BH2 domains of VP5 in Chilean IPNV isolates, respectively. Numbers shown below the sequence represent the amino acid position in the domains. The alignment was performed using MUSCLE software. The Logo was built using the WebLogo software (<http://weblogo.berkeley.edu/logo.cgi>)

Table 6 Highly conserved Bcl-2 homology (BH) domain 1–4. The table shows amino acids from the conserved BH4, BH3, BH1, and BH2 domains of VP5 for IPNV E1-S (Ab strain) and marine birnavirus Y-6 (MABV Y-6), and Bcl-2 family members

Type	BH 4	BH3	BH1	BH2
VP5 IPNV E1-S	QLISMKDEHKQSNRNLEIHYAS	LRSLGLRKLW	LQMECEPDGTRV R PVAG DVT	HAAGWALCPEDWHQCC
MABV Y-6	EHNKGNRNLEIHYAS	LRGLRLRKLW	LQVESEPDGTGIRPVAGNIT	NPTRW
Bcl-2	REIVMKYIHYKLSQRYEWDAGDV	LRQA GDDFS	VVEELFRDGVNWGRIVAF FEF	HLHTWIQDNGGWDAFV E
Bclx-L	RELVDVFLSYKLSQKGYSW SQFDV	LREA GDEFE	VVNELEFRDGVNWGRIVAF FSF	HLEPWIQENGGWDTFV D
Bax	SEQIMKTGALLLQG	LKRI GDELD	VAADMFS DGNNWGRVVAL FYF	RLLGWIQDQGGWDGLL S
Mcl-1	GLNLYCGGAGL GAGSGG	LRRV GDGVQ	VMIHVFS DGVTWGRIVL ISF	TKRDWLVKQRGWDGFV N
A1		VQKE VEKNL	VMEKEFEDGINWGRIVTI FAF	NTGEWIRQNGGWEFII K
Bak	MASGQGGPPRQC	RAII GDDIN	IATSLFESGINWGRVVAL LGF	HCARWIAQRGGVVAAL E
Bad	AHALGLRKS DP GIRSLGSDAG GRRW	GAGA METRS	SAPPNLWA AQR YGRELRR MS	SAGWTRI IQSWDRN



sources were unavailable, a single egg origin for both locations cannot be ruled out.

Correlation analysis between the geographic location and phylogenetic characteristics of the outbreaks can provide important information to understand viral patterns of spreading or mobility throughout salmonid farms [42]. For the isolates 1096-BC20, 1096-BC21, 1096-BC22, 1096-BC23, and 1096-V10, which constituted a clade of genogroup 5, at least three (1096-BC20, 1096-BC22, and 1096-BC23) had a common geographical origin (Aysén Region; Fig. 2c). These three isolates were obtained in 2009, as was isolate 1096-BC21, which originated from roughly 500 miles (800 Km) away in the Araucania Region (Fig. 2a). The 1096-BC21 isolate came from an *O. kisutch* hatchery, but as there is no public record for salmonid transport between farms, it is unknown if fry from this hatchery were later transported to the Aysén Region for the next phase of growth. The 1096-V10 isolate was detected in the Los Lagos Region, a midway point between the Araucania and Aysén Regions (Fig. 2b). Nevertheless, 1096-V10 was obtained from a *S. salar* fry at a hatchery in 2006. Curiously, 1096-BC20, 1096-BC22, and 1096-BC23 were isolated from juvenile/adult *O. kisutch* and *S. salar*. The other clade within genogroup 5 was composed by 1096-BC2, 1096-BC5, and 1096-BC14. Of these, 1096-BC2 and 1096-BC14 were isolated in 2010 from the Comau Fjord (Los Lagos Region), while 1096-BC5 was isolated in 2009 from the Aysén Region. Once again, uncertainty remains regarding the transport of *S. salar* fry and juveniles between the Aysén Region and Comau Fjord. Regarding the genogroup 5 clade formed by 1096-BC11, 1096-BC13, and 1096-BC16, all isolated in 2010, two came from Cisnes (Aysén Region), while 1096-BC13 came from the Reloncaví Estuary. While there is no certainty that the fish came from the same hatchery,

1096-BC11 and 1096-BC13 did have identical nucleotide sequences. Other clades within genogroup 5 were even more difficult to correlate. Aquaculture activities in Chile have existed for more than three decades, and the massive transport of salmonids has always occurred between regions. This fact could explain, for example, why the 1096-V6 virus, isolated in 2006, was close to the 1096-V5 virus isolated in 2008, since both were from the Los Lagos Region. Similarly, it might clarify how the 1096-V4 virus, isolated in 2008 in the Los Ríos Region, could be closely related to the 1096-BC24 virus isolated in 2009 in the Aysén Region.

On the other hand, the genogroup 1 isolates were represented in most regions, except the Los Ríos and Magallanes Regions, although this could be due to the overall low number of isolates obtained from these areas. Both clades observed in this genogroup showed a clear relatedness to the isolation year rather than to region: Isolates 1096-V1, 1096-V7, and 1096-V9, which constituted one clade, were isolated in 2006, while the clade composed by isolates 1096-BC1, 1096-BC3, 1096-BC8, 1096-BC9, 1096-BC15, and 1096-BC26 was sampled from 2009 to 2010 in the Maule and Aysén Regions, which are separated by more than 700 miles (1130 Km).

In contrast to Tapia et al [30], the present study found genogroup 1 isolates (American origin) not only in *O. mykiss*, but also in *O. kisutch* and *S. salar*. Similarly, genogroup 5 isolates (European origin) were present not only in *S. salar*, but also in the other two salmonid species. These findings suggest that in Chile there is no fixed relationship between salmonid species and genogroups. Nevertheless, most isolates from *S. salar* (over 60%) belonged to genogroup 5, while *O. mykiss* and *O. kisutch* were equally represented (around 20% each) in that genogroup.

Unlike isolates from genogroup 5, the 10 isolates from genogroup 1 were obtained only in samples from

freshwater and estuary fish. However, considering the low number of cases for these isolates, it would be unwise to assume that Chilean genogroup 1 IPNV isolates do not infect seawater salmonids.

In relation to VP2 amino acid residues related to virulence described for Sp strains, the motifs Thr217, Ala221, and, to a lesser extent, Thr247 and His/Tyr500 are associated with high virulence, whereas the motifs Pro217, Ala221, and Ala247 are related to low/moderate virulence [12–14]. In the current study, more than half of the isolates in genogroup 5 presented high virulence motifs (i.e. Thr, Ala, and Thr at positions 217, 221, and 247, respectively), and seven showed Tyr at position 500. These isolates contrasted with eight genogroup 5 isolates that presented Thr at position 221, which is associated with almost avirulent strains and the establishment of a carrier state or persistent IPNV infection in salmonids [13, 14, 18]. Unfortunately, data on mortality percentages in field cases are frequently unavailable, either because the information provided by the farmer is imprecise or because there is a policy to not provide such information. Although the laboratory managers that provided the isolates used in the present study claimed that the isolates came from virulent outbreaks, precise information was obtained only for two: 1096-V2 and 1096-V4, with high (about 80%) and medium (40%) level of mortality in field, respectively. Interestingly, these two Sp isolates presented the motifs associated with high virulence (i.e. Thr217, Ala221, Thr247), and 1096-V4 also presented Tyr at position 500.

On the other hand, possible virulence markers for VP2 amino acids were not found associated with genogroup 1 strains. Nine of ten genogroup 1 isolates classified in the current study presented Ala, Thr, and Ala at positions 217, 221, and 247, respectively. Although these isolates supposedly came from virulent outbreaks that were duly reported for diagnosis and identification, accurate information about resulting mortality percentages was unavailable. This amino acid arrangement has been frequently found, either partially (A217 and T221) or completely, in most previously described genogroup 1 isolates [30, 43]. Interestingly, two isolates (1096-V5 [*S. salar*] and BC25 [*S. salar*]) presented Ser residues at position 221, while the 1096-BC5 (*O. kisutch*) isolate presented this amino acid at positions 217 and 221. The presence of serine at position 221 was previously described for one Chilean IPNV isolate [30]. Moreover, only one isolate presented Pro at the position 217, which is in contrast to that normally described for moderate- to low-virulence Sp isolates obtained from fish or eggs originating from Ireland, Scotland, Norway, or Spain [12, 29, 44–46].

Recently, Mutoloki et al. [47] described two different VP2 amino acid motifs, or fingerprints, for IPNV associated with clinical and subclinical infections:

I64T137T217A221T247V252T281N282A319 and V64A137P217T221A247N252S281D282E319, respectively. A search for these fingerprints in the VP2 amino acid sequences of the isolates evaluated in this study showed that of the 26 isolates belonging to genogroup 5, 13 presented the complete motif associated with the clinical fingerprint. Of the remaining 13, seven presented T221 as the only substitution with respect to the clinical fingerprint, while a single isolate presented A247 as the only substitution. Three isolates had two substitutions, and two isolates had three substitutions. In most cases, the substitutions corresponded to amino acids present in the motif associated with the subclinical fingerprint, except for the two isolates presenting S221 and that of S217S221. In the case of the ten genogroup 1 isolates, all presented T221A247N252 and seven presented V64T221A247N252E319. In both cases, these corresponded to partial subclinical fingerprint sequences. Additionally, all genogroup 1 isolates presented T137 and T281, replacements associated with the clinical fingerprint. Two isolates also had I264, and one had A319, both of which are also associated with the clinical fingerprint. Furthermore, all isolates had A217 and A282, substitutions not associated with any fingerprint. Therefore, the clinical finger was only present in the genogroup 5 isolates, with very few amino acid substitutions. In turn, genogroup 1 isolates presented sequences mostly associated with the subclinical fingerprint, with practically constant amino acid substitutions in all isolates. In accordance with the above, the genogroup 5 isolates 1096-V2 and 1096-V4, with high and medium mortality levels in the field, presented amino acid motifs associated with the clinical fingerprint.

Analysis of VP5 showed that four isolates from genogroup 5 (1096-BC20 [*S. salar*], BC21 [*O. kisutch*], BC22 [*S. salar*], and BC23 [*O. kisutch*]) had undescribed deduced proteins with unusually long amino acid sequences. Although the relatedness among these four isolates was demonstrated by phylogenetic analysis, they were isolated from different regions, fish species, and environments. Therefore, establishing any relationship is unlikely. The role of these longer isoforms in VP5 is unknown. All VP5 amino acid sequences were analyzed using the online tools TMpred and TOPCONS, which detect characteristic topologies in membrane proteins or peptides and determine the presence of transmembrane domains. Results showed an absence of transmembrane sequences, in accordance with previous classification as an anti-apoptotic protein class III [48]. Moreover, to determine the presence of motifs for plasma membrane tropism, the VP5 sequences of the obtained Chilean isolates and of the infectious bursal disease virus were searched for the presence of polycationic regions [36], the signals of which were not found in the Chilean isolates.

The WebLogo analysis established that the deduced amino acid sequences of the VP5 BH4, BH3, and BH1

domains of the Chilean IPNV isolates were highly similar to corresponding BH domains in MABV Y-6 and, to a slightly lesser extent, IPNV E1-S (Ab strain) [39]. However, the principal Logo sequence obtained for BH2 was not similar to any of the selected canonical sequences. Nonetheless, combining the Logo sequences of HTTGR (with higher percentage of conservation, Fig. 6d, big letter) and DPSRW (with lowest percentages of conservation, Fig. 6d, letters below the largest) obtained a partial match with the NPTRW sequence, the BH2 domain described for MABV Y-6 (Table 6). Nevertheless, this pattern was only present in two genogroup 1 isolates and partially in one genogroup 5 isolates, which is very unrepresentative. Clearly, the Logo sequences of the Chilean isolates were unrelated to the canonical BH domain sequences present in class I (Bcl-2, Bcl-xL, and Mcl-1) and class II (Bcl-w and A-1) anti-apoptotic proteins, nor with the BH sequences of pro-apoptotic proteins (Bax, Bak and Bad) [39, 48]. Namely, VP5 sequences of Chilean IPNV isolates were highly similar to the marine birnaviruses MABV Y-6 and IPNV E1-S, an Ab serotype, in addition to being unrelated with anti- or pro-apoptotic proteins.

Shivappa et al. [12] described the presence of an ORF for a putative 25 kDa protein located between the VP2 and VP4 coding regions. When assessing the sequences coding for this putative protein in the Chilean viral isolates, only two of the nine isolates with a fully sequenced segment A coded for the complete sequence of the putative 25 kDa protein. According to the performed nuclear localization signal (NLS) alignment analysis against the putative 25 kDa protein of the IPNV strain Sp103, previously investigated by Shivappa et al. [12], two of the described sequences coincided with the two assessed Chilean isolates, with the first between amino acid residues 46 to 71 and the second predicted between residues 97 to 113. Moreover, prediction analyses using the online tools NLS Mapper and NLStradamus revealed that only the first would correspond to a NLS. These tools also detected two additional putative NLS in both Chilean isolates between residues 143 to 178 and 202 to 218 (Fig. 7). This finding partly coincided with previous descriptions in the nonstructural protein of the *Drosophila* X virus and the IPNV strain Sp103 [12]. The presently identified NLS sequences were characterized by the presence of polycationic regions that spanned from the middle of the protein towards the carboxyl terminal. A wide variety of cellular proteins employ surface-exposed, positively charged domains to interact with anionic lipids anchored to the inner plasmatic membrane leaflet [49, 50]. Moreover, the TargetP 1.1 predictor was used to supplement information regarding other possible subcellular location signals in the putative 25 kDa protein. This analysis predicted the presence of a 61 amino acid N-terminal

presequence for mitochondrial targeting in the isolates 1096-BC2 and 1096-BC16 (data not shown). The presence of such signals has been previously described for several viral proteins. This situation appears to be similar to what happens with the non-structural Borna disease virus protein X, which is targeted to both the nucleus and mitochondria of infected cells and inhibits the aggregation of the mitochondrial antiviral-signaling protein, thus blocking programmed cell death and conferring resistance to Fas-induced cell death [51]. Clearly, more investigative progress is needed for the putative protein of 25 kDa, mainly regarding detection as a protein in infected cells, but also on its role in relation to potential nuclear and mitochondrial targeting.

Conclusions

This study molecularly characterized 36 Chilean IPNV isolates sampled over 6 years from the three most cultured salmonids in Chile. The obtained results represent an important advancement in the analysis of IPNV strains present in Chile, providing information about phylogenetic relationships, geographical areas, tropism for salmon species, and characteristics of the VP2, VP5, and putative 25 kDa proteins. Moreover, the obtained data are of utmost relevance for productive salmonid management, providing information that might help in identifying epidemiological links and in developing specific sanitary tools for preventing IPNV risks in Chilean aquaculture.

Acknowledgments

We thank Samuel Valdebenito and Carlos Sandoval of Veterquímica S.A., as well as Dr. Ricardo Enriquez and Mónica Monrás of the Institute of Animal Pathology at the Universidad Austral de Chile, for generously supplying viral isolates. We also thank Dr. Jorge Dresdner of the Department of Economics at the University of Concepción, Chile, for kindly providing us with information about the salmon farming centers present in Chile.

Funding

This work was supported in part by grants from the Fondo Nacional de Desarrollo Científico y Tecnológico de Chile [Fondecyt 1120903 and 1150934], Fondo de Fomento al Desarrollo Científico y Tecnológico [Fondef D0811096], and Fondo de Financiamiento de Centros de Investigación en Áreas Prioritarias [FONDAP 15110027] - Interdisciplinary Center for Aquaculture Research (INCAR).

Availability of data and materials

The GenBank accession numbers for infectious pancreatic necrosis virus sequences reported in this paper are: KX523824, KX523802, KX523827, KX523803, KX523816, KX523804, KX523818, KX523828, KX523829, KX523801, KX523819, KX523800, KX523833, KX523821, KX523830, KX523805, KX523822, KX523820, KX523809, KX523810, KX523811, KX523799, KX523807, KX523808, KX523825, KX523806, KX523826, KX523823, KX523817, KX523813, KX523812, KX523815, KX523814, KX523832, KX523831 and KX523834.

Authors' contributions

JGC conceived the study, designed some experiments, analyzed the data, guided the experiment process, and wrote the manuscript. RAM, designed some experiments, analyzed the data, and guided the experiment process. TV, MVV, AM, VNV, and AJY contributed in data collection. All authors reviewed and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The viral isolates were supplied by the Veterquímica S.A Enterprise and Institute of Animal Pathology at the Universidad Austral de Chile, who obtained viral isolates from field samples according to the ethical standards of the National Fisheries and Aquaculture Service of Chile (SERNAPESCA. <http://www.sernapesca.cl/>). All work performed in this study was in vitro and was approved by the Ethics Committee of FONDEF (CONICYT-Chile).

Author details

¹Instituto de Bioquímica y Microbiología, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile. ²Centro FONDAP, Interdisciplinary Center for Aquaculture Research (INCAR), Valdivia, Chile. ³Institute of Marine and Environmental Technology, University of Maryland Baltimore County, Baltimore 21202, USA.

Received: 12 August 2016 Accepted: 17 January 2017

Published online: 31 January 2017

References

- http://www.sernapesca.cl/index.php?option=com_remository&Itemid=246&func=fileinfo&id=14754.
- Roberts RJ, Pearson MD. Infectious pancreatic necrosis in Atlantic salmon, *Salmo salar* L. *J Fish Dis*. 2005;28:383-90.
- Guerrero L, Herrera E, Salinas J, Torres J, Montero AB, Barron B. Detection and genotyping of an infectious pancreatic necrosis virus from asymptomatic rainbow trout (*Oncorhynchus mykiss*) facilities in Mexico. *Intervirology*. 2008;51:285-92.
- Reno PW. Infectious pancreatic necrosis and associated aquatic birnaviruses. In: Woo PTK, Bruno DW, editors. *New York: Fish Diseases and Disorders*; 1999. p. 1-54.
- Munro ES, Gahlawat SK, Acosta F, Ellis AE. In infectious pancreatic necrosis virus carrier Atlantic salmon, *Salmo salar* L., post-smolts, almost all kidney macrophages ex vivo contain a low level of non-replicating virus. *J Fish Dis*. 2005;29:43-8.
- Dobos P. The molecular biology of infectious pancreatic necrosis virus (IPNV). *Ann Rev Fish Dis*. 1995;5:25-54.
- Petit S, Lejal N, Huet JC, Delmas B. Active residues and viral substrate cleavage sites of the protease of the birnavirus infectious pancreatic necrosis virus. *J Virol*. 2000;74:2057-66.
- Galloux M, Chevalier C, Henry C, Huet J-CC, Costa BD, Delmas B. Peptides resulting from the pVP2 C-terminal processing are present in infectious pancreatic necrosis virus particles. *J Gen Virol*. 2004;85:2231-6.
- Lejal N, Costa DB, Huet JC. Role of Ser-652 and Lys-692 in the protease activity of infectious bursal disease virus VP4 and identification of its substrate cleavage sites. *J Gen Virol*. 2000;81:983-92.
- Chevalier C, Lepault J, Erk I, Da Costa B, Delmas B. The maturation process of pVP2 requires assembly of infectious bursal disease virus capsids. *J Virol*. 2002;76:2384-92.
- Sánchez AB, Rodríguez JF. Proteolytic processing in infectious bursal disease virus: identification of the polyprotein cleavage sites by site-directed mutagenesis. *Virology*. 1999; 262(1):190-9.
- Shivappa R, Song H, Yao K, Aas-Eng A, Evensen O, Vakharia V. Molecular characterization of Sp serotype strains of infectious pancreatic necrosis virus exhibiting differences in virulence. *Dis Aquat Org*. 2004;61:23-32.
- Song H, Santi N, Evensen O, Vakharia VN. Molecular determinants of infectious pancreatic necrosis virus virulence and cell culture adaptation. *J Virol*. 2005;79:10289-99.
- Santi N, Vakharia VN, Evensen Ø. Identification of putative motifs involved in the virulence of infectious pancreatic necrosis virus. *Virology*. 2004;322:31-40.
- Magyar G, Dobos P. Evidence for the detection of the infectious pancreatic necrosis virus polyprotein and the 17-kDa polypeptide in infected cells and of the NS protease in purified virus. *Virology*. 1994;204:580-9.
- Weber S, Fichtner D, Mettenleiter TC, Mundt E. Expression of VP5 of infectious pancreatic necrosis virus strain VR299 is initiated at the second in-frame start codon. *J Gen Virol*. 2001;82(Pt 4):805-12.
- Song H, Baxter-Roshek JL, Dinman JD, Vakharia VN. Efficient expression of the 15-kDa form of infectious pancreatic necrosis virus VP5 by suppression of a UGA codon. *Virus Res*. 2006;122:61-8.
- Santi N, Song H, Vakharia VN, Evensen Ø. Infectious pancreatic necrosis virus VP5 is dispensable for virulence and persistence. *J Virol*. 2005;79:9206-16.
- Heppell J, Tarrab E, Berthiaume L, Lecomte J, Arella M. Characterization of the small open reading frame on genome segment A of infectious pancreatic necrosis virus. *J Gen Virol*. 1995;76(Pt 8):2091-6.
- Skjesol A, Skjæveland I, Elnaes M, Timmerhaus G, Fredriksen BN, Jorgensen SM, Krasnov A, Jorgensen JB. IPNV with high and low virulence: host immune responses and viral mutations during infection. *Viol J*. 2011;8:396.
- Chao DT, Korsmeyer SJ. BCL-2 family: regulators of cell death. *Ann Rev Immunol*. 1998;16:395-419.
- Carballeda JM, Maroniche G, Lucero MS, Ricchetta M, Gomez E, Zoth SC, Berinstein A. Infectious Bursal Disease Virus non-structural protein VP5 is not a transmembrane protein. *Virology*. 2015;483:312-7.
- Hill BJ, Way K. Serological classification of infectious pancreatic necrosis (IPN) virus and other aquatic birnaviruses. *Ann Rev Fish Dis*. 1995;5:55-77.
- Blake S, Ma J, Caporale D, Jairath S, Nicholson B. Phylogenetic relationships of aquatic birnaviruses based on deduced amino acid sequences of genome segment A cDNA. *Dis Aquat Org*. 2001;45:89-102.
- Nishizawa T, Kinoshita S, Yoshimizu M. An approach for genogrouping of Japanese isolates of aquabirnaviruses in a new genogroup, VII, based on the VP2/NS junction region. *J Gen Virol*. 2005;86:1973-8.
- Zhang CX, Suzuki S. Aquabirnaviruses isolated from marine organisms form a distinct genogroup from other aquabirnaviruses. *J Fish Dis*. 2004;27:633-43.
- McAllister PE, Reyes X. Infectious pancreatic necrosis virus: isolation from rainbow trout, *Salmo gairdneri* Richardson, imported into Chile. *J Fish Dis*. 1984;7(4):319-22.
- Espinoza E, Farias G, Soler M, Kuznar J. Identity between infectious pancreatic necrosis virus VR-299 and a Chilean isolate. *Intervirology*. 1985; 24(1):58-60.
- Mutoloki S, Evensen Ø. Sequence similarities of the capsid gene of Chilean and European isolates of infectious pancreatic necrosis virus point towards a common origin. *J Gen Virol*. 2011;92:1721-6.
- Tapia D, Eissler Y, Torres P, Jorquera E, Espinoza JC, Kuznar J. Detection and phylogenetic analysis of infectious pancreatic necrosis virus in Chile. *Dis Aquat Organ*. 2015;116:173-84.
- Eissler Y, Pavlov MS, Conejeros P, Espinoza JC, Kuznar J. Detection and quantification of Chilean strains of infectious pancreatic necrosis virus by real-time RT-PCR assays using segment B as a target. *Lat Am J Aquat Res*. 2011;39:544-52.
- Calleja F, Godoy MG, Carcamo JG, Bandin I, Yanez AJ, Dopazo CP, Kibenge FS, Avendano-Herrera R. Use of reverse transcription-real time polymerase chain reaction (real time RT-PCR) assays with Universal Probe Library (UPL) probes for the detection and genotyping of infectious pancreatic necrosis virus strains isolated in Chile. *J Virol Methods*. 2012;183:80-5.
- Blake SL, Schill WB, McAllister PE, Lee MK, Singer JT, Nicholson BL. Detection and identification of aquatic birnaviruses by PCR assay. *J Clin Microbiol*. 1995;33:835-9.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731-9.
- Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci*. 1992;8:275-82.
- Mendez F, de Garay T, Rodríguez D, Rodríguez JF. Infectious bursal disease virus VP5 polypeptide: a phosphoinositide-binding protein required for efficient cell-to-cell virus dissemination. *PLoS One*. 2015;10:e0123470.
- Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res*. 2004;32:1792-7.
- Moldoveanu T, Follis AV, Kriwacki RW, Green DR. Many players in BCL-2 family affairs. *Trends Biochem Sci*. 2014;39(3):101-11.
- Lin JG, Zhang CX, Suzuki S. An anti-apoptosis gene of the Bcl-2 family from Marine Birnavirus inhibiting apoptosis of insect cells infected with baculovirus. *Virus Genes*. 2005;31:185-93.
- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A. ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res*. 2003;31:3784-8.
- Ba ANN, Pogoutse A, Provart N, Moses AM. NLStradamus: a simple Hidden Markov Model for nuclear localization signal prediction. *BMC Bioinformatics*. 2009;10(1):202-12.
- Davies KR, McColl KA, Wang LF, Yu M, Williams LM, Crane MS. Molecular characterisation of Australasian isolates of aquatic birnaviruses. *Dis Aquat Organ*. 2010;93(1):1-15.

43. Salgado-Miranda C, Rojas-Anaya E, Garcia-Espinosa G, Loza-Rubio E. Molecular Characterization of the VP2 Gene of Infectious Pancreatic Necrosis Virus (IPNV) Isolates from Mexico. *J Aquat Anim Health*. 2014;26:43–51.
44. Ruane NM, McCleary SJ, McCarthy LJ, Henshilwood K. Phylogenetic analysis of infectious pancreatic necrosis virus in Ireland reveals the spread of a virulent genogroup 5 subtype previously associated with imports. *Arch Virol*. 2015;160:817–24.
45. Dadar M, Peyghan R, Memari HR, Shapouri MRSA, Hasanzadeh R, Goudarzi LM, Vakharia VN. Sequence analysis of infectious pancreatic necrosis virus isolated from Iranian reared rainbow trout (*Oncorhynchus mykiss*) in 2012. *Virus Genes*. 2013;47:574–8.
46. Bain N, Gregory A, Raynard RS. Genetic analysis of infectious pancreatic necrosis virus from Scotland. *J Fish Dis*. 2008;31:37–47.
47. Mutoloki S, Jøssund TB, Ritchie G, Munang'andu HM, Evensen Ø. Infectious pancreatic necrosis virus causing clinical and subclinical infections in atlantic salmon have different genetic fingerprints. *Front Microbiol*. 2016;7:1393.
48. Hong J-R, Gong H-Y, Wu J-L. IPNV VP5, a novel anti-apoptosis gene of the Bcl-2 family, regulates Mcl-1 and viral protein expression. *Virology*. 2002;295:217–29.
49. Goldenberg NM, Steinberg BE. Surface charge: a key determinant of protein localization and function. *Cancer Res*. 2010;70:1277–80.
50. Marfori M, Mynott A, Ellis JJ, Mehdi AM, Saunders NF, Curmi PM, Forwood JK, Boden M, Kobe B. Molecular basis for specificity of nuclear import and prediction of nuclear localization. *Biochim Biophys Acta*. 1813;2011:1562–77.
51. Ferré CA, Davezac N, Thouard A, Peyrin J-M, Belenguer P, Miquel M-C, et al. Manipulation of the N-terminal sequence of the Borna disease virus X protein improves its mitochondrial targeting and neuroprotective potential. *FASEB J*. 2016;30:1523–33.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

