Published in final edited form as:

Clin Trials. 2016 December; 13(6): 582-591. doi:10.1177/1740774516665598.

Health Research Participants' Preferences for Receiving Research Results

Christopher R. Long,

Office of Community Health and Research, University Arkansas for Medical Sciences, Northwest Campus, Fayetteville, AR, USA

M. Kathryn Stewart,

Fay W. Boozman College of Public Health, University Arkansas for Medical Sciences, Little Rock, AR, USA

Thomas V. Cunningham,

Division of Medical Humanities and Department of Internal Medicine, UAMS College of Medicine, University Arkansas for Medical Sciences, Little Rock, AR, USA

T. Scott Warmack, and

College of Pharmacy, University of Arkansas for Medical Sciences, Northwest Campus, Fayetteville, AR, USA

Pearl A. McElfish

Office of Community Health and Research, University Arkansas for Medical Sciences, Northwest Campus, Fayetteville, AR, USA

Abstract

Background—Participants in health research studies typically express interest in receiving results from the studies in which they participate. However, participants' preferences and experiences related to receiving results are not well understood. In general, existing studies have had relatively small sample sizes and typically address specific and often sensitive issues within targeted populations.

Methods—The present study used an online survey to explore attitudes and experiences of registrants in ResearchMatch, a large database of past, present, and potential health research participants. Survey respondents provided information related to whether or not they received research results from studies in which they participated, the methods used to communicate results, their satisfaction with results, and when and how they would like to receive research results from future studies. 70,699 ResearchMatch registrants were notified of the study's topic. Of the 5,207 registrants who requested full information about the study, 3,381 respondents completed the survey.

Corresponding author contact information: Pearl A. McElfish, Office of Community Health and Research, University Arkansas for Medical Sciences, Northwest Campus, 1125 N. College Ave, Fayetteville, AR 72701, pamcelfish@uams.edu 479.713.8680.

Results—Approximately 33% of respondents with previous health research participation reported receiving results. Approximately half of respondents with previous research participation reported no opportunity to request results. However, almost all respondents said researchers should always or sometimes offer results to participants. Respondents expressed particular interest in results related to their (or a loved one's) health, as well as information about studies' purposes and any medical advances based on the results. In general, respondents' most preferred dissemination methods for results were email and website postings. The least desirable dissemination methods for results included Twitter, conference calls, and text messages. Across all results, we compare the responses of respondents with and without previous research participation experience, and those who have worked in research organizations vs. those who have not. Compared to respondents who have previous participation experience, a greater proportion of respondents with no participation experience indicated that results should always be shared with participants. Likewise, respondents with no participation experience placed higher importance on the receipt of each type of results information included in the survey.

Conclusions—We present findings from a survey assessing attitudes and experiences of a broad sample of respondents that addresses gaps in knowledge related to participants' preferences for receiving results. The study's findings highlight the potential for inconsistency between respondents' expressed preferences to receive specific types of results via specific methods and researchers' unwillingness or inability to provide them. We present specific recommendations to shift the approach of new studies to investigate participants' preferences for receiving research results.

Keywords

results dissemination; results communication; disseminating results; participant preferences; research results; dissemination; social media dissemination; dissemination survey; ResearchMatch; participant registry

Members of the research community often express support for communicating research results to research participants and to the general public. ^{1–7} To this end, institutional review boards (IRBs) and research ethics boards often require researchers to address plans for the dissemination of study results in their study protocols. ^{3,8,9} Likewise, research funders—including the United States (U.S.) National Institutes of Health (NIH), Patient-Centered Outcomes Research Institute, and Agency for Healthcare Research and Quality—have emphasized the importance of dissemination of results to nonacademic audiences. ^{10–12}

Despite the recommendations to disseminate results to participants and communities, participant- and community-level dissemination may be rarely implemented by researchers. One survey of oncology physicians and nurses found 72.4% of respondents believed most participants wanted to know the results of clinical trials, yet 62.4% of respondents shared trial results with participants less than 20% of the time. Likewise, among 158 researchers presenting at an American Society of Hematology Annual Meeting, 69% supported returning research results to participants, but only 30% of the researchers had a plan to offer results to participants. Similarly, it is rarely clear from published research articles whether or not results have been communicated to participants or the general public. A review of 101

journal articles reporting on community-based participatory research studies found only 48% indicated further dissemination beyond publication of the articles.²

While many in the research community express support for communication of results, some researchers have raised ethical concerns related to consequences of communicating results to participants. ^{9,13,14} Researchers have expressed concern about participants receiving research results that they find difficult to understand or emotionally troubling. ^{1,15–17} A 2008 review of research on participants' responses to receiving research results found participants report a range of positive and negative reactions, ranging from anxiety, anger, or guilt to relief or pleasure. ¹⁸ In addition, unintended consequences can result from participants' uncertainty with what to do with results or from researchers' uncertainty with how best to present results or which results to share with participants. ¹³

Nevertheless, the 2008 review examined nine studies reporting percentages of participants who indicated preferences for receiving results; it found a median of 90% of participants reported wanting to receive results. ¹⁸ Prominent among participants' reported motives to receive results are that results carry potential clinical implications for themselves or their loved ones and that participants have a general right or need to know results of studies in which they participated. ^{15,18–21}

Taken together, existing studies about communicating research results to participants suggest the following pattern: Participants and research team members typically express support for communicating research results; results are seldom communicated to participants; when results are communicated, participants' responses may include positive or negative emotions (or both).¹⁸

There are limitations to generalizing across the results of existing studies of participants' reactions and preferences related to receiving research results: In general, existing studies have had relatively small sample sizes and typically address specific and often sensitive issues within targeted populations. Representative examples include studies of 20 participants in a United Kingdom trial of prenatal antibiotics¹⁹; 24 potential participants (or parents of potential participants) in a study testing children for neurodevelopmental deficits²²; 44 participants in a United Kingdom dementia study²³; 135 U.S. participants in a phase II trial of breast excision as breast cancer treatment²⁴; 409 Canadian and U.S. parents of children with cancer and 86 adolescents with cancer²⁰; 540 Ugandan participants in observational research on living with HIV²⁵; and 121 potential participants in genetic research, including 60 Seattle members of a health maintenance organization and 61 Yup'ik Alaska Native people.²¹ These studies document sample-specific concerns and preferences; broadly, their results converge most conclusively upon the basic finding that study participants would like to receive research results.

Rather than focusing on a particular subgroup, the present study assessed attitudes and experiences of a broad sample of participants who have signed up to be part of a research registry. Investigating participants' preferences related to the communication of research results, the present study used online survey methodology to explore attitudes and experiences of registrants in a large database of past, present, and potential health research

participants. Survey respondents provided information across a range of topics, including whether or not they received research results from studies in which they participated, their satisfaction with those results, and when and how they would like to receive research results from future studies.

In addition, the study is intended to increase understanding of similarities and differences between respondents from specific groups who have been examined separately across previous studies related to the communication of results. For example, the analytic strategy compares preferences of respondents who have worked for research-focused organizations and those who have not. Likewise, the number of survey respondents was large enough to allow comparison between preferences of respondents with previous research participation experience and respondents who have registered as potential participants but had not yet participated in research.

METHOD

Participants and Recruitment

Supported by the NIH as part of the Clinical Translational Science Award program, ResearchMatch²⁶ is a registry of volunteers from the U.S. who have indicated interest in being contacted to participate in research studies.²⁷ Intended to connect researchers and people interested in participating in health research, ResearchMatch was developed by academic institutions, who have used online and face-to-face methods to register volunteers for the database.²⁸ Although ResearchMatch registrants represent all 50 U.S. states, they are approximately 70% female and 80% white and 90% non-Hispanic/latino.²⁹ Approximately 40% of ResearchMatch registrants report no health conditions, and 40% report no medications.²⁹ Typical projects that recruit participants from ResearchMatch range in scope from brief surveys to clinical trials.²⁷

To recruit respondents, an email notification was sent to all ResearchMatch registrants who were age 18 or older. This notification gave registrants the option to indicate whether they were interested in participating in a study about the dissemination of research results. ResearchMatch then displayed the email addresses of interested registrants to the researchers. The researchers contacted potential participants via e-mail to explain the study and provide the opportunity to consent electronically and participate. The email included a link to an online survey. Respondents were not compensated for completing the survey. Of the 70,699 individuals who were notified of the study's topic, 5,207 expressed interest in the study and were sent a link. Of those who received a link, 3,381 respondents completed the survey between December 9, 2014 and March 20, 2015. This project was determined to be exempt from human protections oversight by the IRB at the University of Arkansas for Medical Sciences (203614) on 14 October 2014.

Survey

The researchers developed a survey targeting the general population of research participants, incorporating adapted items, response options, and concepts originally assessed by Fernandez et al.,^{20,30} Dorsey et al.,³¹ Murphy et al.,³² and Partridge et al.³³ The survey was

administered online with a median completion time of 5.4 minutes. The survey used Likert-type and multiple-response items to capture perceptions and preferences regarding if, when, what, and how research results are disseminated. To examine the possibility that working for organizations that conduct research might influence individuals' beliefs and preferences regarding the dissemination of results, a question was included to identify those individuals who have worked or currently work for research organizations. Also, if respondents indicated they had participated in research in the past, they were asked a series of questions to determine if and how they had received the research results and their satisfaction level with the communication of those results. All respondents (both those who had and had not participated in research) were asked to rate their perceptions about the importance of disseminating research results, the frequency and type of results that should be shared, and the method of dissemination. In addition to traditional dissemination methods, respondents were asked in detail about their preferences for the use of social media, websites, online video, and text messages in the dissemination of research results.

Analytic Strategy

Due to the relatively large number of survey respondents and the large number of potential inferential analyses, the analytic strategy focused on presenting results of descriptive analyses for each topic included in the survey. Where inferential comparisons are presented (e.g., comparisons between respondents with vs. without previous research experience), indicators of effect size are included to aid interpretation of results. Not every statistically significant inferential comparison (where alpha = .05 and p < .05) is reported. We emphasized reporting results for which effect sizes were non-negligible or which could be interpreted in the context of other reported results. For example, post-hoc analyses of differences among respondents as a function of race/ethnicity yielded no differences large enough to report, particularly given the relatively low number of respondents from various race/ethnic groups, the relatively small effect sizes, and the inconsistent patterns among those small differences. All inferential analyses presented below were planned a priori. (Any planned analyses not reported below are presented in a supplemental document at the journal website, including unreported comparisons between respondents with vs. without previous research experience or who have vs. have not worked in research organizations.)

RESULTS

Participant Characteristics

Demographic characteristics of the respondents are presented in Table 1. The majority of respondents were female (79.5%) and white non-Hispanic/latino (87.5%), similar to the overall ResearchMatch population (70% female, 80% white, and 90% non-Hispanic/latino). Over two-thirds of respondents have household incomes of at least \$50,000 (67.9%) and have completed at least a Bachelor's degree (71.7%). Approximately half of the respondents have worked or currently work at research organizations (50.6%).

Comparison of Respondent Demographics

Comparing the demographics of respondents who work or have worked at research organizations (50.6%) with respondents who have never worked at research organizations

(48.5%) yielded several statistically significant differences. Relative to those who have never worked at research organizations, respondents who have worked at research organizations were more likely to be female than male (85.0% vs. 74.1%; X^2 , 1, = 61.64, p<.01, φ = .14), to be younger (M= 44.58 vs. M= 48.34; t, 3289, = 7.30, p<.01, d= .24), to have completed a higher level of education (Mann-Whitney U= 898543.50, p<.01, r= .31), and to have a higher income (Mann-Whitney U= 1110240.50, p<.01, r= .14). Those who have worked at research organizations were also more likely to have participated previously in research either as participants or as guardians of participants, than were those who have never worked at research organizations, 87.4% vs. 61.1%, X^2 (1) = 303.27, p<.01, φ = .30.

Likewise, comparing demographics of respondents who previously participated in research (74.6%), either as participants or as guardians of participants, with those who have never previously participated (25.4%) yielded statistically significant differences similar to those reported above. Relative to those who have never participated, respondents who have participated in research were more likely to be female than male (80.5% vs. 77.1%; X^2 , 1, = 4.49, p = .03, $\varphi = .04$), to be younger (M = 45.84 vs. M = 48.00; t, 3325, = 3.64, p<.01, t = .15), to have completed a higher level of education (Mann-Whitney t = 829243.00, t = .01, t = .32), and to have a higher income (Mann-Whitney t = 904956.50, t < .01, t = .14). For results reported below, we noted where patterns of responses differ between respondents who have and have not worked at research organizations and between respondents or guardians of participants.

Past Experiences

Respondents who had previously been participants or guardians of participants were asked to report on their experiences associated with any one of the previous studies in which they participated. Of these respondents, 51.8% indicated they were given no opportunity to request the study's results, 9.0% chose not to receive the study's results, and 6.2% indicated they never received results they had requested. Only 33.0% reported receiving the study's results.

Among respondents who reported receiving results, satisfaction with both the method by which results were provided and with the information provided was high, M = 6.26 (SD = 1.08) and M = 6.15 (SD = 1.00), respectively, on 7-point scales anchored from 1 = completely dissatisfied to 7 = completely satisfied.

Table 2 shows the respondents who reported receiving results through each of a series of different methods. The most commonly reported methods were letters or fliers distributed via e-mail or postal mail. Among the 548 respondents who reported receiving results through only a single method, the highest level of satisfaction with the communication method was associated with receiving results via a meeting with a researcher whereas the least satisfaction was associated with receiving results via a website posting, M = 6.66 (SD = 0.56) and M = 0.579 (SD = 0.56), respectively.

Perceptions Related to the Communication of Research Results

Desirability of Receiving Study Results—Only eight respondents indicated researchers should never offer research results to participants (0.2%). As shown in Table 3, 63.2% reported researchers should always offer results to participants, and 32.4% reported researchers should sometimes offer results to participants. Among respondents who had experience as participants or as guardians of participants, there was a smaller proportion of respondents who indicated researchers should always share results than among those who did not have such experience; 61.3% vs. 68.5%, $X^2(1) = 13.33$, p < .01, $\varphi = .06$.

General Preferences for Methods of Receiving Results—Respondents were asked to consider a hypothetical situation in which they or a loved one had participated in a research study and to then indicate the desirability of receiving the study's results through each of a series of different methods (i.e., informational sessions at universities or community gathering places; letters/fliers mailed or emailed; Facebook; Twitter; postings or videos on a website; text messages; and phone or conference calls). On a 5-point scale anchored from $1 = very \ undesirable$ to $5 = very \ desirable$, respondents rated letters or fliers distributed via e-mail (M = 4.06; SD = 0.95) and postings on study websites (M = 3.83; SD = 1.01) as the most desirable methods, and Twitter (M = 2.00; SD = 1.05), text messages (M = 2.35; SD = 1.23), and conference calls (M = 2.35; SD = 1.17) as the least desirable methods. Informational meetings—whether at universities (M = 3.20; SD = 1.16) or community gathering places (M = 3.12; SD = 1.12)—were not rated as particularly desirable or undesirable.

Preferences for Types of Results to Be Disseminated—Table 4 shows respondents' relative preferences for receiving results for particular kinds of research. On a 4-point scale anchored from 1 = not *important* to 4 = very *important*, respondents indicated the highest importance for receiving results directly related to their health or the health of a loved one, followed by results from research in which the respondents have participated, M = 3.61 (SD = 0.63) and M = 3.34 (SD = 0.82), respectively. Results not directly relevant to one's health or to the health of loved ones was rated as relatively unimportant, M = 2.22; SD = 0.84.

With respect to rating the importance for receiving results for particular kinds of research, there were no notable differences between respondents who have worked at research organizations compared to those who have not. However, for each kind of research, respondents who have previous participation experience—either as participants or as guardians of participants—rated receipt of results information as less important than did those who have no previous experience, all *ts* 3.44, all *ps* .01, all *ds* from .14 to .38.

Respondents were also asked to indicate specific types of information participants should receive when receiving results in general. Their responses are summarized in Table 5. The most commonly endorsed type of information was the purpose of the study (90.2%), followed by medical treatment advances based on the study's results (85.2%). Although at least 70% of participants endorsed each type of information, the least commonly endorsed types were neutral results of the study (70.3%) and respondents' own personal results (70.6%).

The greatest percentage difference between the responses of those who have worked at research organizations versus those who have not were that relatively more of those who have worked at research organizations indicated participants should receive information about neutral results of the study (74.5% vs. 65.7%). There were no percentage differences of this magnitude found when comparing responses from respondents with previous participation experience—either as participants or as the guardians of participants—and those without such experience.

Preferred Frequency and Milestones for Sending Results to Participants

With respect to respondents' desired frequencies and milestones for sending results to study participants, no option was endorsed by a majority of respondents. As Table 6 indicates, the modal responses indicated study participants should receive results after the final results have been reviewed for accuracy by other researchers (46.9%) or after the study has been published (45.6%).

The greatest percentage difference between the responses of those who have worked at research organizations and those who have not was that a greater percentage of those who have worked at research organizations selected post-publication as a milestone at which researchers should send results to participants (52.6% vs. 38.6%). Compared to respondents without participation experience, a lower proportion of respondents with previous participation experience— either as a participant or as the guardian of a participant—endorsed a frequency of every six months as a recommendation for how often results should be sent to participants (10.3% vs. 21.2%), but a higher proportion of respondents with previous research participation indicated results should be sent post-publication (48.8% vs. 36.4%).

DISCUSSION

While most prior studies of participants' preferences for receiving results focus on small samples of specific patient populations, ^{18–21,24,25,34} the present study assessed the preferences of a much larger sample of respondents selected independent of their health history. The profile of respondents' preferences in the present study is in many ways consistent with prior smaller-sample studies. As in previous studies, ¹⁸ over 90% of survey respondents indicated researchers should offer results to participants at least sometimes. Respondents were particularly interested in receiving results related to their health or the health of a loved one, ^{15,18–21} even if those results are not personalized, individual results. Respondents with research participation experience reported satisfaction with results they have received from studies; however, consistent with prior research, ¹⁸ only 33.0% of participants reported receiving results from a study in which they participated, and over half indicated being given no opportunity to request results.

While replicating previous research findings, the present study moves beyond those findings in important ways. For example, inasmuch as our study found participants want results provided back to them, our respondents also indicated relatively low-cost methods of dissemination such as email or website postings may be acceptable to participants.

It is important to note several potential limitations to our findings' generalizability. These findings are based upon preferences and experiences of respondents from ResearchMatch who expressed a willingness to participate. The present survey's respondents are approximately 80% female and 90% white non-Hispanic/latino and overrepresent particular demographic segments of the U.S. population whose responses may or may not generalize to other groups' responses. However, in post-hoc analyses of differences among respondents as a function of race/ethnicity, we found no differences large enough to report.

A second potential limitation is that no data were collected regarding participants' current or prior health conditions; likewise; for those 50.6% respondents who have worked in research institutions, we are uncertain of the extent to which they were directly involved in research endeavors. For these reasons, it is difficult to determine the extent to which respondents' experiences are reflective of the population of people who might participate in any particular health research study (e.g., respondents with research expertise may be biased toward sharing all results, regardless of outcome, or toward the use of low-cost dissemination methods). Moreover, no data were collected regarding the time elapsed since respondents participated in the studies they described; the possibility exists that the passage of time has introduced biases among some survey responses (e.g., misremembering whether or not results were received from a study).

Another potential limitation relates to questions on the survey asking participants to report whether or not they would like information about results from a hypothetical study. Absent presentation of potential risks or costs associated with receiving results, it is unsurprising that participants would generally report wanting this information. ^{35,36} Participants may also overestimate the practical utility of results of individual studies and therefore place a high value on receiving results. Because of these ambiguities, further large-sample research is needed to specify respondents' motivations for desiring results, both within and across specific populations.

The above limitations notwithstanding, the present study assessed the attitudes and experiences of a broad sample of respondents as a step toward addressing gaps in knowledge related to participants' preferences for receiving results, and it explored differences between participants with and without previous participation experience and those who have and have not worked at research organizations. This study's findings highlight the potential for inconsistency between respondents' expressed preferences to receive types of results via specific methods, and researchers' unwillingness or inability to provide them. Most respondents indicated preferences to receive results, but only 33% of respondents reported receiving results from studies in which they participated. The resolution of this inconsistency is unlikely to be simple, as would be asking participants to check online repositories of research results or of publications after the study's completion. For example, a 2010 analysis showed that fewer than half of NIH-funded trials post results within 30 months after completion, and a third of trials remained unpublished after a median of 51 months after completion.³⁷

We suggest the following approaches to identify and address barriers to dissemination of research results to participants. Current knowledge about participants' preferences and

experiences related to receiving results is based largely upon findings from studies mostly intended to document preferences and experiences of specific populations. A 2013 Agency for Healthcare Research and Quality report reviewing existing research on health research communication and dissemination strategies noted significant gaps in knowledge, some of which were related to the homogeneity of research methodologies often employed in this domain. To address these gaps, the report recommends reliance upon "proven data collection methods that can include, but might go beyond, self-reported attitudes, levels of knowledge, and behaviors." To the specific domain of disseminating results to participants, we echo this recommendation. We advocate shifting the focus away from studies that document preferences and toward approaches that investigate (a) origins of participants' expressed preferences, (b) how participants' preferences are informed by knowledge about particular studies and about the research process in general, (c) whether and how participants intend to use—and then how they actually use—results from studies in which they participate, and (d) variations in satisfaction and impact of different dissemination approaches.

To further identify and address barriers to dissemination of research results to participants, we recommend expanding the scope of existing research^{e.g.,} 4,5,39 on health researchers' intentions related to returning results. Specifically, we advocate for investigations of how researcher intentions; characteristics of particular types of studies; and logistical, economic, and other barriers interact to contribute to the disjunction between participants' preferences to receive results and researchers' inability or unwillingness to provide them. We also recommend involving participants and researchers in a collaborative process to address this disjunction, in order to gain understanding of how to modify participants' preferences, researchers' practices, or both. For example, there are important questions about the extent to which researchers must balance responsiveness to participants' preferences for research results against any obligations to educate participants with respect to issues of generalizability, replicability, or utility of results. ^{6,13} This could be accomplished by convening working groups of relevant stakeholders to identify appropriate research questions and research agendas that are meaningful to researchers, community members, and patients. Collaboration between researchers, patients, and other stakeholders is a necessary step toward an ethical, practical plan for dissemination that satisfies both constituencies.

Acknowledgments

Grant support acknowledgement:

The project described was supported by the Translational Research Institute, grant UL1TR000039 through the United States National Institutes of Health (NIH) National Center for Research Resources and National Center for Advancing Translational Sciences. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

REFERENCES

- 1. Partridge AH, Winer EP. Informing clinical trial participants about study results. JAMA. 2002; 288(3):363–365. [PubMed: 12117402]
- 2. Chen PG, Diaz N, Lucas G, Rosenthal MS. Dissemination of results in community-based participatory research. Am J Prev Med. 2010; 39(4):372–378. [PubMed: 20837290]

3. Fernandez CV, Kodish E, Shurin S, Weijer C. Offering to return results to research participants: attitudes and needs of principal investigators in the Children's Oncology Group. J Pediatr Hematol Oncol. 2003; 25(9):704–708. [PubMed: 12972805]

- 4. Partridge AH, Hackett N, Blood E, et al. Oncology physician and nurse practices and attitudes regarding offering clinical trial results to study participants. J Natl Cancer Inst. 2004; 96(8):629–632. [PubMed: 15100341]
- Rigby H, Fernandez CV. Providing research results to study participants: support versus practice of researchers presenting at the American Society of Hematology annual meeting. Blood. 2005; 106(4):1199–1202. [PubMed: 15878983]
- Shalowitz DI, Miller FG. The search for clarity in communicating research results to study participants. Journal of Medical Ethics. 2008; 34(9):2. [PubMed: 18156510]
- 7. Fernandez CV, Ruccione K, Wells RJ, et al. Recommendations for the return of research results to study participants and guardians: A report from the Children's Oncology Group. J Clin Oncol. 2012; 30(36):4573–4579. [PubMed: 23109703]
- 8. Fernandez CV, Skedgel C, Weijer C. Considerations and costs of disclosing study findings to research participants. Canadian Medical Association Journal. 2004; 170(9):1417–1419. [PubMed: 15111476]
- 9. Markman M. Providing research participants with findings from completed cancer-related clinical trials: Not quite as simple as it sounds. Cancer. 2006; 106(7):1421–1424. [PubMed: 16502435]
- [Accessed May 18, 2015] National Institute on Minority Health and Health Disparities Vision & Mission. 2015. http://www.ncbi.nlm.nih.gov/pubmed/
- 11. Engagement. 2015 [Accessed May 19, 2015] http://www.pcori.org/program/engagement.
- 12. Carpenter, D., Nieva, V., Albaghal, T., Sorra, J. Development of a Planning Tool to Guide Research Dissemination. In: Henriksen, K.Battles, JB.Marks, ES., Lewin, DI., editors. Advances in Patient Safety: From Research to Implementation (Volume 4: Programs, Tools, and Products). Rockville, MD: Agency for Healthcare Research and Quality; 2005. p. 83-91.
- Miller FA, Christensen R, Giacomini M, Robert JS. Duty to disclose what? Querying the putative obligation to return research results to participants. Journal of Medical Ethics. 2008; 34(3):210– 213. [PubMed: 18316466]
- 14. Laurie G. Recognizing the right not to know: Conceptual, professional, and legal implications. J Law Med Ethics. 2014; 42(1):53–63. [PubMed: 26767476]
- 15. Snowdon C, Garcia J, Elbourne D. Reactions of participants to the results of a randomised controlled trial: exploratory study. BMJ. 1998; 317(7150):21–26. [PubMed: 9651262]
- 16. Lorimer K, Gray CM, Hunt K, Wyke S, Anderson A, Benzeval M. Response to written feedback of clinical data within a longitudinal study: A qualitative study exploring the ethical implications. BMC Medical Research Methodology. 2011; 11:10. [PubMed: 21272336]
- 17. Chung W, Kim H, Nam CM. Breast-feeding in South Korea: factors influencing its initiation and duration. Public Health Nutr. 2008; 11(3):225–229. [PubMed: 17683647]
- 18. Shalowitz DI, Miller FG. Communicating the results of clinical research to participants: Attitudes, practices, and future directions. PLOS Medicine. 2008; 5(5):e91. [PubMed: 18479180]
- Dixon-Woods M, Jackson C, Windridge KC, Kenyon S. Receiving a summary of the results of a trial: qualitative study of participants' views. BMJ. 2006; 332(7535):206–210. [PubMed: 16401631]
- Fernandez CV, Gao J, Strahlendorf C, et al. Providing research results to participants: attitudes and needs of adolescents and parents of children with cancer. J Clin Oncol. 2009; 27(6):878–883.
 [PubMed: 19164211]
- Trinidad SB, Ludman EJ, Hopkins S, et al. Community dissemination and genetic research: moving beyond results reporting. Am J Med Genet A. 2015
- 22. Kadimpati S, McCormick JB, Chiu Y, et al. Utilizing focus groups with potential participants and their parents: An approach to inform study design in a large clinical trial. AJOB Empir Bioeth. 2014; 5(3):31–38. [PubMed: 24955380]
- 23. Barnett NA, O'Brien JT. Patient involvement in research: An audit of study feedback to participants. Int Psychogeriatr. 2013; 25(8):1387–1388. [PubMed: 23469808]

24. Partridge AH, Wong JS, Knudsen K, et al. Offering participants results of a clinical trial: sharing results of a negative study. Lancet. 2005; 365(9463):963–964. [PubMed: 15766998]

- 25. Baylor A, Muzoora C, Bwana M, et al. Dissemination of Research Findings to Research Participants Living with HIV in Rural Uganda: Challenges and Rewards. PLOS Medicine. 2013; 10(3):4.
- Harris PA, Scott KW, Lebo L, Hassan N, Lightner C, Pulley J. ResearchMatch: a national registry to recruit volunteers for clinical research. Academic Medicine. 2012; 87(1):66–73. [PubMed: 22104055]
- ResearchMatch. ResearchMatch Volunteer FAQ. 2015 [Accessed August 12, 2015] https://www.researchmatch.org/volunteers/index_faq.php, 2015.
- 28. ResearchMatch. About. 2016 [Accessed April 13, 2016] https://www.researchmatch.org/about/.
- 29. ResearchMatch. RM Metrics as of 11/16/2015. 2015 [Accessed November 16, 2015] https://www.researchmatch.org/index_pubsitemetrics.php.
- 30. Fernandez CV, Santor D, Weijer C, et al. The return of research results to participants: pilot questionnaire of adolescents and parents of children with cancer. Pediatr Blood Cancer. 2007; 48(4):441–446. [PubMed: 16425279]
- 31. Dorsey ER, Beck CA, Adams M, et al. Communicating clinical trial results to research participants. Arch Neurol. 2008; 65(12):1590–1595. [PubMed: 19064746]
- 32. Murphy J, Scott J, Kaufman D, Geller G, LeRoy L, Hudson K. Public expectations for return of results from large-cohort genetic research. Am J Bioeth. 2008; 8(11):36–43.
- 33. Partridge AH, Burstein HJ, Gelman RS, Marcom PK, Winer EP. Do patients participating in clinical trials want to know study results? Journal of the National Cancer Institute. 2003; 95(6): 491–492. [PubMed: 12644548]
- 34. Gikonyo C, Kamuya D, Mbete B, et al. Feedback of research findings for vaccine trials: experiences from two malaria vaccine trials involving healthy children on the kenyan coast. Dev World Bioeth. 2013; 13(1):48–56. [PubMed: 23433355]
- Nease RF, Brooks WB. Patient desire for information and decision-making in health-care decisions
 The Autonomy Preference Index and the Health Opinion Survey. Journal of General Internal Medicine. 1995; 10(11):593–600. [PubMed: 8583261]
- Manson NC. Why do patients want information if not to take part in decision making? Journal of Medical Ethics. 2010; 36(12):834–837. [PubMed: 20826868]
- 37. Ross JS, Tse T, Zarin DA, Xu H, Zhou L, Krumholz HM. Publication of NIH funded trials registered in ClinicalTrials.gov: cross sectional analysis. British Medical Journal. 2012; 344:10.
- 38. McCormack, L., Sheridan, S., Lewis, M. Evidence Report/Technology Assessment No. 213. Rockville, MD: Agency for Healthcare Research and Quality; 2013. Communication and dissemination strategies to facilitate the use of health-related evidence.
- 39. Cox K, Moghaddam N, Bird L, Elkan R. Feedback of trial results to participants: A survey of clinicians' and patients' attitudes and experiences. Eur J Oncol Nurs. 2011; 15(2):124–129. [PubMed: 20674498]

Author Manuscript

Table 1

Characteristics of survey respondents.

	Respondents Who Have Worked for Research Organizations (n = 1704)	Respondents Who Have Not Worked for Research Organizations (n = 1632)	Respondents with Previous Research Experience (n = 2516)	Respondents without Previous Research Experience (n = 857)	All Respondents (n = 3381)
	Number (% of respondents) or Mean ± SD (range)	Number (% of respondents) or Mean ± SD (range)	Number (% of respondents) or Mean ± SD (range)	Number (% of respondents) or Mean ± SD (range)	Number (% of respondents) or Mean ± SD (range)
Graduate degree	845 (49.7%)	401 (24.7%)	1030 (41.1%)	229 (26.8%)	1263 (37.5%)
Household income					
Less than \$20,000	79 (4.8%)	177 (11.2%)	169 (6.9%)	91 (11.0%)	260 (7.9%)
\$20,000 to \$34,999	153 (9.2%)	202 (12.8%)	249 (10.2%)	108 (13.0%)	358 (10.9%)
\$35,000 to \$49,999	208 (12.5%)	221 (14.0%)	303 (12.4%)	131 (15.8%)	434 (13.2%)
\$50,000 to \$74,999	342 (20.6%)	313 (19.8%)	513 (20.9%)	158 (19.0%)	671 (20.4%)
\$75,000 to \$99,999	290 (17.4%)	241 (15.2%)	421 (17.2%)	116 (14.0%)	538 (16.4%)
\$100,000 to \$149,999	348 (20.9%)	260 (16.4%)	479 (19.6%)	129 (15.5%)	611 (18.6%)
\$150,000 or more	243 (14.6%)	168 (10.6%)	315 (12.9%)	98 (11.8%)	414 (12.6%)
Ever worked for a research organization					
Yes	1704 (100.0%)	0 (0.0%)	1488 (59.4%)	214 (25.0%)	1704 (50.6%)
No	0 (0.0%)	632 (100.0%)	994 (39.7%)	632 (73.9%)	1632 (48.5%)
Unsure	0 (0.0%)	0 (0.0%)	23 (0.9%)	9 (1.1%)	32 (1.0%)
Ever participated in research/been guardian of minor participating in research?					
Yes	1488 (87.4%)	994 (61.1%)	2516 (100.0%)	0 (0.0%)	2516 (74.6%)
No	214 (12.6%)	632 (38.9%)	0 (0.0%)	857 (100.0%)	857 (25.4%)

Note. Percentages are based on the number of valid responses for each item.

Table 2

Satisfaction with disseminated results among respondents who had received results from previous participation experiences.

All Respondents Who Indicated Having Received Study Results from Previous Participation (n = 826)

Number	(%	of	survey	resi	ond	ents)

	rumber (70 of survey respondents)
Method(s) researchers used to provide study results	
Group information session	43 (5.2%)
Phone call with researcher	95 (11.5%)
Meeting with researcher	150 (18.2%)
Letter or flyer via mail	273 (33.1%)
Letter or flyer via e-mail	306 (37.0%)
Information posted on website	179 (21.7%)
Other	106 (12.8%)
Satisfaction with method used to provide results	
Completely satisfied	439 (54.9%)
Mostly satisfied	226 (28.2%)
Somewhat satisfied	74 (9.3%)
Neither satisfied nor dissatisfied	37 (4.6%)
Somewhat dissatisfied	14 (1.8%)
Mostly dissatisfied	6 (0.8%)
Completely dissatisfied	4 (0.5%)
Satisfaction with information provided	
Completely satisfied	353 (44.2%)
Mostly satisfied	294 (36.8%)
Somewhat satisfied	92 (11.5%)
Neither satisfied nor dissatisfied	41 (5.1%)
Somewhat dissatisfied	16 (2.0%)
Mostly dissatisfied	2 (0.3%)
Completely dissatisfied	1 (0.1%)

Note. For the methods items, participants could endorse more than one response option, so percentages are based on the number of respondents who indicated having received study results from previous participation. For the satisfaction items, percentages are based on the number of valid responses for each item.

Table 3

Perceived desirability of researchers' offering research results.

	ı					
$\begin{aligned} & \text{All} \\ & \text{Respondents} \\ & \text{(n = 3381)} \end{aligned}$	Number (% of survey respondents)		2062 (63.2%)	1059 (32.4%)	8 (0.2%)	136 (4.2%)
Respondents without Previous Research Experience (n = 857)	Number (% of survey respondents)		562 (68.5%)	223 (27.2%)	3 (0.4%)	33 (4.0%)
Respondents with Previous Research Experience (n = 2516)	Number (% of survey respondents)		1495 (61.3%)	835 (34.3%)	5 (0.2%)	102 (4.2%)
Respondents Who Have Not Worked for Research Organizations (n = 1632)	Number (% of survey respondents)		1010 (64.2%)	485 (30.8%)	5 (0.3%)	73 (4.6%)
Respondents Who Have Worked for Research Organizations (n = 1704)	Number (% of survey respondents)		1024 (62.1%)	561 (34.0%)	3 (0.2%)	60 (3.6%)
		Should researchers offer to give study results to the participants?	Yes, always	Yes, sometimes	No, never	Not sure / undecided

Note. Percentages are based on the number of valid responses for each item.

 Table 4

 Perceived importance of receiving different types of research results.

	Respondents with Previous Research Experience (n = 2516)	Respondents without Previous Research Experience (n = 857)	All Respondents (n = 3381)
Importance of Receiving Results	Number (% of survey respondents)	Number (% of survey respondents)	Number (% of survey respondents)
For studies conducted in my community			
Very important	616 (25.1%)	249 (30.1%)	868 (26.4%)
Important	945 (38.5%)	345 (41.8%)	1293 (39.3%)
Somewhat important	679 (27.7%)	177 (21.4%)	857 (26.1%)
Not important	215 (8.8%)	55 (6.7%)	271 (8.2%)
For studies addressing topics that interest me			
Very important	950 (38.8%)	417 (50.4%)	1368 (41.7%)
Important	1071 (43.8%)	336 (40.6%)	1411 (43.0%)
Somewhat important	366 (15.0%)	68 (8.2%)	435 (13.3%)
Not important	59 (2.4%)	7 (0.8%)	67 (2.0%)
For a study I participated in			
Very important	1202 (49.0%)	548 (66.3%)	1752 (53.3%)
Important	776 (31.6%)	208 (25.2%)	988 (30.0%)
Somewhat important	390 (15.9%)	59 (7.1%)	450 (13.7%)
Not important	85 (3.5%)	12 (1.5%)	98 (3.0%)
For a study a family member or loved one participated in			
Very important	951 (39.1%)	445 (54.1%)	1397 (42.8%)
Important	783 (32.2%)	238 (29.0%)	1025 (31.4%)
Somewhat important	514 (21.1%)	102 (12.4%)	617 (18.9%)
Not important	185 (7.6%)	37 (4.5%)	224 (6.9%)
Directly relevant to my health or a loved one's health			
Very important	1615 (65.9%)	592 (72.1%)	2212 (67.5%)
Important	653 (26.7%)	195 (23.8%)	850 (25.9%)
Somewhat important	156 (6.4%)	32 (3.9%)	188 (5.7%)
Not important	26 (1.1%)	2 (0.2%)	29 (0.9%)
NOT directly relevant to my health or a loved one's health			
Very important	149 (6.1%)	70 (8.4%)	220 (6.7%)
Important	663 (27.2%)	257 (30.9%)	921 (28.1%)
Somewhat important	1117 (45.8%)	362 (43.6%)	1483 (45.2%)
Not important	512 (21.0%)	142 (17.1%)	656 (20.0%)

Note. Percentages are based on the number of valid responses for each item.

Table 5

Types of Information that Should be Sent to Study Participants.

	Respondents Who Have Worked for Research Organizations (n = 1704)	Respondents Who Have Not Worked for Research Organizations (n = 1632)	Respondents with Previous Research Experience (n = 2516)	Respondents without Previous Research Experience (n = 857)	$All \\ Respondents \\ (n = 3381)$
	Number (% of survey respondents)	Number (% of survey respondents)	Number (% of survey respondents)	Number (% of survey respondents)	Number (% of survey respondents)
Type of information that should be included in results sent to participants					
Purpose of the study	1543 (90.6%)	1463 (89.6%)	2226 (90.1%)	774 (90.3%)	3048 (90.2%)
Medical treatment advances based on the study's results	1464 (85.9%)	1379 (84.5%)	2148 (85.4%)	726 (84.7%)	2881 (85.2%)
Good results of the study	1418 (83.2%)	1276 (78.2%)	2049 (81.4%)	679 (79.2%)	2735 (80.9%)
General scientific advances based on the study's results	1432 (84.0%)	1244 (76.2%)	2053 (81.6%)	656 (76.5%)	2715 (80.3%)
Bad results of the study	1403 (82.3%)	1263 (77.4%)	2030 (80.7%)	670 (78.2%)	2707 (80.1%)
How the study was conducted	1357 (79.6%)	1273 (78.0%)	1963 (78.0%)	694 (81.0%)	2665 (78.8%)
Potential new research based on the study's results	1344 (78.9%)	1244 (76.2%)	1956 (77.7%)	661 (77.1%)	2625 (77.6%)
Potential policy changes based on the study's results	1275 (74.8%)	1092 (66.9%)	1817 (72.2%)	577 (67.3%)	2400 (71.0%)
Results that are specific to me	1167 (68.5%)	1186 (72.7%)	1751 (69.6%)	630 (73.5%)	2386 (70.6%)
Neutral results of the study	1270 (74.5%)	1073 (65.7%)	1801 (71.6%)	569 (66.4%)	2377 (70.3%)

Note. Participants could endorse more than one response option. Percentages are based on the total number of survey responses for each group.

Long et al. Page 19

Table 6

Milestones when Study Information Should be Sent to Study Participants.

k Respondents with out Previous Previous Research Research Respondents Experience Experience Respondents (n = 2516) (n = 857) (n = 3381)	f Number (% of Number (% of Survey survey survey survey) respondents) respondents)) 259 (10.3%) 182 (21.2%) 442 (13.1%)) 228 (9.1%) 69 (8.1%) 299 (8.8%)) 927 (36.8%) 347 (40.5%) 1278 (37.8%)) 1172 (46.6%) 407 (47.5%) 1584 (46.9%)) 1229 (48.8%) 312 (36.4%) 1543 (45.6%)
Respondents Who Have Not Worked for Research Organizations (n = 1632)	Number (% of survey respondents)		283 (17.3%)	133 (8.1%)	666 (40.8%)	765 (46.9%)	630 (38.6%)
Respondents Who Have Worked for Research Organizations (n = 1704)	Number (% of survey respondents)		153 (9.0%)	165 (9.7%)	591 (34.7%) ed	795 (46.7%)	896 (52.6%)
		How often should information about a study be sent out?	Progress updates every 6 months	Progress updates every 12 months	After the final study results have been analyzed	After the final study results have been reviewed by other researchers for accuracy	After the study has

Note. Participants could endorse more than one response option. Percentages are based on the total number of survey responses for each group.