

CrossMark
click for updates

Korean J Pediatr

Single-center experience of the Korean-Developmental Screening Test for infants and children

Chae-Ri Suh¹, Su Ye Sohn, MD², Gun-Ha Kim, MD, PhD¹, Seong-Kwan Jung, MD², Baik-Lin Eun, MD, PhD¹¹Department of Pediatrics, Korea University College of Medicine, Seoul, ²Woorisoa Children's Hospital, Seoul, Korea

Purpose: We investigated the number of test takers of the Korean-Developmental Screening Test (K-DST) in a single children's hospital within a year, according to age, referral rate, and follow-up percentage.

Methods: For this study, 4,062 children who visited and received K-DST at Woorisoa Children's Hospital between January and December 2015 were enrolled. Seven test sets were used according to the Korean National Health Screening Program for infants and children in the following age groups: 4 to 6, 9 to 12, 18 to 24, 30 to 36, 42 to 48, 54 to 60, and 66 to 71 months. The results of the K-DST were categorized into 4 groups as follows: further evaluation (<-2 standard deviation [$-2SD$]), follow-up test ($-2SD$ to $-1SD$), peer level ($-1SD$ to $1SD$), and high level ($>1SD$).

Results: The test participants' population and follow-up population were concentrated before the age of 24 months (2,532, 62.3%). The children most commonly referred for further evaluation were those in the 30- to 41-month age group. A mismatch was found between the results of the K-DST and the additional questions. Most of the infants and children with suspicious developmental delays showed catch-up development in their follow-up tests (43 of 55, 78.2%).

Conclusion: The use of K-DST should be encouraged, especially among children aged over 24 months. Multiple-choice question format for the additional questions is recommended to avoid confusion. We suggest a nationwide study to evaluate and revise the K-DST.

Key words: Screening tests, Development, Infant, Child, Korea

Corresponding author: Baik-Lin Eun, MD, PhD
Department of Pediatrics, Korea University Guro Hospital, 148 Gurodong-ro, Guro-gu, Seoul 08308, Korea

Tel: +82-2-2626-1229

Fax: +82-31-405-1224

E-mail: bleun@korea.ac.kr

Received: 22 August, 2016

Revised: 5 October, 2016

Accepted: 18 October, 2016

Introduction

With an aging Korean society coupled with low-birth rate, the country realizes the importance of providing an environment that fosters healthy birth and development of children as well as facilitating the well-being of the people of the nation¹⁻⁶. The government also realizes that earlier detection of developmental delays among infants and children is critical for a better prognosis and earlier rehabilitation therapy^{2,7-9}. To achieve this purpose, the government has started a national health screening program for infants and children in Korea since 2007²⁻⁵. Korean-Ages and Stages Questionnaires (K-ASQ)¹⁰ and Denver Developmental Screening Test II (DDST II) were used for evaluating developmental status^{3,4}. However, these 2 tests had certain specific problems such as unsuitable questions for Korean environments, limited age groups and test categories, no use for web version due to a copyright and low validity^{2,5,9,11-13}.

Korean Developmental Screening Test (K-DST) for infants and children is a recent screening test designed to suit the needs of Korean infants and children^{2,14}. The National health screening program has been using this test questionnaire since September 2014. As compared with the previous tests, K-DST has no test-skipped age; for national health

Copyright © 2016 by The Korean Pediatric Society

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

screening program for infants and children (0-71 months old) and has compensated some inappropriate questions related to problem-solving and personal social section. Finally, all test results are stored and managed in the National Health Insurance Service web server, which allow parents and clinicians to look all serial data when needed¹⁵. The aim of this study is to evaluate the status of the K-DST in investigating data including the number of test takers by age groups, actual referral and follow-up percentage to check the usefulness of this test in primary clinics.

Materials and methods

1. Study populations

All infants and children visiting the Woorisoa Children's Hospital for the National Health Screening Test from January to December 2015 were included in the study.

2. Study design

The National Health Screening Tests are conducted on infants and children in Korea who visited the clinic a total 7 times, that is, when they turn 4-6, 9-12, 18-24, 30-36, 42-48, 54-60, and 66-71 months of age. The K-DST questionnaire sets are administered at intervals of two months from 4-5 to 22-23 months, 3 months from 24-26 to 33-35 months, and 6 months from 36-41 to 66-71 months¹⁵. The participants take the test papers according to their corrected age at the time of their clinic visit. We did not conduct K-DST to 4-6 months of age group infants because of the current national health screening policy. The policy states that the first trial of K-DST starts from 9 months of corrected age⁷. They visited and received physical examinations and nutritional education only.

The K-DST has 6 sections and additional questions. Gross motor, fine motor, cognition, communication, social interaction, and self-control are the 6 sections. Additional questions are designed to take into account clinically important diseases, such as cerebral palsy, language delay, and autism spectrum disorders¹⁵. According to the K-DST manual, there are 4 categorized groups; the further evaluation, the follow-up test, the peer-level, and the high-level groups. Children who scored below -1 standard deviation (-1SD) are categorized as the follow-up test group, and they need short-term check-ups for re-evaluation. Children with a score between -1SD and 1SD are the peer-level group, and a group of children who scored above 1SD is the high-level group. The manual suggests that test participants with any positive additional questions should be referred for further evaluation¹⁵. According to the manual, we referred some cases of the further evaluation group or the follow-up test on K-DST results, or of the positive additional questions.

3. Statistics

We reviewed the chart retrospectively and described the quantitative data of the discrete variables as frequencies with percentile. We used the IBM SPSS Statistics ver. 22.0 (IBM Co., Armonk, NY, USA) for data processing.

4. Ethics statement

The study was approved by the Institutional Review Board of Korea University Guro Hospital (approval number: KUGH16176-001).

Results

1. Study population

Total sample cases were 4,062 out of which 2,081 (51.2%) were male and 1,981 (48.8%) were female (Table 1). The sex distributions were almost similar in each group. The numbers of each segmental group were approximately similar with the range of

Table 1. Number of cases according to age and sex (n=4,062)

Study group age	Age (mo)	Sex		Total
		Male	Female	
1st (4-6 mo)	4-5	374 (49.7)	379 (50.3)	753
	6-7	1 (100)	0 (0)	1
	Total	375 (49.7)	379 (50.3)	754
2nd (9-12 mo)	8-9	51 (54.3)	43 (45.7)	94
	10-11	219 (55.3)	178 (44.7)	397
	12-13	185 (52.6)	167 (47.4)	352
	14-15	4 (80.0)	1 (20.0)	5
	Total	460 (54.2)	389 (45.8)	848
3rd (18-24 mo)	18-19	123 (52.1)	113 (47.9)	236
	20-21	114 (50.7)	111 (49.3)	225
	22-23	144 (50.9)	139 (49.1)	283
	24-26	95 (51.4)	90 (48.6)	185
	Total	476 (51.2)	453 (48.8)	929
4th (30-36 mo)	30-32	113 (48.9)	118 (51.1)	231
	33-35	145 (54.9)	119 (45.1)	264
	36-41	76 (52.4)	69 (47.6)	145
	Total	334 (52.2)	306 (47.8)	640
5th (42-48 mo)	42-47	164 (48.2)	176 (51.8)	340
	48-53	49 (51.0)	47 (49.0)	96
	Total	213 (48.9)	223 (51.1)	436
6th (54-60 mo)	54-59	116 (52.3)	106 (47.7)	222
	60-65	28 (49.1)	29 (50.9)	57
	Total	144 (51.6)	135 (48.4)	279
7th (66-71 mo)	66-71	79 (45.1)	96 (54.9)	175
Total		2,081 (51.2)	1,981 (48.8)	4,062

Values are presented as number of cases (%).

200 to 400. The largest number of test group was the 18- to 24-month-old age group, 929 (22.9%). The number of test participants' increased until 118- to 24-month-old age group, and then declined. The total number of K-DST takers was 3,308, because 754 out of 4,062 cases were the members of 4- to 6-month-old group who did not take K-DST. Twenty children had been already in rehabilitation therapy.

2. Follow-up cases

The total number of follow-up cases was 264 (Table 2). The 4- to 6-month-old age group had the largest follow-up cases, that is, a total of 196 cases (74.2%), followed by 10- to 11-month-old age group with 22 cases (8.3%). Most children took follow-up tests before the age of 22-23 months (225 cases, 85.2%). Among the

Table 2. Number of follow-up test takers in 1 year according to age group (n=264)

Age at first test (mo)	Age at follow-up test (mo)	No. (%)
4-6	8-9	23 (8.7)
	10-11	96 (36.4)
	12-13	78 (29.5)
8-9	18-19	5 (1.9)
	18-19	16 (6.1)
10-11	20-21	6 (2.3)
	22-23	1 (0.4)
	18-19	7 (2.7)
12-13	20-21	4 (1.5)
	22-23	3 (1.1)
	30-32	1 (0.4)
18-19	30-32	2 (0.8)
	30-32	3 (1.1)
22-23	30-32	2 (0.8)
	33-35	1 (0.4)
30-32	42-47	1 (0.4)
	42-47	3 (1.1)
33-35	42-47	6 (2.3)
	54-59	5 (1.9)
36-41	66-71	1 (0.4)
42-47		
60-65		
Total		264

Values are presented as number of cases (%).

Table 3. Results of the Korean-Development Screening Test for Infants and Children

Development level	Gross motor	Fine motor	Cognition	Communication	Social interaction	Self-control
Further evaluation	89 (2.7)	59 (1.8)	59 (1.8)	95 (2.9)	81 (2.4)	40 (1.6)
Follow-up test	387 (11.7)	369 (11.1)	334 (10.1)	496 (15.0)	350 (10.6)	243 (9.9)
Peer-level	1,531 (46.3)	1,621 (49.0)	1,640 (49.6)	1,455 (44.0)	1,614 (48.8)	1,334 (54.2)
High-level	1,301 (39.3)	1,259 (38.1)	1,275 (38.5)	1,262 (38.1)	1,263 (38.2)	843 (34.3)
Total	3,308 (100)	3,308 (100)	3,308 (100)	3,308 (100)	3,308 (100)	2,460 (100)

Values are presented as number of cases (%).

follow-up cases, 3 people were receiving rehabilitation therapy.

3. Results of K-DST

The percentage of each section according to the 4 developmental levels was similar. The percentages of each section of the "Further evaluation" group were around 2% in general (Table 3). The average percentage of the high developmental level was about 40%.

4. The number of referrals

The distribution of further evaluation cases had 2 peak points, 10- to 11-month-old age group (56 cases) and 33- to 35-month-old age group (35 cases) (Table 4A). With respect to sections, communication section had the largest number of cases, 95 (2.9%), followed by gross motor section, 89 (2.7%). The distribution of follow-up test cases also had 2 peak points, 12-13 months (148 cases) and 42-47 months (83 cases) (Table 4B). The follow-up test group had the largest number of cases on communication, 496 (15.0%), followed by fine motor, 412 (12.5%). Total referral rates among positive cases in at least 1 section were 11% for further evaluation group and 4.5% for the follow-up test group. Both level of groups had no referral under the age of 18-19 months, and both groups had concentrated number of referral between the age of 30-32 months and 36-41 months. In the fine motor section of further evaluation level, 37 was the largest number of cases in this concentrated zone, followed by the communication section with 31 cases. Both communication and self-control sections of the follow-up test group had the largest population with 80 cases each.

5. Additional questions

We compared the results between the social interaction section and questions regarding social interaction among the additional questions (Table 5). The number of cases with positive additional questions in the Below -1 SD (the further evaluation group and the follow-up test group) group was only 47 out of 165 cases (28.5%). Additionally, only 1 out of 140 cases, in the Above -1SD (the peer-level group and the high-level group) group with positive additional questions, was referred. Among all 187 cases with positive additional questions, 2 children were already in treatment.

Table 4A. Results of the “needs further evaluation” group

Age (mo) (n)	Gross motor	Fine motor	Cognition	Communication	Social interaction	Self-control	At leastone section	Refer
6–7 (1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	-	0 (0)	0 (0)
8–9 (94)	5 (5.3)	0 (0)	0 (0)	10 (10.6)	4 (4.3)	-	19 (20.2)	0 (0)
10–11 (397)	25 (6.3)	5 (1.3)	7 (1.8)	8 (2.0)	29 (7.3)	-	56 (14.1)	0 (0)
12–13 (352)	15 (4.3)	1 (0.3)	9 (2.6)	15 (4.3)	23 (6.5)	-	46 (13.0)	0 (0)
14–15(5)	0 (0)	1 (20.0)	0 (0)	0 (0)	1 (20.0)	-	1 (20.0)	0 (0)
18–19 (236)	9 (3.8)	1 (0.4)	1 (0.4)	2 (0.8)	1 (0.4)	2 (0.8)	12 (5.1)	3 (1.3)
20–21 (225)	0 (0)	1 (0.4)	7 (3.1)	4 (1.8)	0 (0)	2 (0.9)	12 (5.3)	0 (0)
22–23 (283)	0 (0)	1 (0.4)	1 (0.4)	5 (1.8)	1 (0.4)	0 (0)	7 (2.5)	0 (0)
24–26 (185)	4 (2.2)	5 (2.7)	8 (4.3)	5 (2.7)	3 (1.6)	7 (3.8)	20 (10.8)	0 (0)
30–32 (231)	6 (2.6)	9 (3.9)	2 (0.9)	10 (4.3)	5 (2.2)	11 (4.8)	24 (10.4)	12 (5.2)
33–35 (264)	4 (1.5)	23 (8.7)	6 (2.3)	10 (3.8)	6 (2.3)	8 (3.0)	35 (13.3)	5 (1.9)
36–41 (145)	1 (0.7)	5 (3.4)	6 (4.1)	11 (7.5)	3 (2.1)	3 (2.1)	15 (10.3)	7 (4.8)
42–47 (340)	5 (1.5)	4 (1.2)	3 (0.9)	5 (1.5)	2 (0.6)	1 (0.3)	12 (3.5)	0 (0)
48–53 (96)	2 (2.0)	0 (0)	1 (1.0)	2 (2.0)	1 (1.0)	0 (0)	4 (4.2)	1 (1.0)
54–59 (222)	9 (4.1)	1 (0.5)	6 (2.7)	6 (2.7)	1 (0.5)	5 (2.3)	6 (7.2)	3 (1.4)
60–65 (57)	0 (0)	0 (0)	2 (3.5)	1 (1.8)	1 (1.8)	0 (0)	2 (3.5)	4 (7.0)
66–71 (175)	4 (2.3)	2 (1.1)	0 (0)	1 (0.6)	0 (0)	1 (0.6)	6 (3.4)	0 (0)
Total (3,308)	89 (2.7)	59 (1.8)	59 (1.8)	95 (2.9)	81 (2.4)	40 (1.2)	277 (8.4)	35 (1.1)

Values are presented as number of cases (%).

Table 4B. Results of the follow-up test group

Age (mo) (n)	Gross motor	Fine motor	Cognition	Communication	Social interaction	Self-control	At leastone section	Refer
6–7(1)	1 (100)	0 (0)	0 (0)	1 (100)	0 (0)	-	1 (100)	0 (0)
8–9(94)	7 (7.4)	11 (11.7)	11 (11.7)	16 (17.0)	11 (11.7)	-	24 (25.5)	0 (0)
10–11 (397)	43 (10.9)	32 (8.1)	37 (9.3)	96 (24.2)	75 (18.9)	-	137 (34.6)	0 (0)
12–13 (352)	90 (25.6)	45 (12.8)	38 (10.8)	70 (19.9)	47 (13.4)	-	148 (42.0)	0 (0)
14–15(5)	2 (40.0)	1 (20.0)	1 (20.0)	1 (20.0)	0 (0)	-	1 (20.0)	0 (0)
18–19 (236)	32 (13.6)	45 (19.5)	31 (13.1)	47 (19.9)	30 (12.7)	40 (16.9)	86 (36.4)	1 (0.4)
20–21 (225)	18 (8.0)	24 (10.7)	25 (11.1)	28 (12.4)	13 (5.8)	29 (12.9)	65 (28.9)	0 (0)
22–23 (283)	26 (9.2)	41 (14.5)	35 (12.4)	43 (15.2)	32 (11.3)	25 (8.8)	94 (33.2)	0 (0)
24–26 (185)	18 (9.7)	8 (4.3)	27 (14.6)	46 (24.9)	21 (11.4)	13 (7.0)	57 (30.8)	0 (0)
30–32 (231)	12 (5.2)	13 (5.6)	21 (9.1)	37 (16.0)	21 (9.1)	22 (9.5)	47 (20.3)	8 (3.5)
33–35 (264)	24 (9.1)	29 (11.0)	28 (10.6)	27 (10.2)	29 (11.0)	29 (11.0)	57 (21.6)	13 (4.9)
36–41 (145)	18 (12.3)	29 (19.9)	18 (12.3)	16 (11.0)	15 (10.3)	29 (19.9)	47 (32.4)	10 (6.9)
42–47 (340)	40 (11.7)	29 (19.9)	19 (5.6)	23 (6.7)	22 (6.5)	25 (7.3)	83 (24.4)	3 (0.9)
48–53 (96)	11 (11.2)	17 (17.3)	5 (5.1)	3 (3.1)	7 (7.1)	5 (5.1)	21 (21.9)	5 (5.2)
54–59 (222)	18 (8.2)	18 (8.2)	19 (8.7)	11 (5.0)	11 (5.0)	10 (4.6)	41 (18.5)	3 (1.4)
60–65 (57)	10 (17.5)	8 (14.0)	4 (7.0)	10 (17.5)	2 (3.5)	4 (7.0)	23 (40.4)	1 (1.8)
66–71 (175)	17 (9.7)	19 (10.9)	15 (8.6)	21 (12.0)	14 (8.0)	12 (6.9)	48 (27.4)	0 (0)
Total (3,308)	387 (11.7)	369 (11.2)	334 (10.1)	496 (15.0)	350 (10.6)	243 (7.3)	980 (29.6)	44 (1.3)

Values are presented as number of cases (%).

6. K-DST results of follow-up cases

Among the total follow-up cases (Table 2), 197 cases had their first time visit in their 4- to 6-month-old age group. Only 67 cases had taken 2 serial tests. There were 16 cases in a “Above

-1SD to Below -1SD” group, whose first K-DST results were Above -1SD but the following test results became Below -1SD (Fig. 1). The fine motor section had the largest case numbers (10) on “Above -1SD to Below -1SD” group. Not a single case was referred because all cases on their following study were the follow-

Table 5. Comparison of numbers between the social interaction levels and additional questions in the K-DST

Age (mo) (No. of cases)	Developmental level of the social interaction section (No. of cases)	Positive answers for additional questions referring social interaction (No. of referral)	
		0	More than 1
24–26 (185)	Delayed (24)	17 (0)	7 (0)
	Normal (161)	150 (0)	11 (0)
30–32 (231)	Delayed (26)	16 (0)	10 (2)
	Normal (205)	194 (1)	11 (1)
33–35 (264)	Delayed (36)	25 (0)	11 (1)
	Normal (228)	216 (0)	12 (0)
36–41 (145)	Delayed (18)	11 (0)	7 (2)
	Normal (127)	117 (0)	10 (0)
42–47 (340)	Delayed (24)	21 (1)	3 (1)
	Normal (316)	294 (0)	22 (0)
48–53 (96)	Delayed (8)	7 (0)	1 (0)
	Normal (88)	79 (0)	9 (0)
54–59 (222)	Delayed (12)	8 (0)	4 (0)
	Normal (210)	180 (0)	30 (0)
60–65 (57)	Delayed (3)	3 (0)	0 (0)
	Normal (54)	46 (0)	8 (0)
66–71(175)	Delayed (14)	10 (0)	4 (0)
	Normal (161)	134 (0)	27 (1)

K-DST, Korean-Development Screening Test for Infants and Children; delayed, the further evaluation group + the follow-up test group; normal, the peer-level + the high-level.

up test levels. There were a total of 43 cases whose previous K-DST results indicated suspicious developmental delay and then scored higher in follow-up study. Communication section had the largest number of cases (12). Considering that one child scored below -1SD in multiple sections, total 38 out of 67 follow-up cases scored below -1SD on at least one of the K-DST sections. A more in-depth analysis of each section revealed that 3 children on gross motor were diagnosed as suspicious delayed group on the second test, and their scores were between -2SD and -1SD in the follow-up test group. Ten children were diagnosed as suspicious delayed group, and their scores were between -2SD and -1SD. Only three cases of the further evaluation level in both serial tests were sent to special clinics.

Discussion

With growing realization of the importance to invest in people's welfare and a conducive environment for nurturing children, the Korean government is increasingly putting emphasis on developmental delay prevention and life-long health care^{1,3-5}. Some of the developmental delays could catch up to an almost normal status with early rehabilitation programs^{3,6-9}. Early detection of developmental delay needs comprehensive judgements of medical history, developmental status evaluation, physical and neurological examination along with a developmental screening test^{3,4,6-9}. The developmental screening test is especially very important in early




Fig. 1. Follow-up data of the K-DST scores. The numbers of infants and children show that the follow-up results were different, either normalized or newly screened as delayed, from the previous results. n, number of cases; K-DST, Korean-Development Screening Test for Infants and Children; Below -1SD: the further evaluation group + the follow-up test group, Above -1SD: the peer-level + the high-level.

diagnosis of developmental delay^{3,6-9,15}. The Korean government launched the Korea National Health Screening Program for Infants and Children on November 2007^{1-5,9,14,15}. Since then, every infants and children visits the primary clinic and receives serial developmental screening tests from 4 months to 71 months of age, at least before starting with elementary school^{1-4,9,15}.

The original screening tools were K-ASQ and DDST II. K-ASQ is a modified version of Ages and Stage Questionnaires-II designed in America and is more convenient than DDST II; therefore, most clinicians used K-ASQ as a screening tool^{3,5,9,15}. However, this tool could not exactly reflect Korean culture and infant care environments. Also, several questions related to problem-solving and personal-social sections were inappropriate³. The same questions were included in questionnaires in different age groups⁵. This caused a high false-negative rate, especially in the 60-month-old age group⁹. Therefore, the difficulty levels of some questions were not suitable for suspicious developmental delay detection^{5,9}. This strongly indicated the modification of questionnaires. The last test age was 60 months and children had a 1-year gap before entering elementary schools⁹. To compensate the insufficiency, a new screening tool, K-DST was designed in 2014, and this tool was suitable for Korean infants and children^{1-3,9,14}.

The study population by aging group, over 800 cases took the test in aging groups from 4-6 to 118- to 24-month-old age groups. The study participation rate started to drop from 30- to 36-month-old age group. According to the national statistic data between 2012 and 2014, the participation of the screening test started dropped from 42- to 48-month-old group (71.2%)¹⁶. This drop rate of the test participant number seems to be related to the referral rate since the first referred case appeared at the age of 18 months. Many referred cases seemed to stay at the tertiary hospitals for follow-ups.

Children usually catch up with the peer level as time goes by. Therefore, the follow-up data is very important for evaluation of their precise developmental status. Most cases of delayed group scored higher on their follow-up tests (Fig. 1). All cases of “normal to delayed” group on their follow-up study were the follow-up test levels.

In the additional question section, the screening test questions related to cerebral palsy, language delay, and autism spectrum disorders were very important and the clinicians needed to pay attention to the answers of the questions¹⁵. We found that the actual referral rate was low (3.7%) in positive additional questions, although, the manual suggested to refer all positive additional question cases for further evaluation despite normal K-DST level. Twenty cases among delayed group were already receiving rehabilitation therapy, and this could be one of the reasons for the low referral rate. Another reason could be the answer choices. Parents could answer the questions only in a “yes or no.” Therefore, the parents would answer a “yes” if their child performed the

action once or twice. This could be the reason for high positive rate for the “peer” and “high-level” groups. Therefore, multiple choices for the additional questions are necessary to improve the accuracy.

The limitation of our study was inability to evaluate the referral cases against the final results. This may be correlated with the low revisiting rate of referred cases (3 of 31, 9.7%). The parents of the referral cases choose tertiary hospital for their children’s evaluation. The primary clinician could only know the results of the further evaluation tests only when their parents notified directly. This inaccessibility would get solved with the extended web service to feed information about the test results.

When children between 4 to 6 months of age visited the primary clinic for the first time, they only received physical examination and nutrition and health education. This is because the government had previously decided the first K-DST stage as nine to twelve months in chronological age, whereas the K-DST actually uses corrected age. Therefore, any 9-month-old preterm birth infants were required to take the test questionnaire that was meant for 4- to 5-month-old infants or 6- to 7-month-old infants. Since preterm infants visit preterm clinics in a tertiary hospital regularly, many primary clinics do not need to see the questionnaires. The clinicians can use the questionnaires for 4 to 5 and 6 to 7 months age for evaluation of full-term infants under 8 months old with suspicious developmental delay. For earlier detection of developmental delay, we should expand the national developmental screening program to younger infants and children.

In conclusion, we should encourage the use of K-DST, especially among children who are more than 24 months old. Multiple-choice question formats for the additional questions are recommended to avoid confusion. We suggest a nation-wide study to evaluate and revise the K-DST.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

References

1. Eun BL, Moon JS, Eun SH, Lee HK, Shin SM, Sung IK, et al. The current child and adolescent health screening system: an assessment and proposal for an early and periodic check-up program. *Korean J Pediatr* 2010;53:300-6.
2. Moon JS, Lee SY, Eun BL, Kim SW, Kim YK, Shin SM, et al. One-year evaluation of the national health screening program for infants and children in Korea. *Korean J Pediatr* 2010;53:307-13.
3. Eun BL, Chung HJ, Cho S, Kim JK, Shin SM, Lee JH, et al. The appropriateness of the items of Korean Ages and Stages Question-

- naires (K-ASQ) developmental screening test in Korean infants and children. *J Korean Child Neurol Soc* 2014;22:29-41.
4. Eun BL, Kim SW, Kim YK, Kim JW, Moon JS, Park SK, et al. Overview of the national health screening program for infant and children. *Korean J Pediatr* 2008;51:225-32.
 5. Moon JS. Review of National Health Screening Program for infant and children in Korea. *J Korean Med Assoc* 2010;53:377-85.
 6. Im SH, Han EY, Song J. Clinical application of neurodevelopmental assessment for infants and toddlers. *J Korean Acad Child Adolesc Psychiatry* 2012;23:175-80.
 7. Edwards SL, Sarwark JF. Infant and child motor development. *Clin Orthop Relat Res* 2005;434:33-9.
 8. Yeargin-Allsopp M, Murphy CC, Oakley GP, Sikes RK. A multiple-source method for studying the prevalence of developmental disabilities in children: the Metropolitan Atlanta Developmental Disabilities Study. *Pediatrics* 1992;89(4 Pt 1):624-30.
 9. Chung HJ, Eun BL, Kim HS, Kim JK, Shin SM, Lee JH, et al. The validity of Korean Ages and Stages Questionnaires (K-ASQ) in Korean infants and children. *J Korean Child Neurol Soc* 2014;22:1-11.
 10. Heo KH, Squires J, Lee SY, Lee JS. Korean ages and stage questionnaires. Seoul: Seoul Welfare Center for Persons with Disabilities, 2005.
 11. Kim YJ, Lee JY, Sohn JA, Lee EH, Lee JA, Choi CW, et al. A validity study of the Korean Ages and Stages Questionnaires: screening for developmental delay in preterm infant. *J Korean Soc Neonatol* 2010;17:217-23.
 12. Shin DH, Lee HS, Lee JY, Choi BM, Eun BL, Hong YS, et al. Usefulness of Korean infant developmental screening test in premature infants. *Korean J Pediatr* 2005;48:1337-41.
 13. Kim EY, Sung IK. The ages and stages questionnaire: screening for developmental delay in the setting of a pediatric outpatient clinic. *Korean J Pediatr* 2007;50:1061-6.
 14. Eun BL, Chung HJ, Shin SM, Oh KJ, Kim YA, Kim CT, et al. A study on the development and validation of Korean developmental screening test for infant & children. Cheongju: Korea Centers for Disease Control and Prevention; c2016 [cited 2016 Sep 23]. Available from: http://cdc.go.kr/CDC/info/CdcKrInfo0201.jsp?menuIds=HOME001-MNU1154-MNU0004-MNU1889&fid=28&q_type=&q_value=&cid=61795&pageNum=1.
 15. Eun BL, Chung HJ. Korean-development screening test for infants and children. Cheongju: Korea Centers for Disease Control and Prevention, 2014.
 16. National Health Insurance service. The status of the Korean-Developmental Screening Test for Infants and Children target population and the number of the actual test takers [Internet]. Daejeon: Statistics Korea; 2016 [cited 2016 Nov 15]. Available from: http://stat.kosis.kr/nsieu/view/tree.do?task=branchView&id=350_350_35007_004*MT_OTITLE&hOrg=350.