

Article

Essential Oil Characterization of *Thymus vulgaris* from Various Geographical Locations

Prabodh Satyal^{1,2}, Brittney L. Murray², Robert L. McFeeters² and William N. Setzer^{2,*}

¹ Alchemy Aromatic LLC, 621 Park East Blvd., New Albany, IN 47150, USA; prabodhsatyal@gmail.com

² Department of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899, USA; blm0029@uah.edu (B.L.M.); robert.mcfeeters@uah.edu (R.L.M.)

* Correspondence: wsetzer@chemistry.uah.edu; Tel.: +1-256-824-6519

Academic Editor: Angel A. Carbonell-Barrachina

Received: 1 September 2016; Accepted: 24 October 2016; Published: 27 October 2016

Abstract: Thyme (*Thymus vulgaris* L.) is a commonly used flavoring agent and medicinal herb. Several chemotypes of thyme, based on essential oil compositions, have been established, including (1) linalool; (2) borneol; (3) geraniol; (4) sabinene hydrate; (5) thymol; (6) carvacrol, as well as a number of multiple-component chemotypes. In this work, two different *T. vulgaris* essential oils were obtained from France and two were obtained from Serbia. The chemical compositions were determined using gas chromatography–mass spectrometry. In addition, chiral gas chromatography was used to determine the enantiomeric compositions of several monoterpenoid components. The *T. vulgaris* oil from Nyons, France was of the linalool chemotype (linalool, 76.2%; linalyl acetate, 14.3%); the oil sample from Jablanicki, Serbia was of the geraniol chemotype (geraniol, 59.8%; geranyl acetate, 16.7%); the sample from Pomoravje District, Serbia was of the sabinene hydrate chemotype (*cis*-sabinene hydrate, 30.8%; *trans*-sabinene hydrate, 5.0%); and the essential oil from Richerenches, France was of the thymol chemotype (thymol, 47.1%; *p*-cymene, 20.1%). A cluster analysis based on the compositions of these essential oils as well as 81 additional *T. vulgaris* essential oils reported in the literature revealed 20 different chemotypes. This work represents the first chiral analysis of *T. vulgaris* monoterpenoids and a comprehensive description of the different chemotypes of *T. vulgaris*.

Keywords: chiral gas chromatography; mass spectrometry; hierarchical cluster analysis; antifungal activity; enantiomeric distribution

1. Introduction

Thymus vulgaris L. (Lamiaceae) is an evergreen herb native to the southern Europe and the Mediterranean [1]. The plant has been used since ancient times as a culinary ingredient, to add flavor to cheeses [2,3] and liqueurs [4,5], and to flavor meats such as rabbit, boar, and lamb [6]. Today it is a common component of *bouquet garni* [7] and of *herbes de Provence* [8]. In addition to its use in foods, *T. vulgaris* is a well-known herbal medicine that has been used for thousands of years to treat alopecia, dental plaque, dermatophyte infections, bronchitis, cough, inflammatory skin disorders, and gastrointestinal distress [9]. The major constituents of commercial *T. vulgaris* essential oil are thymol (23%–60%), γ -terpinene (18%–50%), *p*-cymene (8%–44%), carvacrol (2%–8%), and linalool (3%–4%) [10]. *T. vulgaris* oil as well as thymol have shown antibacterial, antifungal, and anti-inflammatory effects, accounting for the medicinal uses of *T. vulgaris* [9]. There are, however, numerous varieties and cultivars of *T. vulgaris*; Tropicos lists nine subspecies and varieties of *T. vulgaris* [11]. As many as 13 different chemotypes of *T. vulgaris*, based on the predominance of particular monoterpenoids in the essential oils, have been identified [12–15]. In this work, we present the essential oil compositions, including monoterpenoid enantiomeric compositions, of four different chemotypes of *T. vulgaris* essential oils from Europe. In addition, a hierarchical cluster analysis has been carried out to elucidate/delineate the

various chemotypes of *T. vulgaris*, and we have examined the antifungal properties of three essential oils from Europe.

2. Materials and Methods

2.1. Plant Material

Thymus vulgaris #1, (linalool chemotype), *T. vulgaris* #3, (geraniol chemotype), *T. vulgaris* #4, (sabinene hydrate chemotype), and *T. vulgaris* #5, (thymol chemotype) were collected in May 2015, from Nyons, France (44°21'37'' N, 5°8'23'' E), Jablanicki, Serbia (42°59'43'' N, 21°55'4'' E), Pomoravje District, Serbia (43°58'0'' N, 21°15'0'' E) and Richerenches, France (44°21'37'' N, 4°54'47'' E), respectively. The plants were identified by Julien Abisset (Natural Essential SA, France). The air-dried aerial parts of each *T. vulgaris* sample were subjected to steam distillation for 3 h. After decanting and drying of the oils over anhydrous sodium sulfate, they were stored under refrigeration (−4 °C) until analysis. Aerial parts of *T. vulgaris* #1 produced 1.6% yield of essential oil (EO), *T. vulgaris* #3 produced 1.2% yield, *T. vulgaris* #4 produced 1.0% yield, and *T. vulgaris* #5 produced 1.5% yield.

2.2. Gas Chromatography—Mass Spectrometry (GC-MS)

The essential oils of *T. vulgaris* chemotypes were analyzed by GC-MS using a Shimadzu GCMS-QP2010 Ultra operated in the electron impact (EI) mode (electron energy = 70 eV), scan range = 40–400 amu, scan rate = 3.0 scans/sec, and GC-MS solution software. The GC column was a ZB-5 fused silica capillary column with a (5% phenyl)-polymethylsiloxane stationary phase and a film thickness of 0.25 µm. The carrier gas was helium with a column head pressure of 80 psi and flow rate of 1.37 mL/min. Injector temperature was 250 °C and the ion source temperature was 200 °C. The GC oven temperature program was programmed for 50 °C initial temperature, temperature increased at a rate of 2 °C/min to 260 °C. A 5% *w/v* solution of the sample in CH₂Cl₂ was prepared and 0.1 µL was injected with a splitting mode (30:1). Identification of the oil components was based on their retention indices determined by reference to a homologous series of *n*-alkanes, and by comparison of their mass spectral fragmentation patterns with those reported in the literature [16], and stored in our in-house MS library.

2.3. Chiral Gas Chromatography—Mass Spectrometry

Chiral analysis of the essential oils was performed on a Shimadzu GCMS-QP2010S operated in the EI mode (electron energy = 70 eV), scan range = 40–400 amu, scan rate = 3.0 scans/s. GC was equipped with a Restek B-Dex 325 capillary column (30 m × 0.25 mm ID × 0.25 µm film). Oven temperature was started at 50 °C, and then gradually raised to 120 °C at 1.5 °C/min. The oven was then raised to 200 °C at 2 °C/min and held for 5 min. Helium was the carrier gas and the flow rate was maintained at 1.8 mL/min. Samples were diluted 3% *w/v* with CH₂Cl₂ and then a 0.1 µL sample was injected in a split mode with a split ratio of 1:45.

2.4. Hierarchical Cluster Analysis

A total of 81 *T. vulgaris* essential oil compositions from the published literature, as well as the four compositions from this study were treated as operational taxonomic units (OTUs). The percentage composition of 33 major essential oil components (thymol, *p*-cymene, γ -terpinene, linalool, carvacrol, geraniol, β -caryophyllene, *cis*-sabinene hydrate, borneol, α -pinene, terpinene-4-ol, myrcene, 1,8-cineole, α -terpineol, camphene, α -terpinyl acetate, α -terpinene, camphor, limonene, β -pinene, geranyl acetate, α -thujene, geranyl formate, β -cyclocitral, *cis*-verbenol, *trans*-sabinene hydrate, 1-octen-3-ol, thymol methyl ether, caryophyllene oxide, carvacrol methyl ether, bornyl acetate, thymoquinone, and α -humulene) was used to determine the chemical relationship between the various *T. vulgaris* essential oil samples by agglomerative hierarchical cluster (AHC) analysis using the XLSTAT

software, version 2015.4.01. Pearson correlation was selected as a measure of similarity, and the unweighted pair-group method with arithmetic average (UPGMA) was used for cluster definition.

2.5. Antifungal Screening

Antifungal activity was carried out using *Candida albicans* (ATCC #18804), *Cryptococcus neoformans* 24067 (serotype D or var. *neoformans*), and *Aspergillus niger* (ATCC #16888). Qualitative assessment of antifungal activity utilized broth macrodilution (for *C. albicans* and *C. neoformans*), whereas minimum inhibitory concentrations (MIC) were determined using microdilution methods. Initially cultures were grown on potato dextrose agar for 48–72 h before a single colony was isolated and grown in potato dextrose broth for 48–72 hours to create initial liquid cultures. Cells were diluted to a final concentration of 2×10^3 cells/mL using MOPS (3-(*N*-morpholino)propanesulfonic acid) buffered RPMI (Roswell Park Memorial Institute) medium and 900 μ L were aliquoted into sterile 12 \times 75 mm tubes. Each *T. vulgaris* essential oil (100 μ L of 1% DMSO solution) was added to each tube, which were then incubated at 37 °C for 72 h in a shaking incubator (175 rpm). For determination of the MIC for *C. albicans* and *C. neoformans*, microdilution in 96-well plates was performed in triplicate. Briefly, serial dilution of the *T. vulgaris* samples was performed by adding 50 μ L of RPMI to each well then an equal volume of sample to be tested to the first row. After mixing, 50 μ L was removed and added to the next row. The procedure was repeated for each row. To this mixture, 50 μ L of cells diluted to 2000 cells/mL in RPMI were added to each well. The plates were incubated for 48 hours at 37 °C before growth was quantitated visually based on turbidity.

For the mold-like Ascomycota *A. niger*, disk diffusion was used to characterize each *T. vulgaris* essential oil. Initial cultures were grown on malt extract agar for 5–7 days before conidia were collected and suspended in potato dextrose broth. The suspension was then filtered into a sterile test tube using cheesecloth to remove hyphae. Conidia suspension was then diluted until it reached an OD₆₂₅ of 0.1–0.2. The suspension (100 μ L) was plated on malt extract agar before a sterile filter paper disk was placed in the center and 50 μ L of *T. vulgaris* essential oil was added. The culture was grown for 4–5 days at 25 °C before zones of inhibition were determined.

3. Results and Discussion

3.1. Essential Oil Compositions

The chemical compositions of *T. vulgaris* essential oils from this study are listed in Table 1. *T. vulgaris* sample #1, a linalool chemotype collected from Nyons, France, was dominated by linalool (76.15%) and linalyl acetate (14.26%). The enantiomeric ratios of these two compounds were 1:99 D/L. Other compounds in the oil were β -caryophyllene (2.27%), camphor (1.79%, 100% D enantiomer), and camphene (1.17%, D/L ratio = 99:1). *T. vulgaris* #3 was a geraniol chemotype collected from Jablanicki, Serbia, and was rich in geraniol (59.75%) and esters of geraniol, geranyl acetate (16.72%) and geranyl propionate (1.26%). Oil #3 also had sizable concentrations of linalool (7.15%, D/L ratio = 97:3) and β -caryophyllene (3.67%). *T. vulgaris* #4 was collected from Pomoravje District, Serbia, and had sabinene hydrates as defining compounds with *cis*-sabinene hydrate (30.77%, D/L ratio = 97:3) and *trans*-sabinene hydrate (4.98%). Other major components in oil #4 were terpinene-4-ol (9.50%, D/L = 30:70), linalool (7.89%, D/L = 3:97), γ -terpinene (4.58%), and myrcene (4.09%). *T. vulgaris* #5, representing the thymol chemotype, was collected from Richerenches, France. The composition of oil #5 was rich in thymol (47.06%) and *p*-cymene (20.07%) with lesser quantities of linalool (5.00%, D/L = 1:99) and carvacrol (3.24%). Interestingly, the L-enantiomer dominated linalool in samples #1, #4, and #5, but was reversed in sample #3. Özek and co-workers have found L-linalool to be the predominant enantiomer in several *Thymus* species, but apparently they did not examine *T. vulgaris* [17]. Similarly, D-terpinen-4-ol was the major enantiomer in sample #3, but a minor one in samples #4 and #5. An enantiomeric composition analysis of *Melaleuca alternifolia* revealed this essential oil to be composed of mostly D-*cis*-sabinene hydrate and D-terpinen-4-ol [18]. D- α -Thujene was the dominant enantiomer

in samples #3 and #4, while L- α -thujene predominated in sample #5. The L-enantiomer of α -thujene was found to be predominant in various *Citrus* essential oils [19]. L- α -Pinene dominated samples #1 and #3, but D- α -pinene was the major enantiomer in #4. Sample #5 showed a nearly racemic mixture of α -pinenes. L-Camphene dominated samples #1 and #3, but D-camphene dominated #4 and #5. Both D- α -pinene and D-camphene were found to be the predominant enantiomers in *Ocimum canum* and *O. kilimandscharicum* essential oils [20]. Remarkably, both D-camphor and D-borneol were the only enantiomers in sample #1, and D-borneol was also the only enantiomer found in samples #3 and #4. However, only L-camphor was found in essential oils #3 and #4. This is in contrast to the results of Tabanca and co-workers, who had exclusively observed the L-enantiomers of borneol along with 99% L-camphor in *Micromeria cristata* subsp. *phrygia* [21]. Thus, not only are the chemical profiles different between the different chemotypes, but the enantiomeric distributions are also profoundly different.

Table 1. Chemical compositions and enantiomeric distributions of *Thymus vulgaris* essential oils.

RI	Compound	#1		#3		#4		#5	
		%	D/L	%	D/L	%	D/L	%	D/L
752	3-Methyl-1-penten-3-ol	tr							
783	Methyl α -methyl butyrate	0.10		0.07		tr		tr	
850	(3Z)-Hexenol							0.05	
922	Tricyclene	0.05		tr				0.09	
924	α -Thujene	tr		tr	91:9	0.71	87:13	0.43	10:90
932	α -Pinene	0.47	1:99	0.21	1:99	1.75	85:15	1.32	52:48
947	α -Fenchene							tr	
948	Camphene	1.17	99:1	0.38	99:1	0.25	0:100	1.19	10:90
971	Sabinene	tr		0.05	98:2	2.03	78:22		
977	1-Octen-3-ol	0.43		0.42					
977	1-Octen-3-ol + β -Pinene					0.97	20:80	0.54	20:80
983	6-Methylhept-5-en-2-one			0.05					
983	3-Octanone					tr		tr	
988	Myrcene	0.09		0.44		4.09		1.59	
996	3-Octanol	tr		0.07		tr			
1004	(3Z)-Hexenyl acetate	tr							
1004	<i>p</i> -Mentha-1(7),8-diene								tr
1006	α -Phellandrene					tr	55:45	0.11	
1008	δ -3-Carene							tr	
1016	α -Terpinene			tr		2.65		1.30	
1019	<i>o</i> -Cymene							0.06	
1024	<i>p</i> -Cymene	0.09		0.18		1.09		20.07	
1028	Limonene	0.05		0.39	85:15	2.85	86:14	0.39	80:20
1030	β -Phellandrene			tr		0.37		0.09	
1031	1,8-Cineole	0.37		0.31		0.30		0.72	
1033	<i>m</i> -Cymene							tr	
1034	Lavender lactone + (Z)- β -Ocimene	tr							
1045	(E)- β -Ocimene	tr						0.05	
1057	γ -Terpinene			0.09		4.58		9.03	
1063	3-Methylbut-2-enyl butanoate	tr							
1069	<i>cis</i> -Sabinene hydrate			0.31		30.77	97:3	0.17	
1070	<i>cis</i> -Linalool oxide (furanoid)	0.28							
1071	Pinol							tr	
1084	Terpinolene					0.97		0.07	
1085	<i>trans</i> -Linalool oxide (furanoid)	0.23						tr	
1089	<i>p</i> -Cymenene							tr	
1099	Linalool	76.15	1:99	7.15	97:3	7.89	3:97	5.00	1:99
1101	<i>trans</i> -Sabinene hydrate			tr		4.98		tr	
1103	Hotrienol	0.07							
1104	Nonanal	0.05				tr			
1106	α -Pinene oxide			tr					
1107	1-Octen-3-yl acetate	tr				tr			
1124	<i>cis-p</i> -Menth-2-en-1-ol					0.65			
1136	<i>trans</i> -Limonene oxide					tr			
1142	<i>trans-p</i> -Menth-2-en-1-ol					0.25			
1147	Camphor	1.79	100:0	0.11	0:100	0.10	0:100	1.42	95:5
1148	α -Cyclogeraniol			tr					
1154	β -Pinene oxide					0.05			

Table 1. Cont.

RI	Compound	#1		#3		#4		#5	
		%	D/L	%	D/L	%	D/L	%	D/L
1531	<i>trans</i> -Cadina-1,4-diene								tr
1547	Elemol			0.96					
1554	Geranyl butanoate	0.05		0.74					
1575	Germacrene-D-4-ol	0.10				tr			
1575	Spathulenol			tr					
1581	Caryophyllene oxide	0.41		0.50		0.10		0.13	
1595	Geranyl isovalerate			0.06					
1631	10- <i>epi</i> - γ -Eudesmol			tr					
1654	α -Eudesmol			0.17					

3.2. Chemotypes of *Thymus vulgaris*

A total of 85 *T. vulgaris* essential oil compositions were used to carry out a hierarchical cluster analysis (Figure 1). The cluster analysis revealed as many as 20 different chemotypes. The chemotype with the most samples (39) was the thymol chemotype, which had been recognized previously [12–15]. Other previously recognized chemotypes were the geraniol chemotype, with three representative samples (including sample #3 from Serbia in this study); the linalool chemotype, with four samples (including sample #1 from France in this study); the carvacrol chemotype, with two samples; the borneol chemotype, only one sample; the sabinene hydrate/terpinene-4-ol chemotype, with two samples (including sample #4 from Serbia in this study); and the cyclocitral/verbenol chemotype, with only one sample. The second most populated chemotype in the cluster analysis was a *p*-cymene/thymol type represented by 18 samples. The α -terpineol chemotype [12] was not found in the present study; only three *T. vulgaris* samples (all from Austria) showed α -terpineol concentrations around 10%, but all three had higher concentrations of other components [14]. Other chemotypes identified include carvacrol/ γ -terpinene/thymol, which was previously described as carvacrol and thymol/carvacrol [14]; linalool/carvacrol/*p*-cymene, previously labeled carvacrol/linalool [14]; sabinene hydrate/geraniol/geranyl acetate/ α -terpineol, previously described as geraniol/ α -terpineol/sabinene hydrate [14]; sabinene hydrate/ α -terpinyl acetate/thymol, previously described as α -terpineol/sabinene hydrate/thymol [14]; sabinene hydrate/linalool [14]; α -terpinyl acetate/carvacrol, previously described as α -terpineol/carvacrol [14]; carvacrol/ α -terpineol/borneol; sabinene hydrate/terpinene-4-ol, which includes one of the samples from France in this study; geranyl formate/geraniol, previously called geraniol [13]; *p*-cymene/thymol/carvacrol; terpinene-4-ol/*p*-cymene; camphor/camphene; and 1,8-cineole/ α -terpinyl acetate.

Thymus vulgaris Dendrogram




Figure 1. Dendrogram obtained from the agglomerative hierarchical cluster analysis of 85 *Thymus vulgaris* essential oil compositions.

3.3. Antifungal Activity

Three of the four *T. vulgaris* chemotypes in this study were tested for inhibition of *Aspergillus niger*, *Cryptococcus neoformans* var. *neoformans*, and *Candida albicans*. Macro-dilution was utilized for the yeast-like Ascomycota, *C. albicans*, and Basidiomycota *C. neoformans*, to determine antifungal minimum inhibitory concentrations (MICs, Table 2). The *T. vulgaris* linalool and geraniol chemotypes both demonstrated some degree of inhibition against these pathogens. For the sporulating mold-like Ascomycota, *A. niger*, a larger surface area was required for hyphal growth. Thus, it was grown on malt extract agar plates with a filter disk impregnated with the *T. vulgaris* chemotype of interest. Disk diffusion showed only slight inhibition of *A. niger* with the only clear zone of inhibition for *T. vulgaris* sample #1. The remaining chemotypes showed growth over the filter disk, indicating no significant antifungal activity against *A. niger*. Because it has been shown that linalool is biotransformed to non-pathogenic compounds and that linalyl acetate increases *A. niger* hyphal growth [64], it is speculated that camphor in sample #1 is responsible for *A. niger* inhibition. Significant levels of camphor are not found in the other *T. vulgaris* samples.

Table 2. Antifungal activities (minimum inhibitory concentrations, MIC) of *Thymus vulgaris* essential oils.

Essential Oil	MIC ($\mu\text{g/mL}$)	
	<i>C. albicans</i>	<i>C. neoformans</i>
#1 (linalool chemotype)	1250	313
#3 (geraniol chemotype)	625	156
#4 (sabinene hydrate chemotype)	>2500	>2500

The differential antifungal activities observed in this study mirror those previously reported by Giordani and co-workers [13]. That is, the sabinene hydrate chemotype showed the lowest antifungal activity, the linalool chemotype was next, then the geraniol chemotype. Giordani and co-workers had found that the thymol chemotype showed much stronger antifungal activity [13]. The *T. vulgaris* thymol chemotype was also found to be the most larvicidal against *Culex quinquefasciatus* [15] and exhibited the most antioxidant properties [14].

4. Conclusions

This work has presented the most comprehensive analysis of *Thymus vulgaris* chemotypes, revealing at least 20 different types based on essential oil composition. In addition, this is the first analysis to characterize the enantiomeric distributions of *T. vulgaris* monoterpenoids. Enantiomers are well known to elicit different odorant responses in insects [65,66] as well as humans [67], and it is reasonable to assume that different enantiomers will have different medicinal biological activities [68]. Thus, for example L-linalool has shown anticonvulsant activity in a mouse model whereas D-linalool was inactive [69]. Similarly, both D- α -pinene and D- β -pinene showed antifungal activity whereas the L-enantiomers were inactive [70], while L- α -pinene was more active than the D-enantiomer against *Listeria monocytogenes* [71]. Therefore, not only is the particular chemotype of a culinary and medicinal herb such as *T. vulgaris* an important consideration, but the enantiomeric distribution may also have a profound influence on its bioactivity, flavor, and aroma profile.

Acknowledgments: We thank Julien Abisset for plant identification, Joshua Thomerson for assistance with steam distillation of the essential oils, and Erin McClelland (Middle Tennessee State University) for *C. neoformans* cells. B.L.M. was supported by a GEM University Fellowship.

Author Contributions: P.S. conceived and designed the experiments; P.S. carried out the GC-MS analysis; B.L.M. did the antifungal assays; W.N.S. performed the cluster analysis; R.L.M. and W.N.S. analyzed the data; all authors contributed to writing the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gillett, M. *Growing and Using Thyme*; Storey Publishing: North Adams, MA, USA, 1998.
2. Aygun, O.; Aslantas, O.; Oner, S. A survey on the microbiological quality of Carra, a traditional Turkish cheese. *J. Food Eng.* **2005**, *66*, 401–404. [[CrossRef](#)]
3. Akarca, G.; Çağlar, A.; Tomar, O. The effects of spicing on quality of mozzarella cheese. *Mljekarstvo* **2016**, *66*, 112–121.
4. Martinez-Francés, V.; Rios, S. Lesser known herbal spirits of the Valencia and Alicante provinces (east-southern Spain). *Ind. Crops Rural Dev.* **2005**, 417–426.
5. Karabegović, I.T.; Vukosavljević, P.V.; Novaković, M.M.; Gorjanović, S.Ž.; Džamić, A.M.; Lazić, M.L. Influence of the storage on bioactive compounds and sensory attributes of herbal liqueur. *Dig. J. Nanomater. Biostruct.* **2012**, *7*, 1587–1598.
6. Cornara, L.; La Rocca, A.; Marsili, S.; Mariotti, M.G. Traditional uses of plants in the eastern Riviera (Liguria, Italy). *J. Ethnopharmacol.* **2000**, *125*, 16–30. [[CrossRef](#)] [[PubMed](#)]
7. Gimeno Gasca, J.M. Tomillo (*Thymus vulgaris* L.). *Med. Natur.* **2001**, *3*, 173–175.
8. Alleman, F.; Gabriel, I.; Dufourcq, V.; Perrin, F.; Gabarrou, J.F. Utilisation des huiles essentielles en alimentation des volailles. 1. Performances de croissance et réglementation. *INRA Prod. Anim.* **2013**, *26*, 3–12.
9. Basch, E.; Ulbricht, C.; Hammerness, P.; Blevins, A.; Sollars, D. Thyme (*Thymus vulgaris* L.), thymol. *J. Herb. Pharmacother.* **2004**, *4*, 49–67. [[CrossRef](#)] [[PubMed](#)]
10. Duke, J.A. *Handbook of Phytochemical Constituents of GRAS Herbs and Other Economic Plants*; CRC Press: Boca Raton, FL, USA, 1992.
11. Tropicos.org. Missouri Botanical Garden. Available online: <http://www.tropicos.org> (accessed on 17 August 2016).
12. Thompson, J.D.; Chalchat, J.C.; Michet, A.; Linhart, Y.B.; Ehlers, B. Qualitative and quantitative variation on monoterpene co-occurrence and composition in the essential oil of *Thymus vulgaris* chemotypes. *J. Chem. Ecol.* **2003**, *29*, 859–880. [[CrossRef](#)] [[PubMed](#)]
13. Giordani, R.; Regli, P.; Kaloustian, J.; Mikail, C.; Abou, L.; Portugal, H. Antifungal effect of various essential oils against *Candida albicans*. Potentiation of antifungal action of amphotericin B by essential oil from *Thymus vulgaris*. *Phytother. Res.* **2004**, *18*, 990–995. [[CrossRef](#)] [[PubMed](#)]
14. Chizzola, R.; Michitsch, H.; Franz, C. Antioxidative properties of *Thymus vulgaris* leaves: Comparison of different extracts and essential oil chemotypes. *J. Agric. Food Chem.* **2008**, *56*, 6897–6904. [[CrossRef](#)] [[PubMed](#)]
15. Pavela, R.; Vrchotová, N.; Tříska, J. Mosquitocidal activities of thyme oils (*Thymus vulgaris* L.) against *Culex quinquefasciatus* (Diptera: Culicidae). *Parasitol. Res.* **2009**, *105*, 1365–1370. [[CrossRef](#)] [[PubMed](#)]
16. Adams, R.P. *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*, 4th ed.; Allured Publishing: Carol Stream, IL, USA, 2007.
17. Özek, T.; Tabanca, N.; Demirci, F.; Wedge, D.E.; Başer, K.H.C. Enantiomeric distribution of some linalool containing essential oils and their biological activities. *Rec. Nat. Prod.* **2010**, *4*, 180–192.
18. Shellie, R.; Marriott, P.; Cornwell, C. Application of comprehensive two-dimensional gas chromatography (GC × GC) to the enantioselective analysis of essential oils. *J. Sep. Sci.* **2001**, *24*, 823–830. [[CrossRef](#)]
19. Bonaccorsi, I.; Sciarrone, D.; Cotroneo, A.; Mondello, L.; Dugo, P.; Dugo, G. Enantiomeric distribution of key volatile components in *Citrus* essential oils. *Rev. Bras. Farmacog.* **2011**, *21*, 841–849. [[CrossRef](#)]
20. Pragadheesh, V.S.; Saroj, A.; Yadav, A.; Samad, A. Chantitiya, C.S. Compositions, enantiomer characterization and antifungal activity of two *Ocimum* essential oils. *Ind. Crops Prod.* **2013**, *50*, 333–337. [[CrossRef](#)]
21. Tabanca, N.; Kirimer, N.; Demirci, B.; Demirci, F.; Başer, K.H.C. Composition and antimicrobial activity of the essential oils of *Micromeria cristata* subsp. *phrygia* and the enantiomeric distribution of borneol. *J. Agric. Food. Chem.* **2001**, *49*, 4300–4303. [[CrossRef](#)] [[PubMed](#)]
22. Imelouane, B.; Amhamdi, H.; Wathelet, J.P.; Ankit, M.; Khedid, K.; El Bachiri, A. Chemical composition and antimicrobial activity of essential oil of thyme (*Thymus vulgaris*) from eastern Morocco. *Int. J. Agric. Biol.* **2009**, *11*, 205–208.
23. Jordán, M.J.; Martínez, R.M.; Goodner, K.L.; Baldwin, E.A.; Sotomayor, J.A. Seasonal variation of *Thymus hyemalis* Lange and Spanish *Thymus vulgaris* L. essential oils composition. *Ind. Crops Prod.* **2006**, *24*, 253–263. [[CrossRef](#)]

24. Bhaskara Reddy, M.V.; Angers, P.; Gosselin, A.; Arul, J. Characterization and use of essential oil from *Thymus vulgaris* against *Botrytis cinerea* and *Rhizopus stolonifer* in strawberry fruits. *Phytochemistry* **1998**, *47*, 1515–1520. [[CrossRef](#)]
25. Sacchetti, G.; Maietti, S.; Muzzoli, M.; Scaglianti, M.; Manfredini, S.; Radice, M.; Bruni, R. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chem.* **2005**, *91*, 621–632. [[CrossRef](#)]
26. Dimitrijević, S.I.; Mihajlovski, K.R.; Antonović, D.G.; Milanović-Stevanović, M.R.; Mijin, D.Ž. A study of the synergistic antilisterial effects of a sub-lethal dose of lactic acid and essential oils from *Thymus vulgaris* L., *Rosmarinus officinalis* L. and *Origanum vulgare* L. *Food Chem.* **2007**, *104*, 774–782. [[CrossRef](#)]
27. Gavahian, M.; Farahnaky, A.; Javidnia, K.; Majzoobi, M. Comparison of ohmic-assisted hydrodistillation with traditional hydrodistillation for the extraction of essential oils from *Thymus vulgaris* L. *Innov. Food Sci. Emerg. Technol.* **2012**, *14*, 85–91. [[CrossRef](#)]
28. Panizzi, L.; Flamini, G.; Cioni, P.L.; Morelli, I. Composition and antimicrobial properties of essential oils of four Mediterranean Lamiaceae. *J. Ethnopharmacol.* **1993**, *39*, 167–170. [[CrossRef](#)]
29. Lucchesi, M.E.; Chemat, F.; Smadja, J. Solvent-free microwave extraction of essential oil from aromatic herbs: Comparison with conventional hydro-distillation. *J. Chromatogr. A* **2004**, *1043*, 323–327. [[CrossRef](#)] [[PubMed](#)]
30. Zambonelli, A.; Zechini D’Aulerio, A.; Severi, A.; Benvenuti, S.; Maggi, L.; Bianchi, A. Chemical composition and fungicidal activity of commercial essential oils of *Thymus vulgaris* L. *J. Essent. Oil Res.* **2004**, *16*, 69–74. [[CrossRef](#)]
31. Hudaib, M.; Speroni, E.; Di Pietra, A.M.; Cavrini, V. GC/MS evaluation of thyme (*Thymus vulgaris* L.) oil composition and variations during the vegetative cycle. *J. Pharmaceut. Biomed. Anal.* **2002**, *29*, 691–700. [[CrossRef](#)]
32. Szczepanik, M.; Zawitowska, B.; Szumny, A. Insecticidal activities of *Thymus vulgaris* essential oil and its components (thymol and carvacrol) against larvae of lesser mealworm, *Alphitobius diaperinus* Panzer (Coleoptera: Tenebrionidae). *Allelopath. J.* **2012**, *30*, 129–142.
33. Bozin, B.; Mimica-Dukic, N.; Simin, N.; Anackov, G. Characterization of the volatile composition of essential oils of some Lamiaceae spices and the antimicrobial and antioxidant activities of the entire oils. *J. Agric. Food Chem.* **2006**, *54*, 1822–1828. [[CrossRef](#)] [[PubMed](#)]
34. Lee, S.J.; Umamo, K.; Shibamoto, T.; Lee, K.G. Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem.* **2005**, *91*, 131–137. [[CrossRef](#)]
35. Sartoratto, A.; Machado, A.L.M.; Delarmelina, C.; Figueira, G.M.; Duarte, C.T.; Rehder, V.L.G. Composition and antimicrobial activity of essential oils from aromatic plants used in Brazil. *Braz. J. Microbiol.* **2004**, *35*, 275–280. [[CrossRef](#)]
36. Dawidowicz, A.L.; Rado, E.; Wianowska, D.; Mardarowicz, M.; Gawdzik, J. Application of PLE for the determination of essential oil components from *Thymus vulgaris* L. *Talanta* **2008**, *76*, 878–884. [[CrossRef](#)] [[PubMed](#)]
37. Karami-Osboo, R.; Khodaverdi, M.; Ali-Akbari, F. Antibacterial effect of effective compounds of *Satureja hortensis* and *Thymus vulgaris* essential oils against *Erwinia amylovora*. *J. Agric. Sci. Technol.* **2010**, *12*, 35–45.
38. Al-Maqtary, M.A.A.; Alghalibi, S.M.; Alhamzy, E.H. Chemical composition and antimicrobial activity of essential oil of *Thymus vulgaris* from Yemen. *Turk. J. Biochem.* **2011**, *36*, 342–349.
39. Rota, M.C.; Herrera, A.; Martínez, R.M.; Sotomayor, J.A.; Jordán, M.J. Antimicrobial activity and chemical composition of *Thymus vulgaris*, *Thymus zygis* and *Thymus hyemalis* essential oils. *Food Control* **2008**, *19*, 681–687. [[CrossRef](#)]
40. Daferera, D.J.; Ziogas, B.N.; Polissiou, M.G. GC-MS analysis of essential oils from some Greek aromatic plants and their fungitoxicity on *Penicillium digitatum*. *J. Agric. Food Chem.* **2000**, *48*, 2576–2581. [[CrossRef](#)] [[PubMed](#)]
41. Nikolić, M.; Glamočlija, J.; Ferreira, I.C.F.R.; Calhelha, R.C.; Fernandes, Â.; Marković, T.; Marković, D.; Giweli, A.; Soković, M. Chemical composition, antimicrobial, antioxidant and antitumor activity of *Thymus serpyllum* L., *Thymus algeriensis* Boiss. and Reut and *Thymus vulgaris* L. essential oils. *Ind. Crops Prod.* **2014**, *52*, 183–190. [[CrossRef](#)]

42. Soković, M.D.; Vukojević, J.; Marin, P.D.; Brkić, D.D.; Vajs, V.; van Griensven, L.J.L.D. Chemical composition of essential oils of *Thymus* and *Mentha* species and their antifungal activities. *Molecules* **2009**, *14*, 238–249. [[CrossRef](#)] [[PubMed](#)]
43. Ghasemi Pirbalouti, A.; Hashemi, M.; Taherian Ghahfarokhi, F. Essential oil and chemical compositions of wild and cultivated *Thymus daenensis* Celak and *Thymus vulgaris* L. *Ind. Crops Prod.* **2013**, *48*, 43–48. [[CrossRef](#)]
44. Baranauskienė, R.; Venskutonis, P.R.; Viškelis, P.; Dambrauskienė, E. Influence of nitrogen fertilizers on the yield and composition of thyme (*Thymus vulgaris*). *J. Agric. Food Chem.* **2003**, *51*, 7751–7758. [[CrossRef](#)] [[PubMed](#)]
45. Shabnum, S.; Wagay, M.G. Essential oil composition of *Thymus vulgaris* L. and their uses. *J. Res. Dev.* **2011**, *11*, 83–94.
46. Asllani, U.; Toska, V. Chemical composition of Albanian thyme oil (*Thymus vulgaris* L.). *J. Essent. Oil Res.* **2003**, *15*, 165–167. [[CrossRef](#)]
47. Porte, A.; Godoy, R.L.O. Chemical composition of *Thymus vulgaris* L. (thyme) essential oil from the Rio de Janeiro State (Brazil). *J. Serbian Chem. Soc.* **2008**, *73*, 307–310. [[CrossRef](#)]
48. Piccaglia, R.; Marotti, M. Composition of the essential oil of an Italian *Thymus vulgaris* L. ecotype. *Flavour Fragr. J.* **1991**, *6*, 241–244. [[CrossRef](#)]
49. Atti-Santos, A.C.; Pansera, M.R.; Paroul, N.; Atti-Serafini, L.; Moyna, P. Seasonal variation of essential oil yield and composition of *Thymus vulgaris* L. (Lamiaceae) from south Brazil. *J. Essent. Oil Res.* **2004**, *16*, 294–295. [[CrossRef](#)]
50. Golmakani, M.T.; Rezaei, K. Comparison of microwave-assisted hydrodistillation with the traditional hydrodistillation method in the extraction of essential oils from *Thymus vulgaris* L. *Food Chem.* **2008**, *109*, 925–930. [[CrossRef](#)] [[PubMed](#)]
51. Dawidowicz, A.L.; Rado, E.; Wianowska, D. Static and dynamic superheated water extraction of essential oil components from *Thymus vulgaris* L. *J. Sep. Sci.* **2009**, *32*, 3034–3042. [[CrossRef](#)] [[PubMed](#)]
52. Vurro, E.; Bruni, R.; Bianchi, A.; di Toppi, L.S. Elevated atmospheric CO₂ decreases oxidative stress and increases essential oil yield in leaves of *Thymus vulgaris* grown in a mini-FACE system. *Environ. Exp. Bot.* **2009**, *65*, 99–106. [[CrossRef](#)]
53. McGimpsey, J.A.; Douglas, M.H.; van Klink, J.W.; Beauregard, D.A.; Perry, N.B. Seasonal variation in essential oil yield and composition from naturalized *Thymus vulgaris* L. in New Zealand. *Flavour Fragr. J.* **1994**, *9*, 347–352. [[CrossRef](#)]
54. Mota, K.S.L.; Pereira, F.O.; de Oliveira, W.A.; Lima, I.O.; Lima, E.O. Antifungal activity of *Thymus vulgaris* L. essential oil and its constituent phytochemicals against *Rhizopus oryzae*: Interaction with ergosterol. *Molecules* **2012**, *17*, 14418–14433. [[CrossRef](#)] [[PubMed](#)]
55. Aziz, E.E.; Hendawi, S.T.; Azza, E.E.D.; Omer, E.A. Effect of soil type and irrigation intervals on plant growth, essential oil yield and constituents of *Thymus vulgaris* plant. *Am.-Eurasian J. Agric. Environ. Sci.* **2008**, *4*, 443–450.
56. Venskutonis, P.R. Effect of drying on the volatile constituents of thyme (*Thymus vulgaris* L.) and sage (*Salvia officinalis* L.). *Food Chem.* **1997**, *59*, 219–227. [[CrossRef](#)]
57. Šegvić Klarić, M.; Kosalec, I.; Masterlić, J.; Piecková, E.; Pepeljnak, S. Antifungal activity of thyme (*Thymus vulgaris* L.) essential oil and thymol against moulds from damp dwellings. *Lett. Appl. Microbiol.* **2007**, *44*, 36–42. [[CrossRef](#)] [[PubMed](#)]
58. Nguefack, J.; Tamgue, O.; Lekagne Dongmo, J.B.; Dakole, C.D.; Leth, V.; Vismer, H.F.; Amvam Zollo, P.H.; Nkengfack, A.E. Synergistic action between fractions of essential oils from *Cymbopogon citratus*, *Ocimum gratissimum* and *Thymus vulgaris* against *Penicillium expansum*. *Food Control* **2012**, *23*, 377–383. [[CrossRef](#)]
59. Grigore, A.; Paraschiv, I.; Colceru-Mihul, S.; Bubueanu, C.; Draghici, E.; Ichim, M. Chemical composition and antioxidant activity of *Thymus vulgaris* L. volatile oil obtained by two different methods. *Rom. Biotechnol. Lett.* **2010**, *15*, 5436–5443.
60. Díaz-Maroto, M.C.; Hidalgo, I.J.D.M.; Sánchez-Palomo, E.; Pérez-Coello, M.S. Volatile components and key odorants of fennel (*Foeniculum vulgare* Mill.) and thyme (*Thymus vulgaris* L.) oil extracts obtained by simultaneous distillation-extraction and supercritical fluid extraction. *J. Agric. Food Chem.* **2005**, *53*, 5385–5389. [[CrossRef](#)] [[PubMed](#)]

61. Fachini-Queiroz, F.C.; Kummer, R.; Estevão-Silva, C.F.; Carvalho, M.D.B.; Cunha, J.M.; Grespan, R.; Bersani-Amado, C.A.; Cuman, R.K.N. Effects of thymol and carvacrol, constituents of *Thymus vulgaris* L. essential oil, on the inflammatory response. *Evid.-Based Complement. Altern. Med.* **2012**, *2012*, 657026. [[CrossRef](#)] [[PubMed](#)]
62. El-Nekeety, A.A.; Mohamed, S.R.; Hathout, A.S.; Hassan, N.S.; Aly, S.E.; Abdel-Wahhab, M.A. Antioxidant properties of *Thymus vulgaris* oil against aflatoxin-induced oxidative stress in male rats. *Toxicol* **2011**, *57*, 984–991. [[CrossRef](#)] [[PubMed](#)]
63. Pina-Vaz, C.; Gonçalves Rodrigues, A.; Pinto, E.; Costa-de-Oliveira, S.; Tavares, C.; Salgueiro, L.; Cavaleiro, C.; Gonçalves, M.J.; Martinez-de-Oliveira, J. Antifungal activity of *Thymus* oils and their major compounds. *J. Eur. Acad. Dermatol. Venereol.* **2004**, *18*, 73–78. [[CrossRef](#)] [[PubMed](#)]
64. Oh, K.; Matsuoka, H.; Sumita, O.; Takatori, K. Kurata, H. Automatic evaluation of antifungal volatile compounds on the basis of the dynamic growth process of a single hypha. *Appl. Microbiol. Biotechnol.* **1993**, *38*, 790–794. [[CrossRef](#)] [[PubMed](#)]
65. Anderson, A.R.; Wanner, K.W.; Trowell, S.C.; Warr, C.G.; Jaquin-Joly, E.; Zagatti, P.; Robertson, H.; Newcomb, R.D. Molecular basis of female odorant responses in *Bombyx mori*. *Insect Biochem. Mol. Biol.* **2009**, *39*, 189–197. [[CrossRef](#)] [[PubMed](#)]
66. Liu, F.; Chen, L.; Appel, A.G.; Liu, N. Olfactory responses of the antennal trichoid sensilla to chemical repellents in the mosquito, *Culex quinquefasciatus*. *J. Insect Physiol.* **2013**, *59*, 1169–1177. [[CrossRef](#)] [[PubMed](#)]
67. Prasse, A.; Siehl, H.U.; Zeller, K.P.; Berger, S.; Sicker, D. Wie unsere Nase die enantiomeren Linaloole unterscheidet. *Chem. Unserer Zeit* **2014**, *48*, 342–353. [[CrossRef](#)]
68. Asztemborska, M.; Ochocka, J.R. Chiral monoterpenoids in plants—Enantioselective chromatographic analysis, and their bioactivity. *Stud. Nat. Prod. Chem.* **2002**, *27*, 361–391. [[CrossRef](#)]
69. De Sousa, D.P.; Nóbrega, F.F.; Santos, C.C.; de Almeida, R.N. Anticonvulsant activity of the linalool enantiomers and racemate: Investigation of chiral influence. *Nat. Prod. Commun.* **2010**, *5*, 1847–1851. [[PubMed](#)]
70. Da Silva, A.C.R.; Lopes, P.M.; de Azevedo, M.M.B.; Costa, D.C.M.; Alviano, C.S.; Alviano, D.S. Biological activities of α -pinene and β -pinene enantiomers. *Molecules* **2012**, *17*, 6305–6316. [[CrossRef](#)] [[PubMed](#)]
71. Lis-Balchin, M.; Ochocka, R.J.; Deans, S.G.; Asztemborska, M.; Hart, S. Differences in bioactivity between the enantiomers of α -pinene. *J. Essent. Oil Res.* **1999**, *11*, 393–397. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).